



Published in final edited form as:

*Neuroimage*. 2017 October 15; 160: 15–31. doi:10.1016/j.neuroimage.2017.01.079.

## Development of large-scale functional networks from birth to adulthood: a guide to neuroimaging literature

David S. Grayson<sup>1,2,3</sup> and Damien A. Fair<sup>3,4,5,\*</sup>

<sup>1</sup>The MIND Institute, University of California Davis, Sacramento, California 95817, USA

<sup>2</sup>Center for Neuroscience, University of California Davis, Davis, California 95616, USA

<sup>3</sup>Department of Behavioral Neuroscience, Oregon Health and Science University, Portland, OR 97239, USA

<sup>4</sup>Department of Psychiatry, Oregon Health and Science University, Portland, OR 97239, USA

<sup>5</sup>Advanced Imaging Research Center, Oregon Health and Science University, Portland, OR 97239, USA

### Abstract

The development of human cognition results from the emergence of coordinated brain activity between distant brain areas. Network science, combined with non-invasive functional imaging, has generated unprecedented insights regarding the adult brain's functional organization, and promises to help elucidate the development of functional architectures supporting complex behavior. Here we review what is known about functional network development from birth until adulthood, particularly as understood through the use of resting-state functional connectivity MRI (rs-fcMRI). We attempt to synthesize rs-fcMRI findings with other functional imaging techniques, with macro-scale structural connectivity, and with knowledge regarding the development of micro-scale structure. We highlight a number of outstanding conceptual and technical barriers that need to be addressed, as well as previous developmental findings that may need to be revisited. Finally, we discuss key areas ripe for future research in order to 1) better characterize normative developmental trajectories, 2) link these trajectories to biologic mechanistic events, as well as component behaviors and 3) better understand the clinical implications and pathophysiological basis of aberrant network development.

### Keywords

Functional connectivity; graph theory; resting-state functional MRI; brain development; connectomics

---

Correspondence: Damien A. Fair, PA-C, Ph.D. Oregon Health and Science University, 3181 SW Sam Jackson Park Road L470, Portland, Oregon 97239, Tel: 503-418-0995, faird@ohsu.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Introduction

### **The human brain is organized into multiple distributed functional brain networks that can be measured at multiple spatiotemporal scales**

Coordinated neuronal activity between anatomically disparate regions is an essential feature of human brain function. Across all stages of postnatal development, brain activity to a large degree is consolidated within so-called “resting-state networks” (RSNs). RSNs are defined as distinct modules of regions that exhibit highly synchronized activity even in the absence of external stimuli. RSNs have proven to be highly reproducible in the adult brain (Doucet et al., 2011; Gordon et al., 2016; Power et al., 2011; Yeo et al., 2011) and have become an influential framework for interpreting functional and structural neuroimaging data. Patterns of functional connectivity (FC) within and between the major RSNs are increasingly understood as intrinsic properties of brain function, given that they strongly predict patterns of interregional co-activation across different tasks (Cole et al., 2014; Smith et al., 2009) and are associated with task-relevant behavioral performance (Cole et al., 2012; Lewis et al., 2009).

Interregional FC arises from anatomical projection strengths (Messe et al., 2014; Shen et al., 2012), correlated gene expression (Richiardi et al., 2015), and synaptic receptor densities (Turk et al., 2016; van den Heuvel et al., 2016b)—properties that, in turn, undergo experience-dependent and activity-dependent modulation over the lifespan (Huttenlocher, 2002; Markham and Greenough, 2004; Scholz et al., 2009). This intricate structural-functional interplay underscores that the ontogeny of functional networks likely reflects programmed neurodevelopmental events (e.g. neurogenesis, cell death, myelination, pruning, synaptic plasticity, and glial development) (Innocenti and Price, 2005; Stiles and Jernigan, 2010) and that FC aberrations may point towards the etiological bases of neuropsychiatric disorders (e.g. (Swartz and Monk, 2014; van den Heuvel et al., 2016a)). Thus, exploring the normal trajectory of functional network development and its relationships with these underlying biological processes should be a central mission of both basic and clinically relevant neurodevelopmental research.

Current views regarding network development have been heavily informed by resting-state functional connectivity MRI (rs-fcMRI), which measures the correlations in spontaneous, low-frequency activity between investigator-defined regions. This review will accordingly provide in-depth discussions of the developmental rs-fcMRI literature, its limitations, and unique insights gained from complementary approaches, especially electroencephalography and magnetoencephalography (EEG/MEG). We will begin by discussing emerging themes in the analysis of functional network data. We will then synthesize current findings regarding FC changes from birth to adulthood, and place them within the context of co-occurring macro- and micro-scale structural modifications. Finally, we will discuss some of the implications with regard to neurodevelopmental disorders and highlight several crucial knowledge gaps that are likely to guide research efforts in the near future.

## Analytic approaches to functional connectivity are evolving rapidly

Correlated resting-state activity was first identified in bilateral sensorimotor cortex by (Biswal et al., 1995), followed soon after by similar findings of bilaterally correlated activity in early visual and early auditory cortex (Cordes et al., 2001; Kiviniemi et al., 2000; Lowe et al., 1998). These discoveries were made using “seed” regions outlined based on *a priori* anatomical boundaries. Correlations were then quantified between the seed region’s fMRI timecourse and all other voxels of the brain, producing an FC map. More data-driven approaches were developed based on the concept of independent components analysis (Beckmann and Smith, 2004), which decomposes brain activity into a set of spatial maps (i.e. components) that minimally overlap. ICA maps reproducibly reveal early visual, early auditory, and primary sensory/motor components, as well as other components comprising distributed portions of the frontal, parietal, and temporal association cortices (Damoiseaux et al., 2006).

Building upon these advances, the neuroimaging community has increasingly focused on interrogating the brain as an integrated complex system. In this context, graph theory has emerged as a powerful new approach. A ‘graph’ is simply a network of things, referred to as nodes, and the connections between those things, referred to as edges. The network’s behavior can thus be modeled as a set of properties that emerge from the network’s unique global structure and local features (Sporns, 2014). Graph theoretic studies typically construct functional brain networks using regions as nodes and correlations in activity as edges. Past reviews have already written eloquently about the rationales for applying this approach to study development (Power et al., 2010; Vertes and Bullmore, 2015), which we will only briefly summarize and expand upon here.

First, most individual cognitive abilities arise not solely from a particular brain area, but from networks of activity spanning multiple distributed regions (Petersen and Sporns, 2015). From social cognition in infancy (Eggebrecht et al., 2017) to cognitive control and decision-making in adolescence (Dwyer et al., 2014; van Duijvenvoorde et al., 2015), specific cognitive capacities co-evolve with complex network effects within and between different RSNs, throughout postnatal development. Second, structural connectivity shapes and constrains functional networks across the lifespan (Betz et al., 2014; Hagmann et al., 2012; Vertes and Bullmore, 2015). Cross-modal network analysis is therefore crucial for elucidating normative mechanisms of cognitive development and the pathophysiology of neurodevelopmental disturbances. As a proof-of-principle, recent studies have found that even purely local, experimentally induced perturbations (e.g. exogenous stimulation or inactivation) result in widespread, complex neurophysiological changes, which are at least partly explainable as local interactions with global network structure (Andoh et al., 2015; Grayson et al., 2016; Gu et al., 2015a; Masic et al., 2015). By extension, one could reason that developmental modifications in brain FC, whether normative or pathological, are also best understood via an emerging network science that merges structural and functional connectivity data. Third, graph-based methods are flexible and generalizable. Networks can be measured at various temporal scales depending on imaging methodology, at various conditions (during rest or task completion), and can be compared with corresponding networks of structural connectivity or gene co-expression. A graph theoretic approach is

therefore well-suited for multimodal investigations linking brain function across temporal scales or examining structural drivers of functional development.

## Graph theory techniques for defining network organization

Below we describe 3 broad themes that have emerged with regard to functional (and structural) network analysis in neuroimaging: the brain's community structure (i.e. the spatial and topological organization of specialized systems), the significance of hub regions that integrate information within and between these systems, and global network properties that facilitate efficient and integrated information transfer.

### Community Structure

A defining feature of human brain activity is “modularity,” which refers to an unexpectedly high level of within-RSN FC relative to between-RSN FC (Lohse et al., 2014). “Community detection” algorithms partition the brain into distinct communities, or modules, by maximizing this modularity quotient. The adult brain has several modules at the coarsest level that are remarkably reproducible despite slight methodological differences across studies (Doucet et al., 2011; Gordon et al., 2016; Power et al., 2011; Yeo et al., 2011). These modules include 1) segregated, non-distributed modules for sensory/motor activity, such as in early visual cortex and in somatomotor cortex, 2) the default-mode network (DMN), comprising the medial prefrontal, posterior cingulate/precuneus, angular gyrus, and medial temporal cortex; the DMN is considered “task-negative” as it has been linked to internal mentation and is known to deactivate during processing of external stimuli (Buckner et al., 2008), and 3) three large, distributed modules for task-positive cognition: a) dorsal attention, involved in eye movements and attentional orienting, b) the cingulo-opercular salience module, involved primarily in attentional maintenance, and c) the frontoparietal executive module, important for task-switching. At finer resolutions, several other modules can be identified (Doucet et al., 2011; Gordon et al., 2016; Power et al., 2011; Yeo et al., 2011), including hierarchical visual subdivisions, topographical somatomotor subdivisions, an auditory module, a limbic module, a ventral attention module, and other smaller subdivisions of the executive, salience, and default-mode modules. See Figure 1A & B for a graphical summary.

The brain's modular composition is believed to help segregate information processing between distinct sensory modalities or cognitive architectures. It remains a fundamental challenge to understand what other network properties help integrate disparate data streams in order to meet complex task demands (Cohen and D'Esposito, 2016; Petersen and Sporns, 2015). As discussed throughout this review, the themes of segregation and integration are core concepts in understanding the brain's maturational processes. Thus, further analytic exploration is needed to isolate the critical network components that facilitate segregation, integration, or efficiency. Below we briefly describe some local network measures toward this end.

## Node Measures

The most straightforward nodal description is its *strength* (Rubinov and Sporns, 2010), defined as the sum of a node's connection weights. Other *centrality* measures quantify higher order measures of influence. For instance, *betweenness centrality* (Rubinov and Sporns, 2010) computes the frequency with which a node lies on the shortest paths that connect any other nodes, a property that theoretically enables efficient global communication. By far, strength and betweenness have been the most popular hub metrics within neuroimaging, although others such as *eigenvector centrality* (Lohmann et al., 2010; Zuo et al., 2012) and node *communicability* (de Reus and van den Heuvel, 2014; Mantzaris et al., 2013) have seen useful application as well. These metrics, however, require careful consideration of what the network's edge weights signify (see (de Reus and van den Heuvel, 2014; Grayson et al., 2016; Misisic et al., 2015) for discussions about the neurobiological relevance of shortest paths). Investigators must especially consider whether connections are statistical or structural in nature. For example, a node with structural links to many different RSNs (and thus, high structural network centrality) may have weak statistical associations overall (and thus, low functional network centrality). It has therefore been proposed to consider functional hubs as nodes with integrative capacity across distinct modules (Power et al., 2013). This goal is accomplished in part by the *participation coefficient* (Guimera and Nunes Amaral, 2005), which computes the between-module connectivity and therefore is particularly useful for functional networks (see Figure 1C).

Another way of identifying nodal integration is to compare functional against structural networks (see Figure 1E). Structural networks in the mammalian brain exhibit an important phenomenon known as “rich-club” organization, whereby hubs (i.e. nodes with high strength) link disproportionately to other hubs. Rich-club nodes tend to span multiple RSNs, thereby facilitating integration between them (Sporns, 2013; van den Heuvel and Sporns, 2013). Similarly, structural hubness has been linked to integration of global brain activity (Grayson et al., 2016).

Finally, node *clustering coefficients* and *local efficiency* (Rubinov and Sporns, 2010) are popular ways of measuring node specialization – i.e., integration specifically within a node's local environment. The clustering coefficient quantifies the combined strength of connectivity between a given node's immediately connected neighbors, whereas the closely related local efficiency computes the average (inverse) path length between a node's neighbors.

## Global Network Properties

A network's capacities for globally efficient communication and integration can be defined formally as properties. The *global efficiency* is measured by averaging the (inverse) shortest path length between all node pairs. Greater global efficiency in structural and functional networks has been linked to higher IQ, attentional capacity, and working memory in healthy young adults (Cohen and D'Esposito, 2016; Kitzbichler et al., 2011; Li et al., 2009; Stanley et al., 2015; van den Heuvel et al., 2009), supporting the notion that it confers advantages for complex cognitive processing. When networks exhibit both high global efficiency and high average clustering coefficients, they exhibit “small-world” organization (Watts and Strogatz,

1998). This property, which is ubiquitous across vastly different scales of investigation in the brain, is thought to emerge from a biological need to balance the benefits of integration against the costs of long-range wiring (Collin et al., 2013; Vertes and Bullmore, 2015). Furthermore, these distinct network effects, i.e. segregated versus integrated neural processing, are interwoven features of cognition. Both effects in tandem are crucially involved in cognitive control (Dwyer et al., 2014) and executive functioning (Reineberg et al., 2015), and are differentially associated with performance on simple motor tasks versus complex working memory tasks (Cohen and D’Esposito, 2016). This review will highlight extensively these competing network influences in the context of development. Finally, network *integration* can also be defined alternatively. Similar to the participation coefficient of a given node, one can compute the network’s global integrative capacity by taking into account the network modules and the interdependencies between them (de Reus and van den Heuvel, 2014; Tononi et al., 1994).

## Functional brain organization in infants and toddlers

### Overall structural and functional organization

The human brain expands at an explosive rate during the first few years of life (see Figure 2A), doubling in volume during the first year and reaching 80–90% of adult volume by age 3 (Courchesne et al., 2000; Knickmeyer et al., 2008). This same time period is marked by rapid and widespread cortical synaptogenesis, followed by a protracted period of synapse elimination and cell loss which carries into adulthood (Huttenlocher, 2002; Innocenti and Price, 2005). The brain’s major fiber pathways become consolidated through myelination, though their presence is largely established prenatally (Stiles and Jernigan, 2010). It is also worth noting that both progressive and regressive changes in structure occur simultaneously during infancy (Gilmore et al., 2012; Scott et al., 2015), despite overall growth in volume and connectivity. These first few years of life, therefore, coincide with the formation of the brain’s foundational anatomical circuitry, paralleling the genesis of an intricate functional architecture.

RSNs are already present during infancy, and potentially to some extent in utero (Thomason et al., 2013). Seed-based and component-based approaches demonstrate the existence of robust, bilateral segregated networks for somatomotor, primary auditory, primary visual, and extrastriate visual cortex (Fransson et al., 2007; Gao et al., 2015a; Kiviniemi et al., 2000; Lin et al., 2008; Liu et al., 2008; Smyser et al., 2010). These sensory networks undergo subtle refinements and strengthening over the first two years of life, and by age two bear substantial resemblance to their adult counterparts (Gao et al., 2015a; Lin et al., 2008). Distinct components throughout heteromodal association cortex have also been described in infants, and are believed to represent prototypical elements of distributed, higher-order cognitive networks identifiable in adults (Graham et al., 2015). For instance, in adults the DMN is a tightly integrated circuit comprising several distinct and distant areas of cortex (as described in *Community Structure*). In neonates and very young infants, these individual compartments demonstrate locally correlated activity, but fail to synchronize into a coherent network (Damaraju et al., 2014; Fransson et al., 2007; Gao et al., 2009; Smyser et al., 2010; Wylie et al., 2014). Over the course of the first two years of life, these regions undergo a



precise evolution to gradually bring the network ‘online’ (Damaraju et al., 2014; Gao et al., 2013; Gao et al., 2009). By two years of age, the network bears adult-like qualities, although coupling between distant anterior and posterior nodes of the network remain relatively low. Similar spatial and temporal properties characterize the emergence of distributed connectivity in the dorsal attention (Gao et al., 2013), salience (Alcauter et al., 2015a; Gao et al., 2015a; Gao et al., 2015b) and frontoparietal executive (Gao et al., 2015a; Gao et al., 2015b) networks. Collectively, these findings suggest that sensory networks become established at a much earlier age relative to those implicated in higher-level cognition (Gao et al., 2016). These trends are consistent with the presence of basic somatosensory and visual functions at birth, and help contextualize findings that primary sensory regions are generally the first cortical areas to reach plateaus or peaks in growth (Geng et al., 2016; Gilmore et al., 2012; Lyall et al., 2015; Scott et al., 2015).

### Development of community structure from birth

Despite these insights, network topology in early development remains poorly understood. What is the collective community structure of the brain at birth? How does it evolve during early and late infancy, and how is it distinguished from the now well-described adult community structure?

The current nascent literature on infant network topology (see Figure 2B) suggests that a robust community structure does exist, albeit in a primitive form. In neonates, two independent rs-fcMRI studies found evidence for modules comprised largely of anatomically proximate areas (De Asis-Cruz et al., 2015; van den Heuvel et al., 2015b). One recent report compared community structure in adults against that in 1–2 year olds using closely matched methods (Eggebrecht et al., 2017), providing the first direct evidence for numerous anatomically localized modules in the infant brain and their relationship to adult functional organization (Figure 2C). Many of these modules clearly reflect anterior/posterior/temporal subdivisions of the adult salience, frontoparietal, DMN, dorsal attention, and ventral attention modules. Another distinction seen is that infant modules tend to group together anatomically proximate ROIs that have divergent roles in adulthood. Community structure assessed at two years of age (Alcauter et al., 2015b; Gao et al., 2015b) indicates the presence of distinct modules encompassing early auditory, early somatomotor, and early visual areas, and potentially, other prototypical modules similar to the canonical salience, frontoparietal, and DMN seen in adults. These latter modules also exhibit substantial growth of distributed synchronization and homotopic coupling from birth through the first two years of life (Gao et al., 2015b; Homae et al., 2010), concomitant with decreasing between-module connectivity (i.e. greater functional specialization). Taken together, these studies point towards several trends that are consistent with results from seed and component based approaches – 1) modular patterns at birth take a coarse, primitive form that is heavily influenced by local anatomy, 2) this structure undergoes rapid postnatal refinement, leading to progressively stronger coupling between distributed links that facilitate modular specialization, and 3) these enhancements result in the establishment of stable, primary sensory networks well before networks involved in higher-level cognition appear substantially adult-like.

## Regional hub metrics

Hub configurations are robustly evident after birth (De Asis-Cruz et al., 2015; Fransson et al., 2011; Gao et al., 2011), but undergo substantial changes postnatally. Multiple reports have found that high strength and betweenness hubs in adulthood are largely located within within the default mode, attentional, sensorimotor, and visual areas (Cao et al., 2014; Fransson et al., 2011; Grayson et al., 2014; Hwang et al., 2013), whereas neonatal hubs primarily involve early sensorimotor, visual, and limbic regions (De Asis-Cruz et al., 2015; Fransson et al., 2011; Gao et al., 2011). The first two years of life reflect a gradual decrease in the hubness of sensorimotor areas and increased involvement of various association cortical areas, especially the medial frontal and parietal portions of the default mode (Gao et al., 2011). This pattern of hub strengthening bears substantial spatial and temporal overlap with the pattern of increasing within-module synchronization. This is consistent with the notion that node strength and betweenness in functional networks may serve, at least to some extent, as proxies for modular size and synchronization (Power et al., 2013). Thus, there continues to be much less known regarding what regions become more involved in the integration of different information processing streams (e.g. sensory, cognitive, and emotional) over time. Such anatomical areas are likely critical for the development of complex cognitive and social abilities, and are potentially uniquely susceptible to early environmental influences (Graham et al., 2015). It is also clear that several prenatal factors (e.g. immune and endocrine systems) strongly affect postnatal network trajectories (Canetta et al., 2016; Scheinost et al., 2016b). Future work examining node participation coefficients in functional networks and other centrality measures in structural networks (see *network measures*) promises to shed greater light on the emergence of network integrators and their potential susceptibility to pre- and post-natal factors.

## Global network metrics

Other important topological properties are also robustly evident at birth. Small-worldness (De Asis-Cruz et al., 2015; Fransson et al., 2011; Gao et al., 2011), as well as its constituent properties of global efficiency and average clustering coefficient, are already high in neonates. These properties only appear to marginally increase during the first year of life, plateauing thereafter until at least the end of the second year (Gao et al., 2011). Concomitantly, long-distance links are preferentially strengthened during the first year of life, leveling off thereafter (Gao et al., 2011). Rich clubs are also evident in both structural and functional networks in neonates along with both high functional modularity and integration capacity between modules (Ball et al., 2014; Scheinost et al., 2016a; van den Heuvel et al., 2015b). These findings suggest that the development of a small-world architecture, as well as the capacity for globally efficient communication and integration, largely occurs during the first year of life and prenatally, a finding that is supported by multimodal structural and functional network analysis in preterm infants (Ball et al., 2014; van den Heuvel et al., 2015b). This (very) early developmental trajectory stands in contrast to the more protracted progression towards increasingly segregated and refined modular architecture which continues into early childhood and beyond.



## Considerations in infant rs-fcMRI

A crucial caveat to the above observations is the potential for unidentified sources of artifact in infant studies (also see *Methodological challenges and recommendations*). In the case of identified networks in infants it is important to note that different community detection algorithms can provide unique results and all are not appropriate for any given experimental condition (Gates et al., 2016). Other considerations include early postnatal changes in cerebral vasculature, and altered state of consciousness when scanning very young participants. fMRI relies on hemodynamic measurements (usually BOLD), and there is good reason to suspect that infancy is accompanied by specific changes in neurovascular coupling (Hagmann et al., 2012). Human and animal studies indeed indicate differences in stimulus-evoked neurovascular coupling between infants and older individuals, but whether these differences impact rs-fcMRI measurements has yet to be determined (Graham et al., 2015). Infant imaging studies are also predominantly done during natural sleep (e.g. (Liu et al., 2008)) or sedation (e.g. (Fransson et al., 2007)). A natural benefit is the attenuation of micromovements and their attendant signal artifacts. However, altered state of arousal and consciousness might pose important confounds when comparing network findings to those seen in conscious children or adults. This concern is partly mitigated by findings of similar global functional network topology in conscious versus unconscious state in adults (Spoormaker et al., 2010), but key differences have been found in corticocortical (Heine et al., 2012; Horovitz et al., 2009; Spoormaker et al., 2010) and thalamocortical (Spoormaker et al., 2010) connectivity. Different sleep stages also correspond with different neurophysiological phenomena, which impact rs-fcMRI measurements (Laumann et al., 2016; Spoormaker et al., 2010). Until additional work can more definitively tease apart artifactual vs. neural contributions to the fMRI signal or awake versus sleep affects in these early postnatal years, developmental changes in modular partitions and other network changes must be viewed with at least some caution.

## Developmental refinements of network structure into adulthood

### General observations

Many of the network changes seen during infancy reflect long-term trajectories that extend into childhood and adolescence. As with the infant literature, much of the literature in children and adolescents has focused on the regions that define the adult DMN. Seed and component approaches have consistently found that connectivity between these areas (and indeed, connectivity within other cognitive RSNs) continues to strengthen from early childhood throughout development, especially with respect to long-range anterior-posterior links (de Bie et al., 2012; Fair et al., 2008; Sato et al., 2014; Sherman et al., 2014; Supekar et al., 2010).

More generally, one of the more reproducible findings of child and adolescent development (until very recently) had been an apparent shift from a local to global organization. Prior to 2012, multiple studies had found that functional connectivity between anatomically proximate ROIs gradually decreased, whereas long-range functional connections, especially anterior-posterior links, gradually increased until adulthood (Dosenbach et al., 2010; Fair et al., 2009; Kelly et al., 2009; Supekar et al., 2009). In addition, it appeared that the modular

assignments of regions shifted. In childhood, regional assignments appeared to strongly depend on local anatomy. With age, regions gradually became more defined by their respective functional roles (Power et al., 2011; Power et al., 2010). These trends pointed to coherent changes that involved a dampening of non-specific local spread of activity and strengthening of long-range links, facilitating cognitive development through modular specialization and selective cross-network integration (Bunge and Wright, 2007; Fair et al., 2009; Uddin, 2011; Uddin et al., 2011). Distance-dependent effects were even strong enough to enable reasonably accurate predictions of an individual subject's age on the basis of a single resting-state scan (Dosenbach et al., 2010).

However, recent concerns about the influence of head-motion artifact have raised some doubts about the interpretability of such broad trends. In particular, it has been revealed that motion increases non-specific local coupling and decreases long-range coupling in ways that are similar to differences found between children and adults (Fair et al., 2012b; Power et al., 2012; Satterthwaite et al., 2012; Yan et al., 2013). When processing techniques intended to remove motion artifact are applied, developmental effects of this type – both in terms of overall local versus distributed coupling and in terms of changing modular assignments – are substantially attenuated, though still present to a limited extent (Fair et al., 2012b; Power et al., 2012; Satterthwaite et al., 2012). As motion-denoising strategies continue to evolve, such findings will need continual re-evaluation. It is likely, however, that functional network maturation follows more precise spatiotemporal trajectories than previously understood. Indeed, there are now at least 3 reports that suggest the network structure in children as defined by community detection (see Figure 3) are largely similar to the adult (Fair et al., 2012b; Marek et al., 2015; Power et al., 2012). Despite this “adult like” level of organization, refinements within and between systems does appear to occur over time, although details to this end are still being worked out (Fair et al., 2012b; Gu et al., 2015b; Marek et al., 2015).

Much of this new work is finding that systems with different functional roles are characterized by substantially different trajectories (Gu et al., 2015b; Marek et al., 2015). For example, (Gu et al., 2015b) found that sensorimotor systems, already well-segregated in childhood, displayed little change into adulthood. On the other hand, networks involved in task-positive cognitive function (e.g. salience, frontoparietal executive, and attentional networks), being generally well-connected with other systems, became increasingly segregated and exhibited increasingly variable spatial structure. The DMN was found to exhibit both increasingly synchronized within-network and between-network connections. The authors contend that their results are consistent with notions that network development facilitates the ability to traverse through more diverse cognitive states in order to accomplish increasingly complex task demands (Bassett et al., 2011; Braun et al., 2015). Similar, albeit not identical developmental findings, are observed by (Marek et al., 2015), who found increased cross-network integration between the cingulo-opercular/salience and somatomotor modules over the age of 10–26, which they showed to be associated with the development of cognitive control. In agreement with these data, two additional reports also had findings that suggested that functional restructuring over this timespan disproportionately involves the cingulo-opercular and somatomotor modules (Fair et al., 2012b; Grayson et al., 2014), and potentially specific links between them (Grayson et al.,

2014) (see Figure 4). Therefore, there is recent convergence of data suggesting that network structure continues to evolve after late childhood in ways that are not primarily distance-dependent; rather, complex but specific changes may coincide with the development of specific cognitive abilities. Additional follow-up work using up-to-date methods for artifact prevention and removal, and more sensitive longitudinal designs, will be crucial for expanding on these important new findings.

### EEG/MEG findings

Resting-state networks obtained using electrophysiological recording (EEG or MEG) provide greater temporal resolution ( $<100\text{Hz}$ ) relative to rs-fMRI ( $<0.1\text{Hz}$ ) and therefore offer complementary insights about functional development. One of the classical and well-reproduced findings is that spectral power (i.e. amplitude of activity within different temporal frequency bands) decreases continually from 5–30 years of age across most frequency bands, but particularly in the slow-wave (0.5–7Hz) (Miskovic et al., 2015; Rodriguez-Martinez et al., 2015; Smit et al., 2012; Whitford et al., 2007). There is also a progressive shift in peak amplitude towards higher frequency activity (Miskovic et al., 2015; Rodriguez-Martinez et al., 2015; Smit et al., 2012; Whitford et al., 2007). Interestingly, frontal and parietal cortical gray matter (but not white matter) volume decreases across this same age bracket, and exhibits curvilinear trajectories that are closely correlated to the loss of slow-wave power in these same areas (Buchmann et al., 2011; Smit et al., 2012; Whitford et al., 2007). These coincident trajectories have provided key support for the hypothesis that developmental changes in baseline electrophysiological signatures may primarily be subserved by reduction of neuropil and progressive synapse elimination (Buchmann et al., 2011; Smit et al., 2012; Whitford et al., 2007).

EEG/MEG functional connectivity networks can also be constructed from synchronized activity within different frequency bands. Reports of resting-state EEG/MEG networks in infancy are scant (although see later section on temporal dynamics), but a number of network studies have been conducted regarding later development. It is important to note that these studies are generally unable to map activity to precise brain areas due to challenges in source localization, and often treat the scalp sensors as the network nodes. Nonetheless, there are several noteworthy findings. From roughly 5 years of age until adulthood, theta ( $\sim 4\text{Hz}$ ), alpha ( $\sim 10\text{Hz}$ ), and beta ( $\sim 20\text{Hz}$ ) oscillation networks exhibit concurrent increases in clustering coefficient (Boersma et al., 2011; Smit et al., 2012) and strength of long-range anterior-posterior and interhemispheric connections (Barry et al., 2004; Miskovic et al., 2015; Smit et al., 2012; Srinivasan, 1999). There is also consensus that network structure becomes increasingly non-random across this timespan (Boersma et al., 2011; Miskovic et al., 2015; Smit et al., 2012). Somewhat unexpectedly, two reports found path length to also increase with age (Boersma et al., 2011; Smit et al., 2012), although more recent work demonstrates age-related reductions (Miskovic et al., 2015). While this work is in its infancy, these studies thus far suggest some consensus between electrophysiological and MRI-based findings, implying that changes in brain organization are at least partially preserved across temporal scales of investigation.

## Regional hub metrics

Returning to the MRI literature, there is convergent evidence that the locations of high-strength and high-betweenness hub regions appear to be stable after 5 years of age (Cao et al., 2014; Grayson et al., 2014; Hwang et al., 2013; Wang et al., 2012; Wu et al., 2013; Zuo et al., 2012). However, there is also evidence for regional refinement of hub properties. Node strength and centrality of subcortical regions have both been reported to decrease in late childhood and adolescence (Sato et al., 2015; Supekar et al., 2009). Among cortical areas, some have reported increasing hubness of frontal nodes (Cao et al., 2014; Wu et al., 2013) while in other lobes there have been mixed reports of increasing (Cao et al., 2014; Grayson et al., 2014) and decreasing (Wu et al., 2013; Zuo et al., 2012), or stable (Hwang et al., 2013) hubness. At first glance, these mixed or non-significant findings might suggest that regional properties of cortical hubs are largely established during very early development. However, given that different networks display varying trajectories for within-network and between-network connectivity (Gu et al., 2015b; Marek et al., 2015), renewed examination of nodal hub-like properties such as the participation coefficient (which quantifies within vs. between module strength) may be warranted (e.g. see (Marek et al., 2015)). More detailed network analyses have revealed that specific connections to cortical hub regions increase during childhood and adolescence (Grayson et al., 2014; Hwang et al., 2013) resulting in increased rich club coefficients in adulthood (Cao et al., 2014; Grayson et al., 2014). Regional properties have also been found to exhibit nonlinear relationships with age (Wang et al., 2012), emphasizing the importance of mapping trajectories with more detailed temporal precision.

## Development of temporal dynamics

Here we will briefly discuss the topic of temporal dynamics in brain function, a rapidly emerging theme in the functional neuroimaging literature. First, we will point out that brain functional dynamics can relate to changes that occur over decades (e.g., myelination) or over fractions of a second (e.g. receptor turnover). Thus far, network changes in infancy, childhood, and adolescence have been discussed. This section discusses the growing literature of dynamics that occur over seconds and minutes measurable via rs-fcMRI.

Mounting evidence suggests that the functional connectome exhibits temporal instability that allows it to flexibly switch between multiple different configurations within a scanning session (Allen et al., 2014; Chang and Glover, 2010; Kang et al., 2011; Shen et al., 2015). There is evidence that this property is important for executive functioning, learning, and switching between challenging task demands (Bassett et al., 2011; Braun et al., 2015; Chen et al., 2016). Over development, rs-fcMRI networks show increased within-subject variability (Hutchison and Morton, 2015; Marusak et al., 2016; Qin et al., 2015), consistent with EEG studies showing that signal complexity increases over development (McIntosh et al., 2008; Vakorin et al., 2011). Recent simultaneous EEG-fMRI work (Fransson et al., 2013) links developmental differences (infants versus adults) in rs-fcMRI network dynamics with differences in EEG power spectra, consistent with the notion that temporal variability in very low frequency correlations is an emergent, hidden property of higher frequency power spectra (Chang et al., 2013; Tagliazucchi et al., 2012), which is known to change

continuously throughout development (Miskovic et al., 2015; Rodriguez-Martinez et al., 2015; Smit et al., 2012; Whitford et al., 2007). Given that different EEG frequency components have unique neurophysiological underpinnings (Miskovic et al., 2015), changing frequency spectra and temporal dynamics may point to clearer mechanistic substrates for network development.

Unfortunately, we have to repeat a recurring theme here and make yet another note of caution. Laumann et al. (Laumann et al., 2016) has recently provided strong evidence that much of the observed dynamics in rs-fcMRI networks primarily reflects noise (especially movement). The authors note that instability is not absent, but that it exists in the form of measurable state transitions (e.g. from awake to sleep, or eyes open versus closed), and that there is considerably less evidence for intrinsic dynamics otherwise. Other work has called into question some of the common statistical approaches to assessing dynamic FC (Hindriks et al., 2016; Leonardi and Van De Ville, 2015), which may not test against null models that properly account for random noise. It is also not clear whether the BOLD measurements are simply less sensitive to dynamic changes because of the sluggish nature of the signal. Altogether, these findings warrant more work, especially using simultaneous EEG-fMRI and animal models, to determine the potential role of network dynamics in development.

## Structure-function relationships over development

What are the underlying substrates of functional network development? This question is informed, in part, by previous studies investigating the correspondence between structural connectivity (SC) and FC. In humans, SC networks can be estimated using diffusion-weighted imaging (DWI), which allows for virtual reconstructions of white-matter tracts (a method known as tractography). There is an abundance of research demonstrating a correlation between DWI-tractography connection weights and FC in adults (Hagmann et al., 2008; Hagmann et al., 2010; Honey et al., 2009). More generally, SC networks can be combined with various modeling approaches to predict FC networks, allowing for detailed exploration of how the structural connectome gives rise to complex neural interactions. Models can involve highly parameterized simulations of regional activity (Hansen et al., 2015; Messe et al., 2014; Sanz-Leon et al., 2015), or sparser analytic predictions of FC using graph metrics of mutual communication (Goni et al., 2014; Grayson et al., 2016) (although, both approaches appear to be comparably effective; see Figure 1D). Predictive models have been extensively validated on both rs-fcMRI and EEG data (Chu et al., 2015), suggesting that SC strongly shapes and constrains activity across temporal scales.

A natural extension to this work would be to examine how structure-function correspondence develops, and whether optimized model parameters might evolve in tandem. Given that macroscale FC is modulated by both interregional connectivity strength as well as local population dynamics (Sanz-Leon et al., 2015), alterations in SC-FC coupling could potentially point to important shifts in the underlying drivers of various network phenomena. At least two reports have found abnormally low SC-FC correlation in epileptic patients (Chiang et al., 2015; Zhang et al., 2011), highlighting that aberrant SC-FC coupling may indeed indicate aberrations in neuronal physiology.

The first foray into developmental work was conducted in 2010, when Hagmann et al. (Hagmann et al., 2010) reported that SC-FC correspondence was higher in older teens versus in young children. Although the cause remains unknown, this result is broadly consistent with the notion that large white-matter bundles increase their capacity for information transfer with age via mechanisms such as increased myelination and axon diameter (Hagmann et al., 2012). Unfortunately, follow-up work on this topic in humans has been relatively scant, and this work was prior to new information regarding tractography artifacts (Calabrese et al., 2015; Donahue et al., 2016; Reveley et al., 2015; Thomas et al., 2014), and prior to systematic motion artifacts being demonstrated in diffusion-weighted MRI (Kong, 2014; Yendiki et al., 2014). The possibility for confounding artifactual contributions should therefore not be ruled out. However, new findings by Van den Heuvel (van den Heuvel et al., 2015b) show similar phenomena in preterm infants, such that neonates at 40 weeks gestational age have substantially greater SC-FC coupling than neonates born at 30 weeks. Another study found that age-related changes in FC disproportionately impacts polysynaptic connections (Betzel et al., 2014), although this study was performed across the lifespan and may reflect an effect of aging.

## The role of animal models

The importance of animal work for understanding structure-function relationships over development, and developmental fMRI findings in general, cannot be understated. The non-invasive measurements used to examine human populations are now readily available in rodent and monkey models where rich methodologies for studying genetic and pharmacologic manipulations exist. While clearly several cognitive domains in humans (e.g. language) and their relation to connectivity profiles are unlikely to be measured directly in animal models, network properties and structure-function relationships supporting sensory and cognitive function (see Figure 5) are attainable across species (Grayson et al., 2016; Miranda-Dominguez et al., 2014b; Stafford et al., 2014).

Why is this important? As is oft repeated in the current review, a very large portion of the current human developmental literature is in dire need of re-examination in light of recent developments regarding systematic artifacts in the imaging signals. Even with the current approaches aimed at optimizing signal versus noise, the question remains: What other lurking land mines are hidden in our data?

For example, structure-function modeling using MRI data suffers from a number of inherent limitations in DWI-tractography that have not been overcome by state-of-the-art acquisition and processing approaches (Calabrese et al., 2015; Donahue et al., 2016; Reveley et al., 2015; Thomas et al., 2014). Invasive tract-tracer injections, a method that is only feasible in animal models, remains the “gold standard” for identifying and quantifying interareal connection weights. In the monkey, whole-brain tract-tracer connection networks and tractography networks demonstrate a significant, but only moderately high correspondence. Currently, the highest reported correlation stands at  $r = 0.59$  (Donahue et al., 2016), though it is often much lower (van den Heuvel et al., 2015a). Future work could probably improve these relationships substantially by imposing rigorous anatomical constraints in the



tractography process (Smith et al., 2015), but such efforts have not seen extensive validation yet.

In the meantime, animal models are proving to be an important and necessary “bridge” for identifying ground-truth structure-to-function principles (Deco et al., 2014; Grayson et al., 2016; Miranda-Dominguez et al., 2014b; Stafford et al., 2014). One interesting aspect of the work in (Grayson et al., 2016) is how tightly functional imaging conditions had to be controlled in order to reveal novel structure-function relationships and in order to maximize model performance (see supplementary materials of (Grayson et al., 2016) for an extensive exposition). First, motion had to be reduced to undetectable levels with head fixation; second, the MR coil was specially optimized to enhance SNR; and third, a contrast agent was used to boost the functional signal. Without these efforts (which are not feasible in the large majority of human studies), critical structure-function relationships were much degraded.

Along the same lines, other factors outside of macro-anatomical connectivity likely affect FC measurements. Even the best-performing SC-FC models generally account for less than 50% of the variance in FC across the brain in both humans and non-human primates (Grayson et al., 2016; Messe et al., 2014; Misisic et al., 2015). Identifying these other hidden factors will be of critical importance to understanding the drivers of FC development in a given age range and why FC networks may go awry in various developmental disorders. More to the point, macroscale FC is known to be modulated by interareal correlated gene expression, areal densities of excitatory and inhibitory receptors, and other features of microscale architecture that vary across brain structures (Richiardi et al., 2015; Scholtens et al., 2014; Turk et al., 2016; van den Heuvel et al., 2016b) and undoubtedly evolve with age. Thus, more empirical mapping of brain structure and integration into a coherent modeling framework is needed. Of course, probing cellular and molecular markers across the whole human brain at high-throughput is exceptionally difficult (Richiardi et al., 2015) and cannot be done under strict experimental conditions. Experimental animal models should therefore remain indispensable for this purpose the foreseeable future.

## Clinical disorders of neurodevelopment

While this review has focused on studies of normative development, there is also a rich rs-fcMRI and EEG/MEG network literature in neurodevelopmental disorders. Complex, distributed disruptions in resting-state networks are consistently implicated in Schizophrenia (Baker et al., 2014; Khadka et al., 2013; Satterthwaite et al., 2015), ADHD (Castellanos and Aoki, 2016; Fair et al., 2012b), and Autism (Kessler et al., 2016; Moseley et al., 2015; Supekar et al., 2013) - though a thorough examination of this literature is beyond the scope of this review. Rather, we turn our attention here to two crucial challenges that are likely to dominate future research efforts into these conditions.

The first challenge is the ever-expanding evidence that Schizophrenia (Meda et al., 2014; Satterthwaite and Baker, 2015), ADHD (Costa Dias et al., 2015; Fair et al., 2012a; Gates et al., 2014; Karalunas et al., 2014), Autism (Jeste and Geschwind, 2014; Rudie et al., 2012), and potentially other DSM defined disorders, each constitutes a highly heterogeneous set of

etiologies. Indeed, recent work suggests that Autism Spectrum Disorder is characterized by increased subject-to-subject variability relative to the general population (Hahamy et al., 2015). Such findings are consistent with the notion that different pathologies in brain function may lead to related clinical outcomes, and underscore the importance of identifying individualized abnormalities.

Some recent work has already aimed at addressing this by attempting to identify brain-based subtypes within a diagnostic category, mostly through the application of various analytic clustering methods to multivariate neuroimaging or behavioral data. For an excellent review of recent progress, as well as promises and pitfalls in this area, we direct the reader to (Marquand et al., 2016). Another related approach to tackling heterogeneity would be to focus on deviations from typical brain organization at the individual-subject level, rather than performing classical between-group analyses. Cutting-edge work in resting-state fMRI acquisition and network analysis suggests that individual brains, even among typical subjects, are distinguishable via their functional connectomes (Finn et al., 2015; Laumann et al., 2015; Miranda-Dominguez et al., 2014b). Future work exploring the notion of the ‘functional fingerprint’ or ‘connectotype’ (Miranda-Dominguez et al., 2014a) may help to identify what deviations from the canonical functional connectome lead to atypical behavioral outcomes, versus what deviations may be considered to fall within the typical range. This would also be strongly in line with the National Institute of Health’s new initiative for precision medicine (Collins and Varmus, 2015).

The second challenge is in understanding the pathophysiological basis of network-level disturbances that are distributed and complex. In other words, if one sees atypical functional profiles amongst a specific set of brain regions or connections, does that suggest that the substance of the pathology is co-located? There are several reasons from a theoretical standpoint (Pessoa, 2014), which are largely supported empirically, why this is not necessarily the case. As an example, simulation experiments using structure-function models have predicted that even purely focal and transient perturbations in activity should result in diffuse network reorganization that affects distant brain systems (Alstott et al., 2009; Honey and Sporns, 2008; Misisic et al., 2015). In our recent paper, we presented experimental validation of these ideas through focal pharmacogenetic inactivation (Grayson et al., 2016), and similar findings were presented by (Andoh et al., 2015) using cortical transcranial magnetic stimulation. Both works demonstrate a causal effect on distributed resting-state connectivity and other neurophysiological signatures, driven by cascading neuronal interactions. These data empirically suggest that in a broad context, disease processes localized to one part of the brain can cause widespread neurophysiological disturbances in areas distant and unconnected from the pathology. This work also provides evidence that site-specific pathology can be deduced from distributed functional network disturbances (Grayson et al., 2016). However, in the absence of strong hypotheses regarding a ground-truth, one must be mindful that multiple structural network configurations can theoretically give rise to the same apparent aberrations in function – this is essentially the ‘many-to-many mappings’ problem very nicely described in (Pessoa, 2014). In summary, reducing these sources of complexity in neuropsychiatric case-control studies is likely to require integrated multimodal imaging approaches, more widespread application of SC-FC modeling, and

breakthrough innovations in imaging tissue structural and functional properties that relate to functional connectivity.

## Methodological challenges and recommendations

### FMRI signal artifacts

Rs-fcMRI investigators must be mindful that the fMRI signal is susceptible to a number of spurious sources of variance due to hardware instabilities, respiratory and cardiac signals, and head motion artifacts (Jo et al., 2010; Power et al., 2016). Effects of respiratory and cardiac activity can be attenuated through the use of temporal filtering, ICA-denoising, or directly acquiring these signals and removing them post-hoc. There is ongoing debate regarding the adequacy of these techniques (Jo et al., 2010; Power et al., 2016), and more work may be warranted to determine optimal ways to account for age-related physiological confounds. In addition, head motion has become a substantial concern since 2012 when a trio of papers pointed to systematic motion-related artifacts in rs-fcMRI (Power et al., 2012; Satterthwaite et al., 2012; Van Dijk et al., 2012). Chief among these findings was distance-dependent effects; head motion increases functional correlations between spatially proximal nodes and decreases correlations between distant nodes (Power et al., 2012). Motion tends to be greater in patient populations and younger children, posing a fundamentally important challenge for developmental research. Considerable work and debate is now ongoing to determine the optimal ‘denoising’ approaches during data processing (Patriat et al., 2016; Power et al., 2014). The most contentious debate probably concerns the use of global signal regression (GSR) – arguably the most effective motion-denoising technique currently available (Burgess et al., 2016; Power et al., 2015; Yan et al., 2013). Recent data reinforces concerns about artifactual contributions to the global signal in conventional 3T rs-fcMRI acquisitions sensitive to BOLD contrast (Grayson et al., 2016; Power et al., 2016), but also partly substantiates concerns regarding GSR under less conventional, high signal-to-noise scenarios where motion can be physically restricted (Grayson et al., 2016). Thus, GSR is likely to be a useful tool in developmental rs-fcMRI studies, where good control of such artifact is exceptionally important, but this debate may need to be revisited as methodologies in acquisition continue to advance. Moving forward, there is strong consensus about the importance of preventing head motion during acquisition and rigorously quantifying motion effects in study data (Goto et al., 2016; Power et al., 2015; Siegel et al., 2016).

We also wish to emphasize that future work using strict artifact removal are an increasing need for neuroimaging studies quite broadly – well beyond functional connectivity studies. Strong artifacts due to motion, in addition to their relevance for fMRI, have been noted in diffusion weighted imaging (Roalf et al., 2016; Yendiki et al., 2014) and cortical thickness measurements (Reuter et al., 2015; Savalia et al., 2017). Motion artifacts appear to be tightly related to clinical factors (Fair et al., 2012b) and a whole host of behavioral phenotypes and metrics (Siegel et al., 2016). Some of the techniques currently being used to remove these artifacts are non-optimal (Burgess et al., 2016; Goto et al., 2016). Thus, close examination of many of the developmental MR imaging findings is still warranted.

## Parcellation choice

Developmental network researchers also must make important decisions when defining the nodes (i.e. functional areas) under investigation. There is currently no consensus regarding an optimal set of criteria for defining an areal parcellation (Craddock et al., 2013; Glasser et al., 2016a; Gordon et al., 2016), and even less known about the stability of these parcellations at younger ages.

Parcellation density (i.e. number of regions) is a crucial consideration. Global topological metrics for structural and functional networks show stability across different parcellation schema of similar density, but vary considerably as a function of network size (de Reus and van den Heuvel, 2013; Fornito et al., 2010; Zalesky et al., 2010). In addition, with functional connectivity data some parcellation schemes can obscure known community structure. For example, regions or parcels that are overly large may encompass multiple distinct functional areas in ways that make community/network organization non-recognizable (Power et al., 2011). Functional network analyses that strongly relate to known network organization may therefore find signal homogeneity within parcels to be a useful optimization criterion for choosing a parcellation (e.g. see (Gordon et al., 2016) and (Glasser et al., 2016a).

In our opinion, the primary guiding factors in choosing a parcellation are based in fundamental principles of neuroscience. The cerebral cortex is both anatomically and functionally organized at many physical scales - starting at the level of single neurons and extending up to functional systems (Churchland and Sejnowski, 1992). At one scale are discrete regions of the cortex, known as functional areas. Functional areas possess unique internal structure (e.g. architectonics and topography). They also contain distinct combinations of inputs and outputs, which in turn group related areas into specialized networks (Felleman and Van Essen, 1991). These areal boundaries appear very early in development (Kostovic et al., 1995) and each area is thought to make a distinct contribution to information processing (Cohen et al., 2008; Felleman and Van Essen, 1991; Fuster, 2002; O'Leary and Nakagawa, 2002; Sur and Rubenstein, 2005). Ideally, parcellations for functional connectivity studies would closely reflect these areal boundaries in order to more clearly link network effects to the brain's underlying information processing architecture. In animal studies (rodents and non-human primates), this is feasible through the use of comprehensive, histologically and functionally defined whole-brain areal parcellations (Bezgin et al., 2012; Lein et al., 2007; Paxinos et al., 2000; Saleem and Logothetis, 2007; Swanson, 2015; Van Essen et al., 2012). In humans, there is much less clarity regarding potential areal boundaries (outside of elementary cytoarchitectonic divisions described, for example, by (Brodmann, 1909) and (Economo and Koskinas, 1925)) since a comparable amount of comprehensive data has not been readily available in a common reference space. Several groups, however, are finding ways to utilize today's non-invasive imaging tools to bring us closer to realizing these goals (Behrens et al., 2003; Glasser et al., 2016a; Gordon et al., 2016; Wig et al., 2014a; Wig et al., 2014b; Zhang et al., 2010).

## Atlas or template choice

A topic related to parcellation choice is how to spatially normalize individual subjects onto a common reference template. Cortical folding patterns change continually from birth into

adulthood, and especially during infancy. Infancy is also a time during which the image contrasts in various MR acquisitions changes dramatically. These changes pose challenges during template-based data processing steps such as spatial normalization and image segmentation, and are particularly problematic in studies that use template brains derived from only one age group and therefore rely on assumptions regarding similarity in shape characteristics of the brain across participant groups. The use of high-contrast, age-specific and study-specific template brain images for image registration is a well-validated approach to help overcome some of these confounds (Avants et al., 2010; Fonov et al., 2011), and is increasingly adopted in both structural and functional studies of (especially early) development (Eggebrecht et al., 2017; Fonov et al., 2011; Gao et al., 2015a; Scott et al., 2015). Surface based registrations that take into account the sulcal and gyral anatomy also help mitigate these confounds, and may be the future direction of the field (Fischl, 2012; Van Essen, 2012).

On top of these issues, functional areal boundaries (see *Parcellation Choice*) of heteromodal association areas are variable with respect to sulcal and gyral anatomy (Fischl et al., 2008). New multimodal approaches that also anchor surface-based registrations to other structural or functional characteristics of an individual are likely to improve subject to subject registrations, and hence reduce intersubject variability, even more (e.g. (Glasser et al., 2016b; Robinson et al., 2014)). Unfortunately, these advanced methods require more data acquisition time than is typically afforded for developmental studies. Nonetheless, a high quality anatomical image that is capable of being segmented properly into grey and white matter by itself is enough for the most basic surface-based registration tools.

### Test-retest reliability

A final point regarding the use of rs-fcMRI for network mapping concerns its reliability as a measurement tool. Multiple studies have found evidence for limited test-retest reliability using a multitude of conventional rs-fcMRI network characteristics and analytic approaches (Braun et al., 2012; Wang et al., 2011; Wisner et al., 2013; Zuo et al., 2010; Zuo and Xing, 2014). Recent evidence suggests that these effects may largely result from insufficient data acquisition. For instance, FC networks in one highly sampled individual converged as a function of the timeseries length used to extract correlations and were extremely reproducible using approximately 1 hour of motion-free data (Laumann et al., 2015). However, this length of scan time for each individual is likely prohibitive in most developmental rs-fcMRI studies, underscoring the importance of internal study replication. In addition, more work could be done to develop analytic methods with both high sensitivity to individual variation and high test-retest reliability (see (Miranda-Dominguez et al., 2014a)).

### Conclusions

This review has surveyed the development of correlated, resting-state activity and the emergence of network organization from birth until adulthood. Across all stages of postnatal development, functional brain networks exhibit non-trivial properties such as community structure, hub organization, cross-module integration, and global efficiency. Community

structure at birth is coarse and primitive, but gradually evolves into a denser, more spatially structured, and distributed organization. The most dramatic changes appear to occur in the first two years of life, though network refinement continues through childhood and adolescence, strengthening connections particularly among functional hubs. While the past decade has seen remarkable growth of network neuroscience and its application to development, there is still ample work that needs to be done. The field will benefit from resolving technical issues with acquisition and artifact removal, and from gaining a better understanding of the heterogeneity that exists across typical and atypical populations. Finally, we believe there is an urgent need to establish stronger links between different functional imaging modalities, between structural and functional connectivity, and between micro- and macro-scales of investigation via animal models.

## Acknowledgments

DSG was supported by an Autism Speaks Dennis Weatherstone Predoctoral Fellowship (#9591). This work was supported by the DeStefano Family Foundation, Gates Foundation, MacArthur foundation, and National Institutes of Health (UG3 OD023349, R01 MH105538, R01 MH096773, R01MH107508, P60-AA010760, R01 MH086654, R00MH091238).

## References

- Alcauter S, Lin W, Keith Smith J, Gilmore JH, Gao W. Consistent anterior-posterior segregation of the insula during the first 2 years of life. *Cereb Cortex*. 2015a; 25:1176–1187. [PubMed: 24248433]
- Alcauter S, Lin W, Smith JK, Goldman BD, Reznick JS, Gilmore JH, Gao W. Frequency of spontaneous BOLD signal shifts during infancy and correlates with cognitive performance. *Dev Cogn Neurosci*. 2015b; 12:40–50. [PubMed: 25459875]
- Allen EA, Damaraju E, Plis SM, Erhardt EB, Eichele T, Calhoun VD. Tracking whole-brain connectivity dynamics in the resting state. *Cereb Cortex*. 2014; 24:663–676. [PubMed: 23146964]
- Alstott J, Breakspear M, Hagmann P, Cammoun L, Sporns O. Modeling the impact of lesions in the human brain. *PLoS Comput Biol*. 2009; 5:e1000408. [PubMed: 19521503]
- Andoh J, Matsushita R, Zatorre RJ. Asymmetric Interhemispheric Transfer in the Auditory Network: Evidence from TMS, Resting-State fMRI, and Diffusion Imaging. *J Neurosci*. 2015; 35:14602–14611. [PubMed: 26511249]
- Avants BB, Yushkevich P, Pluta J, Minkoff D, Korczykowski M, Detre J, Gee JC. The optimal template effect in hippocampus studies of diseased populations. *NeuroImage*. 2010; 49:2457–2466. [PubMed: 19818860]
- Baker JT, Holmes AJ, Masters GA, Yeo BT, Krienen F, Buckner RL, Ongur D. Disruption of cortical association networks in schizophrenia and psychotic bipolar disorder. *JAMA Psychiatry*. 2014; 71:109–118. [PubMed: 24306091]
- Ball G, Aljabar P, Zebari S, Tusor N, Arichi T, Merchant N, Robinson EC, Ogundipe E, Rueckert D, Edwards AD, et al. Rich-club organization of the newborn human brain. *Proc Natl Acad Sci U S A*. 2014; 111:7456–7461. [PubMed: 24799693]
- Barry RJ, Clarke AR, McCarthy R, Selikowitz M, Johnstone SJ, Rushby JA. Age and gender effects in EEG coherence: I. Developmental trends in normal children. *Clin Neurophysiol*. 2004; 115:2252–2258. [PubMed: 15351366]
- Bassett DS, Wymbs NF, Porter MA, Mucha PJ, Carlson JM, Grafton ST. Dynamic reconfiguration of human brain networks during learning. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:7641–7646. [PubMed: 21502525]
- Beckmann CF, Smith SM. Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans Med Imaging*. 2004; 23:137–152. [PubMed: 14964560]
- Behrens TE, Johansen-Berg H, Woolrich MW, Smith SM, Wheeler-Kingshott CA, Boulby PA, Barker GJ, Sillery EL, Sheehan K, Ciccarelli O, et al. Non-invasive mapping of connections between



- human thalamus and cortex using diffusion imaging. *Nat Neurosci.* 2003; 6:750–757. [PubMed: 12808459]
- Betzel RF, Byrge L, He Y, Goni J, Zuo XN, Sporns O. Changes in structural and functional connectivity among resting-state networks across the human lifespan. *NeuroImage.* 2014; 102(Pt 2):345–357. [PubMed: 25109530]
- Bezgin G, Vakorin VA, van Opstal AJ, McIntosh AR, Bakker R. Hundreds of brain maps in one atlas: registering coordinate-independent primate neuro-anatomical data to a standard brain. *NeuroImage.* 2012; 62:67–76. [PubMed: 22521477]
- Biswal B, Yetkin FZ, Haughton VM, Hyde JS. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine/Society of Magnetic Resonance in Medicine.* 1995; 34:537–541.
- Boersma M, Smit DJ, de Bie HM, Van Baal GC, Boomsma DI, de Geus EJ, Delemarre-van de Waal HA, Stam CJ. Network analysis of resting state EEG in the developing young brain: structure comes with maturation. *Hum Brain Mapp.* 2011; 32:413–425. [PubMed: 20589941]
- Braun U, Plichta MM, Esslinger C, Sauer C, Haddad L, Grimm O, Mier D, Mohnke S, Heinz A, Erk S, et al. Test-retest reliability of resting-state connectivity network characteristics using fMRI and graph theoretical measures. *NeuroImage.* 2012; 59:1404–1412. [PubMed: 21888983]
- Braun U, Schafer A, Walter H, Erk S, Romanczuk-Seiferth N, Haddad L, Schweiger JI, Grimm O, Heinz A, Tost H, et al. Dynamic reconfiguration of frontal brain networks during executive cognition in humans. *Proc Natl Acad Sci U S A.* 2015; 112:11678–11683. [PubMed: 26324898]
- Brodmann, K. Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Leipzig: Barth; 1909.
- Buchmann A, Ringli M, Kurth S, Schaerer M, Geiger A, Jenni OG, Huber R. EEG sleep slow-wave activity as a mirror of cortical maturation. *Cereb Cortex.* 2011; 21:607–615. [PubMed: 20624840]
- Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci.* 2008; 1124:1–38. [PubMed: 18400922]
- Bunge SA, Wright SB. Neurodevelopmental changes in working memory and cognitive control. *Curr Opin Neurobiol.* 2007; 17:243–250. [PubMed: 17321127]
- Burgess GC, Kandala S, Nolan D, Laumann TO, Power JD, Adeyemo B, Harms MP, Petersen SE, Barch DM. Evaluation of Denoising Strategies to Address Motion-Related Artifacts in Resting-State Functional Magnetic Resonance Imaging Data from the Human Connectome Project. *Brain Connect.* 2016; 6:669–680. [PubMed: 27571276]
- Calabrese E, Badea A, Cofer G, Qi Y, Johnson GA. A Diffusion MRI Tractography Connectome of the Mouse Brain and Comparison with Neuronal Tracer Data. *Cereb Cortex.* 2015; 25:4628–4637. [PubMed: 26048951]
- Canetta S, Bolkan S, Padilla-Coreano N, Song LJ, Sahn R, Harrison NL, Gordon JA, Brown A, Kellendonk C. Maternal immune activation leads to selective functional deficits in offspring parvalbumin interneurons. *Mol Psychiatry.* 2016; 21:956–968. [PubMed: 26830140]
- Cao M, Wang JH, Dai ZJ, Cao XY, Jiang LL, Fan FM, Song XW, Xia MR, Shu N, Dong Q, et al. Topological organization of the human brain functional connectome across the lifespan. *Dev Cogn Neurosci.* 2014; 7:76–93. [PubMed: 24333927]
- Castellanos FX, Aoki Y. Intrinsic Functional Connectivity in Attention-Deficit/Hyperactivity Disorder: A Science in Development. *Biol Psychiatry Cogn Neurosci Neuroimaging.* 2016; 1:253–261. [PubMed: 27713929]
- Chang C, Glover GH. Time-frequency dynamics of resting-state brain connectivity measured with fMRI. *NeuroImage.* 2010; 50:81–98. [PubMed: 20006716]
- Chang C, Liu Z, Chen MC, Liu X, Duyn JH. EEG correlates of time-varying BOLD functional connectivity. *NeuroImage.* 2013; 72:227–236. [PubMed: 23376790]
- Chen T, Cai W, Ryali S, Supekar K, Menon V. Distinct Global Brain Dynamics and Spatiotemporal Organization of the Salience Network. *PLoS Biol.* 2016; 14:e1002469. [PubMed: 27270215]
- Chiang S, Stern JM, Engel J Jr, Haneef Z. Structural-functional coupling changes in temporal lobe epilepsy. *Brain Res.* 2015; 1616:45–57. [PubMed: 25960346]

- Chu CJ, Tanaka N, Diaz J, Edlow BL, Wu O, Hamalainen M, Stufflebeam S, Cash SS, Kramer MA. EEG functional connectivity is partially predicted by underlying white matter connectivity. *NeuroImage*. 2015; 108:23–33. [PubMed: 25534110]
- Churchland, PS., Sejnowski, TJ. *The computational brain*. Cambridge, Mass: MIT Press; 1992.
- Cohen AL, Fair DA, Dosenbach NU, Miezin FM, Dierker D, Van Essen DC, Schlaggar BL, Petersen SE. Defining functional areas in individual human brains using resting functional connectivity MRI. *NeuroImage*. 2008; 41:45–57. [PubMed: 18367410]
- Cohen JR, D’Esposito M. The Segregation and Integration of Distinct Brain Networks and Their Relationship to Cognition. *J Neurosci*. 2016; 36:12083–12094. [PubMed: 27903719]
- Cole MW, Bassett DS, Power JD, Braver TS, Petersen SE. Intrinsic and task-evoked network architectures of the human brain. *Neuron*. 2014; 83:238–251. [PubMed: 24991964]
- Cole MW, Yarkoni T, Repovs G, Anticevic A, Braver TS. Global connectivity of prefrontal cortex predicts cognitive control and intelligence. *J Neurosci*. 2012; 32:8988–8999. [PubMed: 22745498]
- Collin G, Sporns O, Mandl RCW, van den Heuvel MP. *Structural and Functional Aspects Relating to Cost and Benefit of Rich Club Organization in the Human Cerebral Cortex*. Cerebral cortex (New York, NY: 1991). 2013
- Collins FS, Varmus H. A new initiative on precision medicine. *N Engl J Med*. 2015; 372:793–795. [PubMed: 25635347]
- Cordes D, Haughton VM, Arfanakis K, Carew JD, Turski PA, Moritz CH, Quigley MA, Meyerand ME. Frequencies contributing to functional connectivity in the cerebral cortex in “resting-state” data. *AJNR Am J Neuroradiol*. 2001; 22:1326–1333. [PubMed: 11498421]
- Costa Dias TG, Iyer SP, Carpenter SD, Cary RP, Wilson VB, Mitchell SH, Nigg JT, Fair DA. Characterizing heterogeneity in children with and without ADHD based on reward system connectivity. *Dev Cogn Neurosci*. 2015; 11:155–174. [PubMed: 25660033]
- Courchesne E, Chisum HJ, Townsend J, Cowles A, Covington J, Egaas B, Harwood M, Hinds S, Press GA. Normal brain development and aging: quantitative analysis at in vivo MR imaging in healthy volunteers. *Radiology*. 2000; 216:672–682. [PubMed: 10966694]
- Craddock RC, Jbabdi S, Yan CG, Vogelstein JT, Castellanos FX, Di Martino A, Kelly C, Heberlein K, Colcombe S, Milham MP. Imaging human connectomes at the macroscale. *Nat Methods*. 2013; 10:524–539. [PubMed: 23722212]
- Damaraju E, Caprihan A, Lowe JR, Allen EA, Calhoun VD, Phillips JP. Functional connectivity in the developing brain: a longitudinal study from 4 to 9 months of age. *NeuroImage*. 2014; 84:169–180. [PubMed: 23994454]
- Damoiseaux JS, Rombouts SA, Barkhof F, Scheltens P, Stam CJ, Smith SM, Beckmann CF. Consistent resting-state networks across healthy subjects. *Proc Natl Acad Sci U S A*. 2006; 103:13848–13853. [PubMed: 16945915]
- De Asis-Cruz J, Bouyssi-Kobar M, Evangelou I, Vezina G, Limperopoulos C. Functional properties of resting state networks in healthy full-term newborns. *Sci Rep*. 2015; 5:17755. [PubMed: 26639607]
- de Bie HM, Boersma M, Adriaanse S, Veltman DJ, Wink AM, Roosendaal SD, Barkhof F, Stam CJ, Oostrom KJ, Delemarre-van de Waal HA, et al. Resting-state networks in awake five- to eight-year old children. *Hum Brain Mapp*. 2012; 33:1189–1201. [PubMed: 21520347]
- de Reus MA, van den Heuvel MP. The parcellation-based connectome: limitations and extensions. *NeuroImage*. 2013; 80:397–404. [PubMed: 23558097]
- de Reus MA, van den Heuvel MP. Simulated rich club lesioning in brain networks: a scaffold for communication and integration? *Front Hum Neurosci*. 2014; 8:647. [PubMed: 25191259]
- Deco G, McIntosh AR, Shen K, Hutchison RM, Menon RS, Everling S, Hagmann P, Jirsa VK. Identification of optimal structural connectivity using functional connectivity and neural modeling. *J Neurosci*. 2014; 34:7910–7916. [PubMed: 24899713]
- Donahue CJ, Sotiropoulos SN, Jbabdi S, Hernandez-Fernandez M, Behrens TE, Dyrby TB, Coalson T, Kennedy H, Knoblauch K, Van Essen DC, et al. Using Diffusion Tractography to Predict Cortical Connection Strength and Distance: A Quantitative Comparison with Tracers in the Monkey. *J Neurosci*. 2016; 36:6758–6770. [PubMed: 27335406]

- Dosenbach NU, Nardos B, Cohen AL, Fair DA, Power JD, Church JA, Nelson SM, Wig GS, Vogel AC, Lessov-Schlaggar CN, et al. Prediction of individual brain maturity using fMRI. *Science*. 2010; 329:1358–1361. [PubMed: 20829489]
- Doucet G, Naveau M, Petit L, Delcroix N, Zago L, Crivello F, Jobard G, Tzourio-Mazoyer N, Mazoyer B, Mellet E, et al. Brain activity at rest: a multiscale hierarchical functional organization. *J Neurophysiol*. 2011; 105:2753–2763. [PubMed: 21430278]
- Dwyer DB, Harrison BJ, Yucel M, Whittle S, Zalesky A, Pantelis C, Allen NB, Fornito A. Large-scale brain network dynamics supporting adolescent cognitive control. *J Neurosci*. 2014; 34:14096–14107. [PubMed: 25319705]
- Economo, C., Koskinas, GN. Die cytoarchitektonik der hirnrinde des erwachsenen menschen. Wien und Berlin: J. Springer; 1925.
- Eggebrecht AT, Elison JT, Feczko E, Todorov A, Wolff JJ, Kandala S, Adams CM, Snyder AZ, Lewis JD, Estes AM, et al. Joint Attention and Brain Functional Connectivity in Infants and Toddlers. *Cereb Cortex*. 2017
- Fair DA, Bathula D, Nikolas MA, Nigg JT. Distinct neuropsychological subgroups in typically developing youth inform heterogeneity in children with ADHD. *Proc Natl Acad Sci U S A*. 2012a; 109:6769–6774. [PubMed: 22474392]
- Fair DA, Cohen AL, Dosenbach NU, Church JA, Miezin FM, Barch DM, Raichle ME, Petersen SE, Schlaggar BL. The maturing architecture of the brain's default network. *Proc Natl Acad Sci U S A*. 2008; 105:4028–4032. [PubMed: 18322013]
- Fair DA, Cohen AL, Power JD, Dosenbach NUF, Church JA, Miezin FM, Schlaggar BL, Petersen SE. Functional brain networks develop from a “local to distributed” organization. *PLoS Comp Biol*. 2009;5.
- Fair, Da, Nigg, JT., Iyer, S., Bathula, D., Mills, KL., Dosenbach, NUF., Schlaggar, BL., Mennes, M., Gutman, D., Bangaru, S., et al. Distinct neural signatures detected for ADHD subtypes after controlling for micro-movements in resting state functional connectivity MRI data. *Frontiers in systems neuroscience*. 2012b; 6:80. [PubMed: 23382713]
- Felleman DJ, Van Essen DC. Distributed hierarchical processing in the primate cerebral cortex. *Cereb Cortex*. 1991; 1:1–47. [PubMed: 1822724]
- Finn ES, Shen X, Scheinost D, Rosenberg MD, Huang J, Chun MM, Papademetris X, Constable RT. Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat Neurosci*. 2015; 18:1664–1671. [PubMed: 26457551]
- Fischl B. *FreeSurfer*. *NeuroImage*. 2012; 62:774–781. [PubMed: 22248573]
- Fischl B, Rajendran N, Busa E, Augustinack J, Hinds O, Yeo BT, Mohlberg H, Amunts K, Zilles K. Cortical folding patterns and predicting cytoarchitecture. *Cereb Cortex*. 2008; 18:1973–1980. [PubMed: 18079129]
- Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL. Unbiased average age-appropriate atlases for pediatric studies. *NeuroImage*. 2011; 54:313–327. [PubMed: 20656036]
- Fornito A, Zalesky A, Bullmore ET. Network scaling effects in graph analytic studies of human resting-state FMRI data. *Front Syst Neurosci*. 2010; 4:22. [PubMed: 20592949]
- Fransson P, Aden U, Blennow M, Lagercrantz H. The functional architecture of the infant brain as revealed by resting-state fMRI. *Cereb Cortex*. 2011; 21:145–154. [PubMed: 20421249]
- Fransson P, Metsaranta M, Blennow M, Aden U, Lagercrantz H, Vanhatalo S. Early development of spatial patterns of power-law frequency scaling in FMRI resting-state and EEG data in the newborn brain. *Cereb Cortex*. 2013; 23:638–646. [PubMed: 22402348]
- Fransson P, Skiold B, Horsch S, Nordell A, Blennow M, Lagercrantz H, Aden U. Resting-state networks in the infant brain. *Proc Natl Acad Sci U S A*. 2007; 104:15531–15536. [PubMed: 17878310]
- Fuster JM. Frontal lobe and cognitive development. *J Neurocytol*. 2002; 31:373–385. [PubMed: 12815254]
- Gao W, Alcauter S, Elton A, Hernandez-Castillo CR, Smith JK, Ramirez J, Lin W. Functional Network Development During the First Year: Relative Sequence and Socioeconomic Correlations. *Cereb Cortex*. 2015a; 25:2919–2928. [PubMed: 24812084]

- Gao W, Alcauter S, Smith JK, Gilmore JH, Lin W. Development of human brain cortical network architecture during infancy. *Brain Struct Funct*. 2015b; 220:1173–1186. [PubMed: 24469153]
- Gao W, Gilmore JH, Giovanello KS, Smith JK, Shen D, Zhu H, Lin W. Temporal and spatial evolution of brain network topology during the first two years of life. *PLoS One*. 2011; 6:e25278. [PubMed: 21966479]
- Gao W, Gilmore JH, Shen D, Smith JK, Zhu H, Lin W. The synchronization within and interaction between the default and dorsal attention networks in early infancy. *Cereb Cortex*. 2013; 23:594–603. [PubMed: 22368080]
- Gao W, Lin W, Grewen K, Gilmore JH. Functional Connectivity of the Infant Human Brain: Plastic and Modifiable. *Neuroscientist*. 2016
- Gao W, Zhu H, Giovanello KS, Smith JK, Shen D, Gilmore JH, Lin W. Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old healthy pediatric subjects. *Proc Natl Acad Sci U S A*. 2009; 106:6790–6795. [PubMed: 19351894]
- Gates KM, Henry T, Steinley D, Fair DA. A Monte Carlo Evaluation of Weighted Community Detection Algorithms. *Front Neuroinform*. 2016; 10:45. [PubMed: 27891087]
- Gates KM, Molenaar PC, Iyer SP, Nigg JT, Fair DA. Organizing heterogeneous samples using community detection of GIMME-derived resting state functional networks. *PLoS One*. 2014; 9:e91322. [PubMed: 24642753]
- Geng X, Li G, Lu Z, Gao W, Wang L, Shen D, Zhu H, Gilmore JH. Structural and Maturational Covariance in Early Childhood Brain Development. *Cereb Cortex*. 2016
- Gilmore JH, Shi F, Woolson SL, Knickmeyer RC, Short SJ, Lin W, Zhu H, Hamer RM, Styner M, Shen D. Longitudinal development of cortical and subcortical gray matter from birth to 2 years. *Cereb Cortex*. 2012; 22:2478–2485. [PubMed: 22109543]
- Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson J, Beckmann CF, Jenkinson M, et al. A multi-modal parcellation of human cerebral cortex. *Nature*. 2016a; 536:171–178. [PubMed: 27437579]
- Glasser MF, Smith SM, Marcus DS, Andersson JL, Auerbach EJ, Behrens TE, Coalson TS, Harms MP, Jenkinson M, Moeller S, et al. The Human Connectome Project's neuroimaging approach. *Nat Neurosci*. 2016b; 19:1175–1187. [PubMed: 27571196]
- Goni J, van den Heuvel MP, Avena-Koenigsberger A, Velez de Mendizabal N, Betzel RF, Griffa A, Hagmann P, Corominas-Murtra B, Thiran JP, Sporns O. Resting-brain functional connectivity predicted by analytic measures of network communication. *Proc Natl Acad Sci U S A*. 2014; 111:833–838. [PubMed: 24379387]
- Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex*. 2016; 26:288–303. [PubMed: 25316338]
- Goto M, Abe O, Miyati T, Yamasue H, Gomi T, Takeda T. Head Motion and Correction Methods in Resting-state Functional MRI. *Magn Reson Med Sci*. 2016; 15:178–186. [PubMed: 26701695]
- Graham AM, Pfeifer JH, Fisher PA, Lin W, Gao W, Fair DA. The potential of infant fMRI research and the study of early life stress as a promising exemplar. *Dev Cogn Neurosci*. 2015; 12:12–39. [PubMed: 25459874]
- Grayson DS, Bliss-Moreau E, Machado CJ, Bennett J, Shen K, Grant KA, Fair DA, Amaral DG. The Rhesus Monkey Connectome Predicts Disrupted Functional Networks Resulting from Pharmacogenetic Inactivation of the Amygdala. *Neuron*. 2016; 91:453–466. [PubMed: 27477019]
- Grayson DS, Ray S, Carpenter S, Iyer S, Dias TGC, Stevens C, Nigg JT, Fair Da. Structural and functional rich club organization of the brain in children and adults. *PLoS one*. 2014; 9:e88297. [PubMed: 24505468]
- Gu S, Pasqualetti F, Cieslak M, Telesford QK, Yu AB, Kahn AE, Medaglia JD, Vettel JM, Miller MB, Grafton ST, et al. Controllability of structural brain networks. *Nat Commun*. 2015a; 6:8414. [PubMed: 26423222]
- Gu S, Satterthwaite TD, Medaglia JD, Yang M, Gur RE, Gur RC, Bassett DS. Emergence of system roles in normative neurodevelopment. *Proc Natl Acad Sci U S A*. 2015b; 112:13681–13686. [PubMed: 26483477]

- Guimera R, Nunes Amaral LA. Functional cartography of complex metabolic networks. *Nature*. 2005; 433:895–900. [PubMed: 15729348]
- Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey CJ, Van Wassenhove J, Sporns O. Mapping the structural core of human cerebral cortex. *PLoS Biol*. 2008; 6:1479–1493.
- Hagmann P, Grant PE, Fair DA. MR connectomics: a conceptual framework for studying the developing brain. In *Frontiers in Systems Neuroscience*. 2012
- Hagmann P, Sporns O, Madan N, Cammoun L, Pienaar R, Wedeen VJ, Meuli R, Thiran JP, Grant PE. White matter maturation reshapes structural connectivity in the late developing human brain. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:19067–19072. [PubMed: 20956328]
- Hahamy A, Behrmann M, Malach R. The idiosyncratic brain: distortion of spontaneous connectivity patterns in autism spectrum disorder. *Nat Neurosci*. 2015; 18:302–309. [PubMed: 25599222]
- Hansen EC, Battaglia D, Spiegler A, Deco G, Jirsa VK. Functional connectivity dynamics: modeling the switching behavior of the resting state. *NeuroImage*. 2015; 105:525–535. [PubMed: 25462790]
- Heine L, Soddu A, Gomez F, Vanhauwenhuysse A, Tshibanda L, Thonnard M, Charland-Verville V, Kirsch M, Laureys S, Demertzi A. Resting state networks and consciousness: alterations of multiple resting state network connectivity in physiological, pharmacological, and pathological consciousness States. *Front Psychol*. 2012; 3:295. [PubMed: 22969735]
- Hindriks R, Adhikari MH, Murayama Y, Ganzetti M, Mantini D, Logothetis NK, Deco G. Can sliding-window correlations reveal dynamic functional connectivity in resting-state fMRI? *NeuroImage*. 2016; 127:242–256. [PubMed: 26631813]
- Homae F, Watanabe H, Otobe T, Nakano T, Go T, Konishi Y, Taga G. Development of global cortical networks in early infancy. *J Neurosci*. 2010; 30:4877–4882. [PubMed: 20371807]
- Honey CJ, Sporns O. Dynamical consequences of lesions in cortical networks. *Hum Brain Mapp*. 2008; 29:802–809. [PubMed: 18438885]
- Honey CJ, Sporns O, Cammoun L, Gigandet X, Thiran JP, Meuli R, Hagmann P. Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106:2035–2040. [PubMed: 19188601]
- Horovitz SG, Braun AR, Carr WS, Picchioni D, Balkin TJ, Fukunaga M, Duyn JH. Decoupling of the brain's default mode network during deep sleep. *Proc Natl Acad Sci U S A*. 2009; 106:11376–11381. [PubMed: 19549821]
- Hutchison RM, Morton JB. Tracking the Brain's Functional Coupling Dynamics over Development. *J Neurosci*. 2015; 35:6849–6859. [PubMed: 25926460]
- Huttenlocher, PR. *Neural plasticity: the effects of environment on the development of the cerebral cortex*. Cambridge, MA: Harvard University Press; 2002.
- Hwang K, Hallquist MN, Luna B. The development of hub architecture in the human functional brain network. *Cerebral cortex (New York, NY: 1991)*. 2013; 23:2380–2393.
- Innocenti GM, Price DJ. Exuberance in the development of cortical networks. *Nat Rev Neurosci*. 2005; 6:955–965. [PubMed: 16288299]
- Jeste SS, Geschwind DH. Disentangling the heterogeneity of autism spectrum disorder through genetic findings. *Nat Rev Neurol*. 2014; 10:74–81. [PubMed: 24468882]
- Jo HJ, Saad ZS, Simmons WK, Milbury LA, Cox RW. Mapping sources of correlation in resting state FMRI, with artifact detection and removal. *NeuroImage*. 2010; 52:571–582. [PubMed: 20420926]
- Kang J, Wang L, Yan C, Wang J, Liang X, He Y. Characterizing dynamic functional connectivity in the resting brain using variable parameter regression and Kalman filtering approaches. *NeuroImage*. 2011; 56:1222–1234. [PubMed: 21420500]
- Karalunas SL, Fair D, Musser ED, Aykes K, Iyer SP, Nigg JT. Subtyping attention-deficit/hyperactivity disorder using temperament dimensions: toward biologically based nosologic criteria. *JAMA Psychiatry*. 2014; 71:1015–1024. [PubMed: 25006969]
- Kelly AM, Di Martino A, Uddin LQ, Shehzad Z, Gee DG, Reiss PT, Margulies DS, Castellanos FX, Milham MP. Development of anterior cingulate functional connectivity from late childhood to early adulthood. *Cereb Cortex*. 2009; 19:640–657. [PubMed: 18653667]



- Kessler K, Seymour RA, Rippon G. Brain oscillations and connectivity in autism spectrum disorders (ASD): new approaches to methodology, measurement and modelling. *Neurosci Biobehav Rev.* 2016; 71:601–620. [PubMed: 27720724]
- Khadka S, Meda SA, Stevens MC, Glahn DC, Calhoun VD, Sweeney JA, Tamminga CA, Keshavan MS, O'Neil K, Schretlen D, et al. Is aberrant functional connectivity a psychosis endophenotype? A resting state functional magnetic resonance imaging study. *Biol Psychiatry.* 2013; 74:458–466. [PubMed: 23746539]
- Kitzbichler MG, Henson RN, Smith ML, Nathan PJ, Bullmore ET. Cognitive effort drives workspace configuration of human brain functional networks. *J Neurosci.* 2011; 31:8259–8270. [PubMed: 21632947]
- Kiviniemi V, Jauhiainen J, Tervonen O, Paakko E, Oikarinen J, Vainionpaa V, Rantala H, Biswal B. Slow vasomotor fluctuation in fMRI of anesthetized child brain. *Magn Reson Med.* 2000; 44:373–378. [PubMed: 10975887]
- Knickmeyer RC, Gouttard S, Kang C, Evans D, Wilber K, Smith JK, Hamer RM, Lin W, Gerig G, Gilmore JH. A structural MRI study of human brain development from birth to 2 years. *J Neurosci.* 2008; 28:12176–12182. [PubMed: 19020011]
- Kong XZ. Association between in-scanner head motion with cerebral white matter microstructure: a multiband diffusion-weighted MRI study. *PeerJ.* 2014; 2:e366. [PubMed: 24795856]
- Kostovic I, Judas M, Petanjek Z, Simic G. Ontogenesis of goal-directed behavior: anatomo-functional considerations. *Int J Psychophysiol.* 1995; 19:85–102. [PubMed: 7622411]
- Laumann TO, Gordon EM, Adeyemo B, Snyder AZ, Joo SJ, Chen MY, Gilmore AW, McDermott KB, Nelson SM, Dosenbach NU, et al. Functional System and Areal Organization of a Highly Sampled Individual Human Brain. *Neuron.* 2015; 87:657–670. [PubMed: 26212711]
- Laumann TO, Snyder AZ, Mitra A, Gordon EM, Gratton C, Adeyemo B, Gilmore AW, Nelson SM, Berg JJ, Greene DJ, et al. On the Stability of BOLD fMRI Correlations. *Cereb Cortex.* 2016
- Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, Boe AF, Boguski MS, Brockway KS, Byrnes EJ, et al. Genome-wide atlas of gene expression in the adult mouse brain. *Nature.* 2007; 445:168–176. [PubMed: 17151600]
- Leonardi N, Van De Ville D. On spurious and real fluctuations of dynamic functional connectivity during rest. *NeuroImage.* 2015; 104:430–436. [PubMed: 25234118]
- Lewis CM, Baldassarre A, Committeri G, Romani GL, Corbetta M. Learning sculpts the spontaneous activity of the resting human brain. *Proc Natl Acad Sci U S A.* 2009; 106:17558–17563. [PubMed: 19805061]
- Li G, Lin W, Gilmore JH, Shen D. Spatial Patterns, Longitudinal Development, and Hemispheric Asymmetries of Cortical Thickness in Infants from Birth to 2 Years of Age. *J Neurosci.* 2015; 35:9150–9162. [PubMed: 26085637]
- Li Y, Liu Y, Li J, Qin W, Li K, Yu C, Jiang T. Brain anatomical network and intelligence. *PLoS Comput Biol.* 2009; 5:e1000395. [PubMed: 19492086]
- Lin W, Zhu Q, Gao W, Chen Y, Toh CH, Styner M, Gerig G, Smith JK, Biswal B, Gilmore JH. Functional connectivity MR imaging reveals cortical functional connectivity in the developing brain. *AJNR Am J Neuroradiol.* 2008; 29:1883–1889. [PubMed: 18784212]
- Liu WC, Flax JF, Guise KG, Sukul V, Benasich AA. Functional connectivity of the sensorimotor area in naturally sleeping infants. *Brain Res.* 2008; 1223:42–49. [PubMed: 18599026]
- Lohmann G, Margulies DS, Horstmann A, Pleger B, Lepsien J, Goldhahn D, Schloegl H, Stumvoll M, Villringer A, Turner R. Eigenvector centrality mapping for analyzing connectivity patterns in fMRI data of the human brain. *PLoS One.* 2010; 5:e10232. [PubMed: 20436911]
- Lohse C, Bassett DS, Lim KO, Carlson JM. Resolving anatomical and functional structure in human brain organization: identifying mesoscale organization in weighted network representations. *PLoS Comput Biol.* 2014; 10:e1003712. [PubMed: 25275860]
- Lowe MJ, Mock BJ, Sorenson JA. Functional connectivity in single and multislice echoplanar imaging using resting-state fluctuations. *NeuroImage.* 1998; 7:119–132. [PubMed: 9558644]
- Lyall AE, Shi F, Geng X, Woolson S, Li G, Wang L, Hamer RM, Shen D, Gilmore JH. Dynamic Development of Regional Cortical Thickness and Surface Area in Early Childhood. *Cereb Cortex.* 2015; 25:2204–2212. [PubMed: 24591525]



- Mantzaris AV, Bassett DS, Wymbs NF, Estrada E, Porter MA, Mucha PJ, Grafton ST, Higham DJ. Dynamic network centrality summarizes learning in the human brain. *Journal of Complex Networks*. 2013; 1:83–92.
- Marek S, Hwang K, Foran W, Hallquist MN, Luna B. The Contribution of Network Organization and Integration to the Development of Cognitive Control. *PLoS Biol*. 2015; 13:e1002328. [PubMed: 26713863]
- Markham JA, Greenough WT. Experience-driven brain plasticity: beyond the synapse. *Neuron Glia Biol*. 2004; 1:351–363. [PubMed: 16921405]
- Marquand AF, Wolfers T, Mennes M, Buitelaar J, Beckmann CF. Beyond Lumping and Splitting: A Review of Computational Approaches for Stratifying Psychiatric Disorders. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2016; 1:433–447. [PubMed: 27642641]
- Marusak HA, Calhoun VD, Brown S, Crespo LM, Sala-Hamrick K, Gotlib IH, Thomason ME. Dynamic functional connectivity of neurocognitive networks in children. *Hum Brain Mapp*. 2016
- McIntosh AR, Kovacevic N, Itier RJ. Increased brain signal variability accompanies lower behavioral variability in development. *PLoS Comput Biol*. 2008; 4:e1000106. [PubMed: 18604265]
- Meda SA, Ruano G, Windemuth A, O’Neil K, Berwise C, Dunn SM, Boccaccio LE, Narayanan B, Kocherla M, Sprooten E, et al. Multivariate analysis reveals genetic associations of the resting default mode network in psychotic bipolar disorder and schizophrenia. *Proc Natl Acad Sci U S A*. 2014; 111:E2066–2075. [PubMed: 24778245]
- Messe A, Rudrauf D, Benali H, Marrelec G. Relating structure and function in the human brain: relative contributions of anatomy, stationary dynamics, and non-stationarities. *PLoS Comput Biol*. 2014; 10:e1003530. [PubMed: 24651524]
- Miranda-Dominguez O, Mills BD, Carpenter SD, Grant KA, Kroenke CD, Nigg JT, Fair DA. Connectotyping: model based fingerprinting of the functional connectome. *PLoS One*. 2014a; 9:e111048. [PubMed: 25386919]
- Miranda-Dominguez O, Mills BD, Grayson D, Woodall a, Grant Ka, Kroenke CD, Fair Da. Bridging the Gap between the Human and Macaque Connectome: A Quantitative Comparison of Global Interspecies Structure-Function Relationships and Network Topology. *J Neurosci*. 2014b; 34:5552–5563. [PubMed: 24741045]
- Misc B, Betzel RF, Nematzadeh A, Goni J, Griffa A, Hagmann P, Flammini A, Ahn YY, Sporns O. Cooperative and Competitive Spreading Dynamics on the Human Connectome. *Neuron*. 2015; 86:1518–1529. [PubMed: 26087168]
- Miskovic V, Ma X, Chou CA, Fan M, Owens M, Sayama H, Gibb BE. Developmental changes in spontaneous electrocortical activity and network organization from early to late childhood. *NeuroImage*. 2015; 118:237–247. [PubMed: 26057595]
- Moseley RL, Ypma RJ, Holt RJ, Floris D, Chura LR, Spencer MD, Baron-Cohen S, Suckling J, Bullmore E, Rubinov M. Whole-brain functional hypoconnectivity as an endophenotype of autism in adolescents. *Neuroimage Clin*. 2015; 9:140–152. [PubMed: 26413477]
- O’Leary DD, Nakagawa Y. Patterning centers, regulatory genes and extrinsic mechanisms controlling arealization of the neocortex. *Curr Opin Neurobiol*. 2002; 12:14–25. [PubMed: 11861160]
- Patriat R, Reynolds RC, Birn RM. An improved model of motion-related signal changes in fMRI. *NeuroImage*. 2016
- Paxinos, G., Huang, XF., Toga, AW. The rhesus monkey brain in stereotaxic coordinates. San Diego, CA: Academic Press; 2000.
- Pessoa L. Understanding brain networks and brain organization. *Phys Life Rev*. 2014; 11:400–435. [PubMed: 24819881]
- Petersen SE, Sporns O. Brain Networks and Cognitive Architectures. *Neuron*. 2015; 88:207–219. [PubMed: 26447582]
- Power J, Schlaggar B, Lessov-Schlaggar C, Petersen S. Evidence for hubs in human functional brain networks. *Neuron*. 2013; 79:798–813. [PubMed: 23972601]
- Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*. 2012; 59:2142–2154. [PubMed: 22019881]

- Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, Vogel AC, Laumann TO, Miezin FM, Schlaggar BL, et al. Functional Network Organization of the Human Brain. *Neuron*. 2011; 72:665–678. [PubMed: 22099467]
- Power JD, Fair DA, Schlaggar BL, Petersen SE. The development of human functional brain networks. *Neuron*. 2010; 67:735–748. [PubMed: 20826306]
- Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*. 2014; 84:320–341. [PubMed: 23994314]
- Power JD, Plitt M, Laumann TO, Martin A. Sources and implications of whole-brain fMRI signals in humans. *NeuroImage*. 2016
- Power JD, Schlaggar BL, Petersen SE. Recent progress and outstanding issues in motion correction in resting state fMRI. *NeuroImage*. 2015; 105:536–551. [PubMed: 25462692]
- Qin J, Chen SG, Hu D, Zeng LL, Fan YM, Chen XP, Shen H. Predicting individual brain maturity using dynamic functional connectivity. *Front Hum Neurosci*. 2015; 9:418. [PubMed: 26236224]
- Reineberg AE, Andrews-Hanna JR, Depue BE, Friedman NP, Banich MT. Resting-state networks predict individual differences in common and specific aspects of executive function. *NeuroImage*. 2015; 104:69–78. [PubMed: 25281800]
- Reuter M, Tisdall MD, Qureshi A, Buckner RL, van der Kouwe AJ, Fischl B. Head motion during MRI acquisition reduces gray matter volume and thickness estimates. *NeuroImage*. 2015; 107:107–115. [PubMed: 25498430]
- Reveley C, Seth AK, Pierpaoli C, Silva AC, Yu D, Saunders RC, Leopold DA, Ye FQ. Superficial white matter fiber systems impede detection of long-range cortical connections in diffusion MR tractography. *Proc Natl Acad Sci U S A*. 2015; 112:E2820–2828. [PubMed: 25964365]
- Richiardi J, Altmann A, Milazzo AC, Chang C, Chakravarty MM, Banaschewski T, Barker GJ, Bokde AL, Bromberg U, Buchel C, et al. BRAIN NETWORKS. Correlated gene expression supports synchronous activity in brain networks. *Science*. 2015; 348:1241–1244. [PubMed: 26068849]
- Roalf DR, Quarmley M, Elliott MA, Satterthwaite TD, Vandekar SN, Ruparel K, Gennatas ED, Calkins ME, Moore TM, Hopson R, et al. The impact of quality assurance assessment on diffusion tensor imaging outcomes in a large-scale population-based cohort. *NeuroImage*. 2016; 125:903–919. [PubMed: 26520775]
- Robinson EC, Jbabdi S, Glasser MF, Andersson J, Burgess GC, Harms MP, Smith SM, Van Essen DC, Jenkinson M. MSM: a new flexible framework for Multimodal Surface Matching. *NeuroImage*. 2014; 100:414–426. [PubMed: 24939340]
- Rodriguez-Martinez EI, Barriga-Paulino CI, Rojas-Benjumea MA, Gomez CM. Co-maturation of theta and low-beta rhythms during child development. *Brain Topogr*. 2015; 28:250–260. [PubMed: 24793861]
- Rubinov M, Sporns O. Complex network measures of brain connectivity: Uses and interpretations. *NeuroImage*. 2010; 52:1059–1069. [PubMed: 19819337]
- Rudie JD, Hernandez LM, Brown JA, Beck-Pancer D, Colich NL, Gorrindo P, Thompson PM, Geschwind DH, Bookheimer SY, Levitt P, et al. Autism-associated promoter variant in MET impacts functional and structural brain networks. *Neuron*. 2012; 75:904–915. [PubMed: 22958829]
- Saleem, KS., Logothetis, N. A combined MRI and histology atlas of the rhesus monkey brain in stereotaxic coordinates. London; Burlington, MA: Academic; 2007.
- Sanz-Leon P, Knock SA, Spiegler A, Jirsa VK. Mathematical framework for large-scale brain network modeling in The Virtual Brain. *NeuroImage*. 2015; 111:385–430. [PubMed: 25592995]
- Sato JR, Salum GA, Gadelha A, Picon FA, Pan PM, Vieira G, Zugman A, Hoexter MQ, Anes M, Moura LM, et al. Age effects on the default mode and control networks in typically developing children. *J Psychiatr Res*. 2014; 58:89–95. [PubMed: 25085608]
- Sato JR, Salum GA, Gadelha A, Vieira G, Zugman A, Picon FA, Pan PM, Hoexter MQ, Anes M, Moura LM, et al. Decreased centrality of subcortical regions during the transition to adolescence: a functional connectivity study. *NeuroImage*. 2015; 104:44–51. [PubMed: 25290886]

- Satterthwaite TD, Baker JT. How can studies of resting-state functional connectivity help us understand psychosis as a disorder of brain development? *Curr Opin Neurobiol.* 2015; 30:85–91. [PubMed: 25464373]
- Satterthwaite TD, Vandekar SN, Wolf DH, Bassett DS, Ruparel K, Shehzad Z, Craddock RC, Shinohara RT, Moore TM, Gennatas ED, et al. Connectome-wide network analysis of youth with Psychosis-Spectrum symptoms. *Mol Psychiatry.* 2015; 20:1508–1515. [PubMed: 26033240]
- Satterthwaite TD, Wolf DH, Loughead J, Ruparel K, Elliott MA, Hakonarson H, Gur RC, Gur RE. Impact of in-scanner head motion on multiple measures of functional connectivity: relevance for studies of neurodevelopment in youth. *NeuroImage.* 2012; 60:623–632. [PubMed: 22233733]
- Savalia NK, Agres PF, Chan MY, Feczko EJ, Kennedy KM, Wig GS. Motion-related artifacts in structural brain images revealed with independent estimates of in-scanner head motion. *Hum Brain Mapp.* 2017; 38:472–492. [PubMed: 27634551]
- Scheinost D, Kwon SH, Shen X, Lacadie C, Schneider KC, Dai F, Ment LR, Constable RT. Preterm birth alters neonatal, functional rich club organization. *Brain Struct Funct.* 2016a; 221:3211–3222. [PubMed: 26341628]
- Scheinost D, Sinha R, Cross SN, Kwon SH, Sze G, Constable RT, Ment LR. Does prenatal stress alter the developing connectome? *Pediatr Res.* 2016b
- Scholtens LH, Schmidt R, de Reus MA, van den Heuvel MP. Linking macroscale graph analytical organization to microscale neuroarchitectonics in the macaque connectome. *J Neurosci.* 2014; 34:12192–12205. [PubMed: 25186762]
- Scholz J, Klein MC, Behrens TE, Johansen-Berg H. Training induces changes in white-matter architecture. *Nat Neurosci.* 2009; 12:1370–1371. [PubMed: 19820707]
- Scott JA, Grayson D, Fletcher E, Lee A, Bauman MD, Schumann CM, Buonocore MH, Amaral DG. Longitudinal analysis of the developing rhesus monkey brain using magnetic resonance imaging: birth to adulthood. *Brain Structure and Function.* 2015:1–25. [PubMed: 24248427]
- Shen K, Bezgin G, Hutchison RM, Gati JS, Menon RS, Everling S, McIntosh AR. Information processing architecture of functionally defined clusters in the macaque cortex. *The Journal of neuroscience: the official journal of the Society for Neuroscience.* 2012; 32:17465–17476. [PubMed: 23197737]
- Shen K, Hutchison RM, Bezgin G, Everling S, McIntosh AR. Network structure shapes spontaneous functional connectivity dynamics. *J Neurosci.* 2015; 35:5579–5588. [PubMed: 25855174]
- Sherman LE, Rudie JD, Pfeifer JH, Masten CL, McNealy K, Dapretto M. Development of the default mode and central executive networks across early adolescence: a longitudinal study. *Dev Cogn Neurosci.* 2014; 10:148–159. [PubMed: 25282602]
- Siegel JS, Mitra A, Laumann TO, Seitzman BA, Raichle M, Corbetta M, Snyder AZ. Data Quality Influences Observed Links Between Functional Connectivity and Behavior. *Cereb Cortex.* 2016
- Smit DJ, Boersma M, Schnack HG, Micheloyannis S, Boomsma DI, Hulshoff Pol HE, Stam CJ, de Geus EJ. The brain matures with stronger functional connectivity and decreased randomness of its network. *PLoS One.* 2012; 7:e36896. [PubMed: 22615837]
- Smith RE, Tournier JD, Calamante F, Connelly A. The effects of SIFT on the reproducibility and biological accuracy of the structural connectome. *NeuroImage.* 2015; 104:253–265. [PubMed: 25312774]
- Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE, Filippini N, Watkins KE, Toro R, Laird AR, et al. Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci U S A.* 2009; 106:13040–13045. [PubMed: 19620724]
- Smyser CD, Inder TE, Shimony JS, Hill JE, Degnan AJ, Snyder AZ, Neil JJ. Longitudinal analysis of neural network development in preterm infants. *Cereb Cortex.* 2010; 20:2852–2862. [PubMed: 20237243]
- Spoormaker VI, Schroter MS, Gleiser PM, Andrade KC, Dresler M, Wehrle R, Samann PG, Czisch M. Development of a large-scale functional brain network during human non-rapid eye movement sleep. *J Neurosci.* 2010; 30:11379–11387. [PubMed: 20739559]
- Sporns O. Network attributes for segregation and integration in the human brain. *Curr Opin Neurobiol.* 2013; 23:162–171. [PubMed: 23294553]

- Sporns O. Contributions and challenges for network models in cognitive neuroscience. *Nat Neurosci.* 2014
- Srinivasan R. Spatial structure of the human alpha rhythm: global correlation in adults and local correlation in children. *Clin Neurophysiol.* 1999; 110:1351–1362. [PubMed: 10454270]
- Stafford JM, Jarrett BR, Miranda-Dominguez O, Mills BD, Cain N, Mihalas S, Lahvis GP, Lattal KM, Mitchell SH, David SV, et al. Large-scale topology and the default mode network in the mouse connectome. *Proc Natl Acad Sci U S A.* 2014; 111:18745–18750. [PubMed: 25512496]
- Stanley ML, Simpson SL, Dagenbach D, Lyday RG, Burdette JH, Laurienti PJ. Changes in brain network efficiency and working memory performance in aging. *PLoS One.* 2015; 10:e0123950. [PubMed: 25875001]
- Stiles J, Jernigan TL. The basics of brain development. *Neuropsychol Rev.* 2010; 20:327–348. [PubMed: 21042938]
- Supekar K, Musen M, Menon V. Development of large-scale functional brain networks in children. *PLoS Biol.* 2009; 7:e1000157. [PubMed: 19621066]
- Supekar K, Uddin LQ, Khouzam A, Phillips J, Gaillard WD, Kenworthy LE, Yerys BE, Vaidya CJ, Menon V. Brain hyperconnectivity in children with autism and its links to social deficits. *Cell Rep.* 2013; 5:738–747. [PubMed: 24210821]
- Supekar K, Uddin LQ, Prater K, Amin H, Greicius MD, Menon V. Development of functional and structural connectivity within the default mode network in young children. *NeuroImage.* 2010; 52:290–301. [PubMed: 20385244]
- Sur M, Rubenstein JL. Patterning and plasticity of the cerebral cortex. *Science.* 2005; 310:805–810. [PubMed: 16272112]
- Swanson LW. Brain Maps Online: Toward Open Access Atlases and a Pan-mammalian Nomenclature. *J Comp Neurol.* 2015; 523:2272–2276. [PubMed: 25879783]
- Swartz JR, Monk CS. The role of corticolimbic circuitry in the development of anxiety disorders in children and adolescents. *Curr Top Behav Neurosci.* 2014; 16:133–148. [PubMed: 23950019]
- Tagliazucchi E, von Wegner F, Morzelewski A, Brodbeck V, Laufs H. Dynamic BOLD functional connectivity in humans and its electrophysiological correlates. *Front Hum Neurosci.* 2012; 6:339. [PubMed: 23293596]
- Thomas C, Ye FQ, Irfanoglu MO, Modi P, Saleem KS, Leopold DA, Pierpaoli C. Anatomical accuracy of brain connections derived from diffusion MRI tractography is inherently limited. *Proc Natl Acad Sci U S A.* 2014; 111:16574–16579. [PubMed: 25368179]
- Thomason ME, Dassanayake MT, Shen S, Katkuri Y, Alexis M, Anderson AL, Yeo L, Mody S, Hernandez-Andrade E, Hassan SS, et al. Cross-hemispheric functional connectivity in the human fetal brain. *Sci Transl Med.* 2013; 5:173ra124.
- Tononi G, Sporns O, Edelman GM. A measure for brain complexity: relating functional segregation and integration in the nervous system. *Proc Natl Acad Sci U S A.* 1994; 91:5033–5037. [PubMed: 8197179]
- Turk E, Scholtens LH, van den Heuvel MP. Cortical chemoarchitecture shapes macroscale effective functional connectivity patterns in macaque cerebral cortex. *Hum Brain Mapp.* 2016; 37:1856–1865. [PubMed: 26970255]
- Uddin LQ. Resting-state fMRI and developmental systems neuroscience. *Front Neurosci.* 2011; 5:14. [PubMed: 21390286]
- Uddin LQ, Supekar KS, Ryali S, Menon V. Dynamic reconfiguration of structural and functional connectivity across core neurocognitive brain networks with development. *J Neurosci.* 2011; 31:18578–18589. [PubMed: 22171056]
- Vakorin VA, Lippe S, McIntosh AR. Variability of brain signals processed locally transforms into higher connectivity with brain development. *J Neurosci.* 2011; 31:6405–6413. [PubMed: 21525281]
- van den Heuvel MP, de Reus MA, Feldman Barrett L, Scholtens LH, Coopmans FM, Schmidt R, Preuss TM, Rilling JK, Li L. Comparison of diffusion tractography and tract-tracing measures of connectivity strength in rhesus macaque connectome. *Hum Brain Mapp.* 2015a; 36:3064–3075. [PubMed: 26058702]

- van den Heuvel MP, Kersbergen KJ, de Reus MA, Keunen K, Kahn RS, Groenendaal F, de Vries LS, Benders MJ. The Neonatal Connectome During Preterm Brain Development. *Cereb Cortex*. 2015b; 25:3000–3013. [PubMed: 24833018]
- van den Heuvel MP, Scholtens LH, de Reus MA, Kahn RS. Associated Microscale Spine Density and Macroscale Connectivity Disruptions in Schizophrenia. *Biol Psychiatry*. 2016a; 80:293–301. [PubMed: 26632269]
- van den Heuvel MP, Scholtens LH, Turk E, Mantini D, Vanduffel W, Feldman Barrett L. Multimodal analysis of cortical chemoarchitecture and macroscale fMRI resting-state functional connectivity. *Hum Brain Mapp*. 2016b; 37:3103–3113. [PubMed: 27207489]
- van den Heuvel MP, Sporns O. An anatomical substrate for integration among functional networks in human cortex. *The Journal of neuroscience*. 2013; 33:14489–14500. [PubMed: 24005300]
- van den Heuvel MP, Stam CJ, Kahn RS, Hulshoff Pol HE. Efficiency of functional brain networks and intellectual performance. *J Neurosci*. 2009; 29:7619–7624. [PubMed: 19515930]
- Van Dijk KR, Sabuncu MR, Buckner RL. The influence of head motion on intrinsic functional connectivity MRI. *NeuroImage*. 2012; 59:431–438. [PubMed: 21810475]
- van Duijvenvoorde AC, Huizenga HM, Somerville LH, Delgado MR, Powers A, Weeda WD, Casey BJ, Weber EU, Figner B. Neural correlates of expected risks and returns in risky choice across development. *J Neurosci*. 2015; 35:1549–1560. [PubMed: 25632132]
- Van Essen DC. Cortical cartography and Caret software. *NeuroImage*. 2012; 62:757–764. [PubMed: 22062192]
- Van Essen DC, Glasser MF, Dierker DL, Harwell J. Cortical parcellations of the macaque monkey analyzed on surface-based atlases. *Cereb Cortex*. 2012; 22:2227–2240. [PubMed: 22052704]
- Vertes PE, Bullmore ET. Annual research review: Growth connectomics--the organization and reorganization of brain networks during normal and abnormal development. *J Child Psychol Psychiatry*. 2015; 56:299–320. [PubMed: 25441756]
- Wang JH, Zuo XN, Gohel S, Milham MP, Biswal BB, He Y. Graph theoretical analysis of functional brain networks: test-retest evaluation on short- and long-term resting-state functional MRI data. *PLoS One*. 2011; 6:e21976. [PubMed: 21818285]
- Wang L, Su L, Shen H, Hu D. Decoding lifespan changes of the human brain using resting-state functional connectivity MRI. *PLoS One*. 2012; 7:e44530. [PubMed: 22952990]
- Watts DJ, Strogatz SH. Collective dynamics of ‘small-world’ networks. *Nature*. 1998; 393:440–442. [PubMed: 9623998]
- Whitford TJ, Rennie CJ, Grieve SM, Clark CR, Gordon E, Williams LM. Brain maturation in adolescence: concurrent changes in neuroanatomy and neurophysiology. *Hum Brain Mapp*. 2007; 28:228–237. [PubMed: 16767769]
- Wig GS, Laumann TO, Cohen AL, Power JD, Nelson SM, Glasser MF, Miezin FM, Snyder AZ, Schlaggar BL, Petersen SE. Parcellating an individual subject’s cortical and subcortical brain structures using snowball sampling of resting-state correlations. *Cereb Cortex*. 2014a; 24:2036–2054. [PubMed: 23476025]
- Wig GS, Laumann TO, Petersen SE. An approach for parcellating human cortical areas using resting-state correlations. *NeuroImage*. 2014b; 93(Pt 2):276–291. [PubMed: 23876247]
- Wisner KM, Atluri G, Lim KO, Macdonald AW 3rd. Neurometrics of intrinsic connectivity networks at rest using fMRI: retest reliability and cross-validation using a meta-level method. *NeuroImage*. 2013; 76:236–251. [PubMed: 23507379]
- Wu K, Taki Y, Sato K, Hashizume H, Sassa Y, Takeuchi H, Thyreau B, He Y, Evans AC, Li X, et al. Topological organization of functional brain networks in healthy children: differences in relation to age, sex, and intelligence. *PLoS One*. 2013; 8:e55347. [PubMed: 23390528]
- Wylie KP, Rojas DC, Ross RG, Hunter SK, Maharajh K, Cornier MA, Tregellas JR. Reduced brain resting-state network specificity in infants compared with adults. *Neuropsychiatr Dis Treat*. 2014; 10:1349–1359. [PubMed: 25092980]
- Yan CG, Cheung B, Kelly C, Colcombe S, Craddock RC, Di Martino A, Li Q, Zuo XN, Castellanos FX, Milham MP. A comprehensive assessment of regional variation in the impact of head micromovements on functional connectomics. *NeuroImage*. 2013; 76:183–201. [PubMed: 23499792]

- Yendiki A, Koldewyn K, Kakunoori S, Kanwisher N, Fischl B. Spurious group differences due to head motion in a diffusion MRI study. *NeuroImage*. 2014; 88:79–90. [PubMed: 24269273]
- Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, Roffman JL, Smoller JW, Zöllei L, Polimeni JR, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*. 2011; 106:1125–1165. [PubMed: 21653723]
- Zalesky A, Fornito A, Harding IH, Cocchi L, Yucel M, Pantelis C, Bullmore ET. Whole-brain anatomical networks: does the choice of nodes matter? *NeuroImage*. 2010; 50:970–983. [PubMed: 20035887]
- Zhang D, Snyder AZ, Shimony JS, Fox MD, Raichle ME. Noninvasive functional and structural connectivity mapping of the human thalamocortical system. *Cerebral cortex (New York, NY: 1991)*. 2010; 20:1187–1194.
- Zhang Z, Liao W, Chen H, Mantini D, Ding JR, Xu Q, Wang Z, Yuan C, Chen G, Jiao Q, et al. Altered functional-structural coupling of large-scale brain networks in idiopathic generalized epilepsy. *Brain*. 2011; 134:2912–2928. [PubMed: 21975588]
- Zuo XN, Ehmke R, Mennes M, Imperati D, Castellanos FX, Sporns O, Milham MP. Network centrality in the human functional connectome. *Cereb Cortex*. 2012; 22:1862–1875. [PubMed: 21968567]
- Zuo XN, Kelly C, Adelstein JS, Klein DF, Castellanos FX, Milham MP. Reliable intrinsic connectivity networks: test-retest evaluation using ICA and dual regression approach. *NeuroImage*. 2010; 49:2163–2177. [PubMed: 19896537]
- Zuo XN, Xing XX. Test-retest reliabilities of resting-state FMRI measurements in human brain functional connectomics: a systems neuroscience perspective. *Neurosci Biobehav Rev*. 2014; 45:100–118. [PubMed: 24875392]



### Highlights


Reviews development of functional connectivity networks from birth until adulthood.

Reviews trends in resting-state functional MR imaging (rs-fMRI) and network analysis.

Synthesizes developmental rs-fMRI findings with structural connectivity and EEG/MEG.

Suggests strategies to overcome limitations of rs-fMRI in developmental studies.


Suggests approaches to interrogate neurodevelopmental disorders.



**Figure 1. Crucial network properties of resting-state activity in the normal adult brain**


**A)** Group-averaged community structure of resting state brain activity, densely sampled across the cortical surface, in three independent studies (Gordon et al., 2016; Power et al., 2011; Yeo et al., 2011). Different colors correspond to different modules with highly correlated activity, i.e. functional connectivity (FC). Several canonical modules appear in all three studies, including early visual, early somatomotor, default mode, dorsal attention, frontoparietal, and cingulo-opercular modules. Community structure is highly reproducible across these studies. The color scheme in Power and Gordon matches the legend, but differs in Yeo. A novel areal parcellation is also defined in Gordon, as evidenced by interareal boundaries. **B)** Community structure and areal parcellation of an individual subject (Laumann et al., 2015), obtained via repeated scanning sessions in the same individual. Community structure strongly resembles that of the group, although idiosyncracies are also clearly observable (see (Laumann et al., 2015)). **C)** Group-average resting-state network organization defined using 264 spherical nodes situated within different functional modules. On the left, the location and modular assignment of these nodes are pictured on the brain, next to a spring-embedded layout of the thresholded functional matrix. The network layout depicts nodes with stronger links (i.e. stronger correlations in activity) as closer together. The layout illustrates that network organization is heterogeneous – some modules are highly segregated from the rest of the network, whereas others are more integrated. On the right, the participation coefficient is quantified for each node. The participation coefficient

signifies the extent to which a node integrates activity across multiple modules. Comparing the network layouts, integrator nodes tend to exist within the modules reflecting task-positive cognition, i.e. the frontoparietal, cingulo-opercular, and dorsal attention modules. See (Power et al., 2013). **D**) Structural connectivity shapes and constrains FC. As an example, this plot illustrates the correspondence between empirical FC (y-axis) and predicted FC (x-axis) based on modeling communication via the structural connectome. Dots represent region pairs that either are (blue) or are not (red) directly connected via fiber pathways. See (Goni et al., 2014). **E**) Structural hubs tend to interlink to each other and across functional modules, providing an anatomical substrate for integration between otherwise segregated domains of information processing. The schematic on the left illustrates this hypothesis, where the modules on top are functionally defined and the connections shown on the bottom reflect neuroanatomical links. The “rich club” nodes (shown in blue) reflect structural hubs that disproportionately connect to each other. On the right, evidence that the rich club serves an integrative function, as functional network nodes with high participation coefficient (i.e. high between-module integration) overlap significantly with the brain’s structural rich club. See (van den Heuvel and Sporns, 2013).




**Figure 2. Infant brain structural and functional development**

**A)** Visualizations of infant brain cortical surfaces at birth, 1 year, and 2 years of age. Large growth is clearly visible in terms of total brain size, cortical surface area, and cortical thickness Adapted from (Li et al., 2015). **B)** Resting-state functional connectivity network visualizations at birth, 1 year, and 2 years of age. The different lobes containing each region are labeled. Increasing separation of regions into functional modules spanning multiple lobes is apparent over this timespan. Adapted from (Gao et al., 2011). **C)** Community structure of 230 functional ROIs in 1–2 year olds (top) and in adults (bottom) using closely matched methods. Labeling of infant modules was informed via the adult set: Vis (visual), tDMN (temporal default mode network), pcDMN (posterior cingulate DMN), aDMN (anterior DMN), SMN (somato-motor network), SMN2 (somato-motor network 2), DAN (dorsal attention network), pFPC (posterior frontal parietal control network), aFPC (anterior frontal parietal control network), SubCtx (subcortex), CO (cingulo-opercular), pCO (posterior CO), and Sal (salience). Adapted from (Eggebrecht et al., 2017).



**Figure 3. Development of functional architecture from childhood to adulthood: similarities in community structure across age**


**A)** Illustration shows community structure of functional networks in childhood (top) and adulthood (bottom), with (right) or without (left) denoising of scans via removal of high head-motion frames (i.e. “scrubbing”). Nodes are 264 spherical ROIs, colored according to community assignments. Circles illustrate areas that show apparent age-related differences prior to motion denoising, but which do not demonstrate age effects after denoising. Numbers next to arrows indicate the mutual information (a graph theoretic measure of similarity) in the two community structures. Motion exaggerates age-related differences in community structure. From (Power et al., 2012). **B)** Similar community structure is obtained in adults and children when using strict criteria to minimize the influence of motion artifact. Overall network structure looks remarkably similar as well. From (Fair et al., 2012b). **C)** Similar community structures are identified in late childhood, early adolescence, late adolescence, and adulthood. On the other hand, there is evidence for refinement within this modular framework (see Figure 4). From (Marek et al., 2015).



**Figure 4. Evidence for disproportionate involvement of the cingulo-opercular and somatomotor systems during development from late childhood into adulthood**

**A)** Regions that are most predictive of age, from late childhood to young adulthood, via age-related changes in functional connectivity. Nodes are sized according to predictive strength in a support-vector machine. Adapted from (Fair et al., 2012b). **B)** On the left, average functional community structure is shown for healthy young adults. On the right, differences in functional connectivity between adults and older children are shown. Adults have greater functional connectivity of selected links within and between the somatomotor and cingulo-opercular modules. Regions on the cortical surface with high functional connectivity overall are highlighted in warm colors, illustrating that developmental differences also involve hub regions. Adapted from (Grayson et al., 2014). **C)** Developmental trajectories are illustrated for five networks. Average participation coefficient for nodes within each network are plotted over age. The cingulo-opercular network exhibited the most substantial increase over development. These changes mediated age-related increases in cognitive control and were especially driven by increased connectivity between cingulo-opercular and somatomotor nodes. From (Marek et al., 2015).





**Figure 5. Functional network organization, hierarchies, and structure-function relationships in the monkey brain gleaned through contrast-enhanced resting-state imaging**

**A)** Community detection performed on an 80-region parcellation of the rhesus monkey brain. Resting-state networks were obtained under anesthesia using enhanced imaging methodologies that included exogenous contrast, a surface coil with high SNR, and head fixation. Reported modules show clear homology with those seen in the human literature. **B)** Spring-embedded graph layout visualizing correlations between individual regions. Integrated versus segregated activity is visible, as are hierarchies within different sensory modalities. For instance, dorsal attention nodes are situated centrally, suggesting globally integrated processing. In contradistinction, primary visual, auditory, and somatosensory cortex are the most peripheral nodes within their respective modules, followed by secondary sensory cortices, suggesting both segregation of sensory streams and hierarchical relationships within them. Nodes are sized by their correlation with the global signal, illustrating that more central nodes have higher global signal correlation. **C)** Plot illustrates the correspondence between empirical FC (y-axis) and predicted FC (x-axis) based on modeling communication in the structural connectome. Dots represent region pairs that either are (blue) or are not (red) directly connected via fiber pathways. **D)** Structure-function relationships are also observable at the node level. Regional correlation with the global signal (y-axis) is associated with total communication capacity to and from the rest of the brain (x-axis). Dots show global signal correlations before (closed) and after (open) regressing out the global signal from each region's timecourse. Adapted from (Grayson et al., 2016).