



Cite this article: Delás MJ, Hannon GJ. 2017 lncRNAs in development and disease: from functions to mechanisms. *Open Biol.* **7**: 170121.
<http://dx.doi.org/10.1098/rsob.170121>

Received: 16 May 2017
Accepted: 3 July 2017

Subject Area:
molecular biology

Keywords:
long non-coding RNAs, development, cancer

Author for correspondence:
Gregory J. Hannon
e-mail: greg.hannon@cr.uk.cam.ac.uk

lncRNAs in development and disease: from functions to mechanisms

M. Joaquina Delás^{1,2} and Gregory J. Hannon^{1,2,3}

¹Watson School of Biological Sciences, Cold Spring Harbor Laboratory, New York, NY 11724, USA

²Cancer Research UK Cambridge Institute, Li Ka Shing Centre, University of Cambridge, Cambridge CB2 0RE, UK

³New York Genome Center, 101 6th Ave, New York, NY 10013, USA

MJD, 0000-0001-9727-9068; GJH, 0000-0003-4021-3898

Differential expression of long non-coding RNAs (lncRNAs) during differentiation and their misregulation in cancer highlight their potential as cell fate regulators. While some example lncRNAs have been characterized in great detail, the functional *in vivo* relevance of others has been called into question. Finding functional lncRNAs will most probably require a combination of complementary approaches that will greatly vary depending on their mode of action. In this review, we discuss the different tools available to dissect genetically lncRNA requirements and how each is best suited to studies in particular contexts. Moreover, we review different strategies used to select candidate lncRNAs and give an overview of lncRNAs described to regulate development and cancer through different mechanisms.

1. Introduction

Regulated gene expression is the basis for the extensive variety of cell types our bodies generate from the same set of DNA instructions. Specific gene programmes are transcribed in particular cells providing them with their molecular identity and the protein products that underlie their functions. Together with coding genes, thousands of long non-coding RNAs (lncRNAs) are also expressed in a cell-type-specific manner during differentiation and in certain cancers. This has been extensively reported in many organisms and cell types [1–4], yet demonstrating that these molecules play functional roles has not been easy.

The now ever-expanding catalogue of lncRNAs first became apparent from efforts to annotate the functional features of the human genome, which showed that the vast majority of the genome was transcribed [5]. Currently, lncRNAs are defined as capped transcripts longer than 200 nucleotides, which coincides with the cut-off for many RNA extraction protocols [6]. They can be spliced and, in most of the published studies, are also polyadenylated. lncRNAs have little or no coding potential, although some do bind to ribosomes [7–9]. They were originally described to have equivalent chromatin features to protein-coding genes [10]. However, more recent work has highlighted differences in the abundance of particular histone marks [11,12] and splicing efficiency [12,13] between lncRNAs and coding genes, as well as subsets of lncRNAs that differ in their chromatin signatures [4,14].

Hundreds of thousands of lncRNAs have been annotated in different species and tissues [15]. Of those, only a handful have been shown to be critical for organism development [16–18] or cancer progression [19], and the mechanisms by which they act have been established in just a few cases. For some, biochemical partners have been carefully identified, yet *in vivo* evidence for their function is missing or questions have been raised regarding the relevance of the previously reported mechanisms of action [20,21]. Bridging this gap is essential for building a solid body of knowledge of how lncRNAs function in cell fate choices and the mechanisms by which they act. In this review, we will focus on the different strategies to select and identify functional lncRNAs and some mechanistic examples of lncRNA acting in differentiation and cancer.

2. Cell-type-specific expression of long non-coding RNAs: cause or consequence?

The cell-type-specific expression observed for lncRNAs has provoked much excitement, as it implies that they might function during cell fate decisions. In accord with this notion, deregulation of lncRNAs has also been widely observed across human cancers [22–25]. Such disease-associated expression changes suggested a potential role for these lncRNAs in driving cancer or at least contributing to maintaining an aberrant transcriptional landscape.

2.1. Long non-coding RNA expression during development

Differential expression of lncRNAs has been reported between regions of the mammalian brain [26,27], and lncRNA dynamics have been analysed in more detail during corticogenesis [28]. Several studies have shown differential expression of lncRNAs in *in vitro* differentiation models of haematopoiesis [29,30] and in freshly isolated cell populations [31,32], as well as during mammalian adipogenesis [33,34]. This tissue and cell-type-specific regulation is observed across species, including during development of zebrafish [2], *Caenorhabditis elegans* [3], and even during the life cell cycle of our close unicellular relative, *Capsaspora ovczarzaki* [4].

If lncRNAs are to regulate key developmental genes, a very appealing possibility is that they do so in *cis*. Correlated expression of lncRNAs and their neighbouring genes has been reported in embryonic stem (ES) cell differentiation to endoderm [35] and to embryoid bodies [36], as well as in human B and T cell lineages [37]. However, neighbour correlation is not a special property of lncRNA–gene pairs, as the expression levels of neighbouring genes are often correlated. This is thought to be due to shared regulatory elements affecting each neighbour [38] and to the general neighbourhood or chromosomal domain around them [39,40].

This leaves us with several possibilities to consider, and a myriad of experimental challenges to distinguish between them. The genomic location of lncRNAs and their neighbouring regulatory elements could determine their cell-type-specific expression, with the RNA being a mere by-product of the regulatory mechanisms already in place [41]. On the other hand, lncRNAs could be critical for expression of those developmental genes, orchestrating chromatin changes by specific RNA–protein interactions, or by increasing the local concentration of transcriptional machinery regardless of the actual RNA sequence transcribed. A general mechanism by which lncRNA transcription in specific cell types could reorganize nuclear architecture, and thus contribute to the new transcription landscape, has even been proposed [15].

2.2. Long non-coding RNA misregulation in cancer

Along the same lines, cancer-specific lncRNA expression could simply be a by-product of aberrant gene expression in cancer. However, genetic mutations can directly affect lncRNA expression, with the lncRNAs themselves playing a causal role in specific scenarios. lncRNA *CCAT2*, for example, encompasses a cancer-associated SNP. The risk allele correlates

with a higher expression of the lncRNA, which in turn promotes proliferation in colorectal cancer [42]. This lncRNA is part of the 8q24.21 region, where many cancer-associated mutations and amplifications have been reported. Several disease-associated SNPs and translocations including the lncRNA *PVT1* have drawn researchers' attention to this very complex locus [43]. One of these amplifications is that of *PVT1* and its neighbouring gene *c-MYC*. In breast cancer models *PVT1* RNA levels correlate with MYC protein, yet *PVT1* and *c-MYC* are not always co-amplified. This suggests that amplification of this lncRNA alone (even without MYC) can promote tumorigenesis in breast cancer by increasing MYC expression [44], which has been proposed to occur by protein stabilization. However, it is unlikely that this is the only mechanism of action for this lncRNA as this multi-exonic transcript encoding over 20 different isoforms is itself under the control of *c-MYC* and harbours multiple microRNAs within its locus [45].

While these are a few well-characterized examples, a much greater number of functional studies will be required to tease apart collaterally expressed lncRNAs from those with important roles both in development and in cancer.

3. Finding functional long non-coding RNAs

Many experimental strategies have been used to dissect genetically lncRNA requirements in differentiation and cancer. Powerful in their own ways, each of these techniques has its own drawbacks. Therefore, a combination of complementary approaches will probably be required to reveal the biological impact of lncRNAs. The choice of approach also strongly depends on the biological question, whether it is the identification of lncRNAs important for a differentiation or disease process, the study of specific types of regulatory mechanisms—*cis* versus *trans*, or an in-depth analysis of a particular lncRNA.

3.1. Different tools for different questions

The main consideration is that, in a particular lncRNA locus, the act of transcription itself could be key to establishing or maintaining the chromatin state of the surrounding area, while, in this scenario, the actual sequence of the RNA would be irrelevant. Or the RNA itself could be the functional unit, having some sequence-dependent interactions with proteins, RNAs or DNA elements. It could even be that both these mechanisms apply for the same locus. Therefore, it is very important to understand each experimental set-up and what it tells us about each particular lncRNA.

Several studies have taken advantage of RNA interference (RNAi) approaches, either transduced shRNAs or transfected siRNAs [46,47]. This strategy has been coupled with a phenotypic readout, such as viability or differentiation, to identify lncRNAs where the RNA molecule itself is important (figure 1). However, many worry about potential off-target effects (though this is no different from shRNA studies with protein-coding genes). There are additional concerns regarding the difficulty of knocking down lncRNAs that are chromatin-associated versus cytoplasmic, given that small RNA loading into the RISC complex takes place in the cytoplasm. While there is some evidence for differences in knockdown efficiency




Figure 1. Different approaches for disrupting lncRNAs. Methods such as knockdown and CRISPRi affect the RNA itself or reduce the transcription of the lncRNA. Knockdown can be achieved in a variety of ways (siRNA, shRNA, LNA, ASO). CRISPRi is most efficient if Cas9 is fused to repressor domains (e.g. KRAB). These methods can also be transient. Insertion of an early terminator sequence or complete deletion of the locus or promoter are achieved via genome engineering and are non-reversible.

depending on subcellular location [48], this concern would apply only to lncRNAs that are never exported to the cytoplasm. lncRNAs that function in the nucleus but in *trans* could very well be exported just like other RNAs and then re-imported. Undoubtedly, the main advantage of knockdown is that it allows for high-throughput screens that could yield a list, though potentially incomplete, of lncRNAs with functions in the phenotypic assay of our choice.

To circumvent possible subcellular localization biases, other researchers have taken advantage of alternative knockdown techniques, independent of the RNAi machinery, that do not require processing in any specific cell compartment. Morpholinos targeting splice junctions or conserved regions have been used in zebrafish for identification of functional lncRNAs *in vivo* [16]. Locked nucleic acids (LNAs) have also been used in mammalian cells [49,50]. Both of these approaches rely on annealing a synthetic nucleic acid to the lncRNA and blocking its function or its splicing. Antisense oligos (ASOs) are another alternative that takes advantage of RNase-H activity. ASOs have been used in a variety of systems, both delivered to cultured cells and administered *in vivo* in mice [19,51]. Several ASOs have now been approved for clinical use and, although their targets so far have been coding genes, this opens up the path towards therapeutic targeting of lncRNAs. These approaches, although incredibly useful because they exclusively target the RNA, can only be deployed where these molecules can be injected or otherwise delivered. This would not allow for pooled high-throughput screening and misses out on the advantages of genetically encoding knockdown, which can be conditionally induced for *in vivo* studies.

The ultimate proof of functionality is a genetic knockout. They allow for the study of *in vivo* function and reduce the possibility of off-target effects. Discrepancies between the hypothesized mechanisms for some lncRNAs based on the *in vitro* data and the absence of or very mild phenotypes observed in knockout animals [20,52–54] have resulted in scepticism regarding broad regulatory roles for lncRNAs [55].

Two recent examples of lncRNA knockouts emphasize how, in some cases, phenotypes might be more context-specific than anticipated. *Malat1* is a very abundant lncRNA that localizes to nuclear speckles. Although it was hypothesized that this RNA was required for speckle or paraspeckle formation

and maintenance and regulated alternative splicing through interaction with SR proteins [56], three independent mouse knockout models showed that *Malat1* was dispensable for viability [52–54]. Furthermore, *Malat1* was shown not to be required for nuclear speckle formation and its deletion did not affect SR protein phosphorylation [54]. However, when crossed with the MMTV-PyMT mouse model of human breast cancer, *Malat1* deletion impaired tumour progression as evidenced by a severe reduction in metastatic burden [19].

LincRNA-EP5 was shown to have an anti-apoptotic role and be required for red blood cell development in tissue culture models of erythroid development [57]. The knockout mouse model for this lncRNA showed no defects in blood development. However, *LincRNA-EP5* controls expression of immune response genes in macrophages and proved essential for the animals to respond to endotoxin challenge [21].

Although their molecular mechanisms still remain to be elucidated, these RNAs are representative examples of how, just as for coding genes, some lncRNAs could play roles under particular stress or disease conditions. This could potentially be the case for other lncRNAs, whose proposed roles have met with controversy, such as *HOTAIR*, where knockout animals seem to be viable and healthy [20,58].

When removing a DNA locus to generate a knockout—especially when dealing with large deletion—any phenotype observed could be due either to loss of an encoded RNA or to deletion of DNA sequences that might include regulatory elements. For this reason, full transcript knockouts can be combined with complementary strategies to dissect roles of RNA from those of DNA elements. The possibilities of genome engineering are not limited to full locus knockout, but instead allow more subtle modifications, such as polyadenylation (poly-A) signal insertions for premature termination (figure 1). Taking advantage of CRISPR/Cas9, a recent study looked at whether a set of 12 lncRNAs regulated the expression of their neighbouring genes in *cis*. Although expression defects were observed upon promoter knockout for five lncRNAs, only one had the same effect when a poly-A signal was inserted, suggesting that only the regulatory elements in the DNA surrounding the promoter and not the RNAs were required for these *cis* effects [59]. An equivalent mechanism has been proposed to explain the differences between knockout

and poly-A insertions for the lncRNA *Lockd* and its neighbour gene *Cdkn1b* [60].

Two independent models for *Fendrr* knockout showed that it was required for mouse development [17,18]. Interestingly, the phenotypes differed, one being embryonic lethal with a presumed requirement for lateral plate mesoderm [17], while the other was perinatal lethal [18]. These differences could be the consequence of the distinct genetic strategies, one being a triple polyadenylation insertion and the other one a whole gene replacement, emphasizing the need for complementary approaches that distinguish between DNA and RNA elements.

The scenario is substantially more complicated when the lncRNA and its target gene have overlapping transcripts. *Airn* overlaps in antisense with the imprinted gene *Igfr2*. Through a series of polyadenylation cassette insertions, it was shown that transcriptional termination of *Airn* only leads to *Igfr2* de-repression when the non-coding transcript no longer overlapped the *Igfr2* promoter. This work concluded that transcription of *Airn*, rather than the final transcript, is responsible for promoter silencing [61]. Combinations of promoter, exon knockouts and termination signals created using CRISPR/Cas9 have helped dissect the relationship between *Haunt* (also known as *linc1547* or *linc-Hoxa5*) and the *HoxA* locus. Knockdown, termination or deletions of the first exons lead to increased expression of *HoxA* genes during retinoid acid-induced differentiation of ES cells, supporting a repressive role for *Haunt* at the *HoxA* genes. However, deletion of the whole *Haunt* locus prevents expression of *HoxA*, presumably due to deletion of some regulatory DNA elements required for *HoxA* induction [62].

While more accessible owing to the advancements in genome engineering, mouse model generation is still not amenable to high-throughput studies, and therefore requires careful selection of lncRNA candidates. Genome-scale strategies for lncRNA CRISPR/Cas9 deletion are being developed [63], and variations in the ever-expanding CRISPR/Cas9 toolkit could help identify functional lncRNAs. Cas9 fused to repressors [64] or activators [65] now allows for the manipulation of expression levels at the loci themselves (figure 1). Using the former approach, researchers have identified human lncRNAs essential for cell growth in a diverse set of cell lines [64]. This technology is scalable and could help identify lncRNAs required in a variety of contexts by performing loss-of-function studies. The main limitation is that altering the chromatin state of the lncRNA promoter could directly affect nearby genes, complicating the interpretation of the phenotype.

3.2. Which long non-coding RNAs to study

The primary approach will depend on the biological question being asked. Some studies directly focus on a particular lncRNA of interest, while others aim at the unbiased identification of lncRNAs important for a process. Different strategies have been employed to select subsets of lncRNAs for study based on their expression level, dynamic regulation, tissue expression and even conservation.

Historically, highly abundant lncRNAs were chosen as representatives of this RNA class, with the hope of identifying possible mechanisms by which lncRNAs might act generally. This was, in part, due to experimental convenience and technical limitations, and this class includes the best characterized lncRNA to date, *Xist*, as well as *Malat1*, and *Neat1*. *Xist* orchestrates X chromosome inactivation. Expressed

from the silenced allele, this lncRNA acts in *cis* to inactivate the expression of that X chromosome copy [66]. Focusing on a particular lncRNA has allowed researchers to channel all their efforts towards a mechanistic understanding of its mode of action, while also developing *in vivo* tools for its study. While *in vivo* functional validation is everyone's dream, placing all eggs in one basket is always risky and can lead to disappointing outcomes [20,52,53].

The opposite approach to studying a single lncRNA is genome-wide unbiased screening of lncRNAs. This can be an excellent filter to identify potential functional lncRNAs, building a resource for further mechanistic studies, although this approach is best suited to easily measured phenotypes, such as proliferation. Genome-wide screens have been used to identify lncRNAs essential for human cancer cell growth or survival [64], or those required in mouse ES cell self-renewal [47]. In ES cells, this approach identified *TUNA*, a lncRNA required for neural specification from ES cells that had already been described to have neural phenotype in zebrafish (named *Megamind*) [16], validating this strategy.

lncRNA annotation can be intersected with expression levels if one wishes to reduce the number of targeted lncRNAs. This can be necessary for more elaborate phenotypic assays that are not as scalable. A straightforward approach is to assess only the lncRNAs expressed in the cell type or tissue by setting some minimal expression cut-off [46]. However, for some strategies further reduction of lncRNA candidates is necessary. By analysing expression in different related tissues, especially in developmental systems, several groups have focused only on differentially regulated lncRNAs. The rationale behind this is that a transcript dynamically induced or silenced during a cell fate transition, for example, is more likely to be important for that process. Differential expression helped in the selection of candidates in epidermal differentiation [67,68], cardiac differentiation [69] and haematopoiesis [32].

Another extremely useful layer of filtering is evolutionary conservation. Although their sequence is not broadly conserved, lncRNAs can often be found in syntenic positions in different species. These 'syntelogs' can even share some small conserved domains [70]. This approach has been used to identify several conserved lncRNAs that act during zebrafish development [16] and drew researchers' attention to *NORAD*, a conserved lncRNA in mammals that modulates Pumilio proteins [71,72]. Combined with differential expression, conservation can help focus the candidate list on lncRNAs with key functions in developmental or disease processes. As 'syntenic conservation' is a rather loose criterion and the presence of a lncRNA does not necessarily indicate that it will have the same function in a different organism, complementary strategies will be very helpful in identifying orthologous lncRNAs. If they are functionally conserved, lncRNA 'syntelogs' might share structure similarities, even if they do not share much sequence identity. Some studies have approached this by looking at predicted secondary RNA structure [73]. Although RNA structure predictions for long RNAs might not be particularly useful, new experimental approaches to globally identify structure features in lncRNAs could aid in this task [74].

When dealing with lncRNA annotation, it is important to be aware of the limitations. Although most assemblies set up stringent coding potential cut-offs, lncRNAs often contain very short open reading frames (ORFs). The functionality of these micropeptides is hard to assess unless one addresses it experimentally. Three different short proteins have been




Figure 2. Themes in lncRNA functions. lncRNAs have been described to play multiple roles affecting gene expression at the transcriptional level via interactions with chromatin remodelling complexes, direct binding to the DNA as an RNA–DNA triplex, or facilitating chromatin looping (a), interacting with RNA processing machinery or affecting mRNA stability (b) or directly regulating protein function (c).

found to play a role in muscle function or regeneration [75–77], which emphasizes the importance of testing for RNA-mediated rather than protein-mediated effects.

Overall, the ability to modulate the expression of lncRNAs or disrupt it altogether now allows an assessment of lncRNA requirements in many developmental and cancer contexts. This, combined with some clever candidate selection strategies, has identified a number of lncRNAs important in these processes. The level of current mechanistic understanding for each of these lncRNAs is variable, yet the techniques available and being developed hint at a promising future.

4. Long non-coding RNAs shape development and cancer

Even for well-studied lncRNAs, our mechanistic understanding has deepened only in the last few years. The poster child for lncRNA researchers, *Xist*, orchestrates X chromosome inactivation. The functional properties of *Xist* and the order of events it directs have been known for decades (reviewed in [66]). However, it has taken until very recently to better understand the X inactivation at the molecular level. We now know the protein partners *Xist* requires for X chromosome silencing [78,79], how *Xist* spreads [80] and takes advantage of the chromosome's three-dimensional structure to initiate silencing [81], and how that chromosomal conformation changes during transcriptional silencing [82,83]. Additionally, this RNA is modified with N6-methyladenosines, which contribute to its transcriptional repressive activity [84]. *Xist* illustrates not only the detailed mechanistic understanding to which we can aspire for other lncRNAs of interest but also the tremendous amount of effort required to understand even a single lncRNA. Of note, *Xist* is also highly abundant when it is expressed, and that induction takes place in a cell type we

can culture in large amounts (ES cells). Greater challenges can be expected with lncRNAs expressed to lower levels in very specific cell types.

Some hints at how other lncRNAs exert their functions in development and cancer have been reported. Although the field is still maturing, lncRNAs have been described to play a myriad of roles, from regulating gene expression to regulating mRNA processing or affecting protein stability (figure 2). There are also several examples of lncRNA loci where the act of transcription but not the RNA itself seems to be of functional relevance [59,85] or where transcription is even dispensable [59]. Here, we focus on the RNAs themselves as the functional units.

4.1. Effects on chromatin and DNA interactions

Following *Xist*'s example, many researchers have focused on potential chromatin regulatory roles of lncRNAs. *Mistral* (*Mira*) is a lncRNA expressed in ES cells that is reported to interact with MLL1 to recruit this protein to *Hoxa6* and *Hoxa7*, leading to their activation. Consequently, siRNA-mediated knockdown of *Mistral* leads to reduced transcription of these genes and negation of the overall germ-cell specification programme [86]. Also expressed in ES cells, *HoxBlinc* binds to the same complex to promote *HoxB* transcription and mesoderm specification [87].

Two lncRNAs have been shown to be required for heart development. *Braveheart*, required for the production of contracting embryoid bodies from ES cells, interacts with the Polycomb factor SUZ12 [69], while *Fendrr*, a lncRNA essential for mouse development [17,18], binds to SUZ12 as well as EZH2 and WDR5 [17]. These interactions with members of the TrxG/MLL and Polycomb complexes place these lncRNAs in a position to direct chromatin modifications to particular DNA loci in a sequence-specific manner (figure 2a).

Following this hypothesized mechanism, it was shown that *Fendrr* interacts, at least *in vitro*, with the promoter regions of *Pitx2* and *Foxf1*, both of which are expressed during heart development [17]. In HeLa cells, *Khps1* regulates the promoter of its antisense gene, the proto-oncogene *SPHK1*, by forming a DNA–RNA triplex at its promoter while recruiting p300/CBP [88] (figure 2a). *TARID* also regulates its antisense gene, *TFC21*, by guiding GADD45A to the locus and promoting demethylation, which leads to gene activation. *TARID* and *TCF21* are silent and heavily covered by DNA methylation in non-small cell lung cancer (NSCLC), head and neck squamous cell carcinomas (HNSCC) and ovarian cancers (OVC) [49]. And although direct RNA–DNA binding has not been shown, lncRNA *SLNCR1* is required for bringing androgen receptor to the *MMP9* promoter, which increases *MMP9* expression and leads to melanoma invasion [89].

The ultraconserved lncRNA *Megamind/TUNA* showed brain development phenotypes upon knockdown in zebrafish [16] and is required for ES cell pluripotency and neuronal differentiation from mouse ES cells [47]. In the mouse, this lncRNA binds three RNA-binding proteins and interacts with the *Sox2* promoter [47]. *Sox2* is a key transcription factor in neuronal differentiation, so its regulation could explain the resulting phenotype. And while the interaction seems to be indirect, *Dali* has been shown to localize globally to active promoters in the N2A neuronal differentiation model [90], as shown by CHART-seq [91].

Interaction with transcription factors themselves is another plausible mechanism to promote expression of specific gene programmes. *RMST*, for example, is a lncRNA up-regulated during neurogenesis that interacts with *Sox2*. Knockdown of *RMST* leads to a reduction in *Sox2* ChIP-seq peaks in this model, which suggests that this lncRNA is somehow facilitating the binding of this transcription factor [92].

Additionally, *LUNAR* and *DEANR1* both seem to function by facilitating DNA looping between the lncRNA locus and their target gene to promote activation. *LUNAR* is a Notch-regulated lncRNA that activates the *IGFR1* gene in T-cell acute lymphoblastic leukaemia [51], while *DEANR1* functions via a similar mechanism in endoderm development, activating *FOXA2* expression through the recruitment of SMAD2/3 [93] (figure 2a).

4.2. Effects on mRNA stability and processing

Conceptually, the next level of regulation would be for lncRNAs to negatively or positively affect the stability or processing of coding mRNAs. By having cell-type or tumour-specific expression, lncRNAs would effectively control the output levels for these genes. In accord with this model, a few exemplar lncRNAs have been shown to interact with heterogeneous nuclear ribonucleoproteins (hnRNPs) (figure 2b). *LincRNA-p21*, induced in response to DNA damage, interacts with hnRNP-K, and it is regulated by p53. Knockdown of this lncRNA leads to up-regulation of genes normally repressed by p53 and also reduces apoptosis similarly to p53 knockdown [94]. This suggests a model whereby this lncRNA acts as a repressor of p53-dependent genes.

In neural differentiation, *Pnky* knockdown leads to progenitor expansion, and mass spectroscopy of lncRNA-interacting proteins followed by immunoblotting validation revealed PTPB1 as one of its interaction partners. Knockdown of this lncRNA leads to misexpression and altered splicing of many

key genes [95]. Both PTPB1 and hnRNP-K also bind *TUNA* during *in vitro* neuronal differentiation [47]. Being highly expressed and broadly acting proteins, it is only reasonable to wonder whether these functions are truly specific. Only more detailed biochemical studies will be able to clarify this.

In a complementary approach to mass spectroscopy, protein microarrays identified STAU1 as the interacting partner of lncRNA *TINCR* [68]. Combined knockdown of *TINCR* and *STAU1* seems to affect the stability of important epidermal differentiation genes such as *Krt80* [68], which would explain its essential role in skin differentiation. During muscle differentiation, *LncMyoD* binds IGF2 mRNA-binding protein 2 (IMP2), which leads to enhanced translation of mRNAs involved in proliferation. Interestingly, this is a conserved lncRNA and the mouse and human sequence can rescue each other's knockdown [96] (figure 2b).

Rather than binding elements of the RNA processing machinery as a way to regulate the fate of coding mRNAs, other lncRNAs have been shown to act as endogenous competitors for microRNAs, thus dampening the silencing of microRNA targets. *LincND*, for example, is a primate-conserved lncRNA expressed in neural progenitors and down-regulated in neurons. This lncRNA competes for miR-143-3p, which would normally target Notch. Relieving Notch silencing promotes neuronal differentiation [97].

In the cancer context, *LINC00152* acts as an oncogenic lncRNA, competing with *HIF1- α* for miR-138. Expression of this lncRNA promotes invasion in gall bladder cancer [98]. As an additional example, *linc-223* would usually bind to miR-125-5p but it is down-regulated in acute myeloid leukaemia, leading to increased repression of *IRF4*, a target of miR-125-5p [99] (figure 2b).

The range of action of *lncARSR* extends even further because, apart from competing with mir-34 and miR-449 thus promoting stability of AXL and c-MET, this lncRNA can be packaged into exosomes to be secreted. Down-regulation of microRNA target genes renders renal cancer cells resistant to sunitinib, and secretion of the lncRNA can disseminate this property to neighbouring cells [50].

4.3. Effects on protein stability and function

Rather than affecting the mRNAs of genes important for differentiation or malignant proliferation, lncRNAs can also directly bind proteins essential for a signalling pathway and modulate their function. *Linc-DC*, for example, is induced during dendritic differentiation from human monocytes. This lncRNA interacts with STAT3 and, when knocked down, leads to a reduction in Y705 phosphorylation of STAT3, decreasing its nuclear translocation. For that reason, expression of *linc-DC* is required for differentiation of dendritic cells [100] (figure 2c).

Similar protein–lncRNA relationships have been observed in different cancer models. *FAL1* and *PVT1* are amplified in ovarian and breast cancer, respectively. *FAL1* associates with Bmi1, and *FAL1* knockdown leads to a reduction in Bmi1 levels and misregulation of large numbers of genes involved in cell cycle progression [101]. *PVT1* has a similar relationship with the oncogene C-MYC, promoting the stability of this protein [44] (figure 2c).

lncRNA-LET and *LINK-A* have opposing effects on *HIF1- α* in hepatocellular carcinoma. Enforced expression of *lncRNA-LET* leads to reduced *HIF1- α* and results in lower metastatic potential [102], while *LINK-A* interacts with tyrosine protein

kinase 6 to promote stabilization of *HIF1- α* in triple negative breast cancer [103].

Other lncRNAs have specific roles in particular pathways, such as *MAYA*, a lncRNA recruited by HER3-ROR1 that then binds directly to NSUN6, preventing methylation of MST1. In breast cancer, MST1 is inactivated by this methylation resulting in YAP signalling target activation and increased bone metastasis [104] (figure 2c). *SAMMSON* is involved in regulating mitochondrial integrity by associating with p32. Knockdown of this lncRNA results in aberrant mitochondrial structures in melanoma, a cancer where *SAMMSON* is amplified [105].

By playing roles in essential cell functions, other lncRNAs also affect cancer progression. *NORAD* is a conserved lncRNA with repetitive regions that binds to PUMILIO proteins [71]. When this lncRNA is absent, PUMILIO proteins carry out their roles as negative regulators of mRNA stability and translation, and this results in aneuploidy [72]. *LINP1* interacts with Ku80 and DNA-PKcs, coordinating the non-homologous end-joining (NHEJ) pathway. Apart from providing essential functions for any cell, this pathway is particularly required in triple negative breast cancer [106].

5. Concluding remarks

lncRNAs are being heavily studied in the context of development and cancer, as their unique properties could allow them to interact with multiple proteins via three-dimensional structures and also recognize other nucleic acids by base pairing. Their specific expression during differentiation and disease places them in an ideal position to play key regulatory roles.

Because of the added complexity in studying lncRNA loci, a combination of genetic approaches is often required to

distinguish between the function of the RNA molecule and the regulatory activity from a DNA element in that locus. Many examples have been described for lncRNAs interacting with chromatin, regulating genes at the RNA or protein level, or interfering globally with splicing. These functions are diverse and expand the original hypothesis of a nuclear-specific function for most lncRNAs. It was also proposed that lncRNAs would mostly act in *cis*, as their expression mirrored that of their neighbour genes [36,37,107,108]. Although that is the case for some examples, it does not seem to be a general rule [59].

lncRNAs are diverse molecules that are not likely to fit in one functional class. Consequently, we should start thinking of them more like proteins, some functioning in the nucleus [109], others acting in the cytoplasm and others supporting the structure of cells [110,111]. What is becoming clearer with the development of *in vivo* models and our expanding mechanistic understanding is that there are lncRNAs with essential functions in development and others required for cancer progression, taking this class of RNAs out of the ‘junk DNA’ category once and for all. Not every annotated lncRNA will have an RNA-mediated function—or a function at all—but identifying the biologically relevant ones and understanding their mechanisms will certainly be a hotbed for future study.

Data accessibility. This article has no additional data.

Competing interests. The authors have no relevant competing interests.

Funding. M.J.D. was funded by a PhD fellowship from Boehringer Ingelheim Fonds, a graduate studies fellowship from ‘la Caixa’ Foundation and by the Watson School of Biological Sciences. G.J.H. is a Wellcome Trust Investigator, was an investigator of the Howard Hughes Medical Institute, and is supported by Cancer Research UK, The Royal Society (Wolfson Professorship) and a generous gift from Kathryn W. Davis.

Acknowledgements. We thank the members of the Hannon laboratory for helpful discussion and their critical reading of the manuscript.

References

- Ponting CP, Oliver PL, Reik W. 2009 Evolution and functions of long noncoding RNAs. *Cell* **136**, 629–641. (doi:10.1016/j.cell.2009.02.006)
- Pauli A *et al.* 2012 Systematic identification of long noncoding RNAs expressed during zebrafish embryogenesis. *Genome Res.* **22**, 577–591. (doi:10.1101/gr.133009.111)
- Nam J-W, Bartel DP. 2012 Long noncoding RNAs in *C. elegans*. *Genome Res.* **22**, 2529–2540. (doi:10.1101/gr.140475.112)
- Sebé-Pedrós A, Ballaré C, Parra-Acero H, Chiva C, Tena JJ, Sabidó E, Gómez-Skarmeta JL, Di Croce L, Ruiz-Trillo I. 2016 The dynamic regulatory genome of *Capsaspora* and the origin of animal multicellularity. *Cell* **165**, 1224–1237. (doi:10.1016/j.cell.2016.03.034)
- ENCODE Project Consortium *et al.* 2007 Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* **447**, 799–816. (doi:10.1038/nature05874)
- Rinn JL, Chang HY. 2012 Genome regulation by long noncoding RNAs. *Annu. Rev. Biochem.* **81**, 145–166. (doi:10.1146/annurev-biochem-051410-092902)
- Ingolia NT, Lareau LF, Weissman JS. 2011 Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell* **147**, 789–802. (doi:10.1016/j.cell.2011.10.002)
- Guttman M, Russell P, Ingolia NT, Weissman JS, Lander ES. 2013 Ribosome profiling provides evidence that large noncoding RNAs do not encode proteins. *Cell* **154**, 240–251. (doi:10.1016/j.cell.2013.06.009)
- Ruiz-Orera J, Messegue X, Subirana JA, Alba MM, Tautz D. 2014 Long non-coding RNAs as a source of new peptides. *Elife* **3**, e03523. (doi:10.7554/elife.03523)
- Guttman M *et al.* 2009 Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* **458**, 223–227. (doi:10.1038/nature07672)
- Alam T, Medvedeva YA, Jia H, Brown JB, Lipovich L, Bajic VB. 2014 Promoter analysis reveals globally differential regulation of human long non-coding RNA and protein-coding genes. *PLoS ONE* **9**, e109443. (doi:10.1371/journal.pone.0109443)
- Melé, M., Mattioli K, Mallard W, Shechner DM, Gerhardinger C, Rinn JL. 2017 Chromatin environment, transcriptional regulation, and splicing distinguish lincRNAs and mRNAs. *Genome Res.* **27**, 27–37. (doi:10.1101/gr.214205.116)
- Schlackow M, Nojima T, Gomes T, Dhir A, Carmo-Fonseca M, Proudfoot NJ. 2017 Distinctive patterns of transcription and RNA processing for human lincRNAs. *Mol. Cell* **65**, 25–38. (doi:10.1016/j.molcel.2016.11.029)
- Marques AC, Hughes J, Graham B, Kowalczyk MS, Higgs DR, Ponting CP. 2013 Chromatin signatures at transcriptional start sites separate two equally populated yet distinct classes of intergenic long noncoding RNAs. *Genome Biol.* **14**, R131. (doi:10.1186/gb-2013-14-11-r131)
- Melé, M, Rinn JL. 2016 ‘Cat’s cradling’ the 3D genome by the act of lncRNA transcription. *Mol. Cell* **62**, 657–664. (doi:10.1016/j.molcel.2016.05.011)
- Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP. 2011 Conserved function of lincRNAs in vertebrate embryonic development despite rapid sequence evolution. *Cell* **147**, 1537–1550. (doi:10.1016/j.cell.2011.11.055)
- Grote P *et al.* 2013 The tissue-specific lncRNA Fendrr is an essential regulator of heart and body wall development in the mouse. *Cell* **24**, 206–214. (doi:10.1016/j.devcel.2012.12.012)
- Sauvageau M *et al.* 2013 Multiple knockout mouse models reveal lincRNAs are required for life and

- brain development. *Elife* **2**, e01749. (doi:10.7554/eLife.01749)
19. Arun G *et al.* 2016 Differentiation of mammary tumors and reduction in metastasis upon Malat1 lncRNA loss. *Genes Dev.* **30**, 34–51. (doi:10.1101/gad.270959.115)
 20. Amândio AR, Necsulea A, Joye E, Mascrez B, Duboule D. 2016 Hotair is dispensable for mouse development. *PLoS Genet.* **12**, e1006232. (doi:10.1371/journal.pgen.1006232)
 21. Atianand MK *et al.* 2016 A long noncoding RNA lincRNA-EP5 acts as a transcriptional brake to restrain inflammation. *Cell* **165**, 1672–1685. (doi:10.1016/j.cell.2016.05.075)
 22. Yan X *et al.* 2015 Comprehensive genomic characterization of long non-coding RNAs across human cancers. *Cancer Cell* **28**, 529–540. (doi:10.1016/j.ccell.2015.09.006)
 23. Iyer MK *et al.* 2015 The landscape of long noncoding RNAs in the human transcriptome. *Nat. Genet.* **47**, 199–208. (doi:10.1038/ng.3192)
 24. Du Z, Fei T, Verhaak RG. W., Su Z, Zhang Y, Brown M, Chen Y, Liu XS. 2013 Integrative genomic analyses reveal clinically relevant long noncoding RNAs in human cancer. *Nat. Struct. Mol. Biol.* **20**, 908–913. (doi:10.1038/nsmb.2591)
 25. Brunner AL, Beck AH, Edris B. 2012 Transcriptional profiling of long non-coding RNAs and novel transcribed regions across a diverse panel of archived human cancers. *Genome Biol.* **13**, R75. (doi:10.1186/gb-2012-13-8-r75)
 26. Mercer TR, Dinger ME, Sunken SM, Mehler MF, Mattick JS. 2008 Specific expression of long noncoding RNAs in the mouse brain. *Proc. Natl Acad. Sci. USA* **105**, 716–721. (doi:10.1073/pnas.0706729105)
 27. Goff LA *et al.* 2015 Spatiotemporal expression and transcriptional perturbations by long noncoding RNAs in the mouse brain. *Proc. Natl Acad. Sci. USA* **112**, 6855–6862. (doi:10.1073/pnas.1411263112)
 28. Molyneux BJ, Goff LA, Brettler AC, Chen H.-H., Brown JR, Hrvatin S, Rinn JL, Arlotta P. 2015 DeCoN: genome-wide analysis of in vivo transcriptional dynamics during pyramidal neuron fate selection in neocortex. *Neuron* **85**, 275–288. (doi:10.1016/j.neuron.2014.12.024)
 29. Alvarez-Dominguez JR, Hu W, Yuan B, Shi J, Park SS, Gromatzky AA, van Oudenaarden A, Lodish HF. 2014 Global discovery of erythroid long noncoding RNAs reveals novel regulators of red cell maturation. *Blood* **123**, 570–581. (doi:10.1182/blood-2013-10-530683)
 30. Paralkar VR *et al.* 2014 Lineage and species-specific long noncoding RNAs during erythro-megakaryocytic development. *Blood* **123**, 1927–1937. (doi:10.1182/blood-2013-12-544494)
 31. Cabezas-Wallscheid N *et al.* 2014 Identification of regulatory networks in HSCs and their immediate progeny via integrated proteome, transcriptome, and DNA methylome analysis. *Cell Stem Cell* **15**, 507–522. (doi:10.1016/j.stem.2014.07.005)
 32. Luo M *et al.* 2015 Long non-coding RNAs control hematopoietic stem cell function. *Cell Stem Cell* **16**, 426–438. (doi:10.1016/j.stem.2015.02.002)
 33. Alvarez-Dominguez JR *et al.* 2015 *De novo* reconstruction of adipose tissue transcriptomes reveals long non-coding RNA regulators of brown adipocyte development. *Cell Metab.* **21**, 764–776. (doi:10.1016/j.cmet.2015.04.003)
 34. Sun L *et al.* 2013 Long noncoding RNAs regulate adipogenesis. *Proc. Natl Acad. Sci. USA* **110**, 3387–3392. (doi:10.1073/pnas.1222643110)
 35. Sigova AA *et al.* 2013 Divergent transcription of long noncoding RNA/mRNA gene pairs in embryonic stem cells. *Proc. Natl Acad. Sci. USA* **110**, 2876–2881. (doi:10.1073/pnas.1221904110)
 36. Dinger ME *et al.* 2008 Long noncoding RNAs in mouse embryonic stem cell pluripotency and differentiation. *Genome Res.* **18**, 1433–1445. (doi:10.1101/gr.078378.108)
 37. Casero D, Sandoval S, Seet CS, Scholes J, Zhu Y, Ha VL, Luong A, Parekh C, Crooks GM. 2015 Long non-coding RNA profiling of human lymphoid progenitor cells reveals transcriptional divergence of B cell and T cell lineages. *Nat. Immunol.* **16**, 1282–1291. (doi:10.1038/ni.3299)
 38. Cohen BA, Mitra RD, Hughes JD, Church GM. 2000 A computational analysis of whole-genome expression data reveals chromosomal domains of gene expression. *Nat. Genet.* **26**, 183–186. (doi:10.1038/79896)
 39. Downen JM *et al.* 2014 Control of cell identity genes occurs in insulated neighborhoods in mammalian chromosomes. *Cell* **159**, 374–387. (doi:10.1016/j.cell.2014.09.030)
 40. Gonzalez-Sandoval A, Gasser SM. 2016 On TADs and LADs: spatial control over gene expression. *Trends Genet.* **32**, 485–495. (doi:10.1016/j.tig.2016.05.004)
 41. van Bakel H, Nislow C, Blencowe BJ, Hughes TR. 2010 Most ‘dark matter’ transcripts are associated with known genes. *PLoS Biol.* **8**, e1000371. (doi:10.1371/journal.pbio.1000371)
 42. Ling H *et al.* 2013 CCAT2, a novel noncoding RNA mapping to 8q24, underlies metastatic progression and chromosomal instability in colon cancer. *Genome Res.* **23**, 1446–1461. (doi:10.1101/gr.152942.112)
 43. Huppi K, Pitt JJ, Wahlberg BM, Caplen NJ. 2012 The 8q24 gene desert: an oasis of non-coding transcriptional activity. *Front Genet.* **3**, 69. (doi:10.3389/fgene.2012.00069)
 44. Tseng Y-Y *et al.* 2014 PVT1 dependence in cancer with MYC copy-number increase. *Nature* **512**, 82–86. (doi:10.1038/nature13311)
 45. Carramusa L, Contino F, Ferro A, Minafra L, Perconti G, Giallongo A, Feo S. 2007 The PVT-1 oncogene is a Myc protein target that is overexpressed in transformed cells. *J. Cell. Physiol.* **213**, 511–518. (doi:10.1002/jcp.21133)
 46. Guttman M *et al.* 2011 lincRNAs act in the circuitry controlling pluripotency and differentiation. *Nature* **477**, 295–300. (doi:10.1038/nature10398)
 47. Lin N *et al.* 2014 An evolutionarily conserved long noncoding RNA TUNA controls pluripotency and neural lineage commitment. *Mol. Cell* **53**, 1005–1019. (doi:10.1016/j.molcel.2014.01.021)
 48. Lennox KA, Behlke MA. 2016 Cellular localization of long non-coding RNAs affects silencing by RNAi more than by antisense oligonucleotides. *Nucleic Acids Res.* **44**, 863–877. (doi:10.1093/nar/gkv1206)
 49. Arab K *et al.* 2014 Long noncoding RNA *TARID* directs demethylation and activation of the tumor suppressor TCF21 via GADD45A. *Mol. Cell* **55**, 604–614. (doi:10.1016/j.molcel.2014.06.031)
 50. Qu L *et al.* 2016 Exosome-transmitted lncARSR promotes sunitinib resistance in renal cancer by acting as a competing endogenous RNA. *Cancer Cell* **29**, 653–668. (doi:10.1016/j.ccell.2016.03.004)
 51. Trimarchi T, Bilal E, Ntziachristos P, Fabbri G, Dalla-Favera R, Tsigiris A, Aifantis I. 2014 Genome-wide mapping and characterization of Notch-regulated long noncoding RNAs in acute leukemia. *Cell* **158**, 593–606. (doi:10.1016/j.cell.2014.05.049)
 52. Nakagawa S, Ip JY, Shioi G, Tripathi V, Zong X, Hirose T, Prasanth KV. 2012 Malat1 is not an essential component of nuclear speckles in mice. *RNA* **18**, 1487–1499. (doi:10.1261/rna.033217.112)
 53. Eißmann M *et al.* 2012 Loss of the abundant nuclear non-coding RNAMALAT1 is compatible with life and development. *RNA Biol.* **9**, 1076–1087. (doi:10.4161/rna.21089)
 54. Zhang B *et al.* 2012 The lncRNA Malat1 is dispensable for mouse development but its transcription plays a cis-regulatory role in the adult. *Cell Rep.* **2**, 111–123. (doi:10.1016/j.celrep.2012.06.003)
 55. Selleri L, Bartolomei MS, Bickmore WA, He L, Stubbs L, Reik W, Barsh GS. 2016 A hox-embedded long noncoding RNA: is it all hot air? *PLoS Genet.* **12**, e1006485. (doi:10.1371/journal.pgen.1006485)
 56. Tripathi V *et al.* 2010 The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol. Cell* **39**, 925–938. (doi:10.1016/j.molcel.2010.08.011)
 57. Hu W, Yuan B, Flygare J, Lodish HF. 2011 Long noncoding RNA-mediated anti-apoptotic activity in murine erythroid terminal differentiation. *Genes Dev.* **25**, 2573–2578. (doi:10.1101/gad.178780.111)
 58. Li L *et al.* 2013 Targeted disruption of hotair leads to homeotic transformation and gene derepression. *Cell Rep.* **5**, 3–12. (doi:10.1016/j.celrep.2013.09.003)
 59. Engreitz JM, Haines JE, Perez EM, Munson G, Chen J, Kane M, McDonel PE, Guttman M, Lander ES. 2016 Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature* **539**, 452–455. (doi:10.1038/nature20149)
 60. Paralkar VR *et al.* 2016 Unlinking an lncRNA from its associated cis element. *Mol. Cell* **62**, 104–110. (doi:10.1016/j.molcel.2016.02.029)
 61. Latos PA *et al.* 2012 Airn transcriptional overlap, but not its lncRNA products, induces imprinted Igf2r silencing. *Science* **338**, 1469–1472. (doi:10.1126/science.1228110)

62. Yin Y *et al.* 2015 Opposing roles for the lncRNA *haunt* and its genomic locus in regulating HOXA gene activation during embryonic stem cell differentiation. *Cell Stem Cell* **16**, 504–516. (doi:10.1016/j.stem.2015.03.007)
63. Zhu S *et al.* 2016 Genome-scale deletion screening of human long non-coding RNAs using a paired-guide RNA CRISPR-Cas9 library. *Nat. Biotechnol.* **34**, 1279–1286. (doi:10.1038/nbt.3715)
64. Liu SJ *et al.* 2017 CRISPRi-based genome-scale identification of functional long noncoding RNA loci in human cells. *Science* **355**, pii. aah7111. (doi:10.1126/science.aah7111)
65. Konermann S *et al.* 2014 Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* **517**, 583–588. (doi:10.1038/nature14136)
66. Augui S, Nora EP, Heard E. 2011 Regulation of X-chromosome inactivation by the X-inactivation centre. *Nat. Rev. Genet.* **12**, 429–442. (doi:10.1038/nrg2987)
67. Kretz M *et al.* 2012 Suppression of progenitor differentiation requires the long noncoding RNA ANCR. *Genes Dev.* **26**, 338–343. (doi:10.1101/gad.182121.111)
68. Kretz M *et al.* 2013 Control of somatic tissue differentiation by the long non-coding RNA TINCR. *Nature* **493**, 231–235. (doi:10.1038/nature11661)
69. Klattenhoff CA *et al.* 2013 Braveheart, a long noncoding RNA required for cardiovascular lineage commitment. *Cell* **152**, 570–583. (doi:10.1016/j.cell.2013.01.003)
70. Ulitsky I. 2016 Evolution to the rescue: using comparative genomics to understand long non-coding RNAs. *Nat. Rev. Genet.* **17**, 601–614. (doi:10.1038/nrg.2016.85)
71. Tichon A, Gil N, Lubelsky Y, Solomon TH, Lemze D, Itzkovitz S, Stern-Ginossar N, Ulitsky I. 2016 A conserved abundant cytoplasmic long noncoding RNA modulates repression by Pumilio proteins in human cells. *Nat. Commun.* **7**, 1–10. (doi:10.1038/ncomms12209)
72. Lee S, Kopp F, Chang T-C, Sataluri A, Chen B, Sivakumar S, Yu H, Xie Y, Mendell JT. 2016 Noncoding RNA NORAD regulates genomic stability by sequestering PUMILIO proteins. *Cell* **164**, 69–80. (doi:10.1016/j.cell.2015.12.017)
73. Quinn JJ, Zhang QC, Georgiev P, Ilik IA, Akhtar A, Chang HY. 2016 Rapid evolutionary turnover underlies conserved lncRNA–genome interactions. *Genes Dev.* **30**, 191–207. (doi:10.1101/gad.272187.115)
74. Lu Z *et al.* 2016 RNA duplex map in living cells reveals higher-order transcriptome structure. *Cell* **165**, 1267–1279. (doi:10.1016/j.cell.2016.04.028)
75. Matsumoto A *et al.* 2016 mTORC1 and muscle regeneration are regulated by the LINC00961-encoded SPAR polypeptide. *Nature* **541**, 228–232. (doi:10.1038/nature21034)
76. Anderson DM *et al.* 2015 A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell* **160**, 595–606. (doi:10.1016/j.cell.2015.01.009)
77. Nelson BR *et al.* 2016 A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in muscle. *Science* **351**, 271–275. (doi:10.1126/science.aad4076)
78. Chu C, Zhang QC, da Rocha ST, Flynn RA, Bharadwaj M, Calabrese JM, Magnuson T, Heard E, Chang HY. 2015 Systematic discovery of Xist RNA binding proteins. *Cell* **161**, 404–416. (doi:10.1016/j.cell.2015.03.025)
79. McHugh CA *et al.* 2015 The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature* **521**, 232–236. (doi:10.1038/nature14443)
80. Simon MD *et al.* 2013 High-resolution Xist binding maps reveal two-step spreading during X-chromosome inactivation. *Nature* **504**, 465–469. (doi:10.1038/nature12719)
81. Engreitz JM *et al.* 2013 The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science* **341**, 1237973. (doi:10.1126/science.1237973)
82. Giorgetti L *et al.* 2016 Structural organization of the inactive X chromosome in the mouse. *Nature* **535**, 575–579. (doi:10.1038/nature18589)
83. Darrow EM *et al.* 2016 Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. *Proc. Natl Acad. Sci. USA* **113**, E4504–E4512. (doi:10.1073/pnas.1609643113)
84. Patil DP, Chen C-K, Pickering BF, Chow A, Jackson C, Guttman M, Jaffrey SR. 2016 m(6)A RNA methylation promotes XIST-mediated transcriptional repression. *Nature* **537**, 369–373. (doi:10.1038/nature19342)
85. Anderson KM, Anderson DM, McAnally JR, Shelton JM, Bassel-Duby R, Olson EN. 2016 Transcription of the non-coding RNA upperhand controls Hand2 expression and heart development. *Nature* **539**, 433–436. (doi:10.1038/nature20128)
86. Bertani S, Sauer S, Bolotin E, Sauer F. 2011 The noncoding RNA *mistral* activates *Hoxa6* and *Hoxa7* expression and stem cell differentiation by recruiting MLL1 to chromatin. *Mol. Cell* **43**, 1040–1046. (doi:10.1016/j.molcel.2011.08.019)
87. Deng C *et al.* 2016 HoxBln1 RNA recruits Set1/MLL complexes to activate Hox gene expression patterns and mesoderm lineage development. *Cell Rep.* **14**, 103–114. (doi:10.1016/j.celrep.2015.12.007)
88. Postepska-Igielska A, Giwojna A, Gasri-Plotnitsky L, Schmitt N, Dold A, Ginsberg D, Grummt I. 2015 lncRNA *Khps1* regulates expression of the proto-oncogene *SPHK1* via triplex-mediated changes in chromatin structure. *Mol. Cell* **60**, 626–636. (doi:10.1016/j.molcel.2015.10.001)
89. Schmidt K, Joyce CE, Buquicchio F, Brown A, Ritz J, Distel RJ, Yoon CH, Novina CD. 2016 The lncRNA *SLNCR1* mediates melanoma invasion through a conserved SRA1-like region. *Cell Rep.* **15**, 2025–2037. (doi:10.1016/j.celrep.2016.04.018)
90. Chalei V, Sansom SN, Kong L, Lee S, Montiel JF, Vance KW, Ponting CP. 2014 The long non-coding RNA *Dali* is an epigenetic regulator of neural differentiation. *Elife* **3**, e04530. (doi:10.7554/eLife.04530)
91. Simon MD, Wang CI, Kharchenko PV, West JA, Chapman BA, Alekseyenko AA, Borowsky ML, Kuroda MI, Kingston RE. 2011 The genomic binding sites of a noncoding RNA. *Proc. Natl Acad. Sci. USA* **108**, 20 497–20 502. (doi:10.1073/pnas.1113536108)
92. Ng S-Y, Bogu GK, Soh B.-S, Stanton LW. 2013 The long noncoding RNA *RMST* interacts with *SOX2* to regulate neurogenesis. *Mol. Cell* **51**, 349–359. (doi:10.1016/j.molcel.2013.07.017)
93. Jiang W, Liu Y, Liu R, Zhang K, Zhang Y. 2015 The lncRNA *DEANR1* facilitates human endoderm differentiation by activating *FOXA2* expression. *Cell Rep.* **11**, 137–148. (doi:10.1016/j.celrep.2015.03.008)
94. Huarte M *et al.* 2010 A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell* **142**, 409–419. (doi:10.1016/j.cell.2010.06.040)
95. Ramos AD *et al.* 2015 The long noncoding RNA *Pnky* regulates neuronal differentiation of embryonic and postnatal neural stem cells. *Cell Stem Cell* **16**, 439–447. (doi:10.1016/j.stem.2015.02.007)
96. Gong C, Li Z, Ramanujan K, Clay I, Zhang Y, Lemire-Brachat S, Glass DJ. 2015 A long non-coding RNA, *LncMyoD*, regulates skeletal muscle differentiation by blocking IMP2-mediated mRNA translation. *Dev. Cell* **34**, 181–191. (doi:10.1016/j.devcel.2015.05.009)
97. Rani N, Nowakowski TJ, Zhou H, Godshalk SE, Lisi V, Kriegstein AR, Kosik KS. 2016 A primate lncRNA mediates Notch signaling during neuronal development by sequestering miRNA. *Neuron* **90**, 1174–1188. (doi:10.1016/j.neuron.2016.05.005)
98. Cai Q, Wang Z, Wang S, Weng M, Zhou D, Li C, Wang J, Chen E, Quan Z. 2017 Long non-coding RNA *LINC00152* promotes gallbladder cancer metastasis and epithelial–mesenchymal transition by regulating HIF-1 α via miR-138. *Open Biol.* **7**, 160247. (doi:10.1002/hep.24563)
99. Mangiavacchi A, Sorci M, Masciarelli S, Larivera S, Legnini I, Iosue I, Bozzoni I, Fazi F, Fatica A. 2016 The miR-223 host non-coding transcript *linc-223* induces IRF4 expression in acute myeloid leukemia by acting as a competing endogenous RNA. *Oncotarget* **7**, 60 155–60 168. (doi:10.18632/oncotarget.11165)
100. Wang P *et al.* 2014 The STAT3-binding long noncoding RNA *Inc-DC* controls human dendritic cell differentiation. *Science* **344**, 310–313. (doi:10.1126/science.1251456)
101. Hu X *et al.* 2014 A functional genomic approach identifies *FAL1* as an oncogenic long noncoding RNA that associates with *BMI1* and represses *p21* expression in cancer. *Cancer Cell* **26**, 344–357. (doi:10.1016/j.ccr.2014.07.009)
102. Yang F, Huo X-S, Yuan S-X, Zhang L, Zhou W-P, Wang F, Sun S-H. 2013 Repression of the long noncoding RNA-*LET* by histone deacetylase 3 contributes to hypoxia-mediated metastasis. *Mol. Cell* **49**, 1083–1096. (doi:10.1016/j.molcel.2013.01.010)

103. Lin A *et al.* 2016 The LINK-A lncRNA activates normoxic HIF1 α signalling in triple-negative breast cancer. *Nat. Cell Biol.* **18**, 213–224. (doi:10.1038/ncb3295)
104. Li C *et al.* 2017 A ROR1–HER3–lncRNA signalling axis modulates the Hippo–YAP pathway to regulate bone metastasis. *Nat. Cell Biol.* **19**, 106–119. (doi:10.1038/ncb3464)
105. Leucci E *et al.* 2016 Melanoma addiction to the long non-coding RNA SAMMSON. *Nature* **531**, 518–522. (doi:10.1038/nature17161)
106. Zhang Y *et al.* 2016 Long noncoding RNA LINP1 regulates repair of DNA double-strand breaks in triple-negative breast cancer. *Nat. Struct. Mol. Biol.* **23**, 522–530. (doi:10.1038/nsmb.3211)
107. Wang KC *et al.* 2011 A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature* **472**, 120–124. (doi:10.1038/nature09819)
108. Amaral PP *et al.* 2016 Genomic positional conservation identifies topological anchor point (tap)RNAs linked to developmental loci. Pre-print at bioRxiv. (doi:10.1101/051052)
109. Engreitz JM, Ollikainen N, Guttman M. 2016 Long non-coding RNAs: spatial amplifiers that control nuclear structure and gene expression. *Nat. Rev. Mol. Cell Biol.* **17**, 756–770. (doi:10.1038/nrm.2016.126)
110. Hacisuleyman E *et al.* 2014 Topological organization of multichromosomal regions by the long intergenic noncoding RNA Firre. *Nat. Struct. Mol. Biol.* **21**, 198–206. (doi:10.1038/nsmb.2764)
111. Clemson CM, Hutchinson JN, Sara SA, Ensminger AW, Fox AH, Chess A, Lawrence JB. 2009 An architectural role for a nuclear noncoding RNA: NEAT1 RNA is essential for the structure of paraspeckles. *Mol. Cell* **33**, 717–726. (doi:10.1016/j.molcel.2009.01.026)