

# Lack of heme synthesis in a free-living eukaryote

Anita U. Rao\*, Lynn K. Carta†, Emmanuel Lesuisse‡, and Iqbal Hamza\*§

\*Department of Animal and Avian Sciences and Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742;

†Nematology Laboratory, United States Department of Agriculture–Agricultural Research Service, Beltsville, MD 20705; and ‡Laboratoire d'Ingénierie des Protéines et Contrôle Métabolique, Institut Jacques Monod, Tour 43, Université Paris 7/Paris 6, 2 Place Jussieu, 75251 Paris Cedex 05, France

Communicated by J. Clark Lagarias, University of California, Davis, CA, February 2, 2005 (received for review September 22, 2004)

In most free-living eukaryotes studied thus far, heme is synthesized from a series of intermediates through a well defined evolutionarily conserved pathway. We found that free-living worms, including the model genetic organism *Caenorhabditis elegans*, and parasitic helminths are unable to synthesize heme *de novo*, even though these animals contain hemoproteins that function in key biological processes. Radioisotope, fluorescence labeling, and heme analog studies suggest that *C. elegans* acquires heme from exogenous sources. Iron-deprived worms were unable to grow in the presence of adequate heme unless rescued by increasing heme levels in the growth medium. These data indicate that although worms use dietary heme for incorporation into hemoproteins, ingested heme is also used as an iron source when iron is limiting. Our results provide a biochemical basis for the dependence of worm growth and development on heme, and they suggest that pharmacologic targeting of heme transport pathways in worms could be an important control measure for helminthic infections.

*Caenorhabditis elegans* | iron | metals | nematode | prophyrin

Helminthic infections are an enormous burden to public health and global agriculture. More than two billion people are affected by helminthiasis, including schistosomiasis, and plant-parasitic nematodes cause an estimated annual crop loss of eighty billion dollars (1, 2). There is an urgent need to find unique vulnerabilities in helminths because drug resistance by nematodes is already prevalent in livestock and other animals, and schistosomes resistant to praziquantel have been documented in places where this antihelminthic drug is copiously used (3, 4). Within a parasitized host, helminths exhibit distinct nutritional adaptations such that they acquire their food unidirectionally from the host to sustain their growth and reproduction (5). Thus, metabolic pathways essential for nutrient acquisition in worms could be exploited as potential drug targets to control helminthic infections.

Phylogenetic analysis of biosynthetic enzymes in the evolutionarily conserved multistep pathway for heme synthesis,  $\delta$ -aminolevulinic acid dehydratases (ALAD) and porphobilinogen deaminases (PBGD), has suggested that *Caenorhabditis elegans* lacks orthologs for these enzymes and therefore may acquire tetrapyrroles nutritionally (6). Correspondingly, the trypanosomatid protozoa, *Leishmania* spp., appear to lack seven of the eight enzymes of the heme pathway with the exception being ferrochelatase (7). This defect in tetrapyrrole synthesis is manifested as a nutritional requirement for heme or its immediate precursor protoporphyrin IX. Early studies demonstrated that *C. elegans*, *Caenorhabditis briggsae*, and *Rhabditis maupasi* require cytochrome *c* or hemoglobin as a heme source for growth and reproduction (8–10). However, it is unclear why these nematodes require heme to grow and whether this nutritional necessity also exists in related helminths.

Hemes are critical cofactors for many biological processes, including oxidative metabolism (11), xenobiotic detoxification (12), the synthesis and sensing of diatomic gases (13), cellular differentiation (14), gene regulation at the level of transcription (15), protein translation (16) and targeting (17), and maintaining protein stability (18). Within cells, protoheme (iron-protoporphyrin IX) is synthesized via well defined intermediates that are highly conserved through evolution. Depending on the organelle and cell type, heme

pathway intermediates are used in the synthesis of other tetrapyrrole compounds, including bilins, chlorophylls, and corrins (19, 20). The first universal precursor for the synthesis of heme is  $\delta$ -aminolevulinic acid. Heme synthesis culminates when ferrochelatase catalyzes insertion of ferrous iron into the protoporphyrin IX ring to form protoheme (21). Protoheme is incorporated into numerous heme proteins or is modified further for the synthesis of other types of heme found in cytochrome *c* and terminal oxidases.

An analysis of worm genomes showed that worms lack orthologs to genes encoding enzymes for heme biosynthesis. Here we demonstrate that *C. elegans*, a free-living multicellular eukaryote, and medically relevant helminths are natural heme auxotrophs and suggest that *C. elegans* may serve as a model system for understanding the mechanisms of heme transport in eukaryotes.

## Materials and Methods

**Worm Culture and Growth Assays.** Free-living worms were cultured in CeHR axenic liquid medium (Eric Clegg, personal communication; composition and preparation available on request). Worm growth (generation time 3.5 days), mobility, and development in CeHR medium were comparable to those of worms cultured on *Escherichia coli* in agar plates. We modified CeHR medium (henceforth called mCeHR medium) to eliminate all sources of exogenously added heme; basal growth medium comprised 20  $\mu$ M hemin chloride (Frontier Scientific, Logan, UT) and 150  $\mu$ M ferrous ammonium sulfate (Sigma). Worm strains were grown in mCeHR medium under aerobic conditions in tissue culture flasks at 20°C. We routinely obtained  $\approx 3.7 \times 10^6$  worms after 3.5–4 days in a T75 flask with 30 ml of basal medium from an initial inoculum of  $\approx 1.5 \times 10^5$  worms. For initial culturing of worms in axenic media, three generations of worms were sequentially bleached [1.1% Clorox bleach (sodium hypochlorite)/0.55 M NaOH] to eliminate any contaminating bacterial carryover from agar plates. In all dose–response experiments, worms were growth synchronized by treating the gravid hermaphrodite worms with bleach to dissolve adult worms. The eggs, resistant to bleach, were liberated from the carcasses and extensively washed with M9 buffer followed by overnight hatching in M9 buffer to synchronize growth. By harvesting at appropriate time intervals, synchronous larval stages and adult staged worms were collected for experimental manipulations. Identical numbers of synchronized L1 larvae were inoculated into media with different heme concentrations in 12- or 24-well culture plates. Worm growth was monitored each day and an aliquot was obtained for counting by microscopy, usually at days 3, 6, and 9. The worms in each well were counted twice and each growth condition was analyzed in triplicate. *P* values for statistical significance were calculated by using a one-way ANOVA with Student–Newman–Keuls multiple comparison test by using INSTAT version 3.01 (GraphPad, San Diego).

Abbreviations: ZnMP, zinc mesoporphyrin; HO, heme oxygenase.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. AY856093).

§To whom correspondence should be addressed at: 2413 ANSC, Building 142, University of Maryland, College Park, MD 20742. E-mail: hamza@umd.edu.

© 2005 by The National Academy of Sciences of the USA

**Preparation of Hemin, Hemin Analogs, and [<sup>59</sup>Fe]Heme.** Fresh stock solutions of hemin or hemin analogs (Frontier Scientific) were prepared immediately before use by dissolving required amounts in 0.3 M ammonium hydroxide. The pH of the stock solution was adjusted to 8.0 with 6 M HCl, and the solution was filter sterilized (pore diameter 0.45 μm). The upper limit of free iron was estimated to be 3.8 nM/1 μM hemin chloride by inductively coupled plasma–mass spectrometry (ICP-MS) analysis. For synthesis of [<sup>59</sup>Fe]heme, 50 ml of glacial acetic acid was stirred under a constant flow of N<sub>2</sub> at 60°C followed by addition of 12 mg of protoporphyrin IX in pyridine for 30 min (22). To this mixture, 0.85 μCi of FeCl<sub>3</sub> (specific activity 35.77 mCi/mg, Perkin–Elmer; 1 μCi = 37 kBq) was stirred in for an additional 3 h. Heme was extracted from this mixture with ethyl acetate followed by extensive washes with 4 M HCl and distilled water to remove unincorporated protoporphyrin IX and iron. The heme thus obtained was concentrated by evaporation of the ethyl acetate by using a RotaVapor and frozen at –20°C until further use. Total amount of [<sup>59</sup>Fe]heme synthesized by this method was measured by using a Packard Gamma Counter (~21% efficiency). The purity of heme was determined by thin-layer chromatography using silica gel 60 matrix in an NH<sub>4</sub>OH-saturated chamber with 2,6-lutidine/water solvent. We estimated the specific activity of [<sup>59</sup>Fe]heme synthesized to be ≈2.8 × 10<sup>6</sup> dpm/nmol.

**Metabolic Labeling, Heme Isolation, and TLC.** Equal numbers of L1 worms were grown at 20°C in mCeHR medium with 20 μM hemin and supplemented with 9.4 × 10<sup>6</sup> dpm of either [<sup>59</sup>Fe]heme or <sup>59</sup>FeCl<sub>3</sub>. Radiolabeled adult worms were harvested by first incubating them in M9 buffer for 30 min to empty the gut contents. Worms were then extensively washed with large amounts of M9 buffer/1 mM EDTA on a Gelman Metrical membrane (pore diameter 0.45 μm) that had been incubated with 20 μM hemin or FeCl<sub>3</sub> to prevent nonspecific absorption of the radiotracer signal to the filter membrane. The experiment was done in parallel in the presence of 1 mM sodium azide (NaN<sub>3</sub>) to account for nonspecific binding of the radiolabel to the biological samples. We experimentally determined that this concentration of NaN<sub>3</sub> was not lethal to worms. To isolate heme, worms were washed with M9 buffer and resuspended in ice-cold 1 M HCl to a final pH of 2.0. The acidified solutions were incubated on ice for 30 min to allow complete protein denaturation, and then an equal volume of ice-cold 2-butanone was added (23). The solutions were shaken and allowed to stand until the ketone (heme) and aqueous (worm debris) phases separated. The upper ketonic phase was removed and the heme was concentrated by rotary evaporation. The heme was resuspended in dimethylformamide, and equal volumes of samples and controls were spotted and resolved on silica gel 60 TLC plates. Plates were exposed to a PhosphorImager and the radiolabel signal corresponding to heme was excised and analyzed by using a γ counter. Counts (dpm) obtained were normalized for protein as determined by the bicinchoninic acid method (Sigma) performed on aliquot samples of intact worms. The specific activity of [<sup>59</sup>Fe]heme added to the growth medium was 4.7 × 10<sup>4</sup> dpm/nmol, and the specific activity of [<sup>59</sup>Fe]heme extracted from worms was 0.69 × 10<sup>4</sup> dpm/nmol. The most plausible explanation for this difference in specific activity of [<sup>59</sup>Fe]heme is dilution of the supplied radiolabel heme with preexisting unlabeled heme endogenous to the worm.

**Pulse–Chase Analysis.** Mixed populations of worms grown in mCeHR medium with 4 μM hemin were labeled for 16 h in the presence of either 40 μM zinc mesoporphyrin (ZnMP) or hemin. We empirically determined that 40 μM ZnMP labels the worms without affecting viability. Fluorescently labeled worms were washed with M9 buffer, dispensed into 12-well plates, and allowed to incorporate unlabeled hemin at concentrations of 40, 400, or 800 μM. At timed intervals, aliquots of worms were removed into the appropriate medium containing 10 mM NaN<sub>3</sub>, mounted on a slide, and immediately analyzed with a 543 He/Ne laser on a Zeiss 410

confocal microscope, or with a Peltier-cooled Retiga 1300 12-bit charge-coupled device camera fitted on to a Leica DMIRE2 epifluorescence/differential interference contrast microscope. Images were further analyzed with SIMPLEPCI version 5.2 software (Compix, Cranberry Township, PA). Sensor gain and exposure times were kept constant during all image acquisition. No loss of fluorescence was observed when labeling experiments were performed in parallel in the presence of 1 mM NaN<sub>3</sub> during the chase period and when worms were incubated in medium without hemin. To account for background autofluorescence in *C. elegans*, the sensor gain of the charge-coupled device camera or the laser was set to subtract any fluorescence obtained from control worms incubated in the presence of 40 μM hemin.

**Other Methods.** Other methods are described in *Supporting Methods*, which is published as supporting information on the PNAS web site.

## Results

Analysis of publicly available worm genome databases revealed that these genomes lack obvious orthologs to heme biosynthesis pathway enzymes (6). We queried genome databases by using sequences of the human enzymes that catalyze the seven sequential steps to synthesize heme from the first universal precursor, δ-aminolevulinic acid. Expect (*E*) threshold values obtained from BLAST searches revealed nonsignificant hits only in the *C. elegans* database (Table 2, which is published as supporting information on the PNAS web site) as compared with genome databases from *Saccharomyces cerevisiae*, *Drosophila melanogaster*, and mouse, thus suggesting that *C. elegans* lacks orthologous genes for enzymes that catalyze heme synthesis (24, 25). These *in silico* observations were confirmed by measuring enzyme activity. Because free-living worms in the laboratory use *E. coli* as food, bacterial metabolites could confound identification of the source of heme and enzyme activity. Therefore, we grew worms axenically in mCeHR liquid medium in lieu of growth on Petri plates containing *E. coli* (see *Materials and Methods*). We used three physiologically distinct but phylogenetically related free-living bacterivorous strains, *C. elegans*, *Panagrellus redivivus*, and *Oscheius myriophila* (Fig. 4, which is published as supporting information on the PNAS web site). Synchronized worms were grown aerobically at 20°C in mCeHR to the gravid adult stage and homogenized to obtain cytosol- and mitochondria-enriched fractions for analysis of heme biosynthesis enzymes. Under the assay conditions used, δ-aminolevulinic acid dehydratase and porphobilinogen deaminase activities were undetectable in worm lysates as compared with wild-type *E. coli* lysates (Table 1).

The protozoan *Leishmania* and certain microorganisms such as *Haemophilus influenzae* contain only part of the heme biosynthetic pathway (7, 19). Thus, it is possible that worms also have retained the ability to synthesize heme by using an intermediate of the heme pathway. To directly address this issue, we measured ferrochelatase activity, an inner mitochondrial membrane enzyme, which catalyzes the final step in the heme biosynthetic pathway. Because ferrochelatase from *S. cerevisiae* has been genetically and biochemically well characterized, we used yeast as a control source of this enzyme (21). Ferrochelatase activity was readily detected in wild-type yeast but was undetectable in combined mitochondria- and cytosol-enriched fractions from all three worm species and a *S. cerevisiae* ferrochelatase mutant with genetic disruption at the *HEM15* locus (Table 1). Activity for succinate dehydrogenase, another inner mitochondrial membrane enzyme, was readily detectable in these fractions, indicating the presence of mitochondrial membranes. However, the inability to detect heme synthesis enzyme activities could be attributed to the presence of endogenous inhibitors in worm extracts. We found no inhibition of δ-aminolevulinic acid dehydratase, porphobilinogen deaminase, or ferrochelatase enzyme activities when *C. elegans* extracts were mixed in equal proportions with extracts from either *E. coli* or *S. cerevisiae* (Table 3, which is published as supporting information on the PNAS web site).

**Table 1. Heme biosynthetic enzyme activities in free-living and parasitic worms**

Organism	Host	Enzyme activity, nmol/min per mg of protein*			
		ALAD	PBGD	FC	SDH
<b>Helminths</b>					
<i>Caenorhabditis elegans</i> <sup>†</sup>	Free	ND	ND	ND	725 ± 160 <sup>‡</sup>
<i>Panagrellus redivivus</i> <sup>†</sup>	Free	ND	ND	ND	3,419.3 ± 424.7 <sup>‡</sup>
<i>Oscheius myriophila</i> <sup>†</sup>	Free	ND	ND	ND	1,567 ± 594.6 <sup>‡</sup>
<i>Paragordius varius</i>	Free <sup>§</sup>	ND	ND	–	715.9 ± 9.2
<i>Schistosoma mansoni</i>	Human	ND	ND	ND	515 ± 80.6
<i>Strongyloides stercoralis</i>	Human	ND	ND	ND	715.2 ± 53.8 <sup>‡</sup>
<i>Ancylostoma caninum</i>	Dog	ND	ND	ND	266.7 ± 45.6 <sup>‡</sup>
<i>Haemonchus contortus</i>	Sheep/goat	ND	ND	ND	30.3 ± 4.6 <sup>‡</sup>
<i>Trichuris suis</i>	Pig	ND	ND	ND	7.6 ± 0.8 <sup>‡</sup>
<i>Ascaris suum</i>	Pig	ND	ND	ND	24.2 ± 3.2 <sup>‡</sup>
<b>Bacteria</b>					
<i>E. coli</i>	NA	1.47 ± 0.03	0.035 ± 0	16.36 ± 2.71	1,865 ± 318
<i>E. coli</i> (ALAD mutant)	NA	0.034 ± 0	–	–	–
<b>Yeast</b>					
<i>S. cerevisiae</i>	NA	0.97 ± 0.04	0.096 ± 0	27.42 ± 2.92 <sup>‡</sup>	2,371.2 ± 32.1 <sup>‡</sup>
<i>S. cerevisiae</i> (FC mutant)	NA	0.04 ± 0	0.032 ± 0	ND	915 ± 177 <sup>‡</sup>

Wild-type and mutant *E. coli* and *S. cerevisiae* were used as positive controls for enzyme activity. ND, enzyme activity not detected under current assay conditions; NA, not applicable; –, not assayed; ALAD,  $\delta$ -aminolevulinic acid dehydratase; PBGD, porphobilinogen deaminase; FC, ferrochelatase; SDH, succinate dehydrogenase.

\*Mean values (triplicates) of product formed ± SD.

<sup>†</sup>Grown in axenic mCeHR medium.

<sup>‡</sup>Average activity found in combined crude mitochondria- and cytosol-enriched fractions.

<sup>§</sup>*P. varius* adults live in fresh water but larvae develop in arthropods.

To address whether the lack of discernable heme enzyme activities also held for other worm species that are phylogenetically diverse, we examined five parasitic nematodes that have different host specificities (*Strongyloides stercoralis*, *Ancylostoma caninum*, *Haemonchus contortus*, *Trichuris suis*, and *Ascaris suum*), a nematomorph (*Paragordius varius*), and a trematode flatworm (*Schistosoma mansoni*) (Fig. 4). Irrespective of their host affiliations, enzyme activities were undetectable in all helminthic extracts in our assay conditions (Table 1). These measurements provide further support for the hypothesis that the helminths examined in this study do not have enzymes for heme synthesis.

*C. elegans* appears to have a large number of hemoprotein orthologs, including globin isoforms (26), guanylate cyclases (27), adenylate cyclases (28), catalases (29), cytochrome P450s (30), and respiratory cytochromes (31). Although hemes have been found in all phyla, certain microbial pathogens such as *Borrelia burgdorferi* and *Treponema pallidum* neither make heme nor contain hemoproteins (19, 32). Reduced minus oxidized absorption spectra of pyridine hemochromes revealed that *C. elegans* has discernable hemoproteins in worm fractions enriched either for cytosol or for membranes, including mitochondrial membranes (Fig. 1A). The purity of these fractions was determined by immunoblots probed with antisera for  $\alpha$ -tubulin and ATP2, the  $\beta$ -subunit of the F<sub>1</sub> sector of the mitochondrial F<sub>1</sub>F<sub>0</sub> ATP synthase (33). With ATP2 used as a marker, <1% of the signal was found in the cytosol-enriched fraction, whereas  $\approx$ 88% of ATP2 protein was found in the membrane-enriched fraction. Ultralow-temperature spectra of total worm homogenates revealed the presence of detectable amounts of respiratory cytochromes *a*, *b*, and *c* (Fig. 1B).

To quantitatively determine the heme requirement of worms, *C. elegans* was cultured in the presence or absence of nutritional heme supplements. Worms grown in the absence of exogenous heme (supplemented as hemin chloride) were growth arrested at the L4 stage, whereas worms grown in the presence of heme grew robustly and reproduced over multiple generations (Fig. 1C). Similar heme-dependent growth was also observed for *P. redivivus* and *O. myriophila* (data not shown). This lack of growth in the absence of

heme was reversible, because replenishing heme to the heme-depleted media resulted in normal growth rates of the arrested larvae (data not shown). However, unlike *Leishmania*, which can exogenously acquire either hemin or its immediate precursor protoporphyrin IX for growth (7), *C. elegans* was unable to use protoporphyrin IX (supplied as free acid or disodium salt) as a substitute for hemin (Fig. 1C). Cytochrome *c* or hemoglobin also sustained worm growth, supporting previous studies that have shown heme and vitamin B12, another tetrapyrrole, as necessary factors for *C. briggsae* development (8, 9, 34). Our observations that worms can develop to the L4 stage in the absence of exogenous heme in the growth medium suggest that either there are trace amounts of heme in the medium or, more plausibly, maternal heme stored in the egg during embryogenesis is able to sustain early larval growth.

Worms responded in a biphasic manner to heme when grown in the presence of various amounts of hemin (Fig. 1D). Although the optimal concentration of hemin for growth was 20  $\mu$ M, worms continued to grow and reproduce at concentrations ranging from 1.5 to 500  $\mu$ M hemin, albeit with significantly longer generation times and smaller brood size. Large amounts of hemin ( $\geq$ 800  $\mu$ M) resulted in growth arrest at the L3 stage, possibly because hemes are cytotoxic because of peroxidase activity. To determine if *C. elegans* acquires heme directly from the growth medium, worms were metabolically labeled in the presence of equivalent amounts of <sup>59</sup>Fe or [<sup>59</sup>Fe]heme, and worm homogenates were analyzed for the respective radiotracer. A specific signal was obtained in samples containing worms labeled with [<sup>59</sup>Fe]heme (Fig. 1E, lane 2). Thus, worms use [<sup>59</sup>Fe]heme directly from the growth medium to fulfill their heme auxotrophy and incorporate the tetrapyrrole into proteins. However, no radiolabeled signal was observed in heme extracted from worms when <sup>59</sup>Fe by itself was used (Fig. 1E, lane 4). This observation corroborates our genomic and biochemical analysis (Tables 1 and 2) that *C. elegans* lack ferrochelatase, the terminal enzyme in heme biosynthesis. [<sup>59</sup>Fe]Heme uptake was negligible in worms that were metabolically inhibited in the presence of NaN<sub>3</sub>, an inhibitor of the mitochondrial respiratory chain.







the ability to synthesize heme. This conclusion is supported by genomic, biochemical, and nutritional analysis (Tables 1 and 2 and Fig. 1 C–E) and by previous studies which show that heme is a growth factor for *C. elegans* development (8, 9). This inability to make heme is surprising, given that other free-living metazoans make endogenous heme, and heme synthesis is catalyzed by enzymes encoded by at least seven separate genes that are not clustered in the genome. It is plausible that the ancestral worm lost the genes responsible for heme biosynthesis due to a lack of selective pressure because the progenitor had access to heme either from a parasitized host or from a symbiotic relationship with another organism. Recent studies have shown that the cattle tick *Boophilus microplus*, a bloodsucking arthropod, relies on blood meals to acquire heme (41), and pathogenic human filarial nematodes as well as certain insects harbor the bacteria *Wolbachia*, an intracellular symbiont that has a mutualistic relationship with its host such that the nematode acquires endobacterial-derived metabolites (42, 43). Indeed, the *Wolbachia* genome contains orthologs of genes for heme biosynthetic enzymes, suggesting that this endosymbiont has the ability to make heme.

The phylogenetic maximum parsimony tree (Fig. 4) shows that the loss of the heme pathway is common rather than exceptional among free-living and parasitic nematodes, and it can be found in higher taxa such as the Nematomorpha and Platyhelminthes, both of which have the potential for free-living and parasitic habits. Although the closest outgroup to nematodes is still controversial, if one accepts the Ecdysozoa theory, in which arthropods are a sister group to nematodes (44), the loss of the heme synthesis pathway would have occurred more frequently than with a more traditional grouping of helminths (45). A loss in the heme pathway would be consistent with a close phylogenetic relationship among helminths. Additional taxon sampling along with genomic and biochemical analyses may help clarify the disputed phylogenetic placement of helminths in the animal kingdom (46, 47). Evaluation of the complexity and phylogenetic importance of different molecular phylogenies versus the loss of the heme synthesis pathway cannot be made here. However, it is interesting to note that some other biochemical pathways also show distinct differences between classically simpler invertebrates (flatworms and nematodes) and higher invertebrates (arthropods and molluscs). These include an increased variety of different neuropeptides in higher invertebrates compared with lower (48) and the absence of a Toll-like pathway for immunity in nematodes that is present in insects (49, 50).

Our studies provide preliminary evidence that adaptation to heme auxotrophy in worms has enabled worms to use heme in its entirety as a prosthetic group under normal conditions and as an iron source when iron is low in the environment. We have been unable to identify any significant orthologs of HOs in the worm genome, although we found two putative orthologs of biliverdin reductase in the *C. elegans* and *C. briggsae* genomes. Because HOs from bacteria to humans are a diverse group of heme-catabolizing enzymes, it is possible that heme degradation in *C. elegans* is either catalyzed by a HO with low sequence homology to known HOs or by an entirely novel enzyme. Enzyme activities for HO and biliverdin reductase in homogenates from the trematode *Schistosoma japonicum* have been reported, thus raising the possibility that *C. elegans* may also have the ability to enzymatically degrade heme to obtain iron (51, 52).

Because worms lack the ability to make heme and therefore rely solely on exogenous heme for metabolic processes, these animals must have evolved specific mechanisms for intestinal heme absorption, trafficking, and sequestration. Perhaps, exceptional to worms, an intercellular heme transport system may be required to provide heme to other cell types beyond intestinal cells. Free heme is hydrophobic and is insoluble in the aqueous cellular milieu, and hemes have peroxidase activity that can damage biological macromolecules (53). Thus, in principle, intracellular pathways must exist for heme trafficking in eukaryotes. *C. elegans* may therefore serve as a unique animal model of an obligate heme auxotroph to genetically and cell-biologically delineate the pathways for heme homeostasis that have heretofore been elusive. Identification of molecules involved in heme homeostasis should permit selective drug-targeting of helminthic heme transport and heme-dependent cellular pathways that discriminate the parasite from its host and significantly reduce chronic morbidity and debilitation in affected individuals.

We thank Ray Fetterer, Ben Hanelt, Dolores Hill, Phil LoVerde, and Gerhard Schad for large quantities of hard-to-obtain worm samples for biochemical analysis; Jerry Bommer for ICP-MS analysis and expert technical advice on porphyrin chemistry; Eric Clegg for invaluable help with the CeHR medium; Carla Koehler for ATP2p antisera; W. Kelley Thomas for help with the SSU 18S rDNA gene sequencing strategy; Mark O'Brian and Jerry Kaplan for *E. coli* and yeast strains; the *Caenorhabditis* Genetics Center for worm strains; and Bruce Bowerman, Jonathan Gitlin, Dan Kosman, Mike Krause, Ian Mather, Mark O'Brian, and Tom Porter for helpful discussions. This work was supported by National Institutes of Health Grant DK68260 (to I.H.).

- Colley, D. G., LoVerde, P. T. & Savioli, L. (2001) *Science* **293**, 1437–1438.
- Chitwood, D. J. (2003) *Pest Manage. Sci.* **59**, 748–753.
- Sangster, N. C. & Gill, J. (1999) *Parasitol. Today* **15**, 141–146.
- Ismail, M., Botros, S., Metwally, A., William, S., Farghally, A., Tao, L. F., Day, T. A. & Bennett, J. L. (1999) *Am. J. Trop. Med. Hyg.* **60**, 932–935.
- Halton, D. W. (1997) *Int. J. Parasitol.* **27**, 693–704.
- Jaffe, E. K. (2003) *Chem. Biol.* **10**, 25–34.
- Sah, J. F., Ito, H., Kolli, B. K., Peterson, D. A., Sassa, S. & Chang, K. P. (2002) *J. Biol. Chem.* **277**, 14902–14909.
- Hieb, W. F., Stokstad, E. L. & Rothstein, M. (1970) *Science* **168**, 143–144.
- Vanfleteren, J. R. (1974) *Nature* **248**, 255–257.
- Brockelman, C. R. & Jackson, G. J. (1978) *J. Parasitol.* **64**, 803–809.
- Wenger, R. H. (2000) *J. Exp. Biol.* **203**, 1253–1263.
- Lewis, D. F. (2004) *Pharmacogenomics* **5**, 305–318.
- Rodgers, K. R. (1999) *Curr. Opin. Chem. Biol.* **3**, 158–167.
- Nakajima, O., Takahashi, S., Harigae, H., Furuyama, K., Hayashi, N., Sassa, S. & Yamamoto, M. (1999) *EMBO J.* **18**, 6282–6289.
- Sun, J., Hoshino, H., Takaku, K., Nakajima, O., Muto, A., Suzuki, H., Tashiro, S., Takahashi, S., Shibahara, S., Alam, J., et al. (2002) *EMBO J.* **21**, 5216–5224.
- Chen, J. J. & London, I. M. (1995) *Trends Biochem. Sci.* **20**, 105–108.
- Lathrop, J. T. & Timko, M. P. (1993) *Science* **259**, 522–525.
- Qi, Z., Hamza, I. & O'Brian, M. R. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 13056–13061.
- Panek, H. & O'Brian, M. R. (2002) *Microbiology* **148**, 2273–2282.
- Papenbrock, J. & Grimm, B. (2001) *Planta* **213**, 667–681.
- Dailley, H. A. (2002) *Biochem. Soc. Trans.* **30**, 590–595.
- Galbraith, R. A., Sassa, S. & Kappas, A. (1985) *J. Biol. Chem.* **260**, 12198–12202.
- Teale, F. W. (1959) *Biochim. Biophys. Acta* **35**, 543.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* **215**, 403–410.
- Altschul, S. F. & Gish, W. (1996) *Methods Enzymol.* **266**, 460–480.
- Kloek, A. P., McCarter, J. P., Setterquist, R. A., Schedl, T. & Goldberg, D. E. (1996) *J. Mol. Evol.* **43**, 101–108.
- Riedl, C. A. & Sokolowski, M. B. (2004) *Curr. Biol.* **14**, R657–8.
- Moorman, C. & Plasterk, R. H. (2002) *Genetics* **161**, 133–142.
- Petriv, O. I. & Rachubinski, R. A. (2004) *J. Biol. Chem.* **279**, 19996–20001.
- Gotoh, O. (1998) *Mol. Biol. Evol.* **15**, 1447–1459.
- Tsang, W. Y., Sayles, L. C., Grad, L. I., Pilgrim, D. B. & Lemire, B. D. (2001) *J. Biol. Chem.* **276**, 32240–32246.
- Posey, J. E. & Gherardini, F. C. (2000) *Science* **288**, 1651–1653.
- Grad, L. I. & Lemire, B. D. (2004) *Hum. Mol. Genet.* **13**, 303–314.
- Lu, N. C., Hieb, W. F. & Stokstad, E. L. (1976) *Proc. Soc. Exp. Biol. Med.* **151**, 701–706.
- Stojiljkovic, I., Kumar, V. & Srinivasan, N. (1999) *Mol. Microbiol.* **31**, 429–442.
- Greenbaum, N. L. & Kappas, A. (1991) *Photochem. Photobiol.* **54**, 183–192.
- Worthington, M. T., Cohn, S. M., Miller, S. K., Luo, R. Q. & Berg, C. L. (2001) *Am. J. Physiol.* **280**, G1172–G1177.
- Stojiljkovic, I. & Perkins-Balding, D. (2002) *DNA Cell Biol.* **21**, 281–295.
- Pendrak, M. L., Yan, S. S. & Roberts, D. D. (2004) *Arch. Biochem. Biophys.* **426**, 148–156.
- Maines, M. D. (2000) *Cell. Mol. Biol. (Noisy-le-grand)* **46**, 573–585.
- Braz, G. R., Coelho, H. S., Masuda, H. & Oliveira, P. L. (1999) *Curr. Biol.* **9**, 703–706.
- Taylor, M. J. & Hoerauf, A. (1999) *Parasitol. Today* **15**, 437–442.
- Henkle-Duhrsen, K., Eckelt, V. H., Wildenburg, G., Blaxter, M. & Walter, R. D. (1998) *Mol. Biochem. Parasitol.* **96**, 69–81.
- Aguinaldo, A. M., Turbeville, J. M., Linford, L. S., Rivera, M. C., Garey, J. R., Raff, R. A. & Lake, J. A. (1997) *Nature* **387**, 489–493.
- Brusca, R. C. & Brusca, G. J. (1990) *Invertebrates* (Sinauer, Sunderland, MA).
- Blair, J. E., Ikco, K., Gojborji, T. & Hedges, S. B. (2002) *BMC Evol. Biol.* **2**, 7.
- Copley, R. R., Aloy, P., Russell, R. B. & Telford, M. J. (2004) *Evol. Dev.* **6**, 164–169.
- Masler, E. P. (2005) in *Encyclopedia of Molecular Cell Biology and Molecular Medicine*, ed. Meyers, R. A. (Wiley-VCH, Weinheim, Germany), Vol. 9, pp. 167–181.
- Hodgkin, J. (2004) *Nat Immunol.* **5**, 471–472.
- Fitzpatrick, M. J. & Sokolowski, M. B. (2004) *Integr. Comp. Biol.* **44**, 28–36.
- Wen-qi, L. & Yong-long, L. (2004) *Chin. J. Parasitol. Parasit. Dis.* **22**, 106–108.
- Liu, W. Q., Li, Y. L. & Ruppel, A. (2001) *Chin. J. Parasitol. Parasit. Dis.* **19**, 84–86.
- Balla, G., Vercellotti, G. M., Muller-Eberhard, U., Eaton, J. & Jacob, H. S. (1991) *Lab. Invest.* **64**, 648–655.