

Received: 2016.12.21
Accepted: 2017.02.03
Published: 2017.08.09

Association Between ABCB1 Gene Polymorphism and Renal Function in Patients with Hypertension: A Case-Control Study

Authors' Contribution:
Study Design A
Data Collection B
Statistical Analysis C
Data Interpretation D
Manuscript Preparation E
Literature Search F
Funds Collection G

BDE 1 Xiaofang Chen
BD 2 Tao Zhou
CDE 2 Deguang Yang
AE 3 Jinyang Lu

1 Department of Nursing, The Third Affiliated Hospital of Southern Medical University, Guangzhou, Guangdong, P.R. China
2 Department of Cardiology, The Third Affiliated Hospital of Southern Medical University, Guangzhou, Guangdong, P.R. China
3 Intervention Catheterization Room of Radiology Department, The Second Xiangya Hospital of Central South University, Changsha, Hunan, P.R. China

Corresponding Author: Jinyang Lu, e-mail: lujinyangxy@126.com
Source of support: Departmental sources

Background: A previous study found that target organ damage in patients with hypertension was related to genetic factors. The aim of our study was to explore the association between the ABCB1 gene and renal function injury induced by hypertension.





Material/Methods: We used a case-control study design. Patients with hypertension were enrolled from our hospital between July 2015 and December 2015. Questionnaire data included personal information, life habits and behavior. Clinical data included blood routine examination and liver and renal function. We used restriction fragment length polymorphism methods for ABCB1 gene polymorphism detection.

Results: There were 306 patients with hypertension included in the final analyses: 170 cases of hypertension and 136 controls. Compared to controls, the cases group had higher: drinking ratio (65.3% versus 52.9%, $p=0.029$), body mass index ($p=0.032$), systolic blood pressure ($p<0.001$), total cholesterol ($p=0.004$), blood urea nitrogen ($p=0.029$), creatinine ($p=0.024$), uric acid ($p=0.011$), estimated glomerular filtration rate level ($p<0.001$), and platelet level ($p=0.003$). There were no significant differences for others parameters. Genotype frequency distributions of rs1045642 were statistically significant between the two groups ($\chi^2=24.966$, $p<0.001$). No differences were observed for the frequency distribution of rs10808072 and rs1922242 ($\chi^2=1.293$, $p=0.524$; $\chi^2=0.065$, $p=0.968$). The multivariable logistics results found that patients with TT genotype had a higher risk for renal function injury for hypertensive patients compared to those with CC genotype (OR=3.47, 95% CI: 1.19–10.07).

Conclusions: Our results suggested that the rs1045642-T allele of the ABCB1 gene may be associated with increased risk for renal function injury in hypertensive patients.

MeSH Keywords: Hypertension, Renovascular • Kidney Failure, Chronic • Polymorphism, Genetic

Full-text PDF: <https://www.medscimonit.com/abstract/index/idArt/902954>

 2085  4  —  26



Background

Hypertension is an important public health issue, and a main risk factor for cardiovascular disease and other diseases. Hypertension is a chronic disease can develop into resistant hypertension, and 30% of human blood pressure variation can be attributed to heredity [1,2]. Hypertension can cause target organ damage, including left ventricular hypertrophy, renal function injury, and cerebrovascular distortion. The rate of left ventricular hypertrophy in hypertensive patients is reported to be 15–20%, and hypertension accounts for 60% of all identified factors leading to cerebral injury [3–5]. In recent years, it has been suggested that target organ damage in hypertensive patients may be associated with gene factors [6–9]. Dong et al. found that methylene tetrahydrofolate reductase-C677T gene polymorphism was related to renal function decline in male hypertensive patients [10]. Renal function of patients with TT decreased by 1.37 mL/min/1.73 m² compared with those with CC genotype. Li et al. reported a higher risk of left ventricular hypertrophy in hypertensive patients with Klotho gene [11]. The ATP-binding cassette subfamily B member 1 gene (ABCB1) is common related to drug metabolism; and animal experiments suggest that it is involved in the process of blood regulation. ABCB1 gene encodes permeability glycoprotein. Permeability glycoprotein is widely expressed in many organs such as kidney, liver, and intestinal tract. In rats, permeability glycoprotein may be associated with aldosterone transportation in brain [12] and involved in aldosterone plasma distribution. In humans, the 3435C/T variant may be related to serum aldosterone concentration. Since PGP is also encoded in capillary endothelial cells, the level of permeability glycoprotein expression regulated by ABCB1 gene may also be related to renal dysfunction [13].

Previous studies mainly have focused on drug metabolism, and few studies have reported on the association between gene and hypertension, and the related target-organ damage. This study explored the association of ABCB1 gene polymorphism and renal function damage with the aim of providing scientific guidelines for hypertension-personalized medicine to prevent target organ damage.

Material and Methods

Study population

Using a case-control design, we enrolled 350 hypertensive patients from the Third Affiliated Hospital of South Medical University between July 2015 and December of 2015. Patients completed a study questionnaire and had health and laboratory examinations. Patients who had incomplete or missing genotype information were excluded. Patients were also excluded

for the following: secondary hypertension, heart failure, diabetes mellitus, severe hepatic or renal injury, tumor, systemic inflammatory disease, hematological system disorder and known coronary artery or cerebrovascular disease, disabling diseases such as dementia, or an inability to cooperate were excluded. The hypertensive patients with renal function injury were treated as case group, and the isolated hypertensive patients were a control group. We used the Power and Sample Size Calculation software to estimate the sample size. A previous study reported that the estimated odds ratio was 2.17 for rs1045642 in the Chinese population and the gene frequency of TT genotype was 30–40%. The sample size was at least 121 ($\alpha=0.05$, $1-\beta=0.85$), which was enough for the present study.

The institutional research ethics committee of the Third Affiliated Hospital Southern Medical University approved this study, and written informed consent was obtained from each participant.

Data collection

The collected data included two parts. The demographic and physical data was collected through a standardized questionnaire, including age, sex, smoking, drinking, exercise, self-reported salt intake, education level, marriage status, and region. Height and weight were measured, and body mass index (BMI) was calculated. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) after five minutes of rest was measured in triplicate with at least 15 seconds between measurements with the mean value reported as the final blood pressure.

We took two blood samples from participants early in the morning via the antecubital vein. The blood routine examinations were completed by automatic biochemistry analyzer (Beckman Coulter, Inc., Fullerton, CA, USA), including red blood cell (RBC), white blood cell (WBC), red blood cell distribution width (RDW), hemoglobin (Hb), platelets, high density lipoprotein-cholesterol (HDL-C), low density lipoprotein-cholesterol (LDL-C), triglyceride, total cholesterol, and fasting blood glucose, blood urea nitrogen (BUN), creatinine, uric acid, alanine aminotransferase (ALT), aspartate aminotransferase, (AST).

The diagnosis of hypertension refers to the criterion: SBP \geq 140 mm Hg and/or DBP \geq 90 mm Hg [14]. Kidney function was estimated by glomerular filtration rate using the following equation: $186 \times \text{SCr}^{-1.154} \times \text{age in years}^{-0.203} \times 1.210$ (if black) $\times 0.742$ (if female) [15]. Microalbuminuria was defined as \geq 30 mg/24 hours; diabetes was defined by a fasting blood glucose \geq 7.0 mmol/L or by drug treatment. BMI \geq 24 was defined as overweight, and BMI \geq 28 was defined as obese [16]. Smoking was defined as at least five cigarettes each day for at least half a year, and drinking was defined as \geq 50 mL each week for at least one year [17,18]. Physical exercise was defined as 30 minutes or more of physical exercise for \geq 3 times per week [19].

Genotyping

Single-nucleotide polymorphisms (SNPs) were selected based on minor allele frequency greater than 5% from Chinese SNP data from HapMap. All three SNPs of participants were tested through Hardy-Weinberg methods. The collected blood samples were placed in containers with anticoagulant and stored at -40°C . DNA was extracted from peripheral whole blood using the Qiagen DNA Isolation Kit (Valencia, CA, USA). The ABCB1 gene polymorphism was detected by the restriction fragment length polymorphism [20]. A randomized duplicate detection (5%) was conducted to guarantee accuracy of SNP genotyping.

Statistical analyses

We used Statistical Product and Service Solutions 23.0 software (SPSS, Inc., USA) for all analyses. For continuous data, the two independent samples *t*-test were used for comparisons of two groups. Chi-square test was used for categorical data. Differences of genotype distributions between controls and cases were evaluated by the Chi-square test. The association between three SNPs and renal function injury in patients with hypertension was analyzed via binary logistic regression analysis with some potential factors adjusted. Odds ratios (ORs) and their 95% CIs were also calculated. A two-tailed $p < 0.05$ was considered statistically significant.

Results

Comparisons of general characteristics between controls and cases

The demographic characteristics of controls and cases are presented in Table 1. There were 306 patients with hypertension entered into the final analyses, including 170 patients with renal function injury and 136 controls. There were no significant differences in the distributions of age, sex, smoking, exercise, salt intake based on self-reports, education level, marriage status, and region ($p > 0.05$). Compared with controls, the cases group had higher drinking ratio (65.3% versus 52.9%, $p = 0.029$). The biochemical parameters of the two groups are shown in Table 2. No significant difference was observed in DBP, fasting blood glucose, triglyceride, HDL-C, or LDL-C ($p > 0.05$). However, the case group had higher body mass index ($p = 0.032$), SBP ($p < 0.001$), total cholesterol ($p = 0.004$), BUN ($p = 0.029$), creatinine ($p = 0.024$), uric acid ($p = 0.011$) and eGFR levels ($p < 0.001$). The ALT, AST, RBC, WBC, Hb, and RDW levels of the control group were almost commensurate with the case group ($p > 0.05$), but the difference in platelet levels was significant ($p = 0.003$).

Frequency distribution of SNPs in cases and controls

The genotype distributions of three SNPs are presented in Table 3. The genotype distributions of SNPs were in accordance with Hardy-Weinberg equilibrium in the control group ($\chi^2 = 0.395$, $p = 0.541$, $\chi^2 = 0.160$, $p = 0.689$, $\chi^2 = 0.580$, $p = 0.446$). Genotype frequencies of rs1045642 were significant differences between the control group and case group, as shown in Tables ($\chi^2 = 24.966$, $p < 0.001$). There were no significant differences in the frequency distributions of rs10808072 and rs1922242 between the control group and the case group ($\chi^2 = 1.293$, $p = 0.524$; $\chi^2 = 0.065$, $p = 0.968$).

Multivariable regression analysis

We further explored the association between rs1045642 polymorphism and renal function injury in patients with hypertension. We performed multivariable analyses of the association between rs1045642 and renal function injury after adjusting some potential parameters, including age, sex, smoking, drinking, exercise, salt intake, body mass index, triglyceride, total cholesterol, LDL-C, HDL-C, creatinine, BUN, and uric acid. The multivariable logistics results found that patients with TT genotype had a higher risk of renal function injury for hypertensive patients compared with those with CC genotype (OR=3.47, 95% CI: 1.19–10.07). The risk of renal function injury did not increase for hypertensive patients with CT genotype (OR=2.30, 95% CI: 0.98–5.43). In addition, elevated SBP, drinking more alcohol, and increased creatinine level also increased the risk of renal function injury for hypertensive patients. The specific results are presented in Table 4.

Discussion

In this study we identified that the T allele of the ABCB1 gene was related to a higher risk of renal injury in patients with hypertension. Our findings provide new evidence supporting the association between the drug metabolism-related gene and targeted-organ damage in hypertensive patients, as well as provide a new explanation regarding the mechanisms of targeted-organ injury induced by high blood pressure, and present a new finding for consideration for early identification and prevent for renal function injury in hypertensive patients.

Our results identified a SNP (rs1045642) associated with renal function injury in hypertensive patients. Bochud et al. reported that rs1045642 was associated with renal function and that compared with the CC genotype, people with the TT genotype had higher eGFR levels [21]. This finding was contrary to our results. This difference in result could be attributed to the following aspects. First, the genetic background was different for the two studies. Our study population was from

Table 1. Comparisons of general characteristics between controls and cases.

Variables	Control (n=136)	Case (n=170)	t/ χ^2	P value
Sex			1.398	0.237
Male	70 (51.5%)	99 (58.2%)		
Female	66 (48.5%)	71 (41.8%)		
Age (y)	47.0±13.3	48.0±12.5	-0.691	0.490
Smoking			0.563	0.453
Yes	16 (11.8%)	25 (14.7%)		
No	120 (88.2%)	145 (85.3%)		
Drinking			4.796	0.029
No	64 (47.1%)	59 (34.7%)		
Yes	72 (52.9%)	111 (65.3%)		
Exercise/week			0.428	0.934
None	64 (47.1%)	86 (50.6%)		
One	30 (22.1%)	36 (21.2%)		
Twice	22 (16.2%)	26 (15.3%)		
Everyday	20 (14.7%)	22 (12.9%)		
Salt intake			2.205	0.332
Low	48 (35.3%)	56 (32.9%)		
Moderate	68 (50.0%)	97 (57.1%)		
High	20 (14.7%)	17 (10.0%)		
Education			0.444	0.979
No	13 (9.6%)	13 (7.6%)		
Primary	25 (18.4%)	31 (18.3%)		
Middle	45 (33.1%)	60 (34.3%)		
High	35 (25.7%)	43 (25.5%)		
Colleague	18 (13.2%)	23 (13.4%)		
Marriage			1.340	0.855
Unmarried	52 (38.2%)	62 (35.6%)		
Married	80 (58.8%)	103 (60.6%)		
Others	4 (2.9%)	5 (3.0%)		
Region			0.080	0.778
Rural	38 (27.9%)	50 (29.4%)		
Urban	98 (72.1%)	120 (70.6%)		

Table 2. Comparisons of biochemical indexes between controls and cases.

Variables	Control (n=136)	Case (n=170)	t	P value
Body mass index, kg/m ²	25.7±3.2	26.6±3.8	-2.155	0.032
Systolic blood pressure, mmHg	142.5±12.5	159.6±13.6	-1.326	<0.001
Diastolic blood pressure, mmHg	100.4±8.8	99.4±9.1	0.969	0.333
Fasting blood glucose, mg/dl	6.2±6.8	6.6±7.1	-0.498	0.618
Total Cholesterol, mg/dl	4.8±1.2	5.2±1.2	-2.897	0.004
Triglyceride, mg/dl	1.6±1.2	1.6±0.9	0.00	0.999
High-density lipoprotein-C, mg/dl	1.3±0.3	1.3±0.3	0.008	0.498
Low-density lipoprotein-C, mg/dl	2.8±1.0	2.7±0.9	0.919	0.358
Blood Urea Nitrogen, mg/dl	5.4±1.6	5.8±2.0	-1.896	0.029
Creatinine, mg/dl	79.7±53.3	88.6±21.8	-1.980	0.024
Uric acid, mg/dl	385.9±102.2	415.5±100.6	-2.539	0.011
Estimated glomerular filtration rate, ml/min	75.0±20.3	54.3±16.7	9.786	<0.001
Alanine aminotransferase, U/L	26.5±39.5	25.6±18.1	1.733	0.083
Aspartate aminotransferase, U/L	21.9±14.3	21.7±23.3	0.088	0.930
Red blood cell, ×10 ¹² /l	4.6±0.7	4.6±0.6	0.002	0.996
White blood cell, ×10 ⁹ /L	7.0±9.1	7.2±9.3	-0.188	0.850
Hemoglobin, g/L	140.2±18.6	143.1±19.5	-1.319	0.188
Platelets, ×10 ⁹ /l	245.2±42.2	259.9±43.6	-2.970	0.003
Red cell distribution width, %	12.9±13.7	13.8±9.3	-0.682	0.495

Table 3. Frequency distribution of ABCB1 genotype in controls and cases.

SNP	Genotype	Control group		Case group		χ ²	P value
		n	%	n	%		
rs1045642	CC	57	41.9	34	20.0	24.966	<0.001
	CT	60	44.1	77	45.3		
	TT	19	14.0	59	34.7		
rs10808072	GG	49	36.0	72	42.4	1.293	0.524
	GA	64	47.1	71	41.8		
	AA	23	16.9	27	15.9		
rs1922242	AA	66	48.5	85	49.3	0.065	0.968
	AT	56	41.2	68	40.5		
	TT	14	10.3	17	10.1		

Table 4. Multivariable logistics regression analysis between rs1045642 and renal function in patients with hypertension.

Variables	β	SE	Wald χ^2	P value	OR value	95% CI
SBP	0.027	0.11	5.579	0.018	1.03	1.01–1.05
Drinking	0.713	0.270	6.988	0.008	2.04	1.20–3.46
Creatinine	0.586	0.287	4.175	0.041	1.80	1.02–3.15
rs1045642						
CC					1.00	
CT	0.843	0.439	3.165	0.057	2.30	0.98–5.43
TT	1.243	0.545	5.209	0.022	3.47	1.19–10.07
Constant	–2.666	1.881	2.010	0.156		

Asian, whereas in the Bochud et al. study the population was African Black. Previous studies reported the frequency of T as 16–27% in African populations, 48–57% in Caucasian populations, and 41–66% in Asian populations [22]. An obvious difference was observed between African and Asian populations, which may affect the study differences found in the association between rs1045642 and renal function. Second, there was a difference in inclusion of hypertensive patients. In the previous study, some patients were hypertension and some were not, whereas, our study population was strictly limited to hypertensive patients who were not receiving drug treatment. Third, the TT genotype mutations have been reported to decrease P-gp expression, indirectly reducing the exogenous and endogenous materials transport [23]. This aforementioned study also found that patients with TT genotype tended to have renal nephrotoxicity, which was related to the reduction of P-gp. The reduction of P-gp expression delayed the transport of exogenous and endogenous poison materials. These aforementioned reasons could explain why hypertensive patients with TT genotype more easily have renal function injury. The specific biological mechanism is not clear and further studies are required.

The process of inducing renal function injury through reducing P-gp expression was considered to be direct. Other indirect effects may contribute to renal function injury. ABCB1 gene can regulate blood pressure. In rats, the elevated blood pressure level induced through an injection of sodium has been shown to be suppressed by spironolactone injection [24,25]. This finding suggests that P-gp may be involved in the process of salt-sensitive hypertension. *In vitro*, P-gp modulator can inhibit the aldosterone secretion regulated by angiotensin II. It seems reasonable to suppose that variants of ABCB1 gene are associated with blood pressure in humans. Blood pressure itself could induce renal function injury. High blood pressure could lead to renal artery sclerosis, and a decline in eGFR.

There were some limitations to the present study. First, only three SNPs of ABCB1 gene were studied. Another SNP (C3435T) polymorphism in the promoter region is considered a common SNP of the ABCB1 gene. Therefore, additional studies are needed to explore the function of this and other SNPs. Second, some parameters collected were based on self-reported results, such as salt intake. This data could have had informational bias; it has been reported that the ABCB1 gene is a kind of salt-sensitive gene [26]. Third, our study population setting was limited to Chinese Han. People in different race groups have different genetic backgrounds. The effects of the ABCB1 gene could be different in other population groups. Finally, previous studies have reported other gene could be associated with renal function in hypertensive patients. Interactions can exist among these genes. However, to our knowledge, this has never been demonstrated.

Conclusions

In summary, our results suggested that the rs1045642-T allele of the ABCB1 gene may be associated with increased risk for renal function injury in hypertensive patients. More research is needed to explore the potential mechanisms and effects of interactions among the ABCB1 gene and others gene on renal function in patients with or without hypertension.

Acknowledgement

We thank all our colleagues working in the Third Affiliated Hospital of South Medical University.

Conflict of interest

All authors wish to declare that they have no conflict of interests in this study or its publication, financial or otherwise.

References:

- Bochud M, Bovet P, Burnier M, Eap CB: CYP3A5 and ABCB1 genes and hypertension. *Pharmacogenomics*, 2009; 10: 477–87
- Spezia L, Perandini S, Augelli R et al: Successful treatment of resistant hypertension by means of chronic renal artery occlusion revascularization in a fragile patient. *Pol J Radiol*, 2016; 81: 532–35
- Vallee JP: [Arterial hypertension and left ventricular hypertrophy: A 40-year follow-up in a Framingham cohort study]. *Presse Med*, 1999; 28: 1421–22 [in French]
- Gathier CS, Dankbaar JW, van der Jagt M et al: Effects of induced hypertension on cerebral perfusion in delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage: A randomized clinical trial. *Stroke*, 2015; 46: 3277–81
- Ruilope LM, Bakris GL: Renal function and target organ damage in hypertension. *Eur Heart J*, 2011; 32: 1599–604
- Hadjadj S, Fumeron F, Roussel R et al: Prognostic value of the insertion/deletion polymorphism of the ACE gene in type 2 diabetic subjects: Results from the Non-insulin-dependent Diabetes, Hypertension, Microalbuminuria or Proteinuria, Cardiovascular Events, and Ramipril (DIABHYCAR), Diabete de type 2, Nephropathie et Genetique (DIAB2NEPHROGENE), and Survie, Diabete de type 2 et Genetique (SURDIAGENE) studies. *Diabetes Care*, 2008; 31: 1847–52
- Chaves FJ, Pascual JM, Rovira E et al: Angiotensin II AT1 receptor gene polymorphism and microalbuminuria in essential hypertension. *Am J Hypertens*, 2001; 14: 364–70
- Kim SK, Chung JH, Kwon OY: Promoter polymorphism (-174, G/C) of interleukin-6 and arterial thromboembolic events: A meta-analysis. *Med Sci Monit*, 2016; 22: 4345–53
- Li L, Pan Y, Dai L et al: Association of genetic polymorphisms on vascular endothelial growth factor and its receptor genes with susceptibility to coronary heart disease. *Med Sci Monit*, 2016; 22: 31–40
- Dong J: [Epidemiology study on the renal function in hypertensive population of rural communities in North Jiangsu]. Nanjing Medical University, 2012 [in Chinese]
- Li L, Li L, Cui T, Zhao D: Association between Klotho gene polymorphism and left ventricular hypertrophy in hypertensive patients. *Chin J Gerontol*, 2013: 2501–3
- Uhr M, Holsboer F, Muller MB: Penetration of endogenous steroid hormones corticosterone, cortisol, aldosterone and progesterone into the brain is enhanced in mice deficient for both mdr1a and mdr1b P-glycoproteins. *J Neuroendocrinol*, 2002; 14: 753–59
- Zolk O, Jacobi J, Pahl A et al: MDR1 genotype-dependent regulation of the aldosterone system in humans. *Pharmacogenet Genomics*, 2007; 17: 137–44
- Jones DW, Hall JE: Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure and evidence from new hypertension trials. *Hypertension*, 2004; 43: 1–3
- Teramoto T, Sasaki J, Ishibashi S et al: Chronic kidney disease. *J Atheroscler Thromb*, 2014; 21: 173–74
- Zhou B: [Predictive values of body mass index and waist circumference to risk factors of related diseases in Chinese adult population]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 2002; 23: 5–10 [in Chinese]
- Li K, Yao C, Di X et al: Smoking and risk of all-cause deaths in younger and older adults: a population-based prospective cohort study among Beijing adults in China. *Medicine (Baltimore)*, 2016; 95: e2438
- Guo L, Deng J, He Y et al: Alcohol use and alcohol-related problems among adolescents in China: A large-scale cross-sectional study. *Medicine (Baltimore)*, 2016; 95: e4533
- Liu L, Chen L, Li Z et al: Association between gene polymorphisms of seven newly identified *loci* and type 2 diabetes and the correlate quantitative traits in Chinese Dong populations. *Iran J Public Health*, 2014; 43: 1345–55
- Kao YL, Donaghue K, Chan A et al: A novel polymorphism in the aldose reductase gene promoter region is strongly associated with diabetic retinopathy in adolescents with type 1 diabetes. *Diabetes*, 1999; 48: 1338–40
- Bochud M, Eap CB, Maillard M et al: Association of ABCB1 genetic variants with renal function in Africans and in Caucasians. *BMC Med Genomics*, 2008; 1: 21
- Bochud M, Bovet P, Burnier M, Eap CB: CYP3A5 and ABCB1 genes and hypertension. *Pharmacogenomics*, 2009; 10: 477–87
- Uhr M, Holsboer F, Muller MB: Penetration of endogenous steroid hormones corticosterone, cortisol, aldosterone and progesterone into the brain is enhanced in mice deficient for both mdr1a and mdr1b P-glycoproteins. *J Neuroendocrinol*, 2002; 14: 753–59
- Huang BS, Cheung WJ, Wang H et al: Activation of brain renin-angiotensin-aldosterone system by central sodium in Wistar rats. *Am J Physiol Heart Circ Physiol*, 2006; 291: H1109–17
- Huang BS, Leenen FH: Brain renin-angiotensin system and ouabain-induced sympathetic hyperactivity and hypertension in Wistar rats. *Hypertension*, 1999; 34: 107–12
- Bello-Reuss E, Ernest S, Holland OB, Hellmich MR: Role of multidrug resistance P-glycoprotein in the secretion of aldosterone by human adrenal NCI-H295 cells. *Am J Physiol Cell Physiol*, 2000; 278: C1256–65