

# Protective role of 17 $\beta$ -estradiol on tumor necrosis factor- $\alpha$ -induced apoptosis in human nucleus pulposus cells

HUAN LIU<sup>1\*</sup>, SI-DONG YANG<sup>1\*</sup>, YING XU<sup>2</sup>, SHENG-HUA NING<sup>1</sup>,  
TAO WANG<sup>1</sup>, DA-LONG YANG<sup>1</sup> and WEN-YUAN DING<sup>1</sup>

<sup>1</sup>Department of Spine Surgery, The Third Hospital of Hebei Medical University, Shijiazhuang, Hebei 050051; <sup>2</sup>Department of Cardiology, The Traditional Chinese Medicine Hospital of Hebei Medical University, Shijiazhuang, Hebei 050011, P.R. China

Received April 27, 2016; Accepted March 23, 2017

DOI: 10.3892/mmr.2017.6690

**Abstract.** The molecular mechanisms underlying protection and pathogenesis in spinal degenerative diseases remain unclear. Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) has been demonstrated to induce apoptosis of intervertebral disc (IVD) cells during IVD degeneration, and 17 $\beta$ -estradiol (17 $\beta$ -E2) has a protective effect against IVD cell apoptosis. However, the underlying molecular mechanism by which 17 $\beta$ -E2 protects nucleus pulposus (NP) cells remains to be investigated. The aim of the present study was to evaluate whether 17 $\beta$ -E2 modulates apoptosis of human NP cells induced by TNF- $\alpha$ . In addition, the concentration-response effect of 17 $\beta$ -E2 on human NP cells was investigated. Human NP cells were cultured in complete medium, which was replaced every three days until the culture was ~80% confluent. Cells were treated with 100 ng/ml TNF- $\alpha$  for 48 h, with or without pretreatment with various concentrations of 17 $\beta$ -E2, and ICI 182,780, for 30 min. Morphologic alterations characteristic of apoptosis were observed by inverted phase-contrast microscopy and Hoechst 33258 staining; the apoptosis rate was analyzed by flow cytometry. A Cell Counting kit-8 assay was used to assess cell proliferation. Furthermore, caspase-3 activity was determined and proteins associated with apoptosis were analyzed by western blotting. The level of apoptosis and caspase-3 activity in human NP cells increased, whereas proliferation and the expression of poly ADP-ribose polymerase decreased following TNF- $\alpha$  treatment. These effects of TNF- $\alpha$  were abolished by pretreatment with 17 $\beta$ -E2 in a concentration-dependent manner. The results of the present

study indicated that 17 $\beta$ -E2 serves a critical role in the survival of degenerative human NP cells.

## Introduction

The intervertebral disc (IVD) consists of three distinct sections: The outer annulus fibrosus (AF) which contains abundant type I collagens; the inner nucleus pulposus (NP) which is a gelatinous substance containing type II collagen and proteoglycans; and the cartilage end plates that connect the adjacent vertebral bodies (1-3). With age, morphological and molecular alterations in the IVD may induce progressive degeneration and pathologic alterations of this tissue (4). IVD degeneration may eventually result in conditions including disc herniation, facet joint osteoarthritis, spinal canal stenosis, segmental instability and spondylolisthesis. The mechanisms underlying IVD degeneration are considered to be associated with decreased cell viability and adhesion, reduced diffusion of nutrients and proteoglycan synthesis (5,6). Accelerated aging of the IVD due to abnormal apoptosis of IVD cells is a primary cytological process (7-9), and apoptosis of IVD cells serves a key role in the process of IVD degeneration.

Tumor necrosis factor (TNF)- $\alpha$  is an important member of the TNF superfamily, and is involved in immunity, cellular remodeling, apoptosis and cell survival (10). It acts primarily via TNF receptor-1 to induce apoptosis by activating caspases through mitochondria-dependent and -independent pathways (11). In addition to secretion by infiltrating immune cells, resident NP and AF cells produce high levels of TNF- $\alpha$  and interleukin (IL)-1 $\beta$  to promote inflammation (12-14). TNF- $\alpha$  has been confirmed to have a central role in degenerative osteoarthritis (15). Additionally, it has been reported that TNF- $\alpha$  may induce apoptosis of NP cells *in vitro* (16). Furthermore, TNF- $\alpha$  released during the inflammatory process induced by the herniated IVD serves a fundamental role in the development of mechanical and thermal hyperalgesia (16-18). The inhibition of TNF- $\alpha$ -induced abnormal apoptosis of human NP cells may delay the degeneration of the IVD, preventing degenerative scoliosis and other spinal degenerative diseases.

17 $\beta$ -estradiol (17 $\beta$ -E2) has been extensively investigated due to its anti-apoptotic activity (19-22). Bozzo *et al* (23) reported that 17 $\beta$ -E2 protects nerve cells from  $\beta$ -amyloid peptide-induced apoptosis, and that 17 $\beta$ -E2 exerts anti-apoptotic

---

*Correspondence to:* Dr Wen-Yuan Ding, Department of Spine Surgery, The Third Hospital of Hebei Medical University, 139 Ziqiang Road, Shijiazhuang, Hebei 050051, P.R. China  
E-mail: dingwenyuan2016@163.com

\*Contributed equally

**Key words:** 17 $\beta$ -estradiol, tumor necrosis factor- $\alpha$ , nucleus pulposus, apoptosis

effects by upregulating the expression of essential cell membrane components including  $\alpha$ 1 $\beta$ 1 integrin, and affecting cell cycle progression. In addition, 17 $\beta$ -E2 has been reported to inhibit caspase-3/9 activity and increase B-cell lymphoma 2, cyclin D1 and survivin expression (24-26). Wang *et al.* (6) and Yang *et al.* (22) demonstrated that 17 $\beta$ -E2 protects rat IVD cells from apoptosis induced by IL-1 $\beta$  and levofloxacin. However, whether 17 $\beta$ -E2 inhibits TNF- $\alpha$ -induced apoptosis in human NP cells, and the concentration-response effect of 17 $\beta$ -E2 on TNF- $\alpha$ -induced human NP cell apoptosis, remains unclear.

Therefore, the aim of the present study was to investigate whether 17 $\beta$ -E2 modulates apoptosis induced by TNF- $\alpha$  in human NP cells, the concentration-response effect and whether 17 $\beta$ -E2 exerts protective effects via the caspase-3/poly (ADP-ribose) polymerase (PARP) pathway.

## Materials and methods

**Reagents.** Human NP cells, NP Cell Growth Supplement and NP cell medium (NPCM) were purchased from ScienCell Research Laboratories, Inc. (Carlsbad, CA, USA), and Dulbecco's modified Eagle's medium (DMEM)/F-12 and fetal bovine serum (FBS) were obtained from HyClone; Thermo Fisher Scientific, Inc. (Waltham, MA, USA). Trypsin, Cell Counting kit-8 (CCK-8) and 17 $\beta$ -E2 were purchased from Sigma-Aldrich; Merck KGaA (Darmstadt, Germany). PBS (phosphate buffered saline) was obtained from Gibco; Thermo Fisher Scientific, Inc. The Annexin V-Fluorescein Isothiocyanate (FITC)/Propidium Iodide (PI) kit was purchased from BD Biosciences (Franklin Lakes, NJ, USA), and the caspase-3 activity kit, Hoechst staining kit and cell lysis buffer for western and immunoprecipitation were obtained from Beyotime Institute of Biotechnology (Shanghai, China). Z-DEVD-FMK was purchased from MedChem Express (Monmouth Junction, NJ, USA), ICI 182,780 was obtained from Sigma-Aldrich, Merck KGaA; and TNF- $\alpha$  was purchased from PeproTech, Inc. (Rocky Hill, NJ, USA).

**Initiation of human NP cell culture.** The purchased cryopreserved human NP cells were originally isolated from the NP of human IVD. NPCM (500 ml) was supplemented with 10 ml FBS, 5 ml NP Cell Growth Supplement and 5 ml penicillin/streptomycin solution (P/S) to make complete human NP cell medium prior to cell recovery. Subsequently, cells were thawed in their original vials in a 37°C water bath and placed in a 50-ml culture flask containing 15 ml complete HNPC as soon as possible and with minimal handling. Cells were cultured in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C. The culture medium was replaced with fresh the next day to remove residual dimethyl sulfoxide and non-adherent cells. The medium was replaced every three days thereafter. Once the culture reached 70% confluence, the medium was replaced every other day until the culture was ~90% confluent.

**Subculture and drug treatment.** When the culture reached  $\geq$ 90% confluence, the cells were digested, plated into appropriate culture plates and cultured as described in the previous section. When the culture reached 80-90% confluence, cells were washed twice with PBS and the medium was replaced with DMEM/F-12 without phenol red, FBS and P/S. Cells

were divided into 7 groups according to treatment, the conditions of which are listed in Table I. All groups were incubated for 48 h in serum-free medium without phenol red.

**Morphological observations.** Human NP cells were plated in 6-well plates at a density of 2x10<sup>5</sup> cells/well in NPCM. A total of 48 h after treatment, the apoptotic morphological alterations in the human NP cells were observed under an inverted phase-contrast microscope (Olympus Corporation, Tokyo, Japan) and imaged using a digital camera (Nikon Corporation, Tokyo, Japan). Cells were subsequently washed twice with PBS and stained with Hoechst 33258 staining solution according to the manufacturer's protocol. Stained nuclei were detected under a fluorescence microscope (Olympus Corporation) with an excitation wavelength of 350 nm and an emission wavelength of 460 nm.

**Proliferation assay.** The effects of 17 $\beta$ -E2 on cell viability and proliferation were evaluated using CCK-8. Human NP cells were seeded into 96-well plates at a density of 2x10<sup>4</sup> cells/well (100  $\mu$ l/well). Following treatment for 48 h, 10  $\mu$ l CCK-8 solution was added to each well of the plate, and the plates were incubated for an additional 3 h in a 5% CO<sub>2</sub> incubator at 37°C. The proliferation of cells was measured using a microplate reader at a wavelength of 450 nm.

**Caspase-3 activity assay.** Caspase-3 activity was measured using a Caspase-3 Activity kit, which measures the conversion by caspase-3 of N-acetyl-Asp-Glu-Val-Asp-p-nitroanilide into a yellow formazan product, p-nitroaniline. Human NP cells were cultured in 6-well plates as aforementioned, and the assay was performed according to the manufacturer's protocol. Briefly, the treated NP cells were washed twice with cold PBS and lysed with lysis buffer (100  $\mu$ l/2x10<sup>6</sup> cells) for 15 min on ice. Following incubation of 10  $\mu$ l cell lysates, 80  $\mu$ l reaction buffer and 10  $\mu$ l caspase-3 substrate (2 mM) in 96-well microtiter plates at 37°C for 2 h, caspase-3 activity was quantified at a wavelength of 405 nm using a microplate spectrophotometer (BioTek Instruments, Inc., Winooski, VT, USA). Caspase-3 activity is expressed as the fold change in enzyme activity over control.

**Flow cytometric analysis of apoptosis.** Human NP cells were seeded into 6-well plates at a density of 2x10<sup>5</sup> cells/well and cultured as aforementioned. Following 48 h treatment, adherent cells and cells in suspension were collected and washed twice with cold PBS. Apoptotic cells were detected by staining with annexin V-FITC and PI according to the manufacturer's protocol. Briefly, cells were resuspended in 100  $\mu$ l 1X binding buffer at a concentration of 1x10<sup>6</sup> cells/ml, and 100  $\mu$ l (1x10<sup>5</sup> cells) were transferred to 5 ml culture tubes. PI (5  $\mu$ l) and 5  $\mu$ l annexin V-FITC was added to each tube and incubated for 15 min at room temperature in the dark, followed by addition of 400 ml 1X binding buffer. Samples were analyzed on a flow cytometer (BD Biosciences) with CellQuest software (version 5.2, BD Biosciences).

**Western blot analysis.** Protein expression levels of PARP and caspase-3 were determined by western blot analysis. Cells were incubated in 100  $\mu$ l lysis buffer containing 1% protease

Table I. Experimental groups.

Group	Treatment
Control	Ethanol (<0.1%)
TNF	100 ng/ml TNF for 48 h
TNF + 0.1 E2	100 ng/ml TNF for 48 h, with pretreatment of 0.1 $\mu$ mol/l E2 for 30 min
TNF + 1 E2	100 ng/ml TNF for 48 h, with pretreatment of 1 $\mu$ mol/l E2 for 30 min
TNF + 10 E2	100 ng/ml TNF for 48 h, with pretreatment of 10 $\mu$ mol/l E2 for 30 min
TNF + 10 E2 + ICI	100 ng/ml TNF for 48 h, with pretreatment of 10 $\mu$ mol/l E2 and 10 $\mu$ mol/l ICI for 30 min
TNF + FMK	100 ng/ml TNF for 48 h, with pretreatment of 20 $\mu$ mol/l FMK for 2 h

TNF, tumor necrosis factor- $\alpha$ ; E2, 17 $\beta$ -estradiol; ICI, ICI 182,780; FMK, Z-DEVD-FMK.

inhibitors. The cell lysate was collected by centrifugation at 4°C for 10 min at 16,000 x g. The protein concentrations were determined using a Bicinchoninic Acid Protein assay kit (Beyotime Institute of Biotechnology). Proteins (50  $\mu$ g) from each sample were separated on 12% SDS-polyacrylamide gels according to molecular weight and transferred onto polyvinylidene difluoride membranes (Bio-Rad Laboratories, Inc., Hercules, CA, USA). The membranes were blocked with 5% non-fat milk powder in TBS for 1 h at room temperature and incubated with primary antibodies against caspase-3 p17 (#9664; rabbit; 1:1,000), PARP (#9532; rabbit; 1:1,000) (both from Cell Signaling Technology, Inc., Danvers, MA, USA), and GAPDH (10494-1-AP; rabbit; 1:5,000; Wuhan Sanying Biotechnology, Wuhan, China) diluted in 5% non-fat milk powder in PBS containing Tween 20 (PBST) at 4°C overnight. Membranes were washed 3 times for 15 min in PBST and incubated with a peroxidase labeled anti-rabbit IgG (H+L) antibody (074-1506; goat; 1:1,000; KPL, Inc., Gaithersburg, MD, USA) with agitation for 1 h at room temperature. Following washing 3 times for 15 min with PBST, protein bands were detected using Enhanced Chemiluminescence reagent (Amercontrol Biosciences, San Francisco, CA, USA) and quantified with Quantity One imaging software version 4.6.2 (Bio-Rad Laboratories, Inc.).

**Statistical analysis.** Statistical analyses were performed using SPSS software version 16.0 (SPSS, Inc., Chicago, IL, USA). Data are presented as the mean  $\pm$  standard deviation. Groups were compared by one-way analysis of variance followed by pairwise comparison using the Student-Newman-Keuls post hoc test. All statistical tests were two-sided.  $P < 0.01$  was considered to indicate a statistically significant difference.

## Results

**Morphology of apoptotic human NP cells.** The morphological characteristics of human NP cells were observed using inverted phase-contrast microscopy (Fig. 1). Cell nuclei were stained using Hoechst 33258 and detected under a fluorescence microscope with an excitation wavelength of 350 nm and an emission wavelength of 460 nm (Fig. 2). Control untreated cells were well-adhered, exhibiting typical human NP cell morphology. By contrast, treatment with TNF- $\alpha$  induced apoptosis, as demonstrated by detachment from the plate,

cell shrinkage, and nuclei that were compact, condensed and brightly stained with Hoechst 33258. Following pretreatment with various concentrations of 17 $\beta$ -E2, the number of apoptotic cells markedly decreased compared with those treated with TNF- $\alpha$  only. The effect of 17 $\beta$ -E2 was blocked by the estrogen receptor (ER) antagonist ICI 182,780.

**17 $\beta$ -E2 promotes proliferation of human NP cells damaged by TNF- $\alpha$ .** The effect of 17 $\beta$ -E2 on proliferation of human NP cells was detected by CCK-8 assay. As presented in Fig. 3, TNF- $\alpha$  inhibited the proliferation of human NP cells ( $P < 0.001$ ). This effect was abrogated when the cells were pretreated with 17 $\beta$ -E2 ( $P < 0.001$ ). The effect of 17 $\beta$ -E2 was blocked by ICI 182,780 ( $P < 0.001$ ). 17 $\beta$ -E2 inhibited the effect of TNF- $\alpha$  in a concentration-dependent manner.

**17 $\beta$ -E2 reduces caspase-3 activity.** The effect of 17 $\beta$ -E2 on caspase-3 activity in human NP cells was measured using a caspase-3 activity assay. As presented in Fig. 4, the caspase-3 activity of cells treated with TNF- $\alpha$  alone was 5-fold greater compared with the control untreated group ( $P < 0.001$ ). However, cells pretreated with 17 $\beta$ -E2 demonstrated reduced caspase-3 activity compared with cells treated with TNF- $\alpha$  only; the effect of 17 $\beta$ -E2 was concentration-dependent. The effect of pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 for 30 min was similar to pretreatment with 20  $\mu$ mol/l Z-DEVD-FMK (a specific, irreversible caspase-3 inhibitor) for 2 h ( $P > 0.05$ ). Addition of the ER antagonist ICI 182,780 abolished the protective effects of 17 $\beta$ -E2 ( $P < 0.001$ ).

**17 $\beta$ -E2 reduces TNF- $\alpha$ -induced human NP cell apoptosis.** The effect of 17 $\beta$ -E2 on apoptosis of human NP cells was examined by flow cytometry following annexin V-FITC and PI double staining. The apoptotic rate is presented as apoptotic cells in the experimental group/apoptotic cells in the control group. As presented in Fig. 5, an increase in the apoptotic rate was observed following treatment with TNF- $\alpha$ . Pretreatment with 17 $\beta$ -E2 significantly decreased the rate of apoptosis in a concentration-dependent manner compared with cells treated with TNF- $\alpha$  only ( $P < 0.001$ ). Addition of the ER antagonist ICI 182,780 blocked the effect of 17 $\beta$ -E2 on TNF- $\alpha$  treatment, further demonstrating the potential contribution of 17 $\beta$ -E2 to TNF- $\alpha$ -induced human NP cell apoptosis.




Figure 1. Morphological observations by phase-contrast microscopy. (A) Control untreated human NP cells. Human NP cells treated with (B) 100 ng/ml TNF- $\alpha$ , (C) 100 ng/ml TNF- $\alpha$  following pretreatment with 0.1  $\mu$ mol/l 17 $\beta$ -E2, (D) 100 ng/ml TNF- $\alpha$  following pretreatment with 1  $\mu$ mol/l 17 $\beta$ -E2, (E) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 and (F) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 and 10  $\mu$ mol/l ICI 182,780. Apoptotic cells were characterized by shedding of smaller fragments from the cells, cell shrinkage and condensed nuclei. Original magnification, x200; scale bar, 200 nm. NP, nucleus pulposus; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; 17 $\beta$ -E2, 17 $\beta$ -estradiol.




Figure 2. Hoechst 33258 staining. (A) Control untreated human NP cells. Human NP cells treated with (B) 100 ng/ml TNF- $\alpha$ , (C) 100 ng/ml TNF- $\alpha$  following pretreatment with 0.1  $\mu$ mol/l 17 $\beta$ -E2, (D) 100 ng/ml TNF- $\alpha$  following pretreatment with 1  $\mu$ mol/l 17 $\beta$ -E2, (E) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 and (F) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 and 10  $\mu$ mol/l ICI 182,780. The nuclei of living cells were a homogeneous blue, whereas apoptotic nuclei were compact, condensed and brightly stained with Hoechst 33258. Original magnification, x200. NP, nucleus pulposus; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; 17 $\beta$ -E2, 17 $\beta$ -estradiol.

**Western blot analysis.** Human NP cells were divided into the following four groups: Control untreated, 100 ng/ml TNF- $\alpha$ , 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 for 30 min and 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l 17 $\beta$ -E2 and 10  $\mu$ mol/l ICI 182,780 for 30 min. Following treatment, protein expression levels of

PARP, caspase-3p17 and GAPDH were determined by western blot analysis. Caspase-3p17 protein expression levels were increased in human NP cells following treatment with TNF- $\alpha$  compared with the control cells ( $P < 0.001$ ); this increased was inhibited by pretreatment with 17 $\beta$ -E2 ( $P < 0.001$ ). By contrast, PARP protein expression levels were decreased in




Figure 3. Proliferation of human NP cells, as assessed by CCK-8 assay. The CCK-8 assay reflects the activity of mitochondrial dehydrogenase. Human NP cells were untreated (control), treated with 100 ng/ml TNF- $\alpha$ , or treated with 100 ng/ml TNF- $\alpha$  following pretreatment with various concentrations of 17 $\beta$ -E2 with or without ICI 182,780. Data are expressed as the mean  $\pm$  standard deviation from three independent experiments. \*\*P<0.001; #P>0.05. NP, nucleus pulposus; CCK-8, Cell Counting kit-8; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; 17 $\beta$ -E2, 17 $\beta$ -estradiol.




Figure 4. Caspase-3 activity in human NP cells was determined using a caspase-3 activity assay. Human NP cells were untreated (control), treated with 100 ng/ml TNF- $\alpha$ , or treated with 100 ng/ml TNF- $\alpha$  following pretreatment with various concentrations of E2 with or without ICI. An additional control group was pretreated with the caspase-3 inhibitor FMK prior to treatment with TNF- $\alpha$ . Data are expressed as the mean  $\pm$  standard deviation from three independent experiments. \*\*P<0.001; #P>0.05. NP, nucleus pulposus; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; E2, 17 $\beta$ -estradiol; ICI, ICI 182,780; FMK, Z-DEVD-FMK.

cells subjected to TNF- $\alpha$  treatment (P<0.001); this decrease was abrogated by pretreatment with 17 $\beta$ -E2 (P<0.001). These effects of 17 $\beta$ -E2 were blocked by ICI 182,780 treatment (P<0.001; Fig. 6).

## Discussion

One of the primary causes of spinal degeneration is IVD degeneration. Apoptosis of NP cells and the loss of ECM are the central features of the aged and degenerated IVD (7,8,27,28). Gruber *et al* (29) reported that the ER gene is expressed in human IVD cells *in vivo* and *in vitro*, and further demonstrated

that 17 $\beta$ -E2 significantly increased proliferation of annulus cells. Previous studies have revealed that TNF- $\alpha$  may induce NP cell apoptosis (16,30), and various *in vitro* studies have demonstrated that 17 $\beta$ -E2 protects against apoptosis in IVD cells (6,22,24). Therefore, the hypothesis of the present study was that 17 $\beta$ -E2 would attenuate TNF- $\alpha$ -induced apoptosis in human NP cells.

The results of the present study suggested that 17 $\beta$ -E2 abolished TNF- $\alpha$ -induced apoptosis in human NP cells by increasing proliferation and reducing apoptosis. A CCK-8 assay revealed that 17 $\beta$ -E2 enhanced the proliferative capacity of human NP cells. Morphological observations and flow




Figure 5. Apoptosis of human NP cells. (A) Annexin V/propidium iodide double staining of human NP cells; (a) Control untreated human NP cells. Human NP cells treated with (b) 100 ng/ml TNF- $\alpha$ , (c) 100 ng/ml TNF- $\alpha$  following pretreatment with 0.1  $\mu$ mol/l E2, (d) 100 ng/ml TNF- $\alpha$  following pretreatment with 1  $\mu$ mol/l E2, (e) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l E2 and (f) 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l E2 and 10  $\mu$ mol/l ICI. The cells in lower right quadrant (FITC+/PI-) were considered to be apoptotic. (B) Data are expressed as the mean  $\pm$  standard deviation from three independent experiments. <sup>\*\*</sup>P<0.001; <sup>#</sup>P>0.05. NP, nucleus pulposus; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; E2, 17 $\beta$ -estradiol; ICI, ICI 182,780.




Figure 6. Protein expression levels of caspase-3p17 and PARP in human NP cells. Human NP cells were untreated (control), or treated with 100 ng/ml TNF- $\alpha$ , 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l E2 for 30 min, or 100 ng/ml TNF- $\alpha$  following pretreatment with 10  $\mu$ mol/l E2 and 10  $\mu$ mol/l ICI for 30 min. Data are expressed as the mean  $\pm$  standard deviation from three independent experiments. <sup>\*\*</sup>P<0.001. PARP, poly (ADP-ribose) polymerase; NP, nucleus pulposus; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; E2, 17 $\beta$ -estradiol; ICI, ICI 182,780.

cytometric analysis demonstrated that TNF- $\alpha$ -induced apoptosis in human NP cells was prevented by 17 $\beta$ -E2. Western blot analysis was used to detect alterations in the protein expression levels of PARP and caspase-3 in human NP cells. Caspase-3 is a crucial executioner protease, which activates a caspase-activated DNase and serves an important role in apoptosis (27,31-33). Previous studies have reported that caspase-3 may be a therapeutic target for regulation of IVD degeneration (34). Pro-caspase-3 may be sheared into two fragments, p17 and p10, upon activation. The protein expression levels of caspase-3 p17 are positively correlated with caspase-3 activity. PARP is a substrate of caspase-3, which may trigger inflammation and cell death (35). Shakibaei *et al* (36) demonstrated that the cleavage of PARP was activated *in vitro* in apoptotic human articular chondrocytes.

The present study investigated whether 17 $\beta$ -E2 inhibited human NP cell apoptosis induced by TNF- $\alpha$ . Estrogen has positive effects on bone structure, chondrocyte and nerve cells. Various recent studies have revealed the anti-apoptotic effect of 17 $\beta$ -E2 in rat IVD cells (6,22,37). However, whether 17 $\beta$ -E2 may prevent TNF- $\alpha$ -induced apoptosis in human NP cells remains to be investigated. In the present study, human NP cells were used to provide an environment approaching the physiological state for the *in vitro* test of the effect of 17 $\beta$ -E2 on human NP. Phenol red may increase cell proliferation (38); therefore, medium without phenol red was used in the present study.

IVD degeneration is demonstrated by alterations in cell proliferation and cell matrix metabolism, which includes inhibition of nuclear proteoglycan synthesis and enhanced matrix degradation. IVD cells have been reported to produce inflammatory cytokines, including IL-1 $\beta$ , IL-6, TNF- $\alpha$ , matrix metalloproteinases (MMPs), nitric oxide and prostaglandin E2 (39). As an important inflammatory molecule, TNF- $\alpha$  may induce the cellular and matrix alterations of IVD degeneration (30), and activate crosstalk between signaling pathways including nuclear factor- $\kappa$ B and phosphoinositide 3-kinase (PI3K)/protein kinase B (Akt) (40,41).

The mechanism underlying the protection of 17 $\beta$ -E2 against TNF- $\alpha$ -induced apoptosis in human NP cells is complex. Various potential mechanisms may be involved in the anti-apoptotic effect of 17 $\beta$ -E2 in human NP cells. Firstly, integrin production was induced in NP cells by 17 $\beta$ -E2. It has been reported that 17 $\beta$ -E2 may protect rat IVD cells by upregulating integrin  $\alpha$  and  $\beta$  (22,42). Secondly, 17 $\beta$ -E2 may inhibit human NP cell apoptosis via the mitochondrial pathway. Yang *et al* (43) demonstrated that 17 $\beta$ -E2 protects against apoptosis by downregulating MMP-3 and -13 via a mitochondrial pathway in rat NP cells. Finally, the PI3K/Akt signaling pathway may be involved in the effect of 17 $\beta$ -E2. A further study by Yang *et al* (37) revealed that 17 $\beta$ -E2 protects against apoptosis via the activation of PI3K/Akt signaling pathway. This signaling pathway is associated with cell migration and invasion (44,45).

The present study investigated the effect of 17 $\beta$ -E2 on human NP cells; however, the findings of the present study are limited. The results of the present study were used as preliminary data for our subsequent study, which demonstrated that 17 $\beta$ -E2 inhibits TNF- $\alpha$ -induced apoptosis in human NP cells via the PI3K/Akt pathway (46). Further signaling mechanisms underlying the anti-apoptotic effects of 17 $\beta$ -E2 in human NP

cells will require further investigation. In addition, 17 $\beta$ -E2 is a systemic hormone, and therefore further studies are required to determine the systemic effect of 17 $\beta$ -E2 in humans, to assess the suitability of 17 $\beta$ -E2 as a therapeutic strategy for the clinical treatment of IVD degeneration.

In conclusion, the results of the present study revealed that 17 $\beta$ -E2 protected against TNF- $\alpha$ -induced apoptosis in human NP cells in a concentration-dependent manner, and that the activation of caspase-3 and inhibition of PARP may participate in this process. The ability of 17 $\beta$ -E2 to protect human NP cells from apoptosis was blocked by ICI 172,780, an ER antagonist. The results of the present study may present a novel route for the prevention and therapy of IVD degeneration.

### Acknowledgements

The present study was supported by the Natural Science Foundation of Hebei Province (grant no. H2014206075).

### References

- Zhao CQ, Jiang LS and Dai LY: Programmed cell death in intervertebral disc degeneration. *Apoptosis* 11: 2079-2088, 2006.
- Buckwalter JA: Aging and degeneration human intervertebral disc. *Spine (Phila Pa 1976)* 20: 1307-1314, 1995.
- Ferguson SJ, Ito K and Nolte LP: Fluid flow and convection transport of solutes within the intervertebral disc. *J Biomech* 37: 213-221, 2004.
- Anderson DG, Li X and Balian G: A fibronectin fragment alters the metabolism by rabbit intervertebral disc cells in vitro. *Spine (Phila Pa 1976)* 30: 1242-1246, 2005.
- Podichetty VK: The aging spine: The role of inflammatory mediators in intervertebral disc degeneration. *Cell Mol Biol (Noisy-le-grand)* 53: 4-18, 2007.
- Wang H, Ding W, Yang D, Gu T, Yang S and Bai Z: Different concentrations of 17 $\beta$ -estradiol modulates apoptosis induced by interleukin-1 $\beta$  in rat annulus fibrosus cells. *Mol Med Rep* 10: 2745-2751, 2014.
- Zhou GQ, Yang F, Leung VVL and Cheung KMC: Molecular and cellular biology of the intervertebral disc and the use of animal models. *Current Orthopaedics* 22: 267-273, 2008.
- Le Maitre CL, Freemont AJ and Hoyland JA: Accelerated cellular senescence in degenerate intervertebral discs: A possible role in the pathogenesis of intervertebral disc degeneration. *Arthritis Res Ther* 9: R45, 2007.
- Boos N, Weissbach S, Rohrbach H, Weiler C, Spratt KF and Nerlich AG: Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science. *Spine (Phila Pa 1976)* 27: 2631-2644, 2002.
- Szlosarek P, Charles KA and Balkwill FR: Tumour necrosis factor-alpha as a tumour promoter. *Eur J Cancer* 42: 745-750, 2006.
- Bharadwaj U, Marin-Muller C, Li M, Chen C and Yao Q: Mesothelin confers pancreatic cancer cell resistance to TNF- $\alpha$ -induced apoptosis through Akt/PI3K/NF- $\kappa$ B activation and IL-6/Mcl-1 overexpression. *Mol Cancer* 10: 106, 2011.
- Le Maitre CL, Freemont AJ and Hoyland JA: The role of interleukin-1 in the pathogenesis of human intervertebral disc degeneration. *Arthritis Res Ther* 7: R732-R745, 2005.
- Wang J, Markova D, Anderson DG, Zheng Z, Shapiro IM and Risbud MV: TNF- $\alpha$  and IL-1 $\beta$  promote a disintegrin-like and metalloprotease with thrombospondin type I motif-5-mediated aggrecan degradation through syndecan-4 in intervertebral disc. *J Biol Chem* 286: 39738-39749, 2011.
- Lee S, Moon CS, Sul D, Lee J, Bae M, Hong Y, Lee M, Choi S, Derby R, Kim BJ, *et al*: Comparison of growth factor and cytokine expression in patients with degenerated disc disease and herniated nucleus pulposus. *Clin Biochem* 42: 1504-1511, 2009.
- Mourão AF, Caetano-Lopes J, Costa P, Canhão H, Santos MJ, Pinto P, Brito I, Nicola P, Cavaleiro J, Teles J, *et al*: Tumor necrosis factor-alpha-308 genotypes influence inflammatory activity and TNF-alpha serum concentrations in children with juvenile idiopathic arthritis. *J Rheumatol* 36: 837-842, 2009.

16. Zhang CC, Cui GP, Hu JG, Xiao YZ, Zhou XS, Shao C, Lin Q and Zhou JS: Effects of adenoviral vector expressing hIGF-1 on apoptosis in nucleus pulposus cells in vitro. *Int J Mol Med* 33: 401-405, 2014.
17. de Souza Grava AL, Ferrari LF and Defino HL: Cytokine inhibition and time-related influence of inflammatory stimuli on the hyperalgesia induced by the nucleus pulposus. *Eur Spine J* 21: 537-545, 2012.
18. Le Maitre CL, Hoyland JA and Freemont AJ: Catabolic cytokine expression in degenerate and herniated human intervertebral discs: IL-1beta and TNFalpha expression profile. *Arthritis Res Ther* 9: R77, 2007.
19. Ackermann S, Hiller S, Osswald H, Lösle M, Grenz A and Hambrock A: 17beta-Estradiol modulates apoptosis in pancreatic beta-cells by specific involvement of the sulfonyleurea receptor (SUR) isoform SUR1. *J Biol Chem* 284: 4905-4913, 2009.
20. Yamaguchi A, Nozawa K, Fujishiro M, Kawasaki M, Takamori K, Ogawa H, Sekigawa I and Takasaki Y: Estrogen inhibits apoptosis and promotes CC motif chemokine ligand 13 expression on synovial fibroblasts in rheumatoid arthritis. *Immunopharmacol Immunotoxicol* 34: 852-857, 2012.
21. Hattori Y, Kojima T, Kato D, Matsubara H, Takigawa M and Ishiguro N: A selective estrogen receptor modulator inhibits tumor necrosis factor- $\alpha$ -induced apoptosis through the ERK1/2 signaling pathway in human chondrocytes. *Biochem Biophys Res Commun* 421: 418-424, 2012.
22. Yang SD, Ma L, Gu TX, Ding WY, Zhang F, Shen Y, Zhang YZ, Yang DL, Zhang D, Sun YP and Song YL: 17 $\beta$ -Estradiol protects against apoptosis induced by levofloxacin in rat nucleus pulposus cells by upregulating integrin  $\alpha$ 2 $\beta$ 1. *Apoptosis* 19: 789-800, 2014.
23. Bozzo C, Graziola F, Chiochetti A and Canonico PL: Estrogen and beta-amyloid toxicity: Role of integrin and PI3-K. *Mol Cell Neurosci* 45: 85-91, 2010.
24. Nilsen J and Diaz Brinton R: Mechanism of estrogen-mediated neuroprotection: Regulation of mitochondrial calcium and Bcl-2 expression. *Proc Natl Acad Sci USA* 100: 2842-2847, 2003.
25. Yu X, Zhang X, Dhakal IB, Beggs M, Kadlubar S and Luo D: Induction of cell proliferation and survival genes by estradiol-repressed microRNAs in breast cancer cells. *BMC Cancer* 12: 29, 2012.
26. Ge C, Yu M and Zhang C: G Protein-Coupled Receptor 30 mediates estrogen-induced proliferation of primordial germ cells via EGFR/Akt/ $\beta$ -catenin signaling pathway. *Endocrinology* 153: 3504-3516, 2012.
27. Ding F, Shao ZW, Yang SH, Wu Q, Gao F and Xiong LM: Role of mitochondrial pathway in compression-induced apoptosis of nucleus pulposus cells. *Apoptosis* 17: 579-90, 2012.
28. Nardinocchi L, Puca R, Sacchi A and D'Orazi G: Inhibition of HIF-1alpha activity by homeodomain-interacting protein kinase-2 correlates with sensitization of chemoresistant cells to undergo apoptosis. *Mol Cancer* 8: 1, 2009.
29. Gruber HE, Yamaguchi D, Ingram J, Leslie K, Huang W, Miller TA and Hanley EN Jr: Expression and localization of estrogen receptor-beta in annulus cells of the human intervertebral disc and the mitogenic effect of 17-beta-estradiol in vitro. *BMC Musculoskelet Disord* 3: 4, 2002.
30. Weber A, Wandinger KP, Mueller W, Aktas O, Wengert O, Grundström E, Ehrlich S, Windemuth C, Kuhlmann T, Wienker T, *et al*: Identification and functional characterization of a highly polymorphic region in the human TRAIL promoter in multiple sclerosis. *J Neuroimmunol* 149: 195-201, 2004.
31. Singh M, Sharma H and Singh N: Hydrogen peroxide induces apoptosis in HeLa cells through mitochondrial pathway. *Mitochondrion* 7: 367-373, 2007.
32. He B, Tao H, Liu S and Wei A: Protective effect of carboxymethylated chitosan on hydrogen peroxide-induced apoptosis in nucleus pulposus cells. *Mol Med Rep* 11: 1629-1638, 2015.
33. Li Z, Jo J, Jia JM, Lo SC, Whitcomb DJ, Jiao S, Cho K and Sheng M: Caspase-3 activation via mitochondria is required for long-term depression and AMPA receptor internalization. *Cell* 141: 859-871, 2010.
34. Sudo H and Minami A: Caspase 3 as a therapeutic target for regulation of intervertebral disc degeneration in rabbits. *Arthritis Rheum* 63: 1648-1657, 2011.
35. Zhao H, Yenari MA, Cheng D, Sapolsky RM and Steinberg GK: Bcl-2 overexpression protects against neuron loss within the ischemic margin following experimental stroke and inhibits cytochrome c translocation and caspase-3 activity. *J Neurochem* 85: 1026-1036, 2003.
36. Shakibaei M, John T, Seifarth C and Mobasher A: Resveratrol inhibits IL-1 beta-induced stimulation of caspase-3 and cleavage of PARP in human articular chondrocytes in vitro. *Ann N Y Acad Sci* 1095: 554-563, 2007.
37. Yang SD, Ma L, Yang DL and Ding WY: Combined effect of 17 $\beta$ -estradiol and resveratrol against apoptosis induced by interleukin-1 $\beta$  in rat nucleus pulposus cells via PI3K/Akt/caspase-3 pathway. *PeerJ* 4: e1640, 2016.
38. Wesierska-Gadek J, Schreiner T, Gueorguieva M and Ranftler C: Phenol red reduces ROSC mediated cell cycle arrest and apoptosis in human MCF-7 cells. *J Cell Biochem* 98: 1367-1379, 2006.
39. Weiler C, Nerlich AG, Bachmeier BE and Boos N: Expression and distribution of tumor necrosis factor alpha in human lumbar intervertebral discs: A study in surgical specimen and autopsy controls. *Spine (Phila Pa 1976)* 30: 44-54, 2005.
40. Wu Y and Zhou BP: TNF-alpha/NF-kappaB/Snail pathway in cancer cell migration and invasion. *Br J Cancer* 102: 639-644, 2010.
41. Sandra F, Matsuki NA, Takeuchi H, Ikebe T, Kanematsu T, Ohishi M and Hirata M: TNF inhibited the apoptosis by activation of Akt serine/threonine kinase in the human head and neck squamous cell carcinoma. *Cell Signal* 14: 771-778, 2002.
42. Zhao CM, Chen Q, Zhang WJ, Huang AB, Zhang W, Yang HL and Zhang ZM: 17 $\beta$ -estradiol protects rat annulus fibrosus cells against apoptosis via  $\alpha$ 1 integrin-mediated adhesion to type I collagen: An in-vitro study. *Med Sci Monit* 22: 1375-1383, 2016.
43. Yang SD, Yang DL, Sun YP, Wang BL, Ma L, Feng SQ and Ding WY: 17 $\beta$ -estradiol protects against apoptosis induced by interleukin-1 $\beta$  in rat nucleus pulposus cells by down-regulating MMP-3 and MMP-13. *Apoptosis* 20: 348-357, 2015.
44. Aggarwal BB, Shishodia S, Sandur SK, Pandey MK and Sethi G: Inflammation and cancer: How hot is the link? *Biochem Pharmacol* 72: 1605-1621, 2006.
45. Schmiegel W, Roeder C, Schmielau J, Rodeck U and Kalthoff H: Tumor necrosis factor alpha induces the expression of transforming growth factor alpha and the epidermal growth factor receptor in human pancreatic cancer cells. *Proc Natl Acad Sci USA* 90: 863-867, 1993.
46. Wang T, Yang SD, Liu S, Wang H, Liu H and Ding WY: 17 $\beta$ -estradiol inhibites tumor necrosis factor-a induced apoptosis of human nucleus pulposus cells via the PI3K/Akt pathway. *Med Sci Monit* 22: 4312-4322, 2016.