

Dysregulation of synaptic and extrasynaptic N-methyl-D-aspartate receptors induced by amyloid- β

Zhi-Cong Wang, Jie Zhao, Shao Li

Department of Physiology, Dalian Medical University, Dalian 116044, China

Corresponding authors: Shao Li and Jie Zhao. E-mail: lishao89@hotmail.com, dlzhaoj@163.com

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2013

The toxicity of amyloid-beta ($A\beta$) is strongly associated with Alzheimer's disease (AD), which has a high incidence in the elderly worldwide. Recent evidence showed that alteration in the activity of N-methyl-D-aspartate receptors (NMDARs) plays a key role in $A\beta$ -induced neurotoxicity. However, the activation of synaptic and extrasynaptic NMDARs has distinct consequences for plasticity, gene regulation, neuronal death, and $A\beta$ production. This review focuses on the dysregulation of synaptic and extrasynaptic NMDARs induced by $A\beta$. On one hand, $A\beta$ downregulates the synaptic NMDAR response by promoting NMDAR endocytosis, leading to either neurotoxicity or neuroprotection. On the other hand, $A\beta$ enhances the activation of extrasynaptic NMDARs by decreasing neuronal glutamate uptake and inducing glutamate spillover, subsequently causing neurotoxicity. In addition, selective enhancement of synaptic activity by low doses of NMDA, or reduction of extrasynaptic activity by memantine, a non-competitive NMDAR antagonist, halts $A\beta$ -induced neurotoxicity. Therefore, future neuroprotective drugs for AD should aim at both the enhancement of synaptic activity and the disruption of extrasynaptic NMDAR-dependent death signaling.

Keywords: amyloid- β ; synaptic NMDA receptor; extrasynaptic NMDA receptor; neurotoxicity

Introduction

Alzheimer's disease (AD) is the most common form of dementia and is pathologically characterized by senile plaques, neurofibrillary tangles, and synaptic loss. Although the pathogenesis of the disease is still not well understood, amyloid-beta ($A\beta$) is widely recognized to be neurotoxic. To date, the downstream signaling pathways of N-methyl-D-aspartate receptors (NMDARs) are thought to be involved in $A\beta$ -induced neurotoxicity. For example, $A\beta$ induces synaptic loss^[1-3] and postsynaptic density-95 (PSD-95) degradation^[4-6], stimulates the production of reactive oxygen species (ROS)^[7-10], disrupts axonal transport^[11], and causes microtubule deregulation^[12] through an NMDAR-dependent mechanism. Furthermore, $A\beta$ not only binds to domains within or near NMDARs *in vitro* and *in vivo*^[3, 7, 9, 13], but also directly activates NMDARs^[14-16].

NMDARs are the major subtype of ligand-gated

ionotropic glutamate receptors expressed widely in the central nervous system. Based on their localization on the cell membrane, NMDARs are distinguished as synaptic or extrasynaptic. Synaptic NMDARs are classically defined as functional receptors that are activated by glutamate released during low-frequency synaptic events, whereas extrasynaptic NMDARs, which are not activated by synaptically-released glutamate from presynaptic vesicles, are found at various locations such as the cell body, the dendritic shaft, and the neck of the dendritic spine. For many years it has been thought that the degree of Ca^{2+} influx through NMDARs is solely responsible for differences in cellular outcome: moderate levels of NMDAR activity are beneficial for neurons, while excessive activation of NMDARs is deleterious due to Ca^{2+} overload. However, synaptic and extrasynaptic NMDARs are known to play opposing roles in some signaling pathways: synaptic NMDARs play a protective role by promoting nuclear

signaling to cAMP response element binding protein (CREB), inducing gene expression of brain-derived neurotrophic factor (BDNF), activating extracellular signal-regulated kinases (ERK) and an anti-apoptotic pathway, whereas extrasynaptic NMDARs antagonize signaling to CREB, block BDNF expression, and cause mitochondrial membrane potential loss and cell death^[17-23]. Also, selective stimulation of extrasynaptic NMDARs triggers excitotoxicity, but Ca²⁺ overload through synaptic NMDARs is not neurotoxic^[19, 21, 24]. This review aims to summarize recent studies on the dysregulation of synaptic and extrasynaptic NMDARs induced by A β , and provide new insights into the development of AD therapies based on the balance between synaptic and extrasynaptic NMDARs.

Activation of Synaptic or Extrasynaptic NMDARs Influences A β Production

NMDAR stimulation has been reported to increase the levels of α -C-terminal fragments (C83) of amyloid precursor protein (APP) and release soluble APP (sAPP), as well as decrease the production and release of A β ₁₋₄₀, all of which are blocked by NMDAR antagonists or α -secretase inhibitors^[25]. However, chronic NMDA exposure increases the expression of neuronal Kunitz protease inhibitory domain-containing APPs (KPI-APPs; isoforms exhibiting important amyloidogenic potential), which subsequently inhibit the α -secretase candidate tumor necrosis factor- α converting enzyme, and increase the production of A β ^[26].

To determine whether this contradiction is explainable in terms of the activation of synaptic or extrasynaptic NMDARs, they were separately activated in a series of experiments. Activation of synaptic NMDARs alone inhibits A β release and increases C83 production and ERK phosphorylation^[25] (Fig. 1). A β is produced primarily within neurons and secreted into the interstitial fluid (ISF). Using a micro-dialysis technique to specifically measure dynamic changes in ISF A β levels *in vivo*, Verges *et al.* revealed that high doses of NMDA or NMDAR agonist activate ERK and reduce the processing of APP into A β , while NMDAR antagonists increase ISF A β levels, suggesting that basal activity of these receptors normally suppresses A β levels^[27]. Similar to these reports, after synaptic NMDAR activation, Bordji *et al.* failed to detect any KPI-APPs, and

A β levels were not modified^[19]. Mechanistically, stimulation of NMDARs up-regulates the genes encoding a disintegrin and metalloproteinase 10 (ADAM10), the constitutive α -secretase that governs the non-amyloidogenic pathway of APP processing, and increases the trafficking of ADAM10 to the postsynaptic membrane^[28, 29]. On the other hand, selectively activating extrasynaptic NMDARs does not stimulate C83 production and ERK phosphorylation^[25], but induces a significant increase in KPI-APP mRNA and A β production, with no change in total APP mRNA expression, implying that a shift from normal APP to KPI-APP expression causes the higher production of A β ^[19] (Fig. 1).

Dysregulation of Synaptic NMDARs Induced by A β

A β Promotes Synaptic NMDAR Endocytosis

To determine the effect of A β on cell surface (both synaptic and extrasynaptic) NMDAR expression, Snyder *et al.* treated cultured cortical neurons with A β for 1 h and found that A β reduces the surface expression of the NR2B and NR1 subunits of NMDARs, but does not change the total levels (including internalized surface receptors) of NR2B, which is observed in the neurons from APP_{Swe} mice that over-express human APP with the familial Swedish mutation. And the effect of A β is completely inhibited by a γ -secretase inhibitor, which substantially reduces the A β level^[30]. Consistent with these data, A β -mediated NMDAR endocytosis also occurs in other mouse cortical or hippocampal neurons that are exposed to A β ^[3, 13, 31-33], and neurons or retinal membrane from transgenic animals with AD^[13, 33, 34].

Specifically, A β significantly reduces NR1 at synaptic sites, further supporting the selective action of A β on synaptic NMDARs^[30]. In neurons from APP [V717I] transgenic mice, a model that mimics the early cognitive impairment in AD, the amounts of postsynaptic NR2B are also decreased^[13]. In addition, Li *et al.* used a biochemical approach to separate synaptic and extrasynaptic fractions from acute hippocampal slices after incubation with 7PA2 CM (medium from cells stably over-expressing a human APP mutation and secreting A β). They found a significant decrease of synaptic NR2B at 6 and 16 h, but no change of the NR2B levels in either the synaptic or extrasynaptic fractions at 30 min^[35].

reduce STEP activity in transgenic AD mice increases the expression of NR2B, leads to significantly improved cognitive function, and facilitates LTP^[40, 41] (Fig. 1). Moderate ethanol consumption has been associated with a reduced risk of AD in several epidemiological studies. Recently, ethanol preconditioning-dependent neuroprotection was found to be associated with early enhancement of synaptic NR2B localization and NMDAR activity^[42]. Also, chronic ethanol exposure increases synaptic NR2B clusters and synaptic NMDA currents, but does not change extrasynaptic NMDAR clusters^[43].

However, downregulation of NMDARs can also be neuroprotective. Pretreating cultured neurons with A β promotes NMDAR endocytosis, decreases Ca²⁺ influx, and protects neurons from NMDA- and glutamate-induced excitotoxicity^[31, 44]. Likewise, the retinas of transgenic mice over-expressing A β have reduced NR2B, and present with less NMDA-induced retinal damage than wild-type mice^[34]. Moreover, donepezil, an acetylcholinesterase inhibitor, decreases glutamate toxicity *via* promoting NMDAR endocytosis and attenuating glutamate-mediated Ca²⁺ entry^[45].

A β Reduces Postsynaptic NMDA Currents

To investigate the effect of A β on synaptic transmission, excitatory postsynaptic currents (EPSCs) were recorded in neurons from mice over-expressing APP. Li *et al.* reported a significant decrease of NMDA-EPSCs in slices treated with 7PA2 CM. Moreover, they suggested that an A β -mediated rise in glutamate levels leads to receptor desensitization and then causes NMDA-EPSC reduction^[46]. Also, it has been reported that A β reduces NMDA postsynaptic currents^[47]. A decrease in evoked-response amplitude could result from negative modulation of the presynaptic release machinery, yet A β does not alter the synaptic release probability and presynaptic vesicle release at presynaptic sites^[2, 47, 48].

Since A β promotes synaptic NMDAR endocytosis and reduces NMDA-evoked currents, A β may attenuate Ca²⁺ influx at synapses. Using two-photon uncaging of glutamate to stimulate individual dendritic spines while monitoring spine head Ca²⁺ transients, Shankar *et al.* showed that A β reduces NMDAR-dependent Ca²⁺ influx into the spine head^[1]. Pre-incubation of cortical and hippocampal neurons with a low (100 nmol/L) or a high concentration (1 μ mol/L)

of A β for 30 min or 3 h inhibits the NMDA or glutamate-mediated increase of cytosolic Ca²⁺^[13, 16, 31, 44]. Also, prolonged exposure (16 h) of cortical neurons to A β causes an attenuation of NMDAR-mediated Ca²⁺ influx^[8].

Dysregulation of Extrasynaptic NMDARs Induced by A β

A β Enhances the Activation of Extrasynaptic NMDARs by Reducing Excitatory Amino-acid Transporters

The glutamate-aspartate transporter (GLAST) and glutamate transporter-1 (GLT-1) are the main proteins responsible for removing excess glutamate from the synaptic cleft. Considerable evidence supports the hypothesis that A β downregulates the uptake of glutamate by decreasing these glutamate transporters in both the cortex and fibroblasts of AD patients or animal models of AD^[49-52], as well as astrocytes both *in vitro* and *in vivo*^[53, 54]. Recently, Bicca *et al.* revealed decreases in GLT-1 and GLAST expression and glutamate uptake in the hippocampus of A β -treated mice, which could be prevented by a selective and competitive NMDAR antagonist^[10]. Interestingly, disturbance of cholesterol metabolism or aging may contribute to a reduction in glutamate transporters^[55, 56]. Moreover, A β promotes extracellular glutamate release from hippocampal neurons^[57], and increases the hippocampal levels of extracellular glutamate^[58]. After selectively blocking synaptic NMDARs, perfusing 7PA2 CM markedly increases NMDA-EPSCs, and a selective NR2B inhibitor strongly inhibits NMDA-EPSCs, suggesting that A β enhances the activation of extrasynaptic NR2B^[35, 46] (Fig. 1). Also, inhibition of glutamate transporters causes glutamate spillover from the synapse and increases extrasynaptic NMDA-EPSCs^[35, 59].

As discussed above, A β can decrease the NMDAR-mediated intracellular Ca²⁺ concentration^[13, 16, 31, 44], but triggers a sustained Ca²⁺ influx mediated by NMDARs^[7, 15, 60]. This paradox may be explained by the activation of extrasynaptic NMDARs. Li *et al.* showed that the change in extrasynaptic NMDAR-mediated cytosolic Ca²⁺ in response to NMDA is significantly greater in 7PA2 CM-treated neurons^[35], and this is significantly attenuated by memantine^[7, 16, 60], consistent with the notion that memantine predominantly blocks extrasynaptic NMDARs^[21, 61, 62].

Neurotoxicity Is Mediated by Extrasynaptic NR2B-NMDARs

NR2B-NMDARs are predominantly located at extrasynaptic sites in mature neurons, and their activation promotes cell death through a series of pathways, which have been described in detail elsewhere^[63]. Synaptic plasticity such as LTP or long-term depression (LTD) is thought to underlie learning and memory. Depending on the stimulation protocol, Li *et al.* reported that A β from several sources (synthetic, cell culture, human brain extracts) facilitates LTD through metabotropic glutamate receptors (300 pulses) or NMDARs (900 pulses). They found that the effect of A β is caused by the activation of extrasynaptic NR2B through the inhibition of glutamate uptake, as it can be prevented by an extracellular glutamate scavenger and is closely mimicked by the inhibition of glutamate uptake^[46]. Likewise, A β -inhibited LTP shares the same mechanism^[35, 64]. Moreover, A β application impairs early neuronal function before major cytotoxic effects, including LTP, baseline synaptic transmission, spontaneous neuronal network activity, retraction of synaptic contacts, and accumulated Jacob (a messenger that couples extrasynaptic NMDAR activity to CREB) dephosphorylation in the nucleus; all these effects are blocked by extrasynaptic NR2B antagonists^[65] (Fig. 1). Consistent with these reports, only NR2B antagonists, but not those of NR2A, protect against NMDA-induced excitotoxicity^[16, 66], A β -mediated inhibition of plasticity^[67], A β -induced PSD-95 and synaptophysin loss^[5, 6], and A β -increased Ca²⁺ influx^[16], as well as A β -induced ROS production and endoplasmic reticulum stress^[68]. In addition, activation of either synaptic or extrasynaptic NR2B results in excitotoxicity and neuronal apoptosis^[66].

The Balance between Synaptic and Extrasynaptic NMDARs for Therapeutic Targeting

When glutamate is used to activate both extrasynaptic and synaptic NMDARs, the extrasynaptic receptors shut off synaptic NMDAR signaling^[17, 18, 21, 23]. Conversely, enhancement of synaptic NMDAR activity protects against extrasynaptic NMDAR-induced neuronal death^[69]. Thus, the selective enhancement of synaptic activity or reduction of extrasynaptic activity may be sufficient to prevent A β -induced neurotoxicity.

A β downregulates PSD-95 and synaptophysin in an NMDAR-dependent manner, and only extrasynaptic NR2B-NMDAR antagonists abrogate this effect, while blockade of synaptic NMDAR activity does not influence these effects^[5, 6]. However, pretreatment with a low dose of NMDA (1 μ mol/L) prevents the actions of A β , and the protective effect is eliminated only by blockade of synaptic NMDARs. In contrast, a high dose of NMDA (10 μ mol/L) potentiates the effect of A β , which is only abolished by ifenprodil^[5], suggesting that the enhancement of synaptic NMDAR activity can halt the manifestation of early-stage events in AD.

Memantine has received marketing authorization from the European Medicines Agency and the Food and Drug Administration (USA) for the treatment of moderate to severe AD. Recently, accumulating evidence indicates that memantine, at therapeutic concentrations, preferentially blocks extrasynaptic over synaptic NMDARs in hippocampal autapses and cortical neurons^[21, 61, 62]. Blocking extrasynaptic NMDARs with memantine inhibits NMDAR-induced KPI-APP expression and decreases A β expression and release both in cellular models and transgenic animals^[19, 70], reduces tau hyperphosphorylation^[71, 72] and microtubule deregulation^[12], blocks excessive formation of ROS^[7] and intracellular or mitochondrial Ca²⁺ overload^[7, 15, 60], and prevents A β -inhibited LTP^[73] and synaptic deterioration^[3]. In addition, memantine inhibits CREB shutoff and rescues neurons from NMDA-mediated toxicity^[15, 23].

Conclusions and Perspective

In summary, A β differentially affects the activity of synaptic and extrasynaptic NMDARs, resulting in neurotoxicity or neuroprotection. Although most recent evidence supports this notion, some issues still need to be addressed. For example, how A β influences NR2A levels and whether there are other potential effects. In addition, it is known that A β activates many receptors both on the cellular surface and in the nucleus, and their combined effects as well as the upstream or downstream signaling pathways are yet to be described.

Despite the fact that NMDARs mediate brain damage in AD, clinical trials of NMDAR antagonists are not therapeutically effective due to blockade of synaptic NMDAR activity. Therefore, a sufficient understanding

of NMDAR dysregulation induced by A β provides a novel conceptual basis for the future development of neuroprotective therapies for AD. And antagonists designed to selectively target extrasynaptic NMDAR signaling while sparing the physiological and neuroprotective roles of synaptic NMDARs could be promising therapies for AD.

ACKNOWLEDGEMENTS

This review was supported by the National Natural Science Foundation of China (81371223 and 81371437), the Research Fund for the Doctoral Program of Higher Education of China (20122105110010) and the Science and Technology Project of Liaoning Province, China (2011226006).

Received date: 2012-12-09; Accepted date: 2013-02-25

REFERENCES

- [1] Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL. Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci* 2007, 27: 2866–2875.
- [2] Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, *et al.* Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 2008, 14: 837–842.
- [3] Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, Wood M, *et al.* Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci* 2007, 27: 796–807.
- [4] Roselli F, Tirard M, Lu J, Hutzler P, Lamberti P, Livrea P, *et al.* Soluble beta-amyloid1-40 induces NMDA-dependent degradation of postsynaptic density-95 at glutamatergic synapses. *J Neurosci* 2005, 25: 11061–11070.
- [5] Liu J, Chang L, Roselli F, Almeida OF, Gao X, Wang X, *et al.* Amyloid-beta induces caspase-dependent loss of PSD-95 and synaptophysin through NMDA receptors. *J Alzheimers Dis* 2010, 22: 541–556.
- [6] Liu J, Wu Y, Chang L, Peng J, Song Y, Lu T, *et al.* Effects of NMDA receptors in synapses and extrasynapses of rat hippocampal neurons on amyloid-beta-induced alterations in PSD-95 expression. *Neuroembryol Aging* 2008, 5: 116–120.
- [7] De Felice FG, Velasco PT, Lambert MP, Viola K, Fernandez SJ, Ferreira ST, *et al.* Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Biol Chem* 2007, 282: 11590–11601.
- [8] He Y, Cui J, Lee JC, Ding S, Chalimoniuk M, Simonyi A, *et al.* Prolonged exposure of cortical neurons to oligomeric amyloid-beta impairs NMDA receptor function via NADPH oxidase-mediated ROS production: protective effect of green tea (-)-epigallocatechin-3-gallate. *ASN Neuro* 2011, 3: e00050.
- [9] Decker H, Jurgensen S, Adrover MF, Brito-Moreira J, Bomfim TR, Klein WL, *et al.* N-methyl-D-aspartate receptors are required for synaptic targeting of Alzheimer's toxic amyloid-beta peptide oligomers. *J Neurochem* 2010, 115: 1520–1529.
- [10] Bicca MA, Figueiredo CP, Piermartiri TC, Meotti FC, Bouzon ZL, Tasca CI, *et al.* The selective and competitive N-methyl-D-aspartate receptor antagonist, (-)-6-phosphonomethyl-deca-hydroisoquinoline-3-carboxylic acid, prevents synaptic toxicity induced by amyloid-beta in mice. *Neuroscience* 2011, 192: 631–641.
- [11] Decker H, Lo KY, Unger SM, Ferreira ST, Silverman MA. Amyloid-beta peptide oligomers disrupt axonal transport through an NMDA receptor-dependent mechanism that is mediated by glycogen synthase kinase 3beta in primary cultured hippocampal neurons. *J Neurosci* 2010, 30: 9166–9171.
- [12] Mota SI, Ferreira IL, Pereira C, Oliveira CR, Rego AC. Amyloid-beta peptide 1-42 causes microtubule deregulation through N-methyl-D-aspartate receptors in mature hippocampal cultures. *Curr Alzheimer Res* 2012, 9: 844–856.
- [13] Dewachter I, Filipkowski RK, Priller C, Ris L, Neyton J, Croes S, *et al.* Deregulation of NMDA-receptor function and downstream signaling in APP[V717I] transgenic mice. *Neurobiol Aging* 2009, 30: 241–256.
- [14] Texido L, Martin-Satue M, Alberdi E, Solsona C, Matute C. Amyloid beta peptide oligomers directly activate NMDA receptors. *Cell Calcium* 2011, 49: 184–190.
- [15] Alberdi E, Sanchez-Gomez MV, Cavaliere F, Perez-Samartin A, Zugaza JL, Trullas R, *et al.* Amyloid beta oligomers induce Ca²⁺ dysregulation and neuronal death through activation of ionotropic glutamate receptors. *Cell Calcium* 2010, 47: 264–272.
- [16] Ferreira IL, Bajouco LM, Mota SI, Auberson YP, Oliveira CR, Rego AC. Amyloid beta peptide 1-42 disturbs intracellular calcium homeostasis through activation of GluN2B-containing N-methyl-d-aspartate receptors in cortical cultures. *Cell Calcium* 2012, 51: 95–106.
- [17] Ivanov A, Pellegrino C, Rama S, Dumalska I, Salyha Y, Ben-Ari Y, *et al.* Opposing role of synaptic and extrasynaptic NMDA receptors in regulation of the extracellular signal-regulated kinases (ERK) activity in cultured rat hippocampal neurons. *J Physiol* 2006, 572: 789–798.
- [18] Hardingham GE, Fukunaga Y, Bading H. Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB

- shut-off and cell death pathways. *Nat Neurosci* 2002, 5: 405–414.
- [19] Bordji K, Becerril-Ortega J, Nicole O, Buisson A. Activation of extrasynaptic, but not synaptic, NMDA receptors modifies amyloid precursor protein expression pattern and increases amyloid-ss production. *J Neurosci* 2010, 30: 15927–15942.
- [20] Stanika RI, Pivovarova NB, Brantner CA, Watts CA, Winters CA, Andrews SB. Coupling diverse routes of calcium entry to mitochondrial dysfunction and glutamate excitotoxicity. *Proc Natl Acad Sci U S A* 2009, 106: 9854–9859.
- [21] Leveille F, El Gaamouch F, Goux E, Lecocq M, Lobner D, Nicole O, *et al.* Neuronal viability is controlled by a functional relation between synaptic and extrasynaptic NMDA receptors. *FASEB J* 2008, 22: 4258–4271.
- [22] Papadia S, Soriano FX, Leveille F, Martel MA, Dakin KA, Hansen HH, *et al.* Synaptic NMDA receptor activity boosts intrinsic antioxidant defenses. *Nat Neurosci* 2008, 11: 476–487.
- [23] Kaufman AM, Milnerwood AJ, Sepers MD, Coquinco A, She K, Wang L, *et al.* Opposing roles of synaptic and extrasynaptic NMDA receptor signaling in cocultured striatal and cortical neurons. *J Neurosci* 2012, 32: 3992–4003.
- [24] Xu J, Kurup P, Zhang Y, Goebel-Goody SM, Wu PH, Hawasli AH, *et al.* Extrasynaptic NMDA receptors couple preferentially to excitotoxicity via calpain-mediated cleavage of STEP. *J Neurosci* 2009, 29: 9330–9343.
- [25] Hoey SE, Williams RJ, Perkinson MS. Synaptic NMDA receptor activation stimulates alpha-secretase amyloid precursor protein processing and inhibits amyloid-beta production. *J Neurosci* 2009, 29: 4442–4460.
- [26] Lesne S, Ali C, Gabriel C, Croci N, MacKenzie ET, Glabe CG, *et al.* NMDA receptor activation inhibits alpha-secretase and promotes neuronal amyloid-beta production. *J Neurosci* 2005, 25: 9367–9377.
- [27] Verges DK, Restivo JL, Goebel WD, Holtzman DM, Cirrito JR. Opposing synaptic regulation of amyloid-beta metabolism by NMDA receptors *in vivo*. *J Neurosci* 2011, 31: 11328–11337.
- [28] Marcello E, Gardoni F, Mauceri D, Romorini S, Jeromin A, Epis R, *et al.* Synapse-associated protein-97 mediates alpha-secretase ADAM10 trafficking and promotes its activity. *J Neurosci* 2007, 27: 1682–1691.
- [29] Wan XZ, Li B, Li YC, Yang XL, Zhang W, Zhong L, *et al.* Activation of NMDA receptors upregulates a disintegrin and metalloproteinase 10 via a Wnt/MAPK signaling pathway. *J Neurosci* 2012, 32: 3910–3916.
- [30] Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, *et al.* Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci* 2005, 8: 1051–1058.
- [31] Goto Y, Niidome T, Akaike A, Kihara T, Sugimoto H. Amyloid beta-peptide preconditioning reduces glutamate-induced neurotoxicity by promoting endocytosis of NMDA receptor. *Biochem Biophys Res Commun* 2006, 351: 259–265.
- [32] Dinamarca MC, Colombres M, Cerpa W, Bonansco C, Inestrosa NC. Beta-amyloid oligomers affect the structure and function of the postsynaptic region: role of the Wnt signaling pathway. *Neurodegener Dis* 2008, 5: 149–152.
- [33] Kurup P, Zhang Y, Xu J, Venkitaramani DV, Haroutunian V, Greengard P, *et al.* Abeta-mediated NMDA receptor endocytosis in Alzheimer's disease involves ubiquitination of the tyrosine phosphatase STEP61. *J Neurosci* 2010, 30: 5948–5957.
- [34] Shimazawa M, Inokuchi Y, Okuno T, Nakajima Y, Sakaguchi G, Kato A, *et al.* Reduced retinal function in amyloid precursor protein-over-expressing transgenic mice via attenuating glutamate-N-methyl-d-aspartate receptor signaling. *J Neurochem* 2008, 107: 279–290.
- [35] Li S, Jin M, Koeglsperger T, Shepardson NE, Shankar GM, Selkoe DJ. Soluble Abeta oligomers inhibit long-term potentiation through a mechanism involving excessive activation of extrasynaptic NR2B-containing NMDA receptors. *J Neurosci* 2011, 31: 6627–6638.
- [36] Nagele RG, D'Andrea MR, Anderson WJ, Wang HY. Intracellular accumulation of beta-amyloid(1–42) in neurons is facilitated by the alpha 7 nicotinic acetylcholine receptor in Alzheimer's disease. *Neuroscience* 2002, 110: 199–211.
- [37] Wang HY, Li W, Benedetti NJ, Lee DH. Alpha 7 nicotinic acetylcholine receptors mediate beta-amyloid peptide-induced tau protein phosphorylation. *J Biol Chem* 2003, 278: 31547–31553.
- [38] Wang HY, Bakshi K, Shen C, Frankfurt M, Trocme-Thibierge C, Morain P. S 24795 limits beta-amyloid-alpha7 nicotinic receptor interaction and reduces Alzheimer's disease-like pathologies. *Biol Psychiatry* 2010, 67: 522–530.
- [39] Dzieczapolski G, Glogowski CM, Masliah E, Heinemann SF. Deletion of the alpha 7 nicotinic acetylcholine receptor gene improves cognitive deficits and synaptic pathology in a mouse model of Alzheimer's disease. *J Neurosci* 2009, 29: 8805–8815.
- [40] Venkitaramani DV, Moura PJ, Picciotto MR, Lombroso PJ. Striatal-enriched protein tyrosine phosphatase (STEP) knockout mice have enhanced hippocampal memory. *Eur J Neurosci* 2011, 33: 2288–2298.
- [41] Zhang Y, Kurup P, Xu J, Carty N, Fernandez SM, Nygaard HB, *et al.* Genetic reduction of striatal-enriched tyrosine phosphatase (STEP) reverses cognitive and cellular deficits in an Alzheimer's disease mouse model. *Proc Natl Acad Sci U S A* 2010, 107: 19014–19019.
- [42] Mitchell RM, Neafsey EJ, Collins MA. Essential involvement of the NMDA receptor in ethanol preconditioning-dependent

- neuroprotection from amyloid-beta *in vitro*. *J Neurochem* 2009, 111: 580–588.
- [43] Carpenter-Hyland EP, Woodward JJ, Chandler LJ. Chronic ethanol induces synaptic but not extrasynaptic targeting of NMDA receptors. *J Neurosci* 2004, 24: 7859–7868.
- [44] Niidome T, Goto Y, Kato M, Wang PL, Goh S, Tanaka N, *et al.* Non-fibrillar amyloid-beta peptide reduces NMDA-induced neurotoxicity, but not AMPA-induced neurotoxicity. *Biochem Biophys Res Commun* 2009, 386: 734–738.
- [45] Shen H, Kihara T, Hongo H, Wu X, Kem WR, Shimohama S, *et al.* Neuroprotection by donepezil against glutamate excitotoxicity involves stimulation of alpha7 nicotinic receptors and internalization of NMDA receptors. *Br J Pharmacol* 2010, 161: 127–139.
- [46] Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D. Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 2009, 62: 788–801.
- [47] Cerpa W, Farias GG, Godoy JA, Fuenzalida M, Bonansco C, Inestrosa NC. Wnt-5a occludes Abeta oligomer-induced depression of glutamatergic transmission in hippocampal neurons. *Mol Neurodegener* 2010, 5: 3.
- [48] Ting JT, Kelley BG, Lambert TJ, Cook DG, Sullivan JM. Amyloid precursor protein overexpression depresses excitatory transmission through both presynaptic and postsynaptic mechanisms. *Proc Natl Acad Sci U S A* 2007, 104: 353–358.
- [49] Zoia CP, Tagliabue E, Isella V, Begni B, Fumagalli L, Brighina L, *et al.* Fibroblast glutamate transport in aging and in AD: correlations with disease severity. *Neurobiol Aging* 2005, 26: 825–832.
- [50] Zoia CP, Riva C, Isella V, Proserpio P, Terruzzi A, Arban S, *et al.* Nonfibrillar Abeta 1-42 inhibits glutamate uptake and phosphorylates p38 in human fibroblasts. *Alzheimer Dis Assoc Disord* 2011, 25: 164–172.
- [51] Chen KH, Reese EA, Kim HW, Rapoport SI, Rao JS. Disturbed neurotransmitter transporter expression in Alzheimer's disease brain. *J Alzheimers Dis* 2011, 26: 755–766.
- [52] Schallier A, Smolders I, Van Dam D, Loyens E, De Deyn PP, Michotte A, *et al.* Region- and age-specific changes in glutamate transport in the AbetaPP23 mouse model for Alzheimer's disease. *J Alzheimers Dis* 2011, 24: 287–300.
- [53] Matos M, Augusto E, Oliveira CR, Agostinho P. Amyloid-beta peptide decreases glutamate uptake in cultured astrocytes: involvement of oxidative stress and mitogen-activated protein kinase cascades. *Neuroscience* 2008, 156: 898–910.
- [54] Piermartiri TC, Figueiredo CP, Rial D, Duarte FS, Bezerra SC, Mancini G, *et al.* Atorvastatin prevents hippocampal cell death, neuroinflammation and oxidative stress following amyloid-beta(1-40) administration in mice: evidence for dissociation between cognitive deficits and neuronal damage. *Exp Neurol* 2010, 226: 274–284.
- [55] Tian G, Kong Q, Lai L, Ray-Chaudhury A, Lin CL. Increased expression of cholesterol 24S-hydroxylase results in disruption of glial glutamate transporter EAAT2 association with lipid rafts: a potential role in Alzheimer's disease. *J Neurochem* 2010, 113: 978–989.
- [56] Potier B, Billard JM, Riviere S, Sinet PM, Denis I, Champeil-Potokar G, *et al.* Reduction in glutamate uptake is associated with extrasynaptic NMDA and metabotropic glutamate receptor activation at the hippocampal CA1 synapse of aged rats. *Aging Cell* 2010, 9: 722–735.
- [57] Brito-Moreira J, Paula-Lima AC, Bomfim TR, Oliveira FB, Sepulveda FJ, De Mello FG, *et al.* Abeta oligomers induce glutamate release from hippocampal neurons. *Curr Alzheimer Res* 2011, 8: 552–562.
- [58] O'Shea SD, Smith IM, McCabe OM, Cronin MM, Walsh DM, O'connor WT. Intracerebroventricular administration of amyloid β -protein oligomers selectively increases dorsal hippocampal dialysate glutamate levels in the awake rat. *Sensors* 2008, 8: 7428–7437.
- [59] Harney SC, Jane DE, Anwyl R. Extrasynaptic NR2D-containing NMDARs are recruited to the synapse during LTP of NMDAR-EPSCs. *J Neurosci* 2008, 28: 11685–11694.
- [60] Kelly BL, Ferreira A. beta-Amyloid-induced dynamin 1 degradation is mediated by N-methyl-D-aspartate receptors in hippocampal neurons. *J Biol Chem* 2006, 281: 28079–28089.
- [61] Xia P, Chen HS, Zhang D, Lipton SA. Memantine preferentially blocks extrasynaptic over synaptic NMDA receptor currents in hippocampal autapses. *J Neurosci* 2010, 30: 11246–11250.
- [62] Okamoto S, Pouladi MA, Talantova M, Yao D, Xia P, Ehrnhoefer DE, *et al.* Balance between synaptic versus extrasynaptic NMDA receptor activity influences inclusions and neurotoxicity of mutant huntingtin. *Nat Med* 2009, 15: 1407–1413.
- [63] Hardingham GE, Bading H. Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nat Rev Neurosci* 2010, 11: 682–696.
- [64] Kervern M, Angeli A, Nicole O, Leveille F, Parent B, Villette V, *et al.* Selective impairment of some forms of synaptic plasticity by oligomeric amyloid-beta peptide in the mouse hippocampus: Implication of extrasynaptic NMDA receptors. *J Alzheimers Dis* 2012, 32: 183–196.
- [65] Ronicke R, Mikhaylova M, Ronicke S, Meinhardt J, Schroder UH, Fandrich M, *et al.* Early neuronal dysfunction by amyloid beta oligomers depends on activation of NR2B-containing NMDA receptors. *Neurobiol Aging* 2010, 32: 2219–2228.

- [66] Liu Y, Wong TP, Aarts M, Rooyackers A, Liu L, Lai TW, *et al.* NMDA receptor subunits have differential roles in mediating excitotoxic neuronal death both *in vitro* and *in vivo*. *J Neurosci* 2007, 27: 2846–2857.
- [67] Hu NW, Klyubin I, Anwyl R, Rowan MJ. GluN2B subunit-containing NMDA receptor antagonists prevent Abeta-mediated synaptic plasticity disruption *in vivo*. *Proc Natl Acad Sci U S A* 2009, 106: 20504–20509.
- [68] Costa RO, Lacor PN, Ferreira IL, Resende R, Auberson YP, Klein WL, *et al.* Endoplasmic reticulum stress occurs downstream of GluN2B subunit of N-methyl-d-aspartate receptor in mature hippocampal cultures treated with amyloid-beta oligomers. *Aging Cell* 2012, 11: 823–833.
- [69] Zhang SJ, Buchthal B, Lau D, Hayer S, Dick O, Schwaninger M, *et al.* A signaling cascade of nuclear calcium-CREB-ATF3 activated by synaptic NMDA receptors defines a gene repression module that protects against extrasynaptic NMDA receptor-induced neuronal cell death and ischemic brain damage. *J Neurosci* 2011, 31: 4978–4990.
- [70] Alley GM, Bailey JA, Chen D, Ray B, Puli LK, Tanila H, *et al.* Memantine lowers amyloid-beta peptide levels in neuronal cultures and in APP/PS1 transgenic mice. *J Neurosci Res* 2009, 88: 143–154.
- [71] Song MS, Rauw G, Baker GB, Kar S. Memantine protects rat cortical cultured neurons against beta-amyloid-induced toxicity by attenuating tau phosphorylation. *Eur J Neurosci* 2008, 28: 1989–2002.
- [72] Chohan MO, Khatoun S, Iqbal IG, Iqbal K. Involvement of I2PP2A in the abnormal hyperphosphorylation of tau and its reversal by Memantine. *FEBS Lett* 2006, 580: 3973–3979.
- [73] Klyubin I, Wang Q, Reed MN, Irving EA, Upton N, Hofmeister J, *et al.* Protection against Abeta-mediated rapid disruption of synaptic plasticity and memory by memantine. *Neurobiol Aging* 2009, 32: 614–623.