

Ptosis after glaucoma surgery

Abraham J Park¹
Babak Eliassi-Rad²
Manishi A Desai²

¹Moyes Eye Center, Department of Ophthalmology, ²Boston Medical Center, Department of Ophthalmology, Boston, MA, USA

Purpose: Evaluate factors contributing to ptosis after glaucoma surgery.

Methods: Three-year retrospective chart review from January 1, 2012, to January 1, 2015, 157 eyes, 3 surgeons, at Boston Medical Center, to determine the incidence of ptosis and the effects of each variable contributing to ptosis at 3 months after surgery. Each variable was analyzed using the chi-square or independent samples *t*-test analysis to determine statistical significance of ptosis compared with above variables.

Results: The *t*-test or chi-square analysis showed that gender, steroid duration, use of mitomycin C, duration of surgery, and prior surgeries were not statistically significant factors for ptosis. There was a statistically significant association between those <70 years of age and ptosis ($P<0.05$), non-combined surgery and ptosis ($P<0.05$), shunting surgery and ptosis ($P<0.05$).

Conclusion: Ptotic changes occurred more often in those who have shunting (Ahmed and Baerveldt) surgeries compared with filtering (trabeculectomy and Express) surgeries.

Keywords: ptosis, trabeculectomy, shunting surgery, filtering surgery, glaucoma, combined surgery

Introduction

Ptosis is a noted complication after eye surgery that can lead to cosmetic and functional defects.¹ Its incidence has been documented to be around 10% after cataract surgery in previous studies¹⁻⁴ while others have shown it to be as high as 44% at 6 weeks, although most resolve without treatment.⁵⁻⁷ Radial keratotomy has shown a rate of 10%, and trabeculectomy has shown a rate of 10.7%–12% without mitomycin C (MMC) and 19% with MMC 6 months after surgery.⁸ The incidence of ptosis following trabeculectomy was not influenced by type of conjunctival flap, combined surgery, or previous intraocular surgery.⁹

Some of the proposed mechanism behind the cause of ptosis after surgery include lid edema from locally administered anesthetic, initial myotoxic effects, and the compression of the upper eyelid against the orbital bones from the eyelid speculum reducing blood flow to the levator muscle contributing to the edema.^{1,2,10} However, another study showed the rates of ptosis after cataract surgery were common in both local and general anesthesia groups.¹¹ Another theory is that the use of a lid speculum, stiffness of the lid speculum, and smaller palpebral fissures may play a greater role.^{3,5,12} Crosby et al¹² have shown that the type of lid speculum can also affect the patients' eyelids during surgery with the greatest compression forces on those with the smallest palpebral apertures.

In this study, we investigated whether shunting procedures, duration of surgery, or use of steroids contributed to ptosis after surgery. We also re-evaluated the relationship of age, race, gender, use of MMC, type of speculum, number of previous eye surgeries, and combined glaucoma and cataract surgery to ptosis.

Correspondence: Manishi A Desai
Boston University School of Medicine,
Boston Medical Center, 85 East Concord,
8th Floor, Boston, MA 02118, USA
Tel +1 617 414 2020
Email manishi.desai@bmc.org

Methods

A retrospective chart review was performed, with Institutional Review Board (Integrated Network for Subject Protection In Research [INSPIR II] at Boston University Medical Campus and Boston Medical Center) approval, of a total of 252 eyes that underwent surgery between January 1, 2012, and January 1, 2015, at Boston Medical Center by one of three different surgeons. Diagnosis included angle recession, aphakic, chronic angle closure, mixed mechanism, normal or low tension glaucoma, ocular hypertension, pigment dispersion glaucoma, primary open angle glaucoma, pseudoexfoliation glaucoma, and uveitic glaucoma. Surgeries included shunting (Ahmed and Baerveldt) and filtering (fornix-based trabeculectomy and Express) surgery, either stand-alone (non-combined surgery) or with cataract surgery (combined surgery). The incidence of ptosis was determined by documented involvement of ptosis >3 months. Therefore, inclusion criteria included those 18–99 years of age with the diagnosis of primary and secondary glaucoma as listed above, and the persistent documentation of “ptosis” 3 months after glaucoma surgery. The exclusion criteria included inadequate documentation, previous history of ptosis, additional surgery within 3 months of primary glaucoma surgery, lack or lost to follow-up. The initial 252 eyes were reduced to 157 eyes after the exclusion criteria was applied. Due to the categorical nature of each variable, statistical analysis was performed using chi-square analysis or *t*-test analysis, when appropriate, to determine the statistical significance of variables on ptosis.

Results

Background data

The total number of eyes with ptosis was 27 (17.2%) and the average time for a case with and without ptosis was 81.44 and 79.82 minutes, respectively. Time of surgery did not show statistical significance ($P=0.794$). The number of females and males were 79 and 78, respectively. The number of left and right eyes were 80 and 77, respectively. Neither gender nor left or right eye was clinically significant for ptosis ($P=0.307$ and $P=0.170$, respectively). About 90% of patients were in the 50–90 years age group and those <70 years of age showed clinical significance for ptosis ($P<0.05$, moderate relationship, Cramer’s $V=0.20$). Out of 157 eyes, the type of speculum was only documented in 50 eyes and no clinically significant data could be determined ($P=0.069$). MMC was exclusively used in 20 trabeculectomy cases, and they did not show any clinically significant data in relation to ptosis ($P=0.122$). Steroid duration was split into categories of <4, <8, <12, and >12 weeks and did not show clinical significance for ptosis ($P=0.753$). Fifty-seven eyes had prior

Table 1 Patient demographics

Demographics	No ptosis (%)	Ptosis (%)	P-value
Age			–
Mean	67.23±12.76	64.11±13.14	
Gender			0.31
Male	63 (79.7)	16 (20.3)	
Female	67 (85.9)	11 (14.1)	
Eye			0.17
Left	63 (78.8)	17 (21.3)	
Right	67 (87.0)	10 (13.0)	
Ptosis			–
No		130 (82.8)	
Yes		27 (17.2)	

surgery with no clinical significance for ptosis ($P=0.598$). Forty-six combined cases and 111 non-combined cases were clinically significant for ptosis in those undergoing glaucoma surgery alone (ie, without combined surgery) ($P<0.05$, weak–moderate relationship, $\Phi=0.18$). Sixty-eight shunting surgeries and 89 filtering surgeries showed clinically significant data for having increased ptosis, being greater in those with shunting surgery over filtering surgery ($P<0.05$, weak relationship, $\Phi=0.16$). General demographic data and relation to ptosis is summarized in Tables 1 and 2.

In summary, variables that showed no clinical significance included gender, operated eye, steroid duration, use of MMC,

Table 2 Variables for ptosis

Variables	No ptosis (%)	Ptosis (%)	P-value
Surgical time			0.79
Mean	79.82±29.50	81.44±28.42	
Duration of steroid			0.75
<4 weeks	7 (77.8)	2 (22.2)	
<8 weeks	58 (86.6)	9 (13.4)	
<12 weeks	49 (80.3)	12 (19.7)	
>12 weeks	16 (80.0)	4 (20.0)	
Type of speculum			0.07
Lieberman	5 (83.3)	1 (16.7)	
Netland	12 (94.4)	1 (5.6)	
Wire	17 (65.4)	9 (34.6)	
Mitomycin C			0.12
No	111 (81.0)	26 (19.0)	
Yes	19 (95.0)	1 (5.0)	
Prior surgery			0.60
No	84 (84.0)	16 (16.0)	
Yes	46 (80.7)	11 (19.3)	
Combined surgery			0.02
No	87 (78.4)	24 (21.6)	
Yes	43 (93.5)	3 (6.5)	
Type of surgery			0.04
Filtering*	61 (89.7)	7 (10.3)	
Shunting**	69 (77.5)	20 (22.5)	

Notes: *Filtering surgery includes trabeculectomy and Express. **Shunting surgery includes Ahmed and Baerveldt.

prior surgeries, and duration of surgery. The type of speculum could not be analyzed due to lack of documentation. Age, non-combined surgery, and type of surgery did show clinical significance for ptosis and will be discussed further.

Age

Due to the small number of patients <50 and >80 years of age, new categories were created to perform a chi-square test. New age groups were categorized into <60 years of age, 60–69 years of age, 70–79 years of age, and >80 years of age. Under the new categories, a chi-square test of independence was calculated by comparing the frequency of ptosis at 3 months after surgery for individuals in various age categories. Results showed that there was an association between age and ptosis at 3 months after surgery ($\chi^2(3)=8.04$, $N=157$, $P\leq 0.05$). About 94% of 70–79-year-old patients did not have ptosis, compared with 74%–82% of the rest of the age groups. The relationship was moderate in strength (Cramer's $V=0.23$; Table 3). As expected, the trend of those with prior surgeries were older (68.6 ± 12.4 years of age) compared with those without prior surgery (65.6 ± 13.5 years of age) but did not show statistical significance. As noted before, prior surgery did not show a statistically significant relationship with ptosis.

Combined surgery

A chi-square test of independence was calculated by comparing the frequency of ptosis at 3 months for those who

did and did not have combined surgery. Results showed that there was a statistically significant association between non-combined surgery and having ptosis at 3 months after surgery ($\chi^2(1)=5.21$, $N=157$, $P<0.05$). Results show that 78% of those who did not have combined surgery and 94% of those who had combined surgery did not develop ptosis. The relationship is weak to moderate in strength ($\Phi=0.18$). Individuals who had combined surgery were less likely to develop ptosis (Table 4). If patients with prior surgery were excluded, the results were similar ($\chi^2(1)=5.30$, $N=100$, $P<0.05$) with an even stronger, moderate relationship ($\Phi=0.23$; Table 3).

As expected, those who had prior surgery were much less likely to have a combined surgery. Of the 57 eyes that had prior surgery, only 1 had combined surgery. A chi-square test of independence showed a statistically significant association between those who did not have combined surgery and those who had prior surgery ($\chi^2(1)=32.78$, $N=157$, $P<0.05$). There was a strong relationship ($\Phi=0.46$). Those who were in the >70 years age group were also more likely to have combined surgery. About 45% of the people in the >70 years age group had combined surgery compared with 8% in the <60 years age group and 21% in the 60–69 years age group. A chi-square test of independence showed a statistically significant association between older age and combined surgery ($\chi^2(2)=19.55$, $N=157$, $P\leq 0.05$) with a strong relationship (Cramer's $V=0.35$).

Table 3 Relationship between age categories and ptosis

Age category	Ptosis (3 months)		Total
	No (%)	Yes (%)	
<60 years of age	30 (76.9)	9 (23.1)	39
60–69 years of age	32 (74.4)	11 (25.6)	43
70–79 years of age	50 (94.3)	3 (5.7)	53
>80 years of age	18 (81.8)	4 (18.2)	22
Total	130	27	157
Chi-square test ¹			
	Value	df	Asymptotic significance(2-sided)
Pearson chi-square	8.038*	3	0.045
Likelihood ratio	9.170	3	0.027
Linear-by-linear association	2.798	1	0.094
N of valid cases	157		
Symmetric measures ²			
	Value		P<0.05
Phi	0.226		0.045
Cramer's V	0.226		0.045
N of valid cases	157		

Notes: *1 cell (12.5%) have expected count <5. The minimum expected count is 3.78. ¹Chi-square test: statistical method assessing the relationship between observed and expected categorical values. ²Symmetric measures determine strength of relationship between the observed and expected categorical values.

Table 4 Relationship between non-combined surgery and ptosis

Combined surgery	Ptosis (3 months)		Total
	No (%)	Yes (%)	
Non-combined	87 (78.4)	24 (21.6)	111
Combined	43 (93.5)	3 (6.5)	46
Total	130 (82.8)	27 (17.2)	157
Chi-square test ¹			
	Value	df	Asymptotic significance (2-sided)
Pearson chi-square	5.207*	1	0.022
Continuity correction ^b	4.201	1	0.040
Likelihood ratio	6.046	1	0.014
Linear-by-linear association	5.174	1	0.023
N of valid cases	157		
Symmetric measures ²			
	Value		P<0.05
Phi	0.182		0.022
Cramer's V	0.182		0.022
N of valid cases	157		

Notes: *0 cells (0.0%) have expected count <5. The minimum expected count is 7.91. ^bComputed only for a 2x2 table. ¹Chi-square test: statistical method assessing the relationship between observed and expected categorical values. ²Symmetric measures determine strength of relationship between the observed and expected categorical values.

Type of surgery

A chi-square test of independence was calculated comparing the frequency of ptosis at 3 months after surgery for those who had a filtering surgery compared with shunting surgery. There was a statistically significant association between shunting surgery and ptosis 3 months after surgery ($\chi^2(1)=4.01$, $N=157$, $P<0.05$). Ptosis developed in 10% of filtering surgery compared with 23% of shunting surgery. Shunting surgeries with Ahmed and Baerveldt were more likely to develop ptosis. Evidence shows a weak relationship ($\Phi=0.16$; Table 5).

Further investigation revealed a relationship between non-combined surgery and shunting surgery. A chi-square test of independence results showed that there is a statistically significant association between shunting surgeries and non-combined surgery ($\chi^2(1)=8.17$, $N=157$, $P<0.01$). This is a moderately strong relationship ($\Phi=0.23$) (Table 6). Our results show that 64% of those without combined surgery were more likely to have a shunting surgery compared with 39% of those that did have a combined surgery. Additionally, there was no noted difference in age group and type of surgery performed (Table 7). The current result of this study shows that non-combined surgery and shunting surgery had a statistically significant relationship with ptosis and this relationship may strengthen the association between shunting surgery

Table 5 Relationship between type of surgery (shunting or filtering) and ptosis

Type of surgery	Ptosis (3 months)		Total
	No (%)	Yes (%)	
Shunting*	61 (89.7)	7 (10.3)	68
Filtering**	69 (77.5)	20 (22.5)	89
Total	130 (82.8)	27 (17.2)	157
Chi-square test ¹			
	Value	df	Asymptotic significance (2-sided)
Pearson chi-square	4.014 ^a	1	0.045
Continuity correction ^b	3.205	1	0.073
Likelihood ratio	4.202	1	0.040
Linear-by-linear association	3.989	1	0.046
N of valid cases	157		
Symmetric measures ²			
	Value		P<0.05
Phi	0.160		0.045
Cramer's V	0.160		0.045
N of valid cases	157		

Notes: *Shunting surgery includes Baerveldt and Ahmed implants. **Filtering surgery includes trabeculectomy and Express. ⁰ cells (0.0%) have expected count <5. The minimum expected count is 11.69. ^bComputed only for a 2x2 table. ¹Chi-square test, statistical method assessing the relationship between observed and expected categorical values. ²Symmetric measures determine strength of relationship between the observed and expected categorical values.

Table 6 Relationship between non-combined surgery and type of surgery

Combined surgery	Type of surgery		Total
	Shunt* (%)	Filter** (%)	
Non-combined	71 (64.0)	40 (36.0)	111
Combined	18 (39.1)	28 (60.9)	46
Total	89 (56.7)	68 (43.3)	157
Chi-square test ¹			
	Value	df	Asymptotic significance (2-sided)
Pearson chi-square	8.169 ^a	1	0.004
Continuity correction ^b	7.189	1	0.007
Likelihood ratio	8.148	1	0.004
Linear-by-linear association	8.117	1	0.004
N of valid cases	157		
Symmetric measures ²			
	Value		P<0.05
Phi	0.228		0.004
Cramer's V	0.228		0.004
N of valid cases	157		

Notes: *Shunting surgery includes Baerveldt and Ahmed implants. **Filtering surgery includes trabeculectomy and Express. ⁰ cells (0.0%) have expected count <5. The minimum expected count is 19.92. ^bComputed only for a 2x2 table. ¹Chi-square test, statistical method assessing the relationship between observed and expected categorical values. ²Symmetric measures determine strength of relationship between the observed and expected categorical values.

and ptosis as more patients with non-combined surgery also had a statistically significant association with ptosis.

Discussion

Rates of ptosis can range from 10% to 44% after eye surgery, although most tend to resolve without treatment after cataract surgery.¹⁻⁶ The large difference in ptosis rates may be due to temporary aponeurotic ptosis that can occur after surgery compared with permanent aponeurotic ptosis. Permanent aponeurotic ptosis is the most common form of ptosis before and after surgery. It is marked by good levator function of 12–15 mm and a high or absent lid crease that does not improve. Paris and Quickert⁴ and others^{13,14} have shown that patients with permanent aponeurotic ptosis had disinsertion of the levator palpebrae aponeurosis from the epitarsus and patients with ptosis after surgery had a weak levator

Table 7 The relationship between type of surgery and age

Type of surgery	Age categories (years)			Total
	<60 (%)	60–69 (%)	>70 (%)	
Filtering*	17 (25.0)	16 (23.5)	35 (51.5)	68
Shunting**	22 (39.0)	27 (30.3)	40 (44.9)	89
Total	39 (24.8)	43 (27.4)	75 (47.8)	157

Notes: *Filtering surgery includes trabeculectomy and Express. **Shunting surgery includes Baerveldt and Ahmed implants. There was no clinical significance between age and type of surgery.

palpebrae aponeurosis. Several factors have been investigated without conclusive evidence that there is a direct correlation to ptosis. These factors include general anesthesia, lid edema, and use of a bridle suture. Ptosis and the type of lid speculum has been evaluated and believed to play a greater role in patients with smaller palpebral fissures.¹² Crosby et al¹² noted that a more rigid speculum can cause greater compression and greatest risk of eyelid malposition involving those with smallest palpebral apertures. Our study did not address the lid speculum secondary to lack of documentation regarding the type of speculum in close to two-thirds of the cases.

The association between patients <70 years of age and ptosis was of statistical significance ($P<0.05$). This was counterintuitive to the expected rates of ptosis in older patients with a weaker or higher likelihood of dehiscence levator palpebrae aponeurosis due to involuntional changes or history of prior surgery. However, this could also be due to the large amount of bilateral involuntional changes seen in those >70 years of age. Ptosis became more noticeable in younger patients with less involuntional change prior to surgery. Subsequently, new ptosis was likely to be more common in those who are <70 years of age, while those who are >70 years of age have minimal or less noticeable changes due to involuntional ptosis preoperatively.

Patients >70 years of age were more likely to have combined surgery ($P<0.05$) due to the very nature of older patients having cataracts. As expected, a statistically significant ($P<0.05$) association was present between those who had combined surgery and those without prior surgery since most prior surgeries usually involved cataract surgery. Although combined surgery can be expected in an older age group due to higher prevalence of visually significant cataract in this age group, one could also assume that combined surgery is longer and can result in ptosis due to longer exposure to the lid speculum. However, our results showed that ptosis was more likely to be seen in those <70 years of age who had non-combined surgery. Even after excluding prior surgeries that could have previously affected the levator palpebrae aponeurosis, there was a statistically significant ($P<0.05$) association between ptosis and non-combined surgery.

Naruo-tsuchisaka et al⁸ have shown that rates of ptosis after trabeculectomy were as high as 19% with MMC. Although our study did not show statistically significant rates for ptosis with the use of MMC, ptosis was seen in 10.3% of those with filtering surgery and 22.5% in those with shunting procedures. The difference was statistically significant ($P<0.05$) for ptosis to be more likely with shunting surgery compared with filtering surgery. There was also no difference

in age group and the type of surgery performed, indicating that there were similar percentages of shunting and filtering surgery per age group. The potential cause of increased ptosis in patients with shunting surgery could be due to the need for increased exposure required during shunting surgery in order to place a glaucoma drainage device 8–10 mm posterior to the limbus. The need for increased exposure possibly resulted in more pressure on the levator palpebrae aponeurosis by the lid speculum compared with the fornix-based filtering surgery performed in our study.

In our study, variables that showed no clinical significance for ptosis included gender, operated eye, steroid duration, use of MMC, prior surgeries, and duration of surgery. In contrast, those <70 years of age appeared to have more non-combined surgery and were more likely to have shunting surgery. These were both statistically significant ($P<0.05$). All three variables of <70 years of age group, non-combined surgery, and shunting surgery showed a statistically significant association ($P<0.05$) with ptosis. The larger percentage of patients undergoing non-combined surgery in those <70 years of age could be due to the current practice pattern or patients without the need to have the cataract removed at the time of glaucoma surgery, which may not be reflected in all practices. The larger number of non-combined, shunting surgery could also be a practice pattern or the desire to perform shunting surgery prior to filtering surgery in more advanced glaucoma, likely to be seen in older patients, and may not be applicable to every practice. However, the statistically significant association between non-combined surgery and ptosis may strengthen the relationship between shunting surgery and ptosis because there was a statistically significant association between non-combined surgery and shunting surgery indicating that non-combined, shunting surgery may contribute to ptosis rather than a possible difference in practice pattern.

Limitations of our study include lack of formal ptosis measurements like those performed in Naruo-tsuchisaka et al⁸ study with the marginal reflex distance (MRD). Therefore, in our study, ptosis is more likely to be documented in severe cases that can inherently decrease the true number of patients with ptosis, especially in the elderly. It is also difficult to determine the true rate of aponeurotic ptosis in older patients due to the lack of documentation. This may have resulted in the higher number of ptosis in the younger age group. The higher number of ptosis in the younger age group could also reflect the new onset of aponeurotic ptosis evidenced by the documentation of ptosis due to the greater change in eyelid position than those of older patients who may have aponeurotic changes already. Although levator

function is considered grossly intact when measuring aponeurotic ptosis, there is the possibility that the levator palpebrae muscle is weaker in older patients as well. One could consider the fact that younger patients may increase compression or stress on the eyelid from the speculum secondary to their ability to squeeze their eyelids shut with more strength than older patients. Therefore, the type of speculum, surgery (shunting or filtering), and the pre-, intra-, and postoperative interpalpebral fissures can be measured in further studies to determine if the amount of tension on the eyelids from the speculum also plays a role in aponeurotic ptosis after filtering or shunting surgery.

Crosby et al¹² had noted that, smaller palpebral fissures appear to play a role in ptosis and eyelid malposition. However, our study did not address whether patients had small or large palpebral fissures. The palpebral fissure height may also be affected by the preoperative drop regimen, which include medication to dilate the pupil and raise the eyelid. Subsequently, the amount of pressure on the eyelids may also be greater in non-combined surgery compared with combined surgery due to effect of the phenylephrine. Therefore, further prospective studies should be conducted to potentially strengthen the finding of shunting surgery causing more ptosis than filtering surgery by including MRD pre- and postoperatively and vertical palpebral fissure pre-, intra-, and postoperative measurements. Type of surgery and emphasis on intraoperative vertical palpebral fissures measurements after the placement of the speculum can also be measured to determine if greater stretch on eyelids had an association with increased aponeurotic ptosis or changes.

The effect of MMC can also be re-evaluated since some physicians are beginning to use MMC with shunting surgery, and the rate of ptosis of trabeculectomy and MMC in Naruo-tsuchisaka et al⁸ study was 19% compared with 22.5% for shunting surgery without MMC in our study.

Finally, the limitation of a retrospective study without proper case and control patients limit the utility of the study. Results are limited to the current population and practice patterns in the study. As mentioned above, the pre- and postoperative measurements of MRD and palpebral fissures can provide better results for ptosis after a non-combined shunting or filtering surgery. An age-matched case-control, prospective study measuring changes for ptosis (pre- and postoperative MRD, levator function, and vertical palpebral fissures), use of MMC, type of lid speculum, and non-combined filtering or shunting will likely produce stronger results than our current retrospective study. By conducting a prospective study on non-combined surgery

with appropriate ptosis measurements, the comparative data can be more accurate without the variables of practice patterns, improve clarity of rates of new or worsening ptosis in patients $>$ or $<$ 60 years, and directly compare the rates of new or worsening aponeurotic ptosis between filtering and shunting surgeries.

Conclusion

There was statistically significant association between shunting surgery being more common than filtering surgery in patients with ptosis without a significant difference in the percentage of shunting or filtering surgery across all our age categories. There was also statistically significant association between ptosis and shunting surgery, regardless of whether the patient had prior surgery. Higher ptosis rates were noted in the younger, $<$ 70 years age group, that underwent more non-combined surgery, which is typically shorter in duration, though one might expect surgeries of longer duration that include combined surgery, patients with prior surgeries, and older age to have to ptosis.

There was no statistically significant association between the following variables and ptosis: gender, operated eye, steroid duration, use of MMC, prior surgeries, and duration of surgery. The type of speculum could not be analyzed due to lack of documentation.

Our findings of shunting surgery with younger patients and the association with ptosis may be secondary to practice patterns and the desire to preserve filtering surgery for more advanced glaucoma often associated with older patients and to avoid the increased risk of bleb-related infections over the lifetime of the bleb. Furthermore, the association between younger age and ptosis may be due to less involuntional changes in younger patients preoperatively, resulting in new or greater changes in ptosis than older patient with pre-existing involuntional aponeurotic ptosis prior to surgery. Further studies including the type of surgery; type of lid speculum; pre- and postoperative MRD; pre-, intra-, and postoperative vertical IPF; and pre- and postoperative LF measurements can help evaluate the true rates of aponeurotic ptosis after glaucoma surgery in those who may have had preexisting involuntional aponeurotic ptosis before glaucoma surgery.

Acknowledgment

Patient consent to review medical records was not required since patient data were de-identified and were in compliance with the Boston Medical Center/Boston University School of Medicine Office of Clinical Research Health Insurance

Portability and Accountability Act (HIPPA) requirements for our retrospective study. Patient identified data were kept in a locked room and destroyed at the end of the study to maintain patient confidentiality.

Disclosure

The authors report no conflicts of interest in this work.

References

- Alpar JJ. Acquired ptosis following cataract and glaucoma surgery. *Glaucoma*. 1982;4:66–68.
- Kaplan LJ, Jaffe NS, Clayman HM. Ptosis and cataract surgery. A multivariate computer analysis of a prospective study. *Ophthalmology*. 1985;92(2):237–242.
- Patel JI, Blount M, Jones C. Surgical blepharoptosis – the bridle suture factor? *Eye (Lond)*. 2002;16(5):535–537.
- Paris GL, Quickert MH. Disinsertion of the aponeurosis of the levator palpebrae superioris muscle after cataract extraction. *Am J Ophthalmol*. 1976;81(3):337–340.
- Singh SK, Sekhar GC, Gupta S. Etiology of ptosis after cataract surgery. *J Cataract Refract Surg*. 1997;23(9):1409–1413.
- Bernardino CR, Rubin PA. Ptosis after cataract surgery. *Semin Ophthalmol*. 2002;17(3–4):144–148.
- Hosal BM, Tekeli O, Gürsel E. Eyelid malpositions after cataract surgery. *Eur J Ophthalmol*. 1998;8(1):12–15.
- Naruo-tsuchisaka A, Maruyama K, Arimoto G, Goto H. Incidence of postoperative ptosis following trabeculectomy with mitomycin C. *J Glaucoma*. 2015;24(6):417–420.
- Song MS, Shin DH, Spoor TC. Incidence of ptosis following trabeculectomy: a comparative study. *Korean J Ophthalmol*. 1996;10(2):97–103.
- Ranin EA, Carlson BM. Postoperative diplopia and ptosis. A clinical hypothesis based on the myotoxicity of local anesthetics. *Arch Ophthalmol*. 1985;103(9):1337–1339.
- Ropo A, Ruusuvaara P, Nikki P. Ptosis following periocular or general anesthesia in cataract surgery. *Acta Ophthalmol (Copenh)*. 1992;70(2):262–265.
- Crosby NJ, Shepherd D, Murray A. Mechanical testing of lid speculae and relationship to postoperative ptosis. *Eye (Lond)*. 2013;27(9):1098–1101.
- Baggio E, Ruban JM. Postoperative ptosis: etiopathogenesis, clinical analysis, and therapeutic management. Apropos of a series of 43 cases. *J Fr Ophthalmol*. 1998;21(5):361–373.
- Parsa FD, Wolff DR, Parsa NN, Elahia EE. Upper eyelid ptosis repair after cataract extraction and the importance of Hering's test. *Plast Reconstr Surg*. 2001;108(6):1527–1536; discussion 1537–1538.

Clinical Ophthalmology

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

Dovepress

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.