

The Relationship between Serum Selenium and Zinc with Gastroesophageal Cancers in the Southeast of Iran

Abstract

Background: Selenium (Se) and zinc (Zn) have antioxidant and anticancer properties. **Objective:** The aim of this study was to evaluate the serum levels of Se and Zn and the correlation between the levels of these two elements with risk of incidence of esophageal cancer (EC) and gastric cancer (GC). **Materials and Methods:** In a case-control study, we selected sixty patients with GC or EC as the intervention group and 120 age-matched individuals as the control group. Exclusion criteria were the individuals with kidney and liver failure and the consumer of dietary supplements such as Se and Zn. Measurement of serum Se was done in a graphite furnace system and atomic absorption device of Varian and of serum Zn was done by a flame photometer system (flame) and atomic absorption device of Varian. **Results:** In thirty patients of ECs, 90% were squamous cell carcinoma and 10% adenocarcinoma, and out of thirty patients of GCs, 90% were intestinal type and 10% diffuse type. The level of two elements in cancer patients was lower than the control group ($P < 0.05$). There was no significant difference between two cancer groups for level of Se and Zn, but there was a significant difference between the control group with two other groups. **Conclusion:** Our study confirmed the findings from previous prospective studies and randomized trials that reducing of lower levels of Se and Zn can effect on incidence of cancer.

Keywords: Case-control, esophageal cancer, gastric cancer, selenium, zinc

Introduction

Esophageal cancer (EC) is the eighth most common cancer with respect to incidence and the sixth most common cancer with respect to mortality worldwide.^[1] This cancer is classified into two main types histologically: esophageal adenocarcinoma (EA) and esophageal squamous cell carcinoma (ESCC), each having different risk factors.^[2] Despite recent decline, gastric cancer (GC) is the fourth most common cancer and the second leading cause of cancer-related death worldwide^[3] that is the most common gastrointestinal (GI) cancer in Iran.^[4] In blood, selenium (Se) is found in plasma as well as in the cellular compartment where concentrations are higher and the concentrations are highest in tissues such as liver, kidneys, and spleen.^[5] Se is from the group of VI elements, which has been studied for antioxidant and anticancer properties,^[6,7] especially against GC.^[8] Zinc (Zn) is trace element found in blood, and most of it is contained in bones, skin, and hair (~70%), with the remainder

mainly in liver, kidneys, and muscle. In plasma, one-third of Zn is tightly bound to α 2-macroglobulin, the remainder more loosely to albumin.^[5] Zn deficiency adversely affects the immune system, increases oxidative stress, increases the generation of inflammatory cytokines,^[9] and also potentiates the effects of certain nitrosamines that act as esophageal carcinogens.^[10]

The aim of this study was to evaluate the serum levels of Se and Zn and the correlation between the levels of these two elements with risk of incidence of EC and GC in Southeastern Iran.

Materials and Methods

Patients

This study was approved by the Ethics Committee of Zahedan University of Medical Sciences, Zahedan, Iran, at Winter 2016 (Code: 6000988). In a case-control study, we selected sixty patients with EC or GC as the intervention group who referred to Clinic of Oncology, Emam Ali Hospital, as the intervention group and also

Seyed Mehdi Hashemi,
Mohamadali Mashhadi¹,
Alireza Ansari Moghaddam²,
Javad Yousefi²,
Ahmad Dizangian Mofrad²,
Masoud Sadeghi³,
Abolghasem Allahyari⁴

Department of Hematology and Medical Oncology, Zahedan University of Medical Sciences, ¹Department of Internal Medicine, Ali-Ebne Abitaleb Hospital, Zahedan University of Medical Sciences, ²Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, ³Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, ⁴Department of Hematology and Medical Oncology, Emam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran

Address for correspondence:

Dr. Abolghasem Allahyari,
Department of Hematology and Medical Oncology, Emam Reza Hospital, Mashhad University of Medical Sciences,
Mashhad, Iran.
E-mail: allahyari.abolghasem@yahoo.com

Access this article online

Website: www.ijmpo.org

DOI: 10.4103/ijmpo.ijmpo_83_16

Quick Response Code:



How to cite this article: Hashemi SM, Mashhadi M, Moghaddam AA, Yousefi J, Mofrad AD, Sadeghi M, *et al.* The relationship between serum selenium and zinc with gastroesophageal cancers in the Southeast of Iran. *Indian J Med Paediatr Oncol* 2017;38:169-72.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

120 age-matched individuals as the control group referred patients to the Orthopedic Ward of Khatam-al-Anbia Hospital who had just trauma or fracture.

Exclusion criteria: The individuals with kidney and liver failure and the consumer of dietary supplements such as Se and Zn.

Methods of measurement

Measurement of serum Se was done by a graphite furnace system and atomic absorption device of Varian (Varian 240FS, Co., USA) in Laboratory of Toxicology, Razavi Hospital, Mashhad, Iran. System wavelength was 196 nm and width of the valve was 0.5 nm. The standard had been made by the Merck Company with concentration between 50 and 500 µg/l. We used a calibration standard edition that palladium chloride was used as a matrix modifier.

Measurement of serum Zn was done by a flame photometer system (flame) and atomic absorption device of Varian (wavelength of 213 nm). The standard for Zn had been made by the Merck Company.

Statistical analysis

Data analysis was done by the SPSS version 16 (SPSS, Inc., Chicago, IL, USA) (Student's *t*-test for comparison between the means and Chi-square test for sex).

Results

The mean age and the prevalence of sex for two groups are presented in Table 1. There was no significant difference for age in two groups ($P > 0.05$), but there was a significant difference for sex ($P < 0.05$). Therefore, the control group was just age-matched with the intervention group.

Table 2 shows the type of pathology for the intervention group. Out of sixty patients, thirty patients (50%) had EC (ESCC and EA in 90% and 10%, respectively) and thirty patients (50%) had GC (intestinal type and diffuse type in 90% and 10%, respectively).

The serum level of the variables in the intervention and control groups is presented in Table 3. There was significant difference between two groups for level of Se and also for the level of Zn ($P < 0.001$). Therefore, the level of two elements in cancer patients is lower than the control group.

The serum levels of Se and Zn is located in Table 4 based on age groups. There was a significant difference between the level of Zn in two groups in all age groups and for level of Se was just in age group of 45–65 years.

The serum level of the variables in cancer groups and control group is given in Table 5. There was no significant different between two cancer groups for level of Se and Zn ($P > 0.05$), but there was a significant difference between control groups with two other groups ($P < 0.05$).

Table 1: The baseline characteristics of the intervention and control groups

Variables	Control (n=120)	Intervention (n=60)	P
Age (years)			
Mean±SD	60.7±11.4	59±11.7	>0.05
Sex, n (%)			
Male	87 (72.5)	34 (56.7)	<0.05
Female	33 (27.5)	26 (43.3)	

SD – Standard deviation

Table 2: Type of pathology for intervention group

Variables	n (%)
EC (n=30)	
ESCC	27 (90)
Adenocarcinoma	3 (10)
GC (n=30)	
Intestinal type	27 (90)
Diffuse type	3 (10)

EC – Esophageal cancer; ESCC – Esophageal squamous cell carcinoma; GC – Gastric cancer

Table 3: The serum level of the variables in intervention and control groups

Variables	Control (n=120)	Intervention (n=60)	P
Selenium (µg/L)	117.3±25.3	100.4±17.6	<0.001
Zinc (µg/L)	1020.2±163.0	639±252.2	<0.001

The serum level of the variables in cancer groups and control group based on sex is presented in Figure 1. The levels of Se and Zn in males are lower than females in EC patients, whereas in two other groups are vice versa; there was no significant difference between serum Se and Zn concentrations and sex in the patient groups ($P > 0.05$).

Discussion

GC is the second most common GI malignancy with a male/female ratio of 2.1/1,^[11] and EC is the most common GI malignancies in Zahedan (Southeastern Iran)^[12] that this pattern is different from other regions in Iran that GC is the most common GI malignancies.^[11,12] The highest incidences for EC were reported in China, the Caspian region of Iran, South Africa, and France.^[13] Therefore, in this study, we checked the levels of Se and Zn in GC and EC patients in Southeastern Iran compared with normal or control group as reported risk factors for incidence of GI malignancies.

The studies have shown that Se may have anti-carcinogenic effects, especially against cancers of the lung, prostate, skin, and GI system.^[14,15] Another study^[16] indicated that the inverse association between Se level and GC may occur only among populations with low Se levels. One study^[17] observed significant low serum levels of Zn and Se, as compared with normal healthy controls. This shows an association of serum Se and Zn with cancer esophagus. Ji *et al.*^[18] reported that the serum Zn level was lower

Table 4: The serum level of the variables in intervention and control groups based on age ranges


Variables based age range	n	Control (n=120)	Intervention (n=60)	P
Selenium (µg/L)				
Age group <45	41	116.2±26.2	145.5±172.6	0.390
Age group =45-65	88	119.2±26.1	98.3±17.8	<0.001
Age group >65	51	114.6±23.6	102.4±17.5	0.070
Zinc (µg/L)				
Age group <45	41	1047±181.7	570.1±145.6	<0.001
Age group =45-65	88	1018.5±149.5	652.3±223.6	<0.001
Age group >65	51	1003.4±17.5	648.4±362.4	<0.001

Table 5: The serum level of the variables in cancer groups and control groups

Variables	Control (n=120)	GC patients (n=30)	EC patients (n=30)
Selenium (µg/L)	117.3±25.3*:#	98.4±17.9	102.5±17.5
Zinc (µg/L)	1020.2±163.0*:#	620.6±234.0	657.4±269.4

* $P < 0.05$ between control with GC. # $P < 0.05$ between control with EC. EC – Esophageal cancer; GC – Gastric cancer

in advanced GC and also the serum Se level was highly correlated with the location of GC. Rhee *et al.*^[19] reported that the mean Se concentration in cancer patient group (GC and colon cancer cases) was significantly lower than the normal controls. In Iran, the average serum Se in Ardabil, Kerman, Mazandaran, and Golestan was 82, 119, 123, and 155 µg/L, respectively. Therefore, it was suggested that the high incidence of GC and preneoplastic gastric lesions in Ardabil Province could be partly due to low level of Se.^[8] Mark *et al.*^[20] found highly significant inverse associations of serum Se levels with the incidence of esophageal and gastric cardia cancers. Abnet *et al.*^[10] reported that concentration of high tissue Zn concentration was strongly associated with a reduced risk of developing ESCC. One study^[21] showed that there was an inverse relationship between Se intake and cancer mortality in humans, but there was no correlation between serum/tissue Se concentration and disease histological type or sex in the patient group. Zhang *et al.*^[22] reported that whatever serum levels of Zn were significantly reduced in gastritis, peptic ulcer, and GC group, compared with healthy controls, but the higher the Zn levels are, the more increased risk of GC. The researchers in a review study^[11] concluded that an inverse association between Zn concentrations and EC incidence was apparent in most of the reviewed observational studies. In our study, the level of Se and Zn was significantly lower in cancer patients compared with control group or normal group that this study confirmed the results of other studies. In addition, there was no significant difference between serum Se and Zn concentrations and sex in the patient groups, but there was a significant difference between the level of Zn in two groups in three age groups and for level of Se was just in age group of 45–65 years. Furthermore, the levels of two elements were similar in GC compared with EC ($P > 0.05$).

**Figure 1: The serum level of the variables in cancer groups and control groups based on sex**

Conclusion

Our study confirmed findings from previous prospective studies and randomized trials that reducing of lower levels of Se and Zn can effect on incidence of cancer. Therefore, consideration should be given to supplements that contain Se and Zn for the general population and also patients with esophageal and GCs.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012;62:10-29.

2. Kamangar F, Chow WH, Abnet CC, Dawsey SM. Environmental causes of esophageal cancer. *Gastroenterol Clin North Am* 2009;38:27-57, vii.
3. Crew KD, Neugut AI. Epidemiology of gastric cancer. *World J Gastroenterol* 2006;12:354-62.
4. Hosseini Nezhad Z, Darvish Moghaddam S, Zahedi MJ, Hayatbakhsh M, Sharififar F, Ebrahimi Meimand F, *et al.* Serum selenium level in patients with gastric non-cardia cancer and functional dyspepsia. *Iran J Med Sci* 2015;40:214-8.
5. Biesalski HK, Grimm P. *Pocket Atlas of Nutrition*. 3rd ed. Stuttgart: Georg Thieme Verlag; 2005.
6. Ramoutar RR, Brumaghim JL. Antioxidant and anticancer properties and mechanisms of inorganic selenium, oxo-sulfur, and oxo-selenium compounds. *Cell Biochem Biophys* 2010;58:1-23.
7. Nelson WG, De Marzo AM, Isaacs WB. Prostate cancer. *N Engl J Med* 2003;349:366-81.
8. Nouarie M, Pourshams A, Kamangar F, Sotoudeh M, Derakhshan MH, Akbari MR, *et al.* Ecologic study of serum selenium and upper gastrointestinal cancers in Iran. *World J Gastroenterol* 2004;10:2544-6.
9. Prasad AS, Beck FW, Snell DC, Kucuk O. Zinc in cancer prevention. *Nutr Cancer* 2009;61:879-87.
10. Abnet CC, Lai B, Qiao YL, Vogt S, Luo XM, Taylor PR, *et al.* Zinc concentration in esophageal biopsy specimens measured by x-ray fluorescence and esophageal cancer risk. *J Natl Cancer Inst* 2005;97:301-6.
11. Mashhadi MA, Nezam K, Abdollahinejad MJ. Gastric cancer in South East of Iran. *Int J Hematol Oncol Stem Cell Res* 2009;3:38-42.
12. Mashhadi MA, Nezam K, Bakhshipour AR, Ttorbati TF, Ansarimoghaddam AR. High prevalence of esophageal cancer in South East of Iran. *Int J Hematol Oncol Stem Cell Res* 2011;5:18-21.
13. Islami F, Kamangar F, Aghcheli K, Fahimi S, Semnani S, Taghavi N, *et al.* Epidemiologic features of upper gastrointestinal tract cancers in Northeastern Iran. *Br J Cancer* 2004;90:1402-6.
14. Rayman MP. Selenium in cancer prevention: A review of the evidence and mechanism of action. *Proc Nutr Soc* 2005;64:527-42.
15. Lee SR, Bar-Noy S, Kwon J, Levine RL, Stadtman TC, Rhee SG. Mammalian thioredoxin reductase: Oxidation of the C-terminal cysteine/selenocysteine active site forms a thioselenide, and replacement of selenium with sulfur markedly reduces catalytic activity. *Proc Natl Acad Sci U S A* 2000;97:2521-6.
16. Koriyama C, Campos FI, Yamamoto M, Serra M, Carrasquilla G, Carrascal E, *et al.* Toenail selenium levels and gastric cancer risk in Cali, Colombia. *J Toxicol Sci* 2008;33:227-35.
17. Goyal MM, Kalwar AK, Vyas RK, Bhati A. A study of serum zinc, selenium and copper levels in carcinoma of esophagus patients. *Indian J Clin Biochem* 2006;21:208-10.
18. Ji JH, Shin DG, Kwon Y, Cho DH, Lee KB, Park SS, *et al.* Clinical correlation between gastric cancer type and serum selenium and zinc levels. *J Gastric Cancer* 2012;12:217-22.
19. Rhee JK, Chung JH, Sakong J, Kang PS, Kim CY, Lee KS, *et al.* Association between cancer and selenium concentration in blood and toenails. *Yeungnam Univ J Med* 1992;9:29-43.
20. Mark SD, Qiao YL, Dawsey SM, Wu YP, Katki H, Gunter EW, *et al.* Prospective study of serum selenium levels and incident esophageal and gastric cancers. *J Natl Cancer Inst* 2000;92:1753-63.
21. Charalabopoulos K, Kotsalos A, Batistatou A, Charalabopoulos A, Peschos D, Vezyraki P, *et al.* Serum and tissue selenium levels in gastric cancer patients and correlation with CEA. *Anticancer Res* 2009;29:3465-7.
22. Zhang WH, Wu XJ, Niu JX, Yan H, Wang XZ, Yin XD, *et al.* Serum zinc status and *Helicobacter pylori* infection in gastric disease patients. *Asian Pac J Cancer Prev* 2012;13:5043-6.