



Published in final edited form as:

Mutat Res. 2017 July ; 773: 274–281. doi:10.1016/j.mrrev.2016.08.006.

Potential of Hydrogen Peroxide Toxicity:

From Catalase Inhibition to Stable DNA-iron Complexes

Tulip Mahaseth and Andrei Kuzminov*

Department of Microbiology, University of Illinois at Urbana-Champaign

Abstract

Hydrogen peroxide (H_2O_2) is unique among general toxins, because it is stable in abiotic environments at ambient temperature and neutral pH, yet rapidly kills any type of cells by producing highly-reactive hydroxyl radicals. This life-specific reactivity follows distribution of soluble iron, Fe(II) (which combines with H_2O_2 to form the famous Fenton's reagent), — Fe(II) is concentrated inside cells, but is virtually absent outside them. Because of the immediate danger of H_2O_2 , all cells have powerful H_2O_2 scavengers, the equally famous catalases, which enable cells to survive thousand-fold higher concentrations of H_2O_2 and, in combination with adequate movement of H_2O_2 across membranes, make the killing H_2O_2 concentrations virtually impractical to generate in vivo. And yet, low concentrations of H_2O_2 are somehow used as an efficient biological weapon. Here we review several examples of how cells potentiate H_2O_2 toxicity with other chemicals. At first, these potentiators were thought to simply inhibit catalases, but recent findings with cyanide suggest that potentiators mostly promote the other side of Fenton's reaction, recruiting iron from cell depots into stable DNA-iron complexes that, in the presence of elevated H_2O_2 , efficiently break duplex DNA, pulverizing the chromosome. This multifaceted potentiation of H_2O_2 toxicity results in robust and efficient killing.

Keywords

hydrogen peroxide; Fenton's reaction; iron metabolism; nitric oxide; cyanide; catastrophic chromosomal fragmentation

Life-specific reactivity of H_2O_2

Hydrogen peroxide, H_2O_2 , is a metastable oxygen species and an important intermediate in the redox pathway linking molecular oxygen, O_2 , to the fully reduced oxide in water, H_2O . The complete pathway (1), $O_2 \rightarrow O_2^- \rightarrow H_2O_2 \rightarrow (2)OH \cdot \rightarrow (2)H_2O$, also includes a relatively unstable superoxide (O_2^-), as well as the extremely reactive hydroxyl radical

*for correspondence: B103 C&LSL, 601 South Goodwin Ave., Urbana IL 61801-3709, USA, tel: (217) 265-0329, FAX: (217) 244-6697, kuzminov@life.illinois.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest Statement

The authors declare that there are no conflicts of interest.

(OH·), which is the species responsible for the “reactivity” of all reactive oxygen species (2). Paradoxically, hydrogen peroxide is generally non-reactive with organic compounds (3–5); in everyday life, 3% (~0.8 M) hydrogen peroxide solution is a common household antiseptic, stored in plastic bottles for years. H₂O₂ does not directly interact with pure biopolymers (nucleic acids, demetallated proteins, polysaccharides, lipids); perhaps, the only exceptions are thiol-based sensor proteins (like OxyR in bacteria), reacting to the presence of micromolar levels of H₂O₂ with formation of disulfide bonds that are used for signaling (6). At the same time, hydrogen peroxide is a surprisingly potent bio-toxin, as the same 3% H₂O₂ solution kills all kinds of cells within several minutes (7, 8), that is as soon as H₂O₂ penetrates the cell wall barrier.

The main reason H₂O₂ is generally stable in abiotic environment, yet becomes so reactive upon contact with life, is the differential availability of soluble iron. Since H₂O₂ by itself does not react with organic compounds (with the above exception of select thiol groups in a few signaling proteins), it would be completely innocuous, if not for the fact that iron is broadly employed by all types of cells in catalysis of many essential metabolic reactions and transitions (9). Iron can be found in two forms: the soluble Fe(II) and the practically insoluble Fe(III) (10). In the current oxidative atmosphere of the Earth, at least in the oxic environments, the trace amounts of soluble Fe(II) iron are rapidly oxidized by usually more abundant H₂O₂ to Fe(III) iron (11), which does not react with the remaining H₂O₂, explaining the general H₂O₂ stability in abiotic environment. But exactly due to this reason, the only soluble iron in the environment outside cells is represented by the (highly variable) trace amounts of Fe(III) complexed with natural organic ligands (12–14). The limited and unpredictable availability of Fe(III) iron forces cells of all types to actively procure and stockpile iron to maintain their metabolism and support multiplication (15), accumulating 0.1–1.0 mM total iron (10, 16–18). But even inside the cell, the availability of free Fe(II) iron is limited and tightly controlled (15), because when free Fe(II) and H₂O₂ meet, Fenton’s reaction occurs.

In this famous reaction (19–21) (Fig. 1), soluble Fe(II) iron donates one electron to a hydrogen peroxide molecule, causing its decomposition: $\text{Fe(II)} + \text{H}_2\text{O}_2 \rightarrow \text{Fe(III)} + \text{OH} \cdot + \text{OH}^-$ that produces hydroxyl radical capable of reacting with any organic compound at diffusion rates (5, 22). Hydroxyl radicals kill via DNA damage, as indicated by the exquisite H₂O₂ sensitivity of DNA repair mutants (23–25). Fenton’s reaction is the reason why otherwise relatively innocuous extracellular H₂O₂ becomes a potent poison once inside the cell. At the same time, due to its small size and lack of charge, H₂O₂ shows substantial permeability through the membrane barrier (26, 27), so exposed cells cannot simply block the entry of this “conditional” poison. This life-specific reactivity in combination with substantial membrane permeability makes hydrogen peroxide a popular weapon in bio-warfare: our immune cells use it to kill invading microbes (28–30), bees use it as a honey preservative (31), lactic acid bacteria generate it to kill off the competition (32) (and are one of the few life forms that learned to do without iron in their metabolism (33)), ants (if given a choice) take it as an anti-fungal medication (34), while plants employ hydrogen peroxide to reduce grazing by herbivores (35).

H₂O₂ is impossible to concentrate in vivo

However, this strategy of production of a diffusible life-specific source of hydroxyl radicals suffers from the problems of targeting and is also undermined by efficient detoxification. Because superoxide and hydrogen peroxide in sub-micromolar concentrations are byproducts of aerobic metabolism (36), and because both H₂O₂ and O₂⁻ are commonly employed in pathogen-defense mechanisms, the cells are equipped with powerful catalases (37) (Fig. 1) and superoxide dismutases (38). Both are rare examples of diffusion-limited enzymes — the fastest enzymes possible (39) — capable of scavenging up to low millimolar concentrations of hydrogen peroxide during acute exposures without adverse consequences for the cell, once H₂O₂ is removed. Even though 3 mM concentrations of H₂O₂ will eventually kill during prolonged exposures (40), the H₂O₂ concentrations that kill cells within minutes start around 30 mM (24, 41). This creates a classic engineering problem: besides the obvious caveat that such high H₂O₂ concentrations will be dangerous to the producing cell itself, no known cells are actually capable of producing a short burst of 30 mM H₂O₂, or of maintaining a several-hour 3 mM levels of H₂O₂ (perhaps with the exception of lactobacilli (42)).

In fact, for our leucocytes, the real problem appears several orders of magnitude greater. Due to the abovementioned substantial H₂O₂ permeability, it is almost impossible to achieve a significantly higher concentration of H₂O₂ in any cellular compartment relative to the rest of the cytoplasm, especially if its production is slow and indirect, which is thought to be the case in the phagosome (43, 44). That is, if a leucocyte targets superoxide production (that generates H₂O₂ by dismutation) exclusively to the phagosome around a captured bacterium, the continuous escape of H₂O₂ from the phagosome will keep the maximal H₂O₂ concentration ~3 μM (43, 45, 46), about 1,000 times lower than that required for slow killing, not to mention the fast killing. So, how do the cells solve this problem of creating lethal local concentrations of a readily-diffusible toxin?

Potentiated toxicity of H₂O₂

Theoretically, this engineering problem has an elegant chemical solution. Indeed, the cellular systems that negate H₂O₂ effectiveness (for example, H₂O₂ scavengers or DNA repair pathways) could be inactivated with a different agent, to either increase the effective intracellular H₂O₂ concentrations or to make DNA damage irreparable. In particular, catalases have heme in their active centers (37), so any simple chemical that binds heme iron tightly (NO, CN, H₂S (47, 48)) will inhibit catalases and thus will reduce the killing concentrations of H₂O₂. In fact, catalases *could have been* the original target of the evolutionary arms race, as the cells have a second hydrogen peroxide scavenging enzyme, called alkylperoxidase, which is effective against low H₂O₂ concentrations and that has reaction chemistry very different from the chemistry of catalase (49), meaning that the same inhibition principle will not work against both enzymes. Catalase poisoning has been offered to explain potentiation of H₂O₂ toxicity by other chemicals (Fig. 1), however the inefficiency of this poisoning limits the power of this explanation (41, 50–52). Important for our discussion, though, is the concept of catalase inhibition by a separate agent, which

highlights a general strategy for solving the *in vivo* hydrogen peroxide concentration problem, the strategy of **potentiated toxicity**.

Potentiated toxicity is one of the two types of a more general phenomenon of **synergistic toxicity**, when two agents in sublethal concentrations that do not kill individually, efficiently kill when used together (53–55). Even though synergistic toxicity conceptually sounds like the bio-analog of the principle behind binary weapons (projectiles loaded with two relatively non-toxic chemicals, which combine into a potent toxin during their short flight toward the target (56)), the mechanisms behind synergistic toxicity are different. One of the two basic scenarios of how two biotoxins synergize is *redundancy*, when both toxins target the same essential cellular process, and their combined action overwhelms the robustness limit of this process, producing “co-toxicity”. According to the co-toxicity rules, either agent alone in high enough concentrations is expected to be toxic. In contrast, the second basic scenario of biotoxin synergy, *potentiated toxicity*, envisions that one of the agents is not capable of killing the cells in any concentration, but inactivates an antidote against the second agent, which is the real toxin (41, 57, 58). A classic example from everyday life is clavulanic acid, an inhibitor of beta-lactamase, that potentiates the action of amoxicillin against resistant bacteria without being toxic itself (59). The helpful aspect of H₂O₂ toxicity potentiation is that if the second agent is a charged molecule, it will not be able to easily cross the membrane barrier and escape from the phagosome, where the cell tries to accumulate H₂O₂. In other words, the concentration of H₂O₂ may be only ten times higher inside the phagosome than outside of it, but it is the concentration of the charged potentiating agent inside the phagosome that will make H₂O₂ toxic only within the compartment. Thus, potentiated toxicity not only dramatically reduces the killing concentrations of the toxin that the cell has to produce, but also provides the means to target specific cellular compartments.

How NO potentiates H₂O₂ toxicity

Of the several examples of H₂O₂ toxicity potentiation by other simple chemicals, the best known case is observed in our immune cells that kill infecting microbes with a mixture of hydrogen peroxide and nitric oxide (NO) (28–30). NO is bacteriostatic for wild type bacteria (28, 60, 61), inhibiting their general metabolism by binding to heme iron and thus blocking respiration (62, 63). For targeted NO production, our leucocytes have a phagosome-associated nitric oxide synthase (64); together with (phox) NADPH oxidase that generates superoxide (65) (which then dismutates to H₂O₂ (44)), this produces the potentially-toxic NO + H₂O₂ mix only within the phagosome.

The mechanism of NO potentiation of H₂O₂ toxicity turns out to be complex, as NO affects many enzymes and processes. By readily binding to heme iron, NO inhibits catalases (66, 67), increasing the effective intracellular H₂O₂ concentrations (52, 68) (Fig. 1, scenario #1). In addition, NO binding to cytochrome oxidases inhibits respiration (69–71) decreasing the cellular ATP pools, which should inhibit any repair of complex DNA lesions that requires ATP hydrolysis (72, 73). Finally, NO directly inhibits Fpg DNA glycosylase, that removes the major types of oxidized bases from DNA (74, 75), which should further increase susceptibility of NO-treated cells to oxidative DNA damage. Thus, NO could potentiate

H₂O₂ toxicity by interfering both with H₂O₂ scavenging and with subsequent repair of oxidative DNA lesions (Fig. 1).

Investigation into the metabolic underpinnings of NO + H₂O₂ co-toxicity in *E. coli* has substantially broadened the picture, though. First, iron chelation blocks the NO + H₂O₂ co-toxicity (30, 76), indicating participation of free intracellular iron. It turns out that NO-induced respiration inhibition causes accumulation of NADH and, indirectly, greatly expands the pool of reduced flavin mononucleotides (FMNs) (76), used by iron-siderophore reductases to recycle insoluble Fe(III) back into soluble Fe(II) (77), which promotes Fenton's reaction (Fig. 1, scenario #2). Remarkably, the proposed mechanism (76) expands the emphasis of NO potentiation of the H₂O₂ toxicity from simple interference with H₂O₂ scavenging and/or repair of oxidative DNA damage to the increased availability of the second component of Fenton's reaction, the free reduced iron (Fig. 1). The mechanistic diversification of the potentiation phenomenon makes it robust and efficient.

Some other common potentiators of H₂O₂

A few other simple chemicals potentiate H₂O₂ toxicity in both bacterial and mammalian cells. Ascorbic acid (AA) potentiates H₂O₂ toxicity, both *in vivo* (78–81) and *in vitro* (82). The enhancement of the intracellular oxidative potential of H₂O₂ by AA can be blocked by intracellular iron chelators, suggesting that AA increases the pools of intracellular free iron (83). Curiously, many cancer cell lines are sensitive to extracellular AA concentrations innocuous for normal cells, and this sensitivity is relieved by extracellular catalase, indicating the AA + H₂O₂ co-toxicity as the underlying cause of their sensitivity (84).

Amino acid L-histidine potentiates H₂O₂ toxicity (85–88) (reviewed in (89)). The His + H₂O₂ co-toxicity is associated with formation of double-strand DNA breaks and is blocked by iron chelation (86, 87, 90, 91). Histidine also potentiates pure DNA nicking by Fenton's reaction *in vitro* (91, 92).

L-cysteine, the only redox-active amino acid, also potentiates hydrogen peroxide toxicity (85, 93, 94). Cystine, the oxidized dimeric form of cysteine, also potentiates H₂O₂ toxicity, but in a transient way, by temporarily increasing the intracellular cysteine levels (93–95). Cysteine potentiation is blocked by *in vivo* chelation of iron (93, 94). Cysteine also potentiates pure DNA nicking by Fenton's reaction *in vitro* (94).

Potentiation by cysteine may work via its intracellular conversion into hydrogen sulfide (H₂S) (85), which is another known potentiator of cell killing by H₂O₂ (50, 51, 85). Treatment with hydrogen sulfide inhibits cellular catalase, suggesting the obvious mechanism of its potentiation of H₂O₂ toxicity (50, 51) (Fig. 1). As with NO or L-Cys above, intracellular iron chelation abolishes H₂S + H₂O₂ co-toxicity, suggesting that H₂S enhances the availability of free intracellular iron (85).

Recently, the findings that NO (96) and H₂S (97) protect *E. coli* against antibiotics were interpreted in terms of protection against oxidative damage, based on another recent proposal that lethal oxidative damage is the common mechanism behind antibiotic action (98). However, this review presents enough evidence that both NO and H₂S *potentiate* H₂O₂

poisoning, rather than reduce it, in contrast to the prediction from NO or H₂S protection against antibiotics (96, 97). In light of the recent strong evidence of no oxidative damage associated with antibiotic treatment (41, 99), the NO and H₂S protection most likely works by slowing down cellular metabolism, which would make bacterial cells more tolerant to a broad spectrum of cell-damaging treatments.

The paradoxes and complexities of CN potentiation of H₂O₂ toxicity

The emerging pattern of H₂O₂ potentiation, while leaving aside the importance of control over the intracellular H₂O₂ concentration, does emphasize the role of available iron. A recent study of yet another strong H₂O₂ potentiator, cyanide (25, 41), while further solidifying the critical role of iron, brings back into focus stable DNA-iron interactions and highlights yet another important aspect of this multifaceted toxicity — timely DNA repair.

Like NO above, cyanide (CN) is a “classic” strong-field isoelectronic diatomic ligand for heme iron and a strong potentiator of H₂O₂ poisoning (100). Investigation of the mechanisms of CN potentiation of H₂O₂ toxicity (101) was a trailblazer for the subsequent study of NO potentiation (76). By inhibiting heme-containing enzymes (the preferred target of CN binding) and thus blocking respiration, CN was found to increase the intracellular pool of NADH, which would translate into a higher pool of reduced FMNs and would, therefore, elevate the pools of free soluble Fe(II) iron (101). In fact, this general scenario would work for all known H₂O₂ potentiators, since NO, CN, H₂S, as well as the amino acids cysteine and histidine, are all heme ligands (47, 62, 102–104). Ascorbate is the only exception in this list, but it acts as an electron donor for some heme-containing enzymes (105), and so could be considered a “transient ligand”.

However, equating H₂O₂ potentiators with heme ligands creates a paradox, as the best diatomic competitor of O₂ for binding heme iron is the notorious carbon monoxide (CO) (48, 102), which, according to the hypothesis above, should be among the strongest potentiators of H₂O₂ toxicity. To the contrary, CO is known to protect mammalian cells against H₂O₂ toxicity (106–111). In other words, binding to heme iron may be a common property of the H₂O₂ potentiators, but it does not have to be the mechanism of potentiation.

In fact, our recent investigation of CN + H₂O₂ co-toxicity was inconsistent with the respiration block scenario of H₂O₂ toxicity potentiation, although we did confirm the involvement of reduced FMNs in the process (41). Instead, we found evidence for a specific multi-stage scenario (Fig. 2): 1) cyanide recruits (releases) iron from intracellular iron depots, ferritin being one such iron donor; 2) the recruited (released) iron is somehow delivered directly to DNA and forms stable iron-DNA complexes; 3) these DNA-bound iron atoms, if not promptly removed (for example, by the ferritin-like Dps proteins), catalyze ROS-generating Fenton’s reaction right on the most sensitive target of the cell, the chromosomal DNA; 4) cyanide further potentiates this DNA-self-targeted Fenton’s reaction in an unknown way; 5) expeditious repair of the resulting single-strand breaks and lesions (with which CN may also interfere) somehow facilitates iron removal from DNA, giving DNA one more chance to avoid imminent double-strand breaks, — otherwise catastrophic

chromosomal fragmentation ensues (25, 41). Below we discuss some aspects of CN potentiation of H₂O₂ toxicity.

Iron depots and lockups, stable iron-DNA complexes, and ligand-dependent Fenton's variations

Ferritins are 24-subunit thick-walled spheres that manage the levels of cytoplasmic Fe(II) iron using molecular oxygen and depositing excess iron as crystals of Fe(III) oxide (15, 112). At the same time, ferritins not only can accumulate up to 4,500 Fe(III) atoms, but they also release the sequestered iron on specific cues to proteins that are to be iron-metallated (15, 112), functioning as iron-depot proteins. Due to ferritin's general affinity for DNA (113–115), cyanide-compromised ferritins should release iron (Fe(II)) in the vicinity of DNA, as a result, stable DNA-Fe(II) complexes will form (116). In the presence of H₂O₂, stable DNA-iron complexes become the sites of formation of hydroxyl radicals on chromosomal DNA. To emphasize the role of these stable DNA-iron complexes in H₂O₂-promoted DNA demise, we introduced the term “DNA-self-targeted Fenton's reaction”, which describes the unique susceptibility of nucleic acids to iron-catalyzed reactive oxygen species.

Ferritin homologs Dps proteins of bacteria are called mini-ferritins, because they assemble into much smaller 12-subunit spheres, packing only 500 Fe(III) atoms (15, 112). Dps mini-ferritins are important for bacterial DNA protection against oxidative damage, especially in stationary cells: in contrast to “maxi-ferritins” that oxidatively deposit iron using O₂, mini-ferritins use H₂O₂ for this purpose. In effect, when H₂O₂ appears in the bacterial cytoplasm, mini-ferritins sequester all free iron, functioning as “iron-lockup” during H₂O₂ stress. Since, in addition, Dps proteins have a high affinity for DNA (117), mini-ferritins represent a perfect “sponge” for mopping up iron from DNA-iron complexes in the presence of H₂O₂. At the same time, Dps does not appear to be the predominant iron storage in stationary cells, as at least half of iron in stationary *E. coli* is still found in ferritin (118). The complete resistance of stationary cells to CN + H₂O₂ treatment could be due to either the efficient lockup of the released iron by Dps, or due to DNA binding by this protein overproduced in stationary phase; the sensitivity of the stationary *dps* mutant cells to CN + H₂O₂ treatment is consistent with either idea (41).

Presumably, a low level of free iron binding to DNA and removal from DNA by specialized proteins happens in the cell continuously (82, 119, 120). Iron has a high affinity to DNA and binds it tightly in solutions. Two binding sites for iron on DNA, one at the backbone, while the other at the bases, have been proposed (121–123). In fact, it was suggested on multiple occasions that when iron has a freedom to bind DNA in the cell, the bulk of lethal oxidative DNA lesions comes from such stable DNA-iron complexes (25, 124–128). Remarkably, the rate of formation of oxidative lesions at such stable DNA-iron complexes in cells, apparently limited by the very low intracellular concentration of H₂O₂, is slow enough to allow DNA repair to expedite iron removal from the damaged DNA, effectively eliminating formation of double-strand breaks. At the same time, when the concentration of H₂O₂ increases during the treatment, it takes only a couple of minutes for such stable DNA-iron complexes to,

figuratively speaking, “burn through” DNA, breaking both strands in the same location (25) (Fig. 2).

The proposed new model of H₂O₂ toxicity potentiation needs rigorous experimental testing, as cyanide by itself is a strong iron ligand (129), and the resulting ferrocyanide complexes, especially ferric ferrocyanide (“Prussian blue”) (130), are notoriously inert compared to the moderately reactive ferrous iron. Because of this, for example, cyanide completely blocks a classic *in vitro* Fenton’s reaction that measures decomposition of an organic dye by hydroxyl radicals (41). At the same time, similar to cysteine (94) or histidine (91, 92), cyanide further accelerates Fenton’s reaction with DNA strand scission as a read-out (41), demonstrating that, in complexes with DNA, iron reacts to cyanide differently from iron in solution lacking DNA. It could be that the stable DNA-iron complexes (121, 122, 131) have a configuration resistant to CN chelation, but capable of accepting CN potentiation of the unique DNA self-targeted Fenton’s reaction.

Conclusion

The truly multifaceted mechanism of CN potentiation of H₂O₂ toxicity (comprising all the processes listed in Figures 1 and 2) makes the resulting cell killing robust and unavoidable, but is it applicable to other potentiators of H₂O₂? Remarkably, the CN route of H₂O₂ potentiation via iron Fe(II) recruitment from ferritins (41) is likely shared with other H₂O₂ potentiators, which are also known to release iron from ferritin, including NO (132), H₂S (133), ascorbate (134, 135), cysteine (135, 136) and flavin mononucleotides (135, 137, 138). In other words, iron recruitment from ferritins may represent a general strategy, rather than a peculiarity of CN potentiation of H₂O₂ toxicity. Moreover, two or more different potentiators could work together, for example, in the same cellular compartment (phagosome), enhancing each other’s action. Such a co-potentiation between NO and CN has been already reported before in *E. coli* cultures and was proposed to occur in our immune cells (60). By pumping several different membrane-impermeable potentiators into the same compartment (phagosome), the cell would be able to significantly reduce the killing concentrations of the poison, H₂O₂, in this compartment.

Potentiated toxicity is a unique phenomenon of one agent inducing a non-lethal change in the cellular metabolism that renders cells vulnerable to lethal poisoning by a different agent and can be used to design two-component biocides to fight infections and cancer. It is a part of the more general phenomenon of synergistic toxicity, when two agents, both in safely sublethal concentrations when used alone, cooperate to cause massive lethality when applied together. The phenomenon of synergistic toxicity has a genetic counterpart in the phenomenon of synthetic lethality, in which a combination of two mutations with small individual effects proves nonviable, revealing a gap in our understanding of cell’s metabolism (139–141).

Synthetic lethality is a highly sought-after phenomenon, because isolation of its suppressors (a third mutation that makes the combination viable again) frequently provides a mechanistic understanding of the lethality (142, 143). Similar to synergistic toxicity, there are two general explanations for synthetic lethality (143). The redundancy explanation posits that the

products of the two inactivated genes perform essentially the same enzymatic function, removing a poisonous substrate or synthesizing an essential product. In this sense, the two missing functions are “metabolically proximal”, and the suppressors of synthetic lethality all act to restore the missing function, typically by activating a cryptic activity that functionally overlaps the two mutated activities. In contrast, the potentiation explanation (also known as “defect-damage-repair cycle” (143) or “avoidance-repair couple”) posits that one mutation inactivates a damage-*avoidance* function, while the other mutation inactivates the damage-*repair* function. The two inactivated functions in this case perform completely unrelated, “metabolically distal” enzymatic reactions, and the suppressors of synthetic lethality in this case inactivate the gene(s) responsible for poisoning.

Synthetic lethality is considered to add a second dimension to an otherwise linear mutation analysis, while isolation of suppressors of synthetic lethals makes genetic analysis three-dimensional. Similarly, the phenomenon of synergistic toxicity adds another dimension to the mechanisms of cell poisoning. In the future, it should be instructive to apply a genetic dimension to the analysis of co-toxicity, by isolating resistant mutants.

Acknowledgments

We would like to thank Glen Cronan for enthusiastic discussion of our results and for copyediting. Experimental research in this laboratory was supported by grant # GM 073115 from the National Institutes of Health.

References

1. Wood PM. The potential diagram for oxygen at pH 7. *Biochem J.* 1988; 253:287–289. [PubMed: 2844170]
2. Imlay JA. Pathways of oxidative damage. *Annu Rev Microbiol.* 2003; 57:395–418. [PubMed: 14527285]
3. Eberhardt, MK. Reactive Oxygen Metabolites – Chemistry and Medical Consequences. CRC Press; Boca Raton, FL: 2001. p. 608
4. Olinescu, R., Smith, T. Free Radicals in Medicine. Nova Science Publishers, Inc.; Huntington, NY: 2002. p. 195
5. Pryor WA. Oxy-radicals and related species: their formation, lifetimes, and reactions. *Annu Rev Physiol.* 1986; 48:657–667. [PubMed: 3010829]
6. Antelmann H, Helmann JD. Thiol-based redox switches and gene regulation. *Antioxid Redox Signal.* 2011; 14:1049–1063. [PubMed: 20626317]
7. Dittmar HR, Baldwin IL, Miller SB. The Influence of Certain Inorganic Salts on the Germicidal Activity of Hydrogen Peroxide. *J Bacteriol.* 1930; 19:203–211. [PubMed: 16559422]
8. Rutala, WA., Weber, DJ., H. I. C. P. A. C. (HICPAC). Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008. Centers for Disease Control and Prevention; Atlanta, GA: 2008.
9. Andreini C, Bertini I, Cavallaro G, Holliday GL, Thornton JM. Metal ions in biological catalysis: from enzyme databases to general principles. *J Biol Inorg Chem.* 2008; 13:1205–1218. [PubMed: 18604568]
10. Crichton RR, Pierre JL. Old iron, young copper: from Mars to Venus. *Biometals.* 2001; 14:99–112. [PubMed: 11508852]
11. Gonzalez-Davila M, Santana-Casiano JM, Millero FJ. Oxidation of iron (II) nanomolar with H₂O₂ in seawater. *Geochim Cosmochim Acta.* 2005; 69:83–93.
12. Liu X, Millero FJ. The solubility of iron in seawater. *Mar Chem.* 2002; 77:43–54.
13. Wells ML, Price NM, Bruland KW. Iron chemistry in seawater and its relationship to phytoplankton: a workshop report. *Marine Chem.* 1995; 48:157–182.

14. Xing W, Liu G. Iron biogeochemistry and its environmental impacts in freshwater lakes. *Fresenius Environ Bull.* 2011; 20:1339–1345.
15. Andrews SC, Robinson AK, Rodríguez-Quiriones F. Bacterial iron homeostasis. *FEMS Microbiol Rev.* 2003; 27:215–237. [PubMed: 12829269]
16. Ganz T. Systemic iron homeostasis. *Physiol Rev.* 2013; 93:1721–1741. [PubMed: 24137020]
17. Hartmann A, Braun V. Iron uptake and iron limited growth of *Escherichia coli* K-12. *Arch Microbiol.* 1981; 130:353–356. [PubMed: 7034667]
18. Outten CE, O'Halloran TV. Femtomolar sensitivity of metalloregulatory proteins controlling zinc homeostasis. *Science.* 2001; 292:2488–2492. [PubMed: 11397910]
19. Fenton HJH. The oxidation of tartaric acid in presence of iron. *J Chem Soc Proc.* 1894; 10:157–158.
20. Koppenol WH. The centennial of the Fenton reaction. *Free Rad Biol Med.* 1993; 15:645–651. [PubMed: 8138191]
21. Winterbourn CC. Toxicity of iron and hydrogen peroxide: the Fenton reaction. *Toxicol Lett.* 1995; 82–83:969–974.
22. Buxton GV, Greenstock CL, Helman WP, Ross AB. Critical Review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot\text{OH}/\cdot\text{O}$) in aqueous solution. *J Phys Chem Ref Data.* 1988; 17:513–886.
23. Ananthaswamy HN, Eisenstark A. Repair of hydrogen peroxide-induced single-strand breaks in *Escherichia coli* deoxyribonucleic acid. *J Bacteriol.* 1977; 130:187–191. [PubMed: 323227]
24. Imlay JA, Linn S. Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. *J Bacteriol.* 1986; 166:519–527. [PubMed: 3516975]
25. Mahaseth T, Kuzminov A. Prompt repair of hydrogen peroxide-induced DNA lesions prevents catastrophic chromosomal fragmentation. *DNA Repair.* 2016; 41:42–53. [PubMed: 27078578]
26. Makino N, Sasaki K, Hashida K, Sakakura Y. A metabolic model describing the H₂O₂ elimination by mammalian cells including H₂O₂ permeation through cytoplasmic and peroxisomal membranes: comparison with experimental data. *Biochim Biophys Acta.* 2004; 1673:149–159. [PubMed: 15279886]
27. Seaver LC, Imlay JA. Hydrogen peroxide fluxes and compartmentalization inside growing *Escherichia coli*. *J Bacteriol.* 2001; 183:7182–7189. [PubMed: 11717277]
28. Brunelli L, Crow JP, Beckman JS. The comparative toxicity of nitric oxide and peroxyntrite to *Escherichia coli*. *Arch Biochem Biophys.* 1995; 316:327–334. [PubMed: 7840633]
29. Klebanoff SJ. Reactive nitrogen intermediates and antimicrobial activity: role of nitrite. *Free Radic Biol Med.* 1993; 14:351–360. [PubMed: 8385644]
30. Pacelli R, Wink DA, Cook JA, Krishna MC, DeGraff W, Friedman N, Tsokos M, Samuni A, Mitchell JB. Nitric oxide potentiates hydrogen peroxide-induced killing of *Escherichia coli*. *J Exp Med.* 1995; 182:1469–1479. [PubMed: 7595217]
31. Kwakman PH, Zaat SA. Antibacterial components of honey. *IUBMB Life.* 2012; 64:48–55. [PubMed: 22095907]
32. Otero MC, Nader-Macías ME. Inhibition of *Staphylococcus aureus* by H₂O₂-producing *Lactobacillus gasseri* isolated from the vaginal tract of cattle. *Anim Reprod Sci.* 2006; 96:35–46. [PubMed: 16386389]
33. Weinberg ED. The *Lactobacillus* anomaly: total iron abstinence. *Perspect Biol Med.* 1997; 40:578–583. [PubMed: 9269745]
34. Bos N, Sundström L, Fuchs S, Freitag D. Ants medicate to fight disease. *Evolution.* 2015; 69:2979–2984. [PubMed: 26283006]
35. Scheler C, Durner J, Astier J. Nitric oxide and reactive oxygen species in plant biotic interactions. *Curr Opin Plant Biol.* 2013; 16:534–539. [PubMed: 23880111]
36. Imlay JA. Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev Biochem.* 2008; 77:755–776. [PubMed: 18173371]
37. Chelikani P, Fita I, Loewen PC. Diversity of structures and properties among catalases. *Cell Mol Life Sci.* 2004; 61:192–208. [PubMed: 14745498]

38. Miller AF. Superoxide dismutases: ancient enzymes and new insights. *FEBS Lett.* 2012; 586:585–595. [PubMed: 22079668]
39. Clark TJ, Shafee T. Wikipedia. 2008; 2015
40. Brudzynski K, Abubaker K, St-Martin L, Castle A. Re-examining the role of hydrogen peroxide in bacteriostatic and bactericidal activities of honey. *Front Microbiol.* 2011; 2 article 213.
41. Mahaseth T, Kuzminov A. Cyanide enhances hydrogen peroxide toxicity by recruiting endogenous iron to trigger catastrophic chromosomal fragmentation. *Mol Microbiol.* 2015; 96:349–367. [PubMed: 25598241]
42. Martín R, Suárez JE. Biosynthesis and degradation of H₂O₂ by vaginal lactobacilli. *Appl Environ Microbiol.* 2010; 76:400–405. [PubMed: 19948869]
43. Winterbourn CC, Hampton MB, Livesey JH, Kettle AJ. Modeling the reactions of superoxide and myeloperoxidase in the neutrophil phagosome: implications for microbial killing. *J Biol Chem.* 2006; 281:39860–39869. [PubMed: 17074761]
44. Winterbourn CC, Kettle AJ. Redox reactions and microbial killing in the neutrophil phagosome. *Antioxid Redox Signal.* 2013; 18:642–660. [PubMed: 22881869]
45. Imlay JA. Oxidative Stress. *EcoSal Plus.* 2009; 3 doi:10.1128/ecosalplus.5.4.4.
46. Slauch JM. How does the oxidative burst of macrophages kill bacteria? Still an open question. *Mol Microbiol.* 2011; 80:580–583. [PubMed: 21375590]
47. Bieza SA, Boubeta F, Feis A, Smulevich G, Estrin DA, Boechi L, Bari SE. Reactivity of inorganic sulfide species toward a heme protein model. *Inorg Chem.* 2015; 54:527–533. [PubMed: 25537304]
48. Milani M, Pesce A, Nardini M, Ouellet H, Ouellet Y, Dewilde S, Bocedi A, Ascenzi P, Guertin M, Moens L, Friedman JM, Wittenberg JB, Bolognesi M. Structural bases for heme binding and diatomic ligand recognition in truncated hemoglobins. *J Inorg Biochem.* 2005; 99:97–109. [PubMed: 15598494]
49. Poole LB. Bacterial defenses against oxidants: mechanistic features of cysteine-based peroxidases and their flavoprotein reductases. *Arch Biochem Biophys.* 2005; 433:240–254. [PubMed: 15581580]
50. Carlsson J, Berglin EH, Claesson R, Edlund MB, Persson S. Catalase inhibition by sulfide and hydrogen peroxide-induced mutagenicity in *Salmonella typhimurium* strain TA102. *Mutat Res.* 1988; 202:59–64. [PubMed: 3054531]
51. Phan TN, Kirsch AM, Marquis RE. Selective sensitization of bacteria to peroxide damage associated with fluoride inhibition of catalase and pseudocatalase. *Oral Microbiol Immunol.* 2001; 16:28–33. [PubMed: 11169136]
52. Rauen U, Li T, Ioannidis I, de Groot H. Nitric oxide increases toxicity of hydrogen peroxide against rat liver endothelial cells and hepatocytes by inhibition of hydrogen peroxide degradation. *Am J Physiol Cell Physiol.* 2007; 292:C1440–1449. [PubMed: 17192286]
53. Fried J, Perez AG, Doblin JM, Clarkson BD. Factors modifying the synergistic toxicity of deoxycytidine in combination with thymidine plus 5-fluorouracil in HeLa cells. *Cell Tissue Kinet.* 1983; 16:539–548. [PubMed: 6627345]
54. Litwin MS, Walter CW, Ejarque P, Reynolds ES. Synergistic toxicity of gram-negative bacteria and free colloidal hemoglobin. *Ann Surg.* 1963; 157:485–493. [PubMed: 13931017]
55. McKenzie D, Brody JI. Synergistic toxicity of triethylene thiophosphoramidate and folic acid in the IRC-741 leukemia. *Cancer Res.* 1963; 23:118–121. [PubMed: 13932078]
56. Pitschmann V. Overall view of chemical and biochemical weapons. *Toxins.* 2014; 6:1761–1784. [PubMed: 24902078]
57. Kuzminov A. Inhibition of DNA synthesis facilitates expansion of low-complexity repeats: is strand slippage stimulated by transient local depletion of specific dNTPs? *BioEssays.* 2013; 35:306–313. [PubMed: 23319444]
58. Son MH, Kang KW, Lee CH, Kim SG. Potentiation of cadmium-induced cytotoxicity by sulfur amino acid deprivation through activation of extracellular signal-regulated kinase1/2 (ERK1/2) in conjunction with p38 kinase or c-jun N-terminal kinase (JNK). Complete inhibition of the potentiated toxicity by U0126 an ERK1/2 and p38 kinase inhibitor. *Biochem Pharmacol.* 2001; 62:1379–1390. [PubMed: 11709198]

59. Brogden RN, Carmine A, Heel RC, Morley PA, Speight TM, Avery GS. Amoxicillin/clavulanic acid: a review of its antibacterial activity, pharmacokinetics and therapeutic use. *Drugs*. 1981; 22:337–362. [PubMed: 7037354]
60. Kuong KJ, Kuzminov A. Cyanide, peroxide and nitric oxide formation in solutions of hydroxyurea causes cellular toxicity and may contribute to its therapeutic potency. *J Mol Biol*. 2009; 390:845–862. [PubMed: 19467244]
61. Spek EJ, Wright TL, Stitt MS, Taghizadeh NR, Tannenbaum SR, Marinus MG, Engelward BP. Recombinational repair is critical for survival of *Escherichia coli* exposed to nitric oxide. *J Bacteriol*. 2001; 183:131–138. [PubMed: 11114909]
62. Borisov VB, Forte E, Giuffrè A, Konstantinov A, Sarti P. Reaction of nitric oxide with the oxidized di-heme and heme-copper oxygen-reducing centers of terminal oxidases: Different reaction pathways and end-products. *J Inorg Biochem*. 2009; 103:1185–1187. [PubMed: 19592112]
63. Mancinelli RL, McKay CP. Effects of nitric oxide and nitrogen dioxide on bacterial growth. *Appl Environ Microbiol*. 1983; 46:198–202. [PubMed: 6351744]
64. Stuehr DJ. Structure-function aspects in the nitric oxide synthases. *Annu Rev Pharmacol Toxicol*. 1997; 37:339–359. [PubMed: 9131257]
65. Nauseef WM. Nox enzymes in immune cells. *Semin Immunopathol*. 2008; 30:195–208. [PubMed: 18449540]
66. Brown GC. Reversible binding and inhibition of catalase by nitric oxide. *Eur J Biochem*. 1995; 232:188–191. [PubMed: 7556149]
67. Brunelli L, Yermilov V, Beckman JS. Modulation of catalase peroxidatic and catalatic activity by nitric oxide. *Free Radic Biol Med*. 2001; 30:709–714. [PubMed: 11275470]
68. Rauen U, Li T, de Groot H. Inhibitory and enhancing effects of NO on H(2)O(2) toxicity: dependence on the concentrations of NO and H(2)O(2). *Free Radic Res*. 2007; 41:402–412. [PubMed: 17454122]
69. Brown GC. Nitric oxide and mitochondrial respiration. *Biochim Biophys Acta*. 1999; 1411:351–369. [PubMed: 10320668]
70. Stevanin TM, Ioannidis N, Mills CE, Kim SO, Hughes MN, Poole RK. Flavohemoglobin Hmp affords inducible protection for *Escherichia coli* respiration, catalyzed by cytochromes bo' or bd, from nitric oxide. *J Biol Chem*. 2000; 275:35868–35875. [PubMed: 10915782]
71. Stevanin TM, Poole RK, Demoncheaux EA, Read RC. Flavohemoglobin Hmp protects *Salmonella enterica* serovar *typhimurium* from nitric oxide-related killing by human macrophages. *Infect Immun*. 2002; 70:4399–4405. [PubMed: 12117950]
72. Krogh BO, Symington LS. Recombination proteins in yeast. *Annu Rev Genet*. 2004; 38:233–271. [PubMed: 15568977]
73. Kuzminov A. Recombinational repair of DNA damage in *Escherichia coli* and bacteriophage λ . *Microbiol Mol Biol Rev*. 1999; 63:751–813. [PubMed: 10585965]
74. Jaiswal M, LaRusso NF, Shapiro RA, Billiar TR, Gores GJ. Nitric oxide-mediated inhibition of DNA repair potentiates oxidative DNA damage in cholangiocytes. *Gastroenterology*. 2001; 120:190–199. [PubMed: 11208728]
75. Wink DA, Laval J. The Fpg protein, a DNA repair enzyme, is inhibited by the biomediator nitric oxide in vitro and in vivo. *Carcinogenesis*. 1994; 15:2125–2129. [PubMed: 7955043]
76. Woodmansee AN, Imlay JA. A mechanism by which nitric oxide accelerates the rate of oxidative DNA damage in *Escherichia coli*. *Mol Microbiol*. 2003; 49:11–22. [PubMed: 12823807]
77. Coves J, Fontecave M. Reduction and mobilization of iron by a NAD(P)H:flavin oxidoreductase from *Escherichia coli*. *Eur J Biochem*. 1993; 211:635–641. [PubMed: 8436123]
78. Ericsson Y, Lundbeck H. Antimicrobial effect in vitro of the ascorbic acid oxidation. I. Effect on bacteria, fungi and viruses in pure cultures. *Acta Pathol Microbiol Scand*. 1955; 37:493–506. [PubMed: 13301729]
79. Jonas SK, Riley PA, Willson RL. Hydrogen peroxide cytotoxicity. Low-temperature enhancement by ascorbate or reduced lipoate. *Biochem J*. 1989; 264:651–655. [PubMed: 2515850]
80. Miller TE. Killing and lysis of gram-negative bacteria through the synergistic effect of hydrogen peroxide, ascorbic acid, and lysozyme. *J Bacteriol*. 1969; 98:949–955. [PubMed: 4892384]

81. Vaughan AT, Gordon DJ. Hydrogen peroxide lethality is associated with a decreased ability to maintain positive DNA supercoiling. *Exp Cell Res.* 1992; 202:376–380. [PubMed: 1397090]
82. Samuni A, Aronovitch J, Godinger D, Chevion M, Czapski G. On the cytotoxicity of vitamin C and metal ions. A site-specific Fenton mechanism. *Eur J Biochem.* 1983; 137:119–124. [PubMed: 6317379]
83. Inai Y, Bi W, Shiraishi N, Nishikimi M. Enhanced oxidative stress by L-ascorbic acid within cells challenged by hydrogen peroxide. *J Nutr Sci Vitaminol.* 2005; 51:398–405. [PubMed: 16521698]
84. McCarty MF, Contreras F. Increasing superoxide production and the labile iron pool in tumor cells may sensitize them to extracellular ascorbate. *Front Oncol.* 2014; 4:249. 0.3389/fonc.2014.00249. [PubMed: 25279352]
85. Berglin EH, Carlsson J. Potentiation by sulfide of hydrogen peroxide-induced killing of *Escherichia coli*. *Infect Immun.* 1985; 49:538–543. [PubMed: 3897055]
86. Cantoni O, Sestili P, Brandi G, Cattabeni F. The L-histidine-mediated enhancement of hydrogen peroxide-induced cytotoxicity is a general response in cultured mammalian cell lines and is always associated with the formation of DNA double strand breaks. *FEBS Lett.* 1994; 353:75–78. [PubMed: 7926027]
87. Sestili P, Cattabeni F, Cantoni O. The L-histidine-mediated enhancement of hydrogen peroxide-induced DNA double strand breakage and cytotoxicity does not involve metabolic processes. *Biochem Pharmacol.* 1995; 50:1823–1830. [PubMed: 8615861]
88. Tachon P, Giacomoni PU. Histidine modulates the clastogenic effect of oxidative stress. *Mutat Res.* 1989; 211:103–109. [PubMed: 2921996]
89. Cantoni O, Giacomoni P. The role of DNA damage in the cytotoxic response to hydrogen peroxide/histidine. *Gen Pharmacol.* 1997; 29:513–516. [PubMed: 9352295]
90. Tachon P. Intracellular iron mediates the enhancing effect of histidine on the cellular killing and clastogenicity induced by H₂O₂. *Mutat Res.* 1990; 228:221–228. [PubMed: 2300070]
91. Tachon P, Deflandre A, Giacomoni PU. Modulation by L-histidine of H₂O₂-mediated damage of cellular and isolated DNA. *Carcinogenesis.* 1994; 15:1621–1626. [PubMed: 8055641]
92. Marrot L, Giacomoni PU. Enhancement of oxidative DNA degradation by histidine: the role of stereochemical parameters. *Mutat Res.* 1992; 275:69–79. [PubMed: 1379340]
93. Berglin EH, Edlund MB, Nyberg GK, Carlsson J. Potentiation by L-cysteine of the bactericidal effect of hydrogen peroxide in *Escherichia coli*. *J Bacteriol.* 1982; 152:81–88. [PubMed: 6749824]
94. Park S, Imlay JA. High levels of intracellular cysteine promote oxidative DNA damage by driving the Fenton reaction. *J Bacteriol.* 2003; 185:1942–1950. [PubMed: 12618458]
95. Cantoni O, Brandi G, Albano A, Cattabeni F. Action of cystine in the cytotoxic response of *Escherichia coli* cells exposed to hydrogen peroxide. *Free Radic Res.* 1995; 22:275–283. [PubMed: 7757202]
96. Gusarov I, Shatalin K, Starodubtseva M, Nudler E. Endogenous nitric oxide protects bacteria against a wide spectrum of antibiotics. *Science.* 2009; 325:1380–1384. [PubMed: 19745150]
97. Shatalin K, Shatalina E, Mironov A, Nudler E. H₂S: a universal defense against antibiotics in bacteria. *Science.* 2011; 334:986–990. [PubMed: 22096201]
98. Kohanski MA, Dwyer DJ, Collins JJ. How antibiotics kill bacteria: from targets to networks. *Nat Rev Microbiol.* 2010; 8:423–435. [PubMed: 20440275]
99. Liu Y, Imlay JA. Cell death from antibiotics without the involvement of reactive oxygen species. *Science.* 2013; 339:1210–1213. [PubMed: 23471409]
100. Imlay JA, Chin SM, Linn S. Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science.* 1988; 240:640–642. [PubMed: 2834821]
101. Woodmansee AN, Imlay JA. Reduced flavins promote oxidative DNA damage in non-respiring *Escherichia coli* by delivering electrons to intracellular free iron. *J Biol Chem.* 2002; 277:34055–34066. [PubMed: 12080063]
102. Conner KP, Woods CM, Atkins WM. Interactions of cytochrome P450s with their ligands. *Arch Biochem Biophys.* 2011; 507:56–65. [PubMed: 20939998]
103. Isaac IS, Dawson JH. Haem iron-containing peroxidases. *Essays Biochem.* 1999; 34:51–69. [PubMed: 10730188]

104. Zhong F, Lisi GP, Collins DP, Dawson JH, Pletneva EV. Redox-dependent stability, protonation, and reactivity of cysteine-bound heme proteins. *Proc Natl Acad Sci U S A*. 2014; 111:E306–315. [PubMed: 24398520]
105. Lu P, Ma D, Yan C, Gong X, Du M, Shi Y. Structure and mechanism of a eukaryotic transmembrane ascorbate-dependent oxidoreductase. *Proc Natl Acad Sci U S A*. 2014; 111:1813–1818. [PubMed: 24449903]
106. Chow JM, Shen SC, Huan SK, Lin HY, Chen YC. Quercetin, but not rutin and quercitrin, prevention of H₂O₂-induced apoptosis via anti-oxidant activity and heme oxygenase 1 gene expression in macrophages. *Biochem Pharmacol*. 2005; 69:1839–1851. [PubMed: 15876423]
107. Cremers NA, Lundvig DM, van Dalen SC, Schelbergen RF, van Lent PL, Szarek WA, Regan RF, Carels CE, Wagener FA. Curcumin-induced heme oxygenase-1 expression prevents H₂O₂-induced cell death in wild type and heme oxygenase-2 knockout adipose-derived mesenchymal stem cells. *Int J Mol Sci*. 2014; 15:17974–17999. [PubMed: 25299695]
108. Han W, Wu L, Chen S, Yu KN. Exogenous carbon monoxide protects the bystander Chinese hamster ovary cells in mixed coculture system after alpha-particle irradiation. *Carcinogenesis*. 2010; 31:275–280. [PubMed: 19945969]
109. Heo JM, Kim HJ, Ha YM, Park MK, Kang YJ, Lee YS, Seo HG, Lee JH, Yun-Choi HS, Chang KC. YS 51, 1-(beta-naphthylmethyl)-6,7-dihydroxy-1,2,3,4-tetrahydroisoquinoline, protects endothelial cells against hydrogen peroxide-induced injury via carbon monoxide derived from heme oxygenase-1. *Biochem Pharmacol*. 2007; 74:1361–1370. [PubMed: 17719563]
110. Lin HY, Shen SC, Lin CW, Yang LY, Chen YC. Baicalein inhibition of hydrogen peroxide-induced apoptosis via ROS-dependent heme oxygenase 1 gene expression. *Biochim Biophys Acta*. 2007; 1773:1073–1086. [PubMed: 17532486]
111. Thom SR, Ischiropoulos H. Mechanism of oxidative stress from low levels of carbon monoxide. *Res Rep Health Eff Inst*. 1997; 80:1–19.
112. Liu X, Theil EC. Ferritins: dynamic management of biological iron and oxygen chemistry. *Acc Chem Res*. 2005; 38:167–175. [PubMed: 15766235]
113. Alkhateeb AA, Connor JR. Nuclear ferritin: A new role for ferritin in cell biology. *Biochim Biophys Acta*. 2010; 1800:793–797. [PubMed: 20347012]
114. Surguladze N, Thompson KM, Beard JL, Connor JR, Fried MG. Interactions and reactions of ferritin with DNA. *J Biol Chem*. 2004; 279:14694–14702. [PubMed: 14734543]
115. Thompson KJ, Fried MG, Ye Z, Boyer P, Connor JR. Regulation, mechanisms and proposed function of ferritin translocation to cell nuclei. *J Cell Sci*. 2002; 115:2165–2177. [PubMed: 11973357]
116. Ouameur AA, Arakawa H, Ahmad R, Naoui M, Tajmir-Riahi HA. A Comparative study of Fe(II) and Fe(III) interactions with DNA duplex: major and minor grooves bindings. *DNA Cell Biol*. 2005; 24:394–401. [PubMed: 15941392]
117. Chiancone E, Ceci P. The multifaceted capacity of Dps proteins to combat bacterial stress conditions: Detoxification of iron and hydrogen peroxide and DNA binding. *Biochim Biophys Acta*. 2010; 1800:798–805. [PubMed: 20138126]
118. Abdul-Tehrani H, Hudson AJ, Chang YS, Timms AR, Hawkins C, Williams JM, Harrison PM, Guest JR, Andrews SC. Ferritin mutants of *Escherichia coli* are iron deficient and growth impaired, and fur mutants are iron deficient. *J Bacteriol*. 1999; 181:1415–1428. [PubMed: 10049371]
119. Enright HU, Miller WJ, Hebbel RP. Nucleosomal histone protein protects DNA from iron-mediated damage. *Nucleic Acid Res*. 1992; 20:3341–3346. [PubMed: 1630905]
120. Henle ES, Linn S. Formation, prevention, and repair of DNA damage by iron/hydrogen peroxide. *J Biol Chem*. 1997; 272:19095–19098. [PubMed: 9235895]
121. Eisinger J, Schulman RG, Szymanski BM. Transition metal binding in DNA solutions. *J Chem Phys*. 1962; 36:1721–1729.
122. Luo Y, Han Z, Chin SM, Linn S. Three chemically distinct types of oxidants formed by iron-mediated Fenton reactions in the presence of DNA. *Proc Natl Acad Sci USA*. 1994; 91:12438–12442. [PubMed: 7809055]

123. Netto LE, Ferreira AM, Augusto O. Iron(III) binding in DNA solutions: complex formation and catalytic activity in the oxidation of hydrazine derivatives. *Chem Biol Interact.* 1991; 79:1–14. [PubMed: 1647885]
124. Chevion M. A site-specific mechanism for free radical induced biological damage: the essential role of redox-active transition metals. *Free Radic Biol Med.* 1988; 5:27–37. [PubMed: 3075945]
125. Gutteridge JM. Copper-phenanthroline-induced site-specific oxygen-radical damage to DNA. Detection of loosely bound trace copper in biological fluids. *Biochem J.* 1984; 218:983–985. [PubMed: 6721843]
126. Lloyd DR, Carmichael PL, Phillips DH. Comparison of the formation of 8-hydroxy-2'-deoxyguanosine and single- and double-strand breaks in DNA mediated by fenton reactions. *Chem Res Toxicol.* 1998; 11:420–427. [PubMed: 9585472]
127. Schweitz H. Dégradation du DNA par H₂O₂ en présence d'ions Cu⁺⁺, Fe⁺⁺ et Fe⁺⁺⁺ Biopolymers. 1969; 8:101–119.
128. Ward JF, Evans JW, Limoli CL, Calabro-Jones PM. Radiation and hydrogen peroxide induced free radical damage to DNA. *Br J Cancer Suppl.* 1987; 8:105–112. [PubMed: 2820457]
129. Griffith WP. Cyanide complexes of the transition metals. *Q Rev Chem Soc.* 1962; 16:188–207.
130. Pearce J. Studies of any toxicological effects of Prussian blue compounds in mammals—a review. *Food Chem Toxicol.* 1994; 32:577–582. [PubMed: 8045465]
131. Floyd RA. DNA-ferrous iron catalyzed hydroxyl free radical formation from hydrogen peroxide. *Biochem Biophys Res Commun.* 1981; 99:1209–1215. [PubMed: 7259775]
132. Reif DW. Ferritin as a source of iron for oxidative damage. *Free Radic Biol Med.* 1992; 12:417–427. [PubMed: 1317328]
133. Cassanelli S, Moulis J. Sulfide is an efficient iron releasing agent for mammalian ferritins. *Biochim Biophys Acta.* 2001; 1547:174–182. [PubMed: 11343803]
134. Boyer RF, McCleary CJ. Superoxide ion as a primary reductant in ascorbate-mediated ferritin iron release. *Free Radic Biol Med.* 1987; 3:389–395. [PubMed: 2828195]
135. Sirivech S, Frieden E, Osaki S. The release of iron from horse spleen ferritin by reduced flavins. *Biochem J.* 1974; 143:311–315. [PubMed: 4462557]
136. Boyer RF, Grabill TW, Petrovich RM. Reductive release of ferritin iron: a kinetic assay. *Anal Biochem.* 1988; 174:17–22. [PubMed: 3218730]
137. Funk F, Lenders JP, Crichton RR, Schneider W. Reductive mobilisation of ferritin iron. *Eur J Biochem.* 1985; 152:167–172. [PubMed: 4043077]
138. Melman G, Bou-Abdallah F, Vane E, Maura P, Arosio P, Melman A. Iron release from ferritin by flavin nucleotides. *Biochim Biophys Acta.* 2013; 1830:4669–4674. [PubMed: 23726988]
139. Guarente L. Synthetic enhancement in gene interaction: a genetic tool come of age. *Trends Genet.* 1993; 9:362–366. [PubMed: 8273152]
140. Hartman JL, Garvik B, Hartwell L. Principles for the buffering of genetic variation. *Science.* 2001; 291:1001–1004. [PubMed: 11232561]
141. Nijman SM. Synthetic lethality: general principles, utility and detection using genetic screens in human cells. *FEBS Lett.* 2011; 585:1–6. [PubMed: 21094158]
142. Budke B, Kuzminov A. Production of clastogenic DNA precursors by the nucleotide metabolism in *Escherichia coli*. *Mol Microbiol.* 2010; 75:230–245. [PubMed: 19943897]
143. Ting H, Kouzminova EA, Kuzminov A. Synthetic lethality with the *dut* defect in *Escherichia coli* reveals layers of DNA damage of increasing complexity due to uracil incorporation. *J Bacteriol.* 2008; 190:5841–5854. [PubMed: 18586941]




Fig. 1. Hydrogen peroxide scavenging by catalases versus “radicalization” by soluble iron (Fe(II)), and the two obvious strategies to potentiate H₂O₂ toxicity

Hydrogen atoms are small gray spheres. Oxygen atoms are spheres of various shades of blue: the darker the blue, the more stable this oxygen atom is. At the top, in black font, the two opposite in vivo fates of hydrogen peroxide are shown as formulas. At the bottom, in red font and symbols, the corresponding potentiation strategies are indicated: 1) inhibition of H₂O₂ decomposition (or DNA repair); 2) increasing the concentration of soluble intracellular iron.




Fig. 2. Cyanide potentiation of hydrogen peroxide toxicity is proposed to happen at least at four separate stages

Small green circles, Fe(II) atoms; small orange circles inside ferritin spheres, Fe(III) atoms. Purple double zig-zag, duplex DNA. The big ferritin depots (purple spheres) release their iron in response to specific signals. Dps lockups (brown-yellow thick-walled spheres) remove iron from the DNA-iron complexes. The distinct opportunities for potentiation, marked by “CN-1, 2, 3, 4” arrows, are: 1) Fe(II) recruitment (“iron release”) from the intracellular iron depots; 2) direct formation of stable DNA-iron complexes; 3) promotion of DNA self-targeted Fenton; 4) Inhibition of one-strand DNA repair.