

# Correction: Immediate Effect of a Single Session of Whole Body Vibration on Spasticity in Children With Cerebral Palsy


Chunung Park, MD, Eun Sook Park, MD, Ja Young Choi, MD,  
Yoona Cho, MD, Dong-wook Rha, MD, PhD

Department of Rehabilitation Medicine and Research Institute of Rehabilitation Medicine,  
Yonsei University College of Medicine, Seoul, Korea


<https://doi.org/10.5535/arm.2017.41.2.273>  
Ann Rehabil Med 2017;41(2):273-278

We noticed that the original version of this paper contains typographical errors in Figs. 2 and 3.

In Fig. 2 (page 275), 'With knee extraction' must be changed into 'With knee extension.'  
In Fig. 3 (page 276), 'With knee extraction' must be changed into 'With knee extension.'  
Corrected versions of both figures appear below.



**Fig. 2.** Changes in Modified Ashworth Scale (MAS) after whole body vibration therapy. \* $p < 0.05$  vs. baseline (post-hoc test).



**Fig. 3.** Changes in Modified Tardieu Scale (MTS) scores after whole body vibration therapy. (A) R1 angle, (B) R2 angle, and (C) R2-R1. \*p<0.05 vs. baseline (post-hoc test).