



# Avibactam Restores the Susceptibility of Clinical Isolates of *Stenotrophomonas maltophilia* to Aztreonam

Maria F. Mojica,<sup>a,b</sup> Krisztina M. Papp-Wallace,<sup>a,b,c</sup> Magdalena A. Taracila,<sup>b,c</sup> Melissa D. Barnes,<sup>b,c</sup> Joseph D. Rutter,<sup>b</sup> Michael R. Jacobs,<sup>d,e</sup> John J. LiPuma,<sup>f</sup> Thomas J. Walsh,<sup>g</sup> Alejandro J. Vila,<sup>h,i</sup> Robert A. Bonomo<sup>a,c,j,k,l</sup>

Department of Biochemistry, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA<sup>a</sup>; Research Service, Louis Stokes Veterans Affairs Medical Center, Cleveland, Ohio, USA<sup>b</sup>; Department of Medicine, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA<sup>c</sup>; Department of Pathology, University Hospitals Cleveland Medical Center, Cleveland, Ohio, USA<sup>d</sup>; Department of Pathology, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA<sup>e</sup>; Department of Pediatrics and Communicable Disease, University of Michigan Medical School, Ann Arbor, Michigan, USA<sup>f</sup>; Transplantation-Oncology Infectious Diseases Program, Weill Cornell Medical Center, New York, New York, USA<sup>g</sup>; Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Rosario, Argentina<sup>h</sup>; Área Biofísica, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina<sup>i</sup>; Departments of Pharmacology, Molecular Biology and Microbiology, Biochemistry, and Proteomics and Bioinformatics, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA<sup>j</sup>; Medical Service and GRECC, Louis Stokes Cleveland Department of Veterans Affairs Medical Center, Cleveland, Ohio, USA<sup>k</sup>; CWRU-Cleveland VAMC Center for Antimicrobial Resistance and Epidemiology (Case VA CARES), Cleveland, Ohio, USA<sup>l</sup>

**ABSTRACT** *Stenotrophomonas maltophilia* is an emerging opportunistic pathogen, classified by the World Health Organization as one of the leading multidrug-resistant organisms in hospital settings. The need to discover novel compounds and/or combination therapies for *S. maltophilia* is urgent. We demonstrate the *in vitro* efficacy of aztreonam-avibactam (ATM-AVI) against *S. maltophilia* and kinetically characterize the inhibition of the L2  $\beta$ -lactamase by avibactam. ATM-AVI overcomes aztreonam resistance in selected clinical strains of *S. maltophilia*, addressing an unmet medical need.

**KEYWORDS** aztreonam, *S. maltophilia*, avibactam

*Stenotrophomonas maltophilia* is a Gram-negative environmental bacillus and an emerging nosocomial pathogen (1). Primarily associated with respiratory tract infections, this bacterium is alarmingly increasing in prevalence among patients with cystic fibrosis, exacerbating an already-compromised respiratory function (2–5). *S. maltophilia* is intrinsically resistant to aminoglycosides, tetracycline, fosfomycin, and the majority of  $\beta$ -lactams.  $\beta$ -Lactam resistance is due to the expression of two  $\beta$ -lactamases: L1 is a B3 metallo- $\beta$ -lactamase (M $\beta$ L) that hydrolyzes all  $\beta$ -lactams with the exception of aztreonam (ATM), and it is resistant to all clinically available  $\beta$ -lactamase inhibitors (1, 6, 7); L2 is a clavulanate-susceptible class A cephalosporinase (1, 8, 9). Mimicking the AmpC cephalosporinases in *Pseudomonas aeruginosa*, L1 and L2 are inducible  $\beta$ -lactamases, and their expression is regulated by a similar mechanism (10, 11).

The growing challenge of treating infections caused by *S. maltophilia* is reflected in increasing reports of acquired resistance to historically effective drugs, like trimethoprim-sulfamethoxazole (SXT), ceftazidime (CAZ), ticarcillin-clavulanate (TIM), and fluoroquinolones, and the documented ability to develop high-level resistance during antibiotic treatment (8, 9, 12, 13). Recently, Mojica et al. described a clinical case in which the avibactam and ceftazidime combination coadministered with ATM (CZA) effectively treated a prolonged bacteremia caused by *S. maltophilia* (13). Notwithstanding the effectiveness of this triple combination, we anticipate that aztreonam-avibactam (ATM-

Received 12 April 2017 Returned for modification 2 May 2017 Accepted 26 July 2017

Accepted manuscript posted online 7 August 2017

**Citation** Mojica MF, Papp-Wallace KM, Taracila MA, Barnes MD, Rutter JD, Jacobs MR, LiPuma JJ, Walsh TJ, Vila AJ, Bonomo RA. 2017. Avibactam restores the susceptibility of clinical isolates of *Stenotrophomonas maltophilia* to aztreonam. Antimicrob Agents Chemother 61:e00777-17. <https://doi.org/10.1128/AAC.00777-17>.

**Copyright** © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Robert A. Bonomo, Robert.Bonomo@va.gov.

**TABLE 1** Steady-state kinetic parameters of L2 with NCF and AVI<sup>a</sup>

Parameter	Value
NCF $K_m$ ( $\mu\text{M}$ )	$62 \pm 4$
NCF $k_{\text{cat}}/K_m$ ( $\mu\text{M}^{-1} \cdot \text{s}^{-1}$ )	$9.14 \pm 0.01$
AVI $K_i$ app ( $\mu\text{M}$ )	$0.66 \pm 0.07$
AVI $k_2/K$ ( $\text{M}^{-1} \cdot \text{s}^{-1}$ )	$47,000 \pm 131$
AVI $k_{\text{off}}$ ( $\text{s}^{-1}$ )	$0.0015 \pm 0.0001$
AVI $k_{\text{off}} t_{1/2}$ (min)	$4.0 \pm 0.2$

<sup>a</sup>NCF, nitrocefin. Values reported are averages  $\pm$  standard deviations from triplicate experiments.


AVI), the planned commercial preparation already in clinical development (ClinicalTrials.gov identifier NCT01689207), would be equally effective against *S. maltophilia*. This dual combination is highly active against *Enterobacteriaceae* producing M $\beta$ Ls (14). Accordingly, we asserted that AVI would inhibit L2, while ATM would bypass L1 to reach its likely target (penicillin-binding protein 3 [PBP-3]).

To test our assertion, we used the reference strain ATCC 51331 (A1) and 27 clinical isolates of *S. maltophilia* from University Hospitals (Cleveland, OH) and from the collection of the *Burkholderia cepacia* Research Laboratory and Repository at the University of Michigan (Ann Arbor, MI). The heterogeneity of these isolates was revealed by pulsed-field electrophoresis (data not shown). The susceptibilities of the isolates were determined by agar dilution MICs according to the Clinical and Laboratory Standards Institute (CLSI) guidelines, (15). Where breakpoints for *S. maltophilia* were not available, those for *Pseudomonas aeruginosa* were used. Table S1 in the supplemental material shows that the ATM-AVI combination fully restored ATM susceptibility in 23/28 (82%) isolates and lowered the MICs to intermediate in 3/5 of the remaining isolates.

Employing analytical isoelectric focusing (aIEF) with a nitrocefin overlay,  $\beta$ -lactamase induction was visualized on gels using crude extract of cells grown with and without 10  $\mu\text{g}/\text{ml}$  imipenem, a known inducer (11). For this experiment, 10 isolates were selected based on their MICs to ATM-AVI (7 susceptible, 2 intermediates, and 1 highly resistant). Three of these 10 strains tested (A2, C1, and E1) were derepressed and demonstrated  $\beta$ -lactamase activity in the absence of imipenem (Fig. S1). We assert that these strains could possess alterations (e.g., amino acid substitutions) in one or more of the proteins involved in the  $\beta$ -lactamase regulatory pathway. However, a direct correlation between derepression of  $bla_{L1}$  or  $bla_{L2}$  and nonsusceptibility to ATM-AVI was not observed. Therefore, to test if ATM-AVI can induce  $\beta$ -lactamase expression, isolate A1 was grown in the presence of either 10  $\mu\text{g}/\text{ml}$  imipenem, 4  $\mu\text{g}/\text{ml}$  avibactam, 16  $\mu\text{g}/\text{ml}$  aztreonam, or 4  $\mu\text{g}/\text{ml}$  avibactam plus 0.5  $\mu\text{g}/\text{ml}$  aztreonam. Interestingly, after 2 h of exposure,  $\beta$ -lactamase expression was induced by imipenem but not by ATM-AVI (Fig. S2). Since L1 activity was not detected by aIEF, even with the addition of 50  $\mu\text{M}$   $\text{ZnSO}_4$  to the nitrocefin overlay, we tested the specific activities of L1 and L2 in the crude lysates. In accordance with the aIEF assay, we found that the activities of both enzymes are enhanced only in lysates obtained after exposure to imipenem (Fig. S3). Taken together, these results suggest that neither ATM, AVI, nor the ATM-AVI combination induces  $\beta$ -lactamase expression.

Steady-state kinetics and electrospray ionization mass spectrometry (ESI-MS) reveal that AVI competitively and reversibly inhibits L2 (Table 1). The carbamylation rates ( $k_2/K$ ) for L2 are comparable with those published for the AVI inactivation of another class A  $\beta$ -lactamase, KPC-2 ( $47,000 \pm 131 \text{ M}^{-1} \cdot \text{s}^{-1}$  for L2 versus  $13,000 \pm 100 \text{ M}^{-1} \cdot \text{s}^{-1}$  for KPC-2 [16]). However, the inhibition of these two enzymes by AVI differs in the decarbamylation rate ( $k_{\text{off}}$ ). For KPC, this rate is  $0.00014 \cdot \text{s}^{-1}$ , compared to  $0.0015 \pm 0.0001 \cdot \text{s}^{-1}$  for L2. Thus, despite similar AVI carbamylation rates for KPC-2 and L2, the AVI  $k_{\text{off}}$  rate of L2 is greater than that of KPC-2. Given the high on-rate of AVI for L2, this higher off-rate most likely will not be clinically significant. In other words, even when AVI decarbamylates off L2, the reformed active AVI will carbamylate L2 rapidly; this is supported by our ESI-MS data.

ESI-MS was used to determine if any intermediates (e.g., desulfation) were formed upon incubation of  $\beta$ -lactamases with AVI (16). Observations taken at



**FIG 1** Avibactam is a potent inhibitor of L2. (A and B) ESI-MS spectrum of unreacted L2 (170 nM) (A) and 1:1 molar ratio of L2 and AVI (170 nM) (B) incubated at room temperature in 10 mM phosphate-buffered saline (PBS) (pH 7.4) for 15 s. Spectra obtained after 1-min, 15-min, 1-h, and 24-h incubations of L2 and AVI were identical to those in panel B. (C) Molecular docking of L2 and AVI. The L2 crystal structure (PDB no. 1N40) is represented as a gray ribbon; AVI is displayed as sticks colored by heteroatoms (pink, carbon; red, oxygen; blue, nitrogen; yellow, sulfur). (D) Interactions of AVI with L2. Interacting residues are displayed in gray, while AVI is represented by pink. Heteroatoms are colored as described before. Interacting waters are shown in blue with their corresponding hydrogen bonds displayed as blue dashed lines; distances between important atoms are highlighted by yellow arrows; hydrogen bonds between residues are indicated by green dashed lines.

several time points (i.e., 15 s, 1 min, and 1 h) reveal a mass of  $29,549 \pm 5$  Da, which corresponds to the carbamylated enzyme (L2 [29,282 Da] plus AVI [265 Da]) (Fig. 1A and B). In addition, the AVI-L2 complex is stable for  $\geq 24$  h, without desulfation or hydrolysis observed. With L1 and AVI, only the apo-enzyme of L1 (29,282 Da) was detected by ESI-MS after a 24-h incubation with AVI, consistent with the lack of inhibition observed for L1-mediated nitrocefin hydrolysis in the presence of up to  $500 \mu\text{M}$  AVI (data not shown).

The *in silico* molecular model of AVI covalently bound with L2 (Fig. 1C and D) suggests that AVI preserves key interactions with conserved residues in class A  $\beta$ -lactamases (16–18). We propose that the C-7 carbamoyl of AVI is present in the oxyanion hole and forms hydrogen bonds (H-bonds) with S70:N and S237:N. Moreover, the carbonyl group of the AVI carboxamide makes H-bonds with N132. The highly polar  $\text{SO}_4$  is proposed to interact with T235, R244, and S237 (forming water-mediated interactions), and possibly with K234 (distance of  $\approx 3.1 \text{ \AA}$ ). These interactions result in a highly stable complex, as observed via ESI-MS.

Given the higher  $k_{\text{off}}$  value (compared to KPC-2), we further assessed our model to identify potential pathways that could lead to recyclization. Recyclization is predicted to be initiated by K73 and S130 via a proton shuttle, resulting in a nucleophilic attack on the carbamate bond by the N-6 of AVI, thus reforming active AVI. In our model, AVI's N-6 is positioned at  $2.8 \text{ \AA}$  away from S130:O and its C-7 is at H-bond distance from a water molecule (close enough for S130 to donate a proton to N-6). Moreover, K73 is within H-bond distance to S70 ( $\approx 2.4 \text{ \AA}$  from S70:O $\delta$ ), and S130 is  $\approx 2.5 \text{ \AA}$  away from S70:O. Thus, a possible proton shuttle pathway for recyclization is present. In addition, the positioning of key catalytic residues in the active site may contribute to the faster recyclization of AVI observed with the L2 enzyme (Fig. S4).

In summary, AVI is a potent inhibitor of L2, and AZT-AVI is an effective *in vitro* combination against multidrug-resistant (MDR) *S. maltophilia*. This combination suggests that a potential therapeutic option can be tested against this MDR threat.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AAC.00777-17>.

**SUPPLEMENTAL FILE 1**, PDF file, 2.0 MB.

## ACKNOWLEDGMENTS

We thank Allergan for supplying avibactam powder for this work through an investigator-initiated trial.

The research reported in this publication was supported in part by funds and/or facilities provided by the Cleveland Department of Veterans Affairs, the Veterans Affairs Merit Review Program BX002872 (K.M.P.-W.) and BX001974 (R.A.B.) from the U.S. Department of Veterans Affairs Biomedical Laboratory Research and Development Service of the VA Office of Research and Development, the Geriatric Research Education and Clinical Center VISN 10 (R.A.B.), and the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under awards R21AI114508, R01AI100560, R01AI063517, and R01AI072219 (R.A.B.). This work was also supported by funding from the Cystic Fibrosis Foundation (J.J.L.).

The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, the U.S. Department of Veterans Affairs, or the U.S. government.

## REFERENCES

- Brooke JS. 2012. *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen. *Clin Microbiol Rev* 25:2–41. <https://doi.org/10.1128/CMR.00019-11>.
- Pompilio A, Crocetta V, Ghosh D, Chakrabarti M, Gherardi G, Vitali LA, Fiscarelli E, Di Bonaventura G. 2016. *Stenotrophomonas maltophilia* phenotypic and genotypic diversity during a 10-year colonization in the lungs of a cystic fibrosis patient. *Front Microbiol* 7:1551. <https://doi.org/10.3389/fmicb.2016.01551>.
- Salsgiver EL, Fink AK, Knapp EA, LiPuma JJ, Olivier KN, Marshall BC, Saiman L. 2016. Changing epidemiology of the respiratory bacteriology of patients with cystic fibrosis. *Chest* 149:390–400. <https://doi.org/10.1378/chest.15-0676>.
- Waters V, Yau Y, Prasad S, Lu A, Atenafu E, Crandall I, Tom S, Tullis E, Ratjen F. 2011. *Stenotrophomonas maltophilia* in cystic fibrosis: serologic response and effect on lung disease. *Am J Respir Crit Care Med* 183:635–640. <https://doi.org/10.1164/rccm.201009-1392OC>.
- Junge S, Görlich D, den Reijer M, Wiedemann B, Tümmler B, Ellemunter H, Dübbers A, Küster P, Ballmann M, Koerner-Rettberg C, Große-Onnebrink J, Heuer E, Sextro W, Mainz JG, Hammermann J, Riethmüller J, Graepler-Mainka U, Staab D, Wollschläger B, Szczepanski R, Schuster A, Tegtmeyer FK, Sutharsan S, Wald A, Nofer JR, van Wamel W, Becker K, Peters G, Kahl BC. 2016. Factors associated with worse lung function in cystic fibrosis patients with persistent *Staphylococcus aureus*. *PLoS One* 11:e0166220. <https://doi.org/10.1371/journal.pone.0166220>.
- Crossman LC, Gould VC, Dow JM, Vernikos GS, Okazaki A, Sebahia M, Saunders D, Arrowsmith C, Carver T, Peters N, Adlem E, Kerhornou A, Lord A, Murphy L, Seeger K, Squares R, Rutter S, Quail MA, Rajandream MA, Harris D, Churcher C, Bentley SD, Parkhill J, Thomson NR, Avison MB. 2008. The complete genome, comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol* 9:R74. <https://doi.org/10.1186/gb-2008-9-4-r74>.
- Lin C-W, Huang Y-W, Hu R-M, Chiang K-H, Yang T-C. 2009. The role of AmpR in regulation of L1 and L2  $\beta$ -lactamases in *Stenotrophomonas maltophilia*. *Res Microbiol* 160:152–158. <https://doi.org/10.1016/j.resmic.2008.11.001>.
- Chang YT, Lin CY, Chen YH, Hsueh PR. 2015. Update on infections caused by *Stenotrophomonas maltophilia* with particular attention to resistance mechanisms and therapeutic options. *Front Microbiol* 6:693. <https://doi.org/10.3389/fmicb.2015.00893>.
- Perez F, Adachi J, Bonomo RA. 2014. Antibiotic-resistant Gram-negative bacterial infections in patients with cancer. *Clin Infect Dis* 59:S335–S339. <https://doi.org/10.1093/cid/ciu612>.
- Huang Y-W, Wu C-J, Hu R-M, Lin Y-T, Yang T-C. 2015. Interplay among membrane-bound lytic transglycosylase D1, the CreBC two-component regulatory system, the AmpNG-AmpDI-NagZ-AmpR regulatory circuit, and L1/L2  $\beta$ -lactamase expression in *Stenotrophomonas maltophilia*. *Antimicrob Agents Chemother* 59:6866–6872. <https://doi.org/10.1128/AAC.05179-14>.
- Vadlamani G, Thomas MD, Patel TR, Donald LJ, Reeve TM, Stetefeld J, Standing KG, Vocadlo DJ, Mark BL. 2015. The  $\beta$ -lactamase gene regulator AmpR is a tetramer that recognizes and binds the D-Ala-D-Ala motif of its repressor UDP-N-acetylmuramic acid (MurNAc)-pentapeptide. *J Biol Chem* 290:2630–2643. <https://doi.org/10.1074/jbc.M114.618199>.
- Ryan RP, Monchy S, Cardinale M, Taghavi S, Crossman L, Avison MB, Berg G, van der Lelie D, Dow JM. 2009. The versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nat Rev Microbiol* 7:514–525. <https://doi.org/10.1038/nrmicro2163>.
- Mojica MF, Ouellette CP, Leber A, Becknell MB, Ardura MI, Perez F, Shimamura M, Bonomo RA, Aitken SL, Shelburne SA. 2016. Successful treatment of bloodstream infection due to metallo- $\beta$ -lactamase-producing *Stenotrophomonas maltophilia* in a renal transplant patient. *Antimicrob Agents Chemother* 60:5130–5134. <https://doi.org/10.1128/AAC.00264-16>.
- Biedenbach DJ, Kazmierczak K, Bouchillon SK, Sahn DF, Bradford PA. 2015. *In vitro* activity of aztreonam-avibactam against a global collection of Gram-negative pathogens from 2012–2013. *Antimicrob Agents Chemother* 59:4239–4248. <https://doi.org/10.1128/AAC.00206-15>.
- CLSI. 2012. Performance standards for antimicrobial susceptibility testing: 22nd informational supplement. CLSI document M100-S22. Clinical and Laboratory Standards Institute, Wayne, PA.
- Ehmann DE, Jahić H, Ross PL, Gu R-F, Hu J, Durand-Réville TF, Lahiri S, Thresher J, Livchak S, Gao N, Palmer T, Walkup GK, Fisher SL. 2013. Kinetics of avibactam inhibition against class A, C, and D

- $\beta$ -lactamases. *J Biol Chem* 288:27960–27971. <https://doi.org/10.1074/jbc.M113.485979>.
17. Krishnan NP, Nguyen NQ, Papp-Wallace KM, Bonomo RA, van den Akker F. 2015. Inhibition of *Klebsiella*  $\beta$ -lactamases (SHV-1 and KPC-2) by avibactam: a structural study. *PLoS One* 10:e0136813. <https://doi.org/10.1371/journal.pone.0136813>.
  18. Lahiri SD, Mangani S, Durand-Reville T, Benvenuti M, De Luca F, Sanyal G, Docquier J-D. 2013. Structural insight into potent broad-spectrum inhibition with reversible recyclization mechanism: avibactam in complex with CTX-M-15 and *Pseudomonas aeruginosa* AmpC  $\beta$ -lactamases. *Antimicrob Agents Chemother* 57:2496–2505. <https://doi.org/10.1128/AAC.02247-12>.