





Review

Exploring the Role of RGD-Recognizing Integrins in Cancer

Markus Nieberler ^{1,*},[†], Ute Reuning ^{2,†}, Florian Reichart ³, Johannes Notni ⁴ ,
Hans-Jürgen Wester ⁴, Markus Schwaiger ⁵, Michael Weinmüller ³ , Andreas Räder ³,
Katja Steiger ⁶  and Horst Kessler ^{3,*} 

¹ Department of Oral and Maxillofacial Surgery, University Hospital Rechts der Isar, Technische Universität München, Ismaninger Straße 22, 81679 Munich, Germany

² Clinical Research Unit, Department of Obstetrics & Gynecology, Technische Universität München, Ismaninger Straße 22, 81675 Munich, Germany; ute.reuning@mri.tum.de

³ Institute for Advanced Study and Center for Integrated Protein Science (CIPSM), Department Chemie, Technische Universität München, Lichtenbergstraße 4, 85747 Garching, Germany; florian.fr.reichart@tum.de (F.R.); michael.weinmueller@tum.de (M.W.); andreas.raeder@mytum.de (A.R.)

⁴ Pharmaceutical Radiochemistry, Technische Universität München, Walther-Meißner-Straße 3, 85748 Garching, Germany; johannes.notni@tum.de (J.N.); h.j.wester@tum.de (H.-J.W.)

⁵ Department of Nuclear Medicine, Technische Universität München, Ismaninger Straße 22, 81679 Munich, Germany; markus.schwaiger@tum.de

⁶ Institute of Pathology, Technische Universität München, Trogerstraße 18, 81675 Munich, Germany; katja.steiger@tum.de

* Correspondence: markus.nieberler@tum.de (M.N.); kessler@tum.de (H.K.);
Tel.: +49-89-4140-5916 (M.N.) or +49-163-4714-383 (M.N.); Fax: +49-89-4140-4993 (M.N.)

† These authors contributed equally to this work.

Academic Editor: Helen M. Sheldrake

Received: 6 July 2017; Accepted: 31 August 2017; Published: 4 September 2017

Abstract: Integrins are key regulators of communication between cells and with their microenvironment. Eight members of the integrin superfamily recognize the tripeptide motif Arg-Gly-Asp (RGD) within extracellular matrix (ECM) proteins. These integrins constitute an important subfamily and play a major role in cancer progression and metastasis via their tumor biological functions. Such transmembrane adhesion and signaling receptors are thus recognized as promising and well accessible targets for novel diagnostic and therapeutic applications for directly attacking cancer cells and their fatal microenvironment. Recently, specific small peptidic and peptidomimetic ligands as well as antibodies binding to distinct integrin subtypes have been developed and synthesized as new drug candidates for cancer treatment. Understanding the distinct functions and interplay of integrin subtypes is a prerequisite for selective intervention in integrin-mediated diseases. Integrin subtype-specific ligands labelled with radioisotopes or fluorescent molecules allows the characterization of the integrin patterns in vivo and later the medical intervention via subtype specific drugs. The coating of nanoparticles, larger proteins, or encapsulating agents by integrin ligands are being explored to guide cytotoxic reagents directly to the cancer cell surface. These ligands are currently under investigation in clinical studies for their efficacy in interference with tumor cell adhesion, migration/invasion, proliferation, signaling, and survival, opening new treatment approaches in personalized medicine.

Keywords: RGD-recognizing integrins; $\alpha v \beta 3$; $\alpha v \beta 5$; $\alpha v \beta 6$; $\alpha v \beta 8$; $\alpha 5 \beta 1$; $\alpha 8 \beta 1$; integrin adhesion; migration; apoptosis; and signaling; synthetic integrin ligands; cyclic peptide; peptidomimetics; Cilengitide; epithelial-mesenchymal transition (EMT); transforming growth factor- β (TGF- β); metastasis; angiogenesis

1. Introduction

Cellular recognition of the interstitial neighborhood and microenvironment is required within all organisms, which depends on cell-extracellular matrix (ECM) and cell-cell communications for the correct functioning of individual cells as essential features of life [1]. In this respect, the cell adhesion and signaling receptors of the integrin superfamily are thought to play a major role. Integrins are heterodimeric transmembrane glycoproteins, consisting of one α - and one β -subunit. They execute crucial regulatory functions during cell adhesion, migration/invasion, proliferation, survival, and apoptosis. Moreover, integrins are capable of bidirectional signaling across cell membranes, referred to as “outside-in” and “inside-out” signaling, which results in information exchange between the ECM proteins and intracellular molecules [2–5]. Since integrins are linked to cytoskeletal components within the cell and with ECM proteins within the extracellular space, they function as mechanotransducers exhibiting force-sensing capabilities. These functional features of integrins have gained increasing attention in recent years [6,7].

In 1984, Pierschbacher and Ruoslahti described the Arg-Gly-Asp (RGD) peptide motif as a highly conserved minimal integrin recognition sequence within fibronectin [8]. Subsequently, the RGD motif was identified in many other ECM proteins, including vitronectin [9], von Willebrand factor [10], osteopontin [11], and laminin [12]. Although many integrins recognize ECM proteins via the RGD motif, the specificity of integrins governing this interaction turned out to be more complex. Meanwhile, it became clear that ligand specificity for integrins—besides the RGD motif—also depends on the distinct conformational and spatial presentation of this motif in various ECM proteins as well as on other synergistically functioning adjacent molecular regions [13–15]. Based on these findings, synthetic peptides and peptidomimetics, displaying the RGD motif have been developed as potent integrin ligands and antagonists, inhibiting the adhesion of anchorage-dependent cells to ECM proteins and thereby controlling integrin-mediated (tumor) biological functions.

Among the 24 human integrin subtypes known to date, eight integrin dimers, i.e., $\alpha v\beta 1$, $\alpha v\beta 3$, $\alpha v\beta 5$, $\alpha v\beta 6$, $\alpha v\beta 8$, $\alpha 5\beta 1$, $\alpha 8\beta 1$, and $\alpha IIb\beta 3$, recognize the tripeptide RGD motif within ECM proteins. Those members constitute a most important integrin receptor subfamily instrumental in cancer and their metastasis [16]. The exploration of functional differences between distinct members of the integrin subfamily is challenging, because the integrin expression profiles strongly depend on the cell type and the microenvironment within a given temporal and spatial context. To add even more complexity, integrins act in concert with other receptor proteins that are functionally and/or spatially organized within multiprotein networks within so-called focal adhesions (FA) [17,18]. Thus, the elucidation of the specific properties of distinct integrin subtypes governing their *in vivo* activities requires highly active and selective ligands.

The rapidly growing field of integrins as targets for cancer diagnosis and therapy has stimulated us to provide a thorough update on the impact of RGD-recognizing integrins in cancer, their distinct tumor-cell-specific functions and their crosstalk with other membrane receptors in various cancer entities. For previous reviews of integrins in cancer please see [19–28].

2. Integrin Activation upon Conformational Rearrangements

Over the many years following their discovery, integrins have been characterized in great detail. Nevertheless, the exact mechanisms of their molecular activation and bidirectional signaling remained unclear for a long time, mostly because of the missing structural information regarding these non-covalently linked heterodimeric receptors. This changed in 2001, when the structure of the extracellular domain of $\alpha v\beta 3$ in the absence or presence of Cilengitide, a small cyclic peptide ligand, was solved [29,30]. These structural data unraveled that the extracellular domains of $\alpha v\beta 3$ exist in a resting state in a conformation in which the integrin head groups are bent towards the cell membrane. Ligands may even bind to the kinked integrin in the resting state in which the transmembrane domains of each subunit are closely associated [31,32]. Upon ligand binding, however, structural rearrangements in the hinge region of each integrin subunit occur via the so-called “switchblade”

mechanism. Meanwhile, several structural studies have elucidated the details of the conformational alterations during integrin activation [2]. Key elements in the outside-in activation are a shift of the $\alpha 7$ -helix in the β I-domain and a dissociation of the transmembrane helices of the α - and the β -subunit [31,33]. Homo-oligomerization of both subunits then activates the intracellular part by the binding of intracellular proteins, such as talin and kindlin, to the cytoplasmic domain of the $\beta 3$ -subunit, which is crucial for outside-in signaling [34,35] (Figure 1). In turn, the binding of talin results in changes within the integrin conformation to an activated and erected state, accompanied by the separation of the integrin transmembrane and cytoplasmic domains (inside-out). The headpieces either remain in the closed and bent state or in the open conformation, which then results in high ligand binding affinity.




Figure 1. Schematic illustration of integrin activation and the “inside-out” and “outside-in” signaling mechanism. Integrins in the bent resting conformation reveal low affinity binding to their ECM ligands. The inside-out signaling involves disruption of the intracellular salt bridge, which is established between the cytoplasmic subunits. This induces dissociation of the transmembrane helices, followed by the reorganization and generation of a high affinity binding integrin, plus multimerization in focal adhesions. Conformational changes of the resting integrin state are induced by the integrin binding of ECM ligands causing stronger binding at the focal adhesions. The outside-in signaling requires integrin oligomerization [35].

The elucidation of the molecular mechanisms of integrin activation reveals that integrins can harbor “on” and “off” states upon conformational rearrangements by switching from a low to a high affinity state. The binding of ECM ligands with high affinity then triggers increased adhesiveness and signal transduction to the cytoplasm in the outside-in direction. To this end, phosphorylation events occur to activate downstream signaling kinases, such as the mitogen-activated protein kinases (MAPK), phosphoinositide kinase (PI3K)/Akt, and extracellular signal regulated kinase (ERK) [31,36–39], which in turn have an impact on cell proliferation, migration/invasion, and cell survival. Modulations of those integrin-mediated cell biological events are instrumental in malignant cellular transformation and tumor metastasis [40–42]. Figure 2 summarizes the various functions of specific integrins in the tumor biological contexts.




Figure 2. Integrin functions are instrumental in tumor biological contexts. Integrin functions are involved in various tumor biological processes, including cell adhesion, proliferation, inhibition of apoptosis/anoikis, induction of angiogenesis, cell invasion and migration. Adhesion: Cell adhesion is mediated by integrin binding to the respective recognition motif of RGD-ligands within the ECM, e.g., fibronectin, osteopontin and vitronectin. Binding of RGD ligands to integrins enables communication between the ECM and intracellular components, such as the cytoskeleton. Proliferation: Cell proliferation, mediated by integrin subtypes, such as $\alpha v \beta 3$, $\alpha v \beta 6$, and $\alpha v \beta 8$ may also be induced upon the binding of the RGD-containing latency associated peptide (LAP) of the inactive TGF- β to integrins. Force between the TGF- β binding proteins (LTBP) and the TGF- β molecule results in the activation of the latent TGF- β . Subsequent binding of TGF- β to the TGF- β receptor (TGF- β -R) induces epithelial-mesenchymal transition (EMT) and cell proliferation. Apoptosis/anoikis: Further, integrin expression allows cells to bind to ECM molecules within a mesenchymal tissue context, thereby inhibiting apoptosis/anoikis of invasive carcinoma cells. Angiogenesis: Upon tumor growth, the expression of hypoxia induced factors and vascular endothelial growth factor (VEGF) results in the induction of integrin-associated vessel sprouting and angiogenesis. Invasion/migration: The integrin switch from $\alpha v \beta 1$ and/or $\alpha v \beta 5$ to $\alpha v \beta 6$ allows cells to migrate and cross the basement membrane (BM) to invade the surrounding ECM as invasive cancer cells. This process is associated with EMT-like alterations of the cell phenotype. The EMT process involves the downregulation of E-cadherin and cell adhesion molecules (CAMs), promoting cell mobility, cancer progression and metastasis.

3. Integrins in Cancer Progression and Metastasis

In a series of various cancer entities, the expression of $\alpha v \beta 3$, $\alpha v \beta 5$, $\alpha 5 \beta 1$, $\alpha 6 \beta 4$, $\alpha 4 \beta 1$, $\alpha v \beta 6$ and $\alpha v \beta 8$ has been shown to correlate well with metastasis and poor patient prognosis. Here, $\alpha v \beta 3$, $\alpha v \beta 5$, and $\alpha 5 \beta 1$ are among the most prominent and well-studied integrin subtypes [21]. Table 1 provides

an overview of RGD-binding integrins, their expression in various human cancer entities and their prognostic impact.

As such, the enhanced expression of $\alpha v\beta 3$ favors tumor growth and invasion and metastasis to the bones, especially when ligated by its major ECM ligands vitronectin and/or osteopontin [43,44]. In addition to αv -integrins, $\alpha 5\beta 1$ expression levels are strongly induced upon hypoxia and promote tumor metastasis in breast cancer. Thus, high $\alpha 5$ expression in clinical biopsies is associated with an increased risk of mortality [45].

Moreover, integrin $\alpha v\beta 6$ has been proposed to serve as a marker for tumor cell invasiveness of gastric, pancreatic, cholangiocellular, breast, ovarian, colon, and head and neck cancer. Clinical data have demonstrated that $\alpha v\beta 6$ expression levels correlated with poor patient outcome [46–55]. In accordance, high $\alpha v\beta 6$ expression seems to be associated with a more aggressive disease and, consequently, poor outcome and prognosis for patients afflicted with colon cancer [56,57]. The clinical outcome is mainly attributable to the ability of colon cancer cells to metastasize to the liver. In gastric cancer, $\alpha v\beta 6$ expression was present in 37% of cases and was predictive of the reduced survival of cancer patients. The analysis of the relationship between $\alpha v\beta 6$ expression and the progression of cancer revealed that induced expression of matrix metalloproteases (MMP) and activation of the TGF- $\beta 1$ pathway are reiteratively activated [53,58,59]. In oral squamous cell carcinoma (OSCC) cells, $\alpha v\beta 6$ is highly expressed at the invasive tumor front. In gastroenteropancreatic adenocarcinomas, however, only weak $\alpha v\beta 6$ expression was noticed at the cancer cell membranes [51]. In gastroenteropancreatic malignancies, $\alpha v\beta 6$ was predominantly expressed in pancreatic ductal adenocarcinomas, followed by gastric carcinomas of the intestinal type, and intestinal adenocarcinomas, without a correlation of $\alpha v\beta 6$ expression and its ligands, such as fibronectin and tenascin. Most interestingly, $\alpha v\beta 6$ expression was elevated in well to moderately differentiated as compared to poorly differentiated tumors. This observation indicates a possible additional role of $\alpha v\beta 6$ during cancer differentiation of gastroenteropancreatic adenocarcinomas [51].

Table 1. Overview table of RGD-binding integrins expressed in various human cancers at the tissue level with respect to the expressing cell type and, if available, the prognostic impact.

Integrin	Cancer Type	Cell Type	Main Expression Feature	Reference
$\alpha v\beta 3$	gastric cancer	tumor, endothelial and stromal cells	low to moderate expression frequency in tumor cells, high frequency in stroma and endothelia, correlates with phenotype, endothelial expression correlates with survival	[60]
	glioma	endothelial and tumor cells	expression correlates with grade	[61]
	lung cancer brain metastases	endothelial and tumor cells	high expression frequency in endothelial, low frequency in tumor cells	[62]
	non small cell lung cancer	endothelial and tumor cells	high expression frequency in endothelia, low frequency in tumor cells, no correlation with survival	[63]
	oral squamous cell carcinoma	endothelial cells	higher expression in intratumoral endothelia compared with control tissue	[64]
	pancreatic cancer	tumor cells	moderate expression frequency, involved in lymph node metastasis	[65]
	prostate cancer	endothelial cells	high expression frequency peritumoral	[66]
$\alpha v\beta 5$	gastric cancer	tumor, endothelial and stromal cells	moderate (to high) frequency in tumor cells, high frequency in stroma and endothelial cells, independent prognostic factor in intestinal-type	[60]
	lung cancer (with brain metastases)	vessel endothelia and tumor cells	high expression frequency in endothelia, low frequency in tumor cells	[62]
	non small cell lung cancer	tumor cells and stroma	high frequency in tumor and stroma cells, no correlation with survival	[63]
	oral squamous cell carcinoma	tumor cells and stroma		[64]
	prostate cancer	tumor cells	expression influenced by differentiation	[66]

Table 1. Cont.

Integrin	Cancer Type	Cell Type	Main Expression Feature	Reference
$\alpha v\beta 6$	basal cell carcinoma	tumor cells	higher expression frequency in infiltrative subtype	[67]
	breast cancer		expression correlates with prognosis	[68]
	colon cancer		upregulated at invasive front and in budding tumor cells	[69]
	endometrial cancer		often overexpressed without correlation with occurrence of lymph node metastasis	[70]
	gastric cancer		potential prognostic marker in early stage carcinoma	[53]
	liver		differentiates cholangiocarcinoma from hepatocellular carcinoma	[52]
	non small cell lung cancer		high expression frequency with intratumoral heterogeneity, no correlation with survival	[71]
	lung cancer brain metastases		high expression frequency	[62]
	oral squamous cell carcinoma		expression at invasive front	[72]
	ovarian cancer		expression correlates with grade	[55]
	pancreatic cancer		high expression frequency	[51] [73]
prostate cancer	not/weakly expressed	[66]		
$\alpha v\beta 8$	non small cell lung cancer	tumor cells	low to moderate expression frequency, no correlation with survival	[63]
	prostate cancer		not expressed	[66]
$\alpha 5\beta 1$	oral squamous cell carcinoma	tumor, endothelial cells, stroma	strong expression in stroma, expressed also in tumor and endothelial cells	[64]
	ovarian cancer	tumor cells	moderate expression frequency, correlates with survival	[74]

In addition, we wish to stress here, that the integrins $\alpha v\beta 3$ and $\alpha 5\beta 1$ work in concert with growth factors and their receptors in order to influence differentiation processes in cancer stem cells in several cancer entities [75–77]. Cancer stem cells display increased expression of $\alpha v\beta 3$ on their membranes. The ligation by vitronectin results in the loss of nuclear β -catenin and the downregulation of genes that contribute to cancer stem cell maintenance [78]. To this end, strategies to target $\alpha v\beta 3$ and downstream signaling molecules, such as K-Ras are currently under investigation with the aim of interfering with cancer stemness [79]. Moreover, $\alpha v\beta 3$ represents an indicator for chemosensitivity predicting the response of tumor cells to chemotherapeutics [80,81].

As some malignancies are associated with viral infection, $\alpha v\beta 6$ and $\alpha v\beta 8$ interestingly plays a crucial role during the entry of the foot-and-mouth-disease virus, the Herpes simplex virus (HSV), and the Epstein-Barr viruses. Virus entry into the cells depends on activated viral glycoproteins. HSV infect cells via species-specific glycoproteins and the conserved apparatus gH/gL and gB. Here, HSV uses $\alpha v\beta 6$ or $\alpha v\beta 8$ as gH/gL receptors [82]. Furthermore, the fusion of epithelial cells with Epstein-Barr virus proteins can be triggered by the binding of viral glycoproteins gH/gL to $\alpha v\beta 8$ [83,84].

In the subsequent sections, we will address the characteristics and tumor biologically relevant functions of specific integrin subtypes and their interactions with distinct signaling networks.

3.1. Integrin-Mediated Cell Adhesion, Migration, and Invasion

Epithelial-type, anchorage-dependent tumor cells need to establish adhesive cell-cell and cell-ECM contacts with their microenvironment in order to evade anoikis, a special form of apoptosis attributable to the loss of cell-cell and cell-ECM contacts.

Normal epithelial cells acquire migratory capacities exclusively during embryonic development, tissue renewal and during tissue remodeling, e.g., during wound healing. In contrast, during transformation, tumor cells gain the ability to cross tissue boundaries and to invade into the surrounding ECM and the adjacent vasculature by loosening cell-cell and cell-ECM contacts. This

is a prerequisite for the ability of cells to disseminate to distant body compartments and form metastases [85,86]. Hereby, effective cell migration of individual cancer cell types is influenced by the adhesive strength of their cell-cell and cell-ECM contacts [87]. As such, by altering the adhesive strength, the fine-tuned balance between increased and decreased cell motility might be tipped in order to move a cell's body forward [88]. Hereby, integrins are also involved in cell shape alterations, which, at least in part, depend on integrin clustering and actin filament polymerization at the leading cell front of the migrating tumor cell. This is achieved by integrin cooperation with small guanosine-5'-triphosphate (GTP)-binding proteins, such as Rho and Rac [89].

The acquirement of persistent adhesive and migratory properties involves the epithelial-mesenchymal transition (EMT), which results in the loss of epithelial cell polarity and the formation of an elongated fibroblastoid cell morphology. During EMT, the expression of the cell-cell adhesive protein E-cadherin is decreased concomitant with the elevation of vimentin, desmin, and other mesenchymal markers [90,91].

Breast cancer cells utilize EMT to facilitate their invasion into the vasculature and to establish tumor metastases at distant sites [92]. In the following sections, the most important RGD-recognizing integrin subtypes and their roles in communicating cellular adhesion, migration and invasion, will be considered.

3.1.1. Integrin $\alpha\nu\beta3$ and $\alpha5\beta1$

Integrin $\alpha\nu\beta3$ was first identified by Ruoslahti and coworkers [93]. Because of its predominant ECM ligand, it was originally named vitronectin receptor. In later studies, however, $\alpha\nu\beta3$ turned out to be a highly promiscuous integrin that binds to a plethora of different ECM proteins, among them fibronectin, osteopontin, and laminin, via the RGD motif, thereby triggering integrin signaling. The need for $\alpha\nu\beta3$ -provoked signaling in order to induce cell migration has been reported for many different cell types, including smooth muscle cells [94], endothelial cells [95], and various tumor cells [96]. In human ovarian and breast cancer cells, enhancement of $\alpha\nu\beta3$ /vitronectin-mediated cell adhesion was shown to be indispensable for the provision of a sufficient grip of cells to the underlying ECM and the necessary cytoskeletal rearrangements [97–99]. Indeed, tumor cells exhibiting high $\alpha\nu\beta3$ expression levels are capable of protruding broad lamellipodia, associated with decreased RhoA activity [100,101]. With regard to the regulation of cell contractility, cell/ECM stiffness, and cell movements, $\alpha\nu\beta3$ needs to cooperate with $\alpha5\beta1$ in response to applied tension [87,102]. In prostate cancer cells, $\alpha\nu\beta3$ expression also provoked increased chemokine receptor expression, leading to enhanced migration/invasion during tumor cell metastasis to the bone [103].

Similar observations were made for integrin $\alpha5\beta1$. Breast cancer cells with high $\alpha5\beta1$ levels have revealed a 3-fold increased cell invasiveness, compared with cells exhibiting low $\alpha5\beta1$ expression. However, invasiveness can be reduced by the myosin light chain kinase inhibitor ML-7, only in cells with a high $\alpha5\beta1$ expression. This suggests a crucial role for $\alpha5\beta1$ in the enhancement of cell migration and invasion by transmission and generation of contractile forces [104]. Moreover, the endocytic recycling of $\alpha5\beta1$ also enhances cancer cell invasion, driven by RhoA and filopodial spike-based protrusions [105].

3.1.2. Integrin $\alpha\nu\beta6$

The expression of $\alpha\nu\beta6$ is initiated during embryonic development with high levels being exclusively restricted to epithelial cells, the developing lung tissue, and the kidney epithelia [106,107]. In a physiological context, $\alpha\nu\beta6$ is not constitutively expressed in differentiated epithelial cells, however, it becomes upregulated in the context of tissue remodeling, including wound healing and carcinogenesis [108]. The prevalence of $\alpha\nu\beta6$ expression has been described in several kinds of malignancies [46].

In addition to RGD-containing ECM ligands, such as fibronectin, tenascin-C, osteopontin, and vitronectin, $\alpha\nu\beta6$ binds to the RGD motif contained within the latency associated peptide (LAP) of

TGF- β to activate the inactive, latent form of the TGF- β molecule. TGF- β signaling has a strong impact on EMT during invasive growth and cancer progression and is a crucial physiological process during embryonic development. In differentiated cells, however, EMT-related cellular alterations have also been observed during fibrotic tissue remodeling, wound healing, invasive cancer growth, and tumor metastasis. One hallmark of EMT is the downregulation of E-cadherin, which provokes loss of epithelial cell-cell adherence, followed by the detachment from the epithelial tissue and increased cell motility [109]. Because $\alpha v\beta 6$ functions as a potent activator of TGF- β , a crucial role for $\alpha v\beta 6$ expression during the EMT process has been suggested [46]. The activation of TGF- $\beta 1$ by $\alpha v\beta 6$ at the epithelial-mesenchymal interface seems to be a highly conserved EMT mechanism, first established during embryonic development, and then reactivated in the context of carcinogenesis. Cell alterations associated with EMT include a switch from $\alpha v\beta 1$ and/or $\alpha v\beta 5$ to $\alpha v\beta 6$ and render $\alpha v\beta 6$ an important EMT-marker for invasive cancer cells [42,110,111]. This switch enables epithelial cells to bind to ECM proteins, to cross tissue boundaries, and to invade mesenchymal tissue. Thereby, cancer cells evade apoptosis (anoikis), which under normal conditions would be provoked because of the loss of cell-cell and cell-ECM contacts [112,113]. Increased expression of $\alpha v\beta 6$ in OSCCs triggers EMT, indicated by a decreased E-cadherin expression and the gain of vimentin expression [58]. Among the most common skin cancers, basal cell carcinomas (BCC) are caused by the deregulation of the Sonic hedgehog (Shh) signaling pathway. Although $\alpha v\beta 6$ is expressed in low risk BCC, it is markedly upregulated in infiltrative BCC and promotes invasion by modulating the tumor stroma and activating the TGF- $\beta 1$ pathway [67].

The induction of tumor cell-associated proteolytic activity and the downregulation of E-cadherin, as hallmarks of EMT, promote cancer cell migration, invasion, and tumor metastasis. Hence, the high invasive capacity of $\alpha v\beta 6$ -positive cancer cells is shown by the observation that $\alpha v\beta 6$ expression is correlated with an elevated MMP-9 expression at the invasive cancer front [114]. MMPs are zinc-dependent metallo-endopeptidases involved in the degradation of ECM components [59]. Under physiological conditions, the basement membrane functions as a natural barrier for migrating/invasive cells. Elevated expression of MMP-9 degrades collagen type IV, one of the major components of the basement membrane. In ovarian cancer tissues and cultivated ovarian cancer cells, elevated $\alpha v\beta 6$ expression is associated with the secretion of urokinase-type plasminogen activator (uPA), MMP-2, and MMP-9 [49]. Immunoprecipitation studies have revealed that $\alpha v\beta 6$ directly interacts with the uPA receptor (uPAR) in ovarian cancer cells to promote cell migration and ERK activation. The interaction of $\alpha v\beta 6$ with uPAR is restricted to ovarian cancer cells accompanied with increased TGF- $\beta 1$ [115]. In OSCCs, ligand binding to $\alpha v\beta 6$ leads to the recruitment of the focal adhesion kinase (FAK) and the activation of the Raf-ERK/MAPK pathway, thereby inducing MMPs and promoting cell proliferation and experimental metastasis [116]. Integrin $\alpha v\beta 6$ expression in OSCC also activates MMP-3, a crucial factor for ECM degradation and remodeling, the subsequent activation of further MMPs, cell migration, and invasion upon binding to fibronectin [58]. Thus, $\alpha v\beta 6$ contributes to the remodeling of the ECM as a prerequisite for cell invasion, migration, and tumor metastasis. These cell-cell and cell-ECM interactions in OSCCs also involve EMT-associated events, such as vimentin expression and E-cadherin downregulation [58]. As in OSCC, invasive endometrial carcinomas have been characterized by elevated $\alpha v\beta 6$ at the leading invasive front, concurrently with MMP expression and metastasis formation [70]. Several studies have described the influence of $\alpha v\beta 6$ activation/inhibition on the MMP pathway together with its binding of RGD-containing ligands, such as fibronectin, tenascin-C, vitronectin, and the LAP of TGF- $\beta 1$ or TGF- $\beta 3$, respectively. Deregulated ECM degradation further promotes invasive growth and malignant progression, as abnormal ECM degradation products bind to integrins, leading to the initiation of integrin signal transduction. For example, the cleavage of fibronectin by MMP-9 results in a 120-kDa fragment that promotes the migration of $\alpha v\beta 6$ -positive tumor cells. In turn, the increased expression of MMP-9 and MMP-2 leads to a positive feedback loop that might further promote cancer development [117]. At the invasive front of OSCC, MMP-7, -9, -12, and high levels of $\alpha v\beta 6$ have been described as prognostic markers [72]. The last 11 amino

acids (EKQKVDLSTDC) of the cytoplasmic tail of the $\beta 6$ -subunit seem to be sufficient to activate MMP-9 and promote invasiveness. However, this process depends on specific integrin subunits, because the $\beta 3$ -subunit activates MMP-2 [118]. In OSCC cell lines the migration of $\alpha v\beta 6$ -positive cells attributable to MMP-9 activation depends on integrin binding to insoluble LAP, whereas soluble LAP downregulates the $\alpha v\beta 6$ -induced adhesion and migration of OSCC cells [119]. Accordingly, the overexpression of $\alpha v\beta 6$ enables cells to migrate onto fibronectin with increased invasiveness because of MMP-9 upregulation [50,120]. The main signaling event, associated with $\alpha v\beta 6$ -dependent MMP activation involves the ERK/MAPK pathway [121]. Ligand binding of fibronectin to $\alpha v\beta 6$ recruits the Fyn tyrosin kinase to the $\beta 6$ -subunit. This activates MMP-3, which, in turn, degrades the ECM. Thus, the interrelation of MMP and integrin expression provides a putative molecular mechanism for $\alpha v\beta 6$ enhancement to promote metastasis of OSCC cells. Although the exact molecular mechanisms of the integrin signaling network still remain to be fully elucidated, evidence points to a major role of the ERK/MAPK and TGF- $\beta 1$ pathway in MMP activation by $\alpha v\beta 6$ in various cancer types [116].

A different aspect of integrin function during cancer development is provided by the role of $\alpha v\beta 6$ during endocytosis. Its interaction with the hematopoietic lineage cell-specific protein-1 (HS1) associated protein (HAX-1) is responsible for the clathrin-mediated endocytosis of $\alpha v\beta 6$, an event that promotes the invasive behavior of OSCCs [122]. A recent study further indicates that the clathrin or non-clathrin-mediated endocytosis of integrins affects cell-signaling pathways that control cancer progression [123]. Consequently, the interaction between HAX-1 and $\alpha v\beta 6$ might be the first evidence that integrin-triggered endocytosis plays a crucial role during cancer development and progression.

3.1.3. Integrin $\alpha v\beta 8$

A further RGD-recognizing integrin is represented by $\alpha v\beta 8$. In accordance with its $\beta 6$ counterpart, $\beta 8$ -integrin is a 100 kDa protein that exclusively heterodimerizes with the 130-kDa αv -subunit [124]. Integrin $\alpha v\beta 8$ is far less studied than other members of the integrin αv -subfamily; however, it is not only structurally, but also functionally related to $\alpha v\beta 6$. Nevertheless, its role in adhesion, invasion and migration remains to be clarified.

Its biological functions are mainly associated with its action as a potent activator of TGF- $\beta 1$ [125]. Loss-of-function experiments have revealed that mice lacking $\alpha v\beta 6$ and $\alpha v\beta 8$ activity reproduce the abnormalities observed in TGF- $\beta 1$ and TGF- $\beta 3$ null mice [126]. Site-directed mutagenesis of latent TGF- $\beta 1$ has demonstrated that the high affinity binding of $\alpha v\beta 8$ to latent TGF- $\beta 1$ is defined by Leu-218 following the RGD motif contained within LAP [127]. In contrast to $\alpha v\beta 6$, $\alpha v\beta 8$ has a shorter cytoplasmic tail and does not bind to actin. Therefore, the $\alpha v\beta 8$ -dependent activation of latent TGF- $\beta 1$ from the large latent complex (LLC) have been suggested not to work by force, but to require proteolytic cleavage by co-expressed MMP-14 (or MT1-MMP) [128]. The control of the subsequent intracellular transduction of $\alpha v\beta 6$ signaling is gradually being elucidated. So far, the Band 4.1B and the Rho GDP Dissociation Inhibitor 1 (RhoGDI1) have been identified as $\beta 8$ -binding proteins [129] and the cytoskeletal adapter protein spinophilin has been described to regulate tumor cell invasion in glioblastomas [130]. Interestingly, during wound healing, TGF- β is activated via $\alpha v\beta 8$ and $\alpha v\beta 6$. In contrast to $\alpha v\beta 6$, the inhibition of $\alpha v\beta 8$ enhances the degree of cell motility [131].

Moreover, $\alpha v\beta 8$ plays a dominant role in promoting the migration of astrocytes onto vitronectin [132]. Astrocytic $\alpha v\beta 8$ expression has been proposed to act as a central regulator of brain vessel homeostasis because of the regulation of TGF- β activation and downstream genes that promote vessel differentiation and stabilization, such as the plasminogen activator inhibitor-1 (PAI-1) and thrombospondin-1 [133].

3.2. Impact of Integrins on Cellular Proliferation

In normal cells, distinct mechanisms regulate and terminate integrin-triggered signaling pathways to keep cell biological events under a balanced control and to maintain cells anchored onto the ECM in a quiescent state. Under normal conditions, integrins are essential players for the maintenance

of tissue integrity. In contrast, tumor cells are characterized by uncontrolled behavior and acquire self-sufficiency regarding the expression and action of growth factors and their receptors, cytokines, and the activation of oncogenes, leading to the loss of control over cell proliferative activity in their pericellular microenvironment. Hanahan and Weinberg have defined the hallmarks of cancer as (i) continuous proliferation, (ii) self-sufficiency, (iii) invasion and metastasis, (iv) limitless replicative potential, (v) promotion of angiogenesis, and (vi) evasion of apoptosis [134].

3.2.1. Integrin $\alpha\nu\beta 3$

The growth-promoting effects of $\alpha\nu\beta 3$ on breast and prostate cancer cells and on malignant melanoma have been well characterized [20]. In ovarian cancer, $\alpha\nu\beta 3$ is instrumental in increasing cell proliferation by means of signaling that involves integrin-linked kinase [99]. Consequently, the blocking of $\alpha\nu$ results in drastic cell cycle arrest [135].

During their functions in cell proliferation, integrins crosstalk with growth factors and their respective receptors [136]. In breast and pancreatic cancer as well as glioma cells, $\alpha\nu\beta 3$ physically and functionally cooperates with, for example, the epidermal growth factor receptor (EGF-R), Erb-B2, and the platelet-derived growth factor (PDGF-R) whose activation drives cell proliferation [137–139]. In ovarian cancer cells, $\alpha\nu\beta 3$ expression correlates with enhanced expression and activity of the EGF-R [140]. In this way, even in the absence of respective ligands, growth factor receptor and integrins synergistically induce cellular signaling cascades. In cancer cells, constitutively activated integrin signaling not only leads to dysregulated tumor cell growth, but also avoids apoptosis. The effects of induced cell proliferation by $\alpha\nu\beta 3$ is even further enhanced by the action of anti-apoptotic proteins Bcl-2 and FLIP [90].

TGF- β signaling is well known for its anti-proliferative effects. However, its role in tumor biology is heterogeneous and depends on the oncological context. TGF- β function can switch from inhibition of cell proliferation as a tumor suppressor to activation of a pro-oncogenic EMT-program. The effect of TGF- β signaling is influenced by the abundance of active TGF- β molecules. Here, RGD-binding integrins play a crucial role by binding to LAP and activating the latent form of the TGF- β molecule. The consequences will be explained and exemplified in the following based on the prevailing data for $\alpha\nu\beta 6$ [125].

3.2.2. Integrin $\alpha\nu\beta 6$

The binding of LAP to $\alpha\nu\beta 6$ releases and activates the inactive pro/LAP-TGF- β [141,142]. Hence, $\alpha\nu\beta 6$ -positive cancer cells become self-sufficient in growth signals. In a study of cervical cancer patients, those with positive $\alpha\nu\beta 6$ expression had different expression levels of p53, PCNA, Ki-67, and TIPE2 as proliferation markers. Thus, $\alpha\nu\beta 6$ was described to cause active proliferation but to inhibit apoptosis, at least in cervical cancers [143].

3.2.3. Integrin $\alpha\nu\beta 8$

In contrast to the effects of $\alpha\nu\beta 6$ on pro/LAP-TGF- β , the activation of TGF- β by $\alpha\nu\beta 8$ has interestingly been described to inhibit the proliferation of the airway epithelium in intact bronchial tissue [144]. In this manner, the $\beta 8$ -subunit seems to regulate the growth of epithelial cells and has been suggested to inhibit tumor growth in nude mice [145]. Interestingly, integrins seem to play heterogeneous roles in regulating cell proliferation: growth-promoting integrin signals may be counterbalanced via inhibitory integrin signaling pathways, which can be activated by cytoplasmic integrins domains as the results of either alternative splicing or evolutionary divergence [146].

Integrin $\alpha\nu\beta 8$ has been suggested to regulate proliferation negatively for three main reasons. First, the cytoplasmic part of the $\beta 8$ -subunit differs in sequence, compared with other strongly related cytoplasmic domains of $\alpha\nu$ -associated integrin subunits, such as $\beta 1$, $\beta 3$, $\beta 5$, and $\beta 6$. Second, the $\beta 8$ cytoplasmic domain cannot support stable, high-affinity adhesion to vitronectin, and third, $\beta 8$ expression is restricted and mainly upregulated in non-proliferating cell types [145]. However, the

overexpression of $\alpha v\beta 8$ facilitates the proliferation and invasion of OSCC cells via the MEK/ERK signaling pathway upon the binding of $\alpha v\beta 8$ to collagen type-1 [147].

In the context of chronic obstructive pulmonary disease (COPD), TGF- β activation via $\alpha v\beta 8$ is known to enhance interleukin (IL)-1 β -dependent fibroblast expression of the ccr6 chemokine ligand ccl20, suggesting that $\alpha v\beta 8$, ccl20, and ccr6 interaction leads to the accumulation of dendritic cells (DC) around airways [148]. The convergence of the TGF- β and the IL-1 β signaling pathways on the ccl20 promoter has been defined as a mechanism by which the $\alpha v\beta 8$ -mediated activation of TGF- β regulates IL-1 β -dependent ccl20 expression in COPD patients [149]. The $\alpha v\beta 8$ -mediated activation of TGF- β regulates the chemokine secretion of lung fibroblasts, which directs DC and regulates fibrotic and immune responses in the lung [150]. Interestingly, T regulatory cells (tTregs) lacking $\alpha v\beta 8$ are unable to suppress pathogenic T-cell responses during active inflammation; however, the deletion of $\alpha v\beta 8$ does not result in a spontaneous inflammatory phenotype [151]. For thymus-derived tTregs, $\alpha v\beta 8$ is referred to as a marker protein that mediates the processing of latent TGF- $\beta 1$ from the latent TGF- $\beta 1$ /glycoprotein-A repetitions predominant protein (GARP) complex on the surface of tTregs [152].

3.3. Integrin Effects on Cell Survival and Apoptosis

The integrin-mediated anchorage of epithelial-type cells to the ECM is instrumental for maintenance of cell viability via the activation of FAK-, the PI3K/Akt, and the FAK-MAPK-pathways. In contrast, unligated integrins may eventually lead to cellular apoptosis (integrin-mediated death) [153,154]. Tumor cells derived from epithelia, even losing their ECM contacts, have the ability to overcome controlled cell death (anoikis), e.g., by increasing the expression of receptor tyrosine kinases or small guanosine-5'-triphosphate (GTP)ases in concert with downregulated caspase 8 expression [155,156]. Moreover, tumor cells may circumvent anoikis by modulating the expression pattern and cell surface density of their integrin subtypes and by internalizing activated integrins with sustained signaling in endosomes [111,157,158]. Cancer cells also manage to escape from anoikis by integrin crosstalk with growth factor receptors, resulting in effective survival signaling [159,160]. Interestingly, $\alpha v\beta 3$ -expressing tumor cells, which are detached from the ECM, nevertheless exhibit enhanced tumor growth in vitro in an anchorage-independent fashion. The growth-promoting effect depends on the recruitment of c-Src to the $\beta 3$ -integrin cytoplasmic tail. This, in turn, activates c-Src, leads to the phosphorylation of the Crk-associated substrate (CAS), and promotes tumor cell survival in a mechanism independent of FAK activation. These findings have unraveled an interesting role of $\alpha v\beta 3$ in anchorage-independent tumor cell growth and aggressiveness [21]. Loss of normal integrin action may also arise from abnormal integrin internalization. To this end, $\beta 3$ -integrin has been reported to associate with caveolin-1, indicative of raft/caveolar endocytosis. Following cell detachment, lipid rafts are rapidly internalized. This, in turn, inhibits key integrin signaling molecules such as Erk, PI3K, and Rac. In cancer cells, however, this process appears to be hindered by $\beta 3$ -integrin whereby integrin signaling is maintained [21]. In glioblastoma cells, which overexpress $\alpha v\beta 3$ at the invasive tumor front, fibronectin expression is elevated accompanied by increased cell motility and resistance to apoptosis. Subsequently, it has turned out that apoptosis in glioblastoma cells is not regulated by integrins alone but by their interconnections to the p53 pathways [161,162]. It is observed that integrin $\alpha 5\beta 1$ is more important for inducing apoptosis in glioblastomas than $\alpha v\beta 3$ [161,162]. Loss of anchorage-dependence and consequently, resistance to anoikis, are extremely important features during malignant transformation of tumor cells, enhancing tumor development and metastasis [155,163].

3.4. Integrins $\alpha v\beta 3$, $\alpha v\beta 5$, and $\alpha 5\beta 1$ in Tumor Angiogenesis

Tumor cells, like normal cells, strongly depend on the continuous supply of nutrients and oxygen in order to maintain their vital features. Further outgrowth of tumors above a tumor size of approximately 2 mm in diameter is largely restrained by the absence of functional blood vessels

in the tumor vicinity. Therefore, tumor cells are forced to induce the formation of new vessels, an event involving the three endothelial integrins, $\alpha v\beta 3$, $\alpha v\beta 5$, and $\alpha 5\beta 1$ [164,165]. This is achieved by the secretion of pro-angiogenic molecules, such as the vascular endothelial growth factor (VEGF) and the fibroblast growth factor-2 (FGF2), which upon engagement of their respective receptors leads to the so-called “angiogenic switch” [166]. VEGF-mediated angiogenesis occurs via $\alpha v\beta 3$, whereas FGF-2-mediated angiogenesis is mainly triggered by $\alpha v\beta 3$ and $\alpha 5\beta 1$ [167]. Signaling by these receptor/ligand systems contributes to the formation of new vessels, derived from the surrounding pre-existing vasculature. Hereby, per se quiescent endothelial cells are activated to proliferate and migrate towards the tumor site. This motile activity depends on the remodeling of the ECM, predominantly provoked via proteolytic ECM breakdown by MMPs and uPA. Integrin $\alpha v\beta 3$ co-localizes and assists during the activation of MMPs on the surface of angiogenic blood vessels. This provokes the remodeling of collagen and the basement membrane. The consecutive ECM breakdown then facilitates endothelial cell migration [168,169]. In accordance, pancreatic tumor cells have been shown to exhibit increased $\alpha v\beta 3$ levels that correlated with enhanced MMP-2 activation and lymph node metastasis [65].

One predominant trigger for tumor angiogenesis becomes relevant as soon as tumor growth exceeds blood supply and causes a hypoxic gradient. Hypoxia induces the translocation of the hypoxia-inducible factors-1 α (HIF-1 α), HIF-2 α , and HIF-3 α into the nucleus. Here, they recognize hypoxia response elements (HREs) and induce gene transcription upon binding to the respective promoter regions of their many target genes, including αv , $\beta 3$, $\beta 1$, and $\beta 6$, and the integrin-linked kinase (ILK) and VEGF [170,171]. In turn, these integrins enhance a migratory and invasive tumor cell phenotype that establishes a positive feedback loop and furthers EMT and tumor metastasis [172–179].

In animal models, the blockade of $\alpha v\beta 3$ and $\alpha v\beta 5$ by specific peptide antagonists, such as Cilengitide, or by blocking antibodies, such as Vitaxin, results in the drastic reduction of angiogenesis. This blockade also inhibits downstream signaling via the FAK/Src/Akt-pathway, leading to endothelial and tumor cell apoptosis [180]. Cilengitide is effective in combination with chemotherapy [181]. However, studies in αv -, as well as $\beta 3/\beta 5$ -knock-out mice indicate that $\alpha v\beta 3$ is important but not indispensable for tumor angiogenesis. In addition, $\beta 5$ does not represent an absolute requirement for this process, because extensive angiogenesis and tumor growth were still observed [182–184]. Thus, $\alpha 5\beta 1$ has been proposed also to take over as yet not fully resolved functions during angiogenesis, especially since Hynes and co-workers have shown that αv and $\alpha 5$ cooperate and are capable of functionally backing each other up [185]. Interestingly, low dose administration of the $\alpha v\beta 3$ ligand activated this integrin [182,186,187] and opens a new way in stabilization of the vascular system in the treatment with cytotoxic drugs [188].

As regards $\alpha v\beta 8$ expression, it activates a TGF- β gradient in the brain, a gradient that is angio-suppressive and inhibits endothelial cell sprouting. Consistently, the loss of $\alpha v\beta 8$ in the brain or downstream TGF- $\beta 1$ signaling via ALK5-Smad3 in endothelial cells increases vascular sprouting, branching, and proliferation, resulting in vascular dysplasia and hemorrhage [189]. However, in a developmental context, $\alpha v\beta 8$, βPix , and GIT1 have been suggested to regulate vascular stability, cerebral angiogenesis, and endothelial cell proliferation in the developing embryo [190]. In the developing retina and on astrocytes $\alpha v\beta 8$ expression is essential for neovascularization and regulates blood vessel sprouting [191]. Astrocytoma cells selectively suppress $\alpha v\beta 8$ expression to manipulate their antigenic balance in order to exploit the signaling pathways of developmental brain angiogenesis in adult brain tumors [192]. The reduced expression of the $\beta 8$ -chain had been considered to be instrumental for the pathogenesis of sporadic brain arteriovenous malformations [193].

3.5. TGF- $\beta 1$ and Its Integration with Integrin Signaling

One major inducer of EMT is TGF- β , which is produced at high levels in breast, prostate, and other cancer cells [194]. The TGF- β family comprises three isoforms: TGF- $\beta 1$, TGF- $\beta 2$, and TGF- $\beta 3$. Each member is processed as a homodimeric pro-TGF- β . This cytokine is deposited in the ECM in an inactive

form linked to LAP and one of four latent TGF- β -binding proteins (LTBP) [195]. Upon the binding of $\alpha\nu\beta 3$, $\alpha\nu\beta 6$, or $\alpha\nu\beta 8$ to the RGD motif contained within LAP, a conformational switch is provoked, leading to the exposure of TGF- $\beta 1$ to the neighboring cells where it ligates its respective receptors. This provokes the downregulation of E-cadherin and increases the expression of mesenchymal proteins, such as N-cadherin [196].


Pro-TGF- $\beta 1$ monomers contain an aminoterminal pro-domain of 249 amino acid residues, separated by a pro-protein convertase cleavage site from the 112-amino acid residue carboxy-terminus of the TGF- $\beta 1$ domain. It is cleaved intracellularly by furin-like enzymes and remains non-covalently linked to LAP in order to form the small inactive latent complex (SLC) [197]. The binding of LAP to the mature TGF- β molecule prevents signaling through the TGF- β -receptors [198]. In order to enable receptor binding, TGF- $\beta 1$ has to be activated and released from this complex. However, TGF- $\beta 1$ is mostly secreted as a large latent complex (LLC) formed by the dimerization and covalent disulfide linkage of the LAP-TGF- β to LTBPs or GARP [199]. Each pro-domain forms a ring around TGF- β to keep it inactive/latent [200]. LTBPs belong to the LTBP/fibrillin protein family, comprising fibrillin-1-3 and LTBP1-4 [201]. Except for LTBP3, the LTBP-isoforms LTBP1, LTBP2, and LTBP4 can bind to all TGF- β isoforms. LTBPs have a binding capacity towards fibronectin and vitronectin, thus in the LLC, TGF- β is bound to ECM proteins. This complex building provides in the ECM a pool of latent TGF- β that is activated in a context-dependent manner. In contrast to other mechanisms, such as proteolysis, this way of TGF- $\beta 1$ activation requires cell traction forces and the association of the LLC with the ECM [202]. Thereby, the binding of $\alpha\nu\beta 6$ to LAP directly influences the migration of cancer cells. A recent crystal structure analysis of the $\alpha\nu\beta 6$ head group bound to pro/LAP-TGF- $\beta 1$ has provided insights into the interaction between integrins and their macromolecular ligands. Furthermore, it elucidated the way in which integrin binding to ECM molecules transmits activating forces with biological consequences, such as TGF- $\beta 1$ activation [7]. Interestingly, the resulting effects on the integrin domains by small molecules and RGD-containing peptide binding depend on the force vector and the physiological orientation of the integrin dimer and their ligands.

The activation mechanism of TGF- $\beta 1$ by $\alpha\nu\beta 6$ has been thoroughly studied [119,141,203]. Upon binding of $\alpha\nu\beta 6$, $\alpha\nu\beta 3$, or $\alpha\nu\beta 8$ to the RGD LXX(I/L) motif within LAP of pro-TGF- $\beta 1$, these integrins associate with the actin cytoskeleton. This triggers conformational changes in the LLC complex (LAP-TGF- β -LTBP1), releasing TGF- $\beta 1$ from the LLC and enabling TGF- $\beta 1$ binding to its receptor [198,199]. Interestingly, TGF- $\beta 1$ activation by integrins is LTBP-specific, because $\alpha\nu\beta 6$ can only activate TGF- β molecules that are bound to LTBP1 but not to LTBP3. The various isoforms of the LTBPs exhibit significant differences in the sequence of the hinge region, suggesting a functional role in the specific activation of TGF- $\beta 1$ by the various integrin subtypes [203]. Although LAP-TGF- $\beta 1$ also binds to other integrins, such as $\alpha\nu\beta 1$ and $\alpha\nu\beta 3$, only $\alpha\nu\beta 6$ can release and activate TGF- $\beta 1$ from the LLC [204]. However, the mere binding of integrin to LAP is not sufficient to release and activate TGF- $\beta 1$ in vivo [200]. This is explainable by the different cytoskeleton-binding properties that influence the binding of the cytoplasmic integrin tail to actin and exert traction forces. Recent data have revealed that traction forces by $\alpha\nu\beta 6$ on pro/LAP-TGF- $\beta 1$ is required for TGF- $\beta 1$ activation: truncation of the cytoplasmic domain of the actin-binding $\beta 6$ -subunit or deletion of the binding site of the pro-domain of LAP-TGF- $\beta 1$ to the ECM inhibits the exertion of tensile force across the pro-domain and thus TGF- $\beta 1$ activation [205].

Upon TGF- $\beta 1$ activation, the successive signaling events in turn lead to the additional expression of $\alpha\nu\beta 3$ and $\alpha\nu\beta 6$ and of ECM protein ligands, followed by the activation of the PI3K-, Akt, and NF- κ B pathways. Moreover, growth factor receptors, such as the EGF-R, are upregulated upon TGF- β signaling; this is of note here, since the EGF-R physically and functionally crosstalks with integrins in a synergistic fashion [127,204,206–212].

4. Challenges for the Design of Novel Integrin Ligands and Their Translation into Clinical Applications

The great discovery that the small tripeptide sequence RGD is sufficient to inhibit the interaction of ECM proteins with integrins [8] has stimulated the search for small peptidic or peptidomimetic molecules to selectively address various integrin subtypes and thus their pathophysiological features with high affinity [27,28,213,214]. This culminated in the development of the peptidic compound Cilengitide (4a) (Figure 3) [215,216], which was tested in phase III clinical studies for the treatment of glioblastomas [217,218]. The failure of these studies was a backlash for the development of integrin ligands [219]. However, recent investigations into the mechanism and details of the activities of the various integrin subtypes and their role in apoptosis have shed light on the reasons for this failure and have opened new exciting applications for this molecule [182]. The binding of small ligands can occur to the resting state of an integrin receptor leading to its activation. However, a stronger binding to the focal adhesion complex requires higher concentrations of specific ligands [186].



	$\alpha v \beta 3$ [nM]	$\alpha v \beta 5$ [nM]	$\alpha v \beta 6$ [nM]	$\alpha v \beta 8$ [nM]	$\alpha 5 \beta 1$ [nM]	$\alpha IIb \beta 3$ [nM]	Ref.
1a	>10,000	>10,000	433 ± 101	37 ± 3	2.3 ± 0.02	>10,000	[220]
2a	0.65 ± 0.05	199 ± 21	>10,000	>10,000	108 ± 27.5	>10,000	[221]
3	>10,000	>10,000	23 ± 3.4	8.2 ± 0.52	2.5 ± 0.4	>10,000	[222]
4a	0.61 ± 0.06	8.4 ± 2.1	2050 ± 640	2350 ± 438	14.9 ± 3.1	5400 ± 814	[216]
5a	1200 ± 240	>10,000	0.28 ± 0.019	24 ± 3.1	73 ± 6	>10,000	[223]
6a	1.1 ± 0.1	16.7 ± 2.1	>10,000	>10,000	820 ± 156	>10,000	[224]

Figure 3. Development and testing of selective integrin ligands. Depicted are some integrin-specific ligands according to their selectivity profile (IC_{50} -values) for the subtypes $\alpha v \beta 3$, $\alpha v \beta 5$, $\alpha v \beta 6$, $\alpha v \beta 8$, $\alpha 5 \beta 1$, and $\alpha IIb \beta 3$ [214]. The binding affinities of the ligands have been determined in a cell-free, enzyme-linked immunosorbent assay (ELISA)-like assay for comparability. None of the selected compounds has significant affinity for the platelet integrin $\alpha IIb \beta 3$. Specificity or subtype with the lowest IC_{50} -values are highlighted. The preferential site for modification with diagnostic or therapeutic agents is indicated by an ellipsoid tag.

5. Improving the Activity and Selectivity of Integrin Ligands

Although the above-mentioned integrins all recognize RGD, they differ in their binding activity towards different ECM proteins. Integrins of the α v-family are well known for their strong binding to vitronectin [225]. Very early, the activity and selectivity of peptide ligands to the different integrin subtypes was observed to be controllable by cyclization [226,227], RGD flanking residues [228], the chirality of the amino acids, and the N-methylation of peptide bonds [215,226–229]. The substitution of arginine by lysine, which removes ligand binding to all α v-integrins, retains, however, the activity for the platelet receptor α IIb β 3 [27,230]. This knowledge and the structural data concerning integrin head groups with and without a ligand have enabled the development of many derivatives with improved integrin binding properties, including peptide mimetics and finally have also led to non-peptidic ligands. X-ray structures of α v β 3 [29,30], α IIb β 1/ α IIb β 3 [231], α 5 β 1 [232], and α v β 6 [7] have facilitated the rational design and optimization of ligands. For this aspect of integrin ligand design, please refer to previously published literature [27,213,214]. These selective ligands for distinct integrin subtypes bear the potential of serving as promising new drug candidates for personalized medicine in various cancer entities. Figure 3 depicts a few selected integrin ligands resulting from an extensive effort to develop RGD-peptides or low-molecular RGD-mimicking peptidomimetics, featuring high stability for diagnostic and therapeutic in vivo applications. To this end, the ligands have to be functionalized which is frequently accompanied by a decline in integrin-binding affinity. Hence, after functionalization, the binding affinity of a newly developed integrin ligands has to be carefully controlled and may be minimized by a well-considered selection of the labeling site and an appropriate spacer moiety being placed in between [214,215,220–224,233].

Moreover, during ligand development, we must avoid making the specificity of integrin ligands to be used for therapeutic purposes towards the platelet integrin α IIb β 3 too narrow, since their systemic administration might lead to major hemorrhagic disorders [26–28].

6. In Vivo Targeting of Integrins for Cancer Imaging and Therapy

Because of their different functions in cancer biology and the availability of small molecule ligands, RGD-binding integrins have been identified as attractive in vivo targets for the molecular imaging of tumors [107]. The respective contrast agents are usually based on the peptidic or non-peptidic integrin ligands, which are conjugated to suitable labels, e.g., fluorophores, radionuclides, or MRI contrast agents, such as gadolinium complexes. Moreover, integrin ligands have been incorporated into larger structures, e.g., by grafting them onto the surface of nanomaterials, or by their inclusion into liposomes [234–236].

6.1. Integrin α v β 3 and α 5 β 1

A glance at the currently available literature reveals that the overwhelming majority of pertinent studies is focused on α v β 3. Targeted probes comprising α v β 3-selective cyclic pentapeptides of the RGDxK (x = f, y) type are popular, most likely because the c(RGDxK) motif can be relatively easily synthesized. Thus, they quickly entered the portfolios of producers of fine chemicals as an off-the-shelf building block. Mice models for α v β 3-expressing tumors, such as subcutaneous U87MG glioma or M21 melanoma xenografts, are routinely generated. The imaging of α v β 3 in rodents by using modified RGD peptides has evolved into a standard benchmark scheme for in vivo feasibility studies and the validation of novel bioconjugation and labeling approaches. Hence, a vast number of α v β 3-targeting probes has been described, whereas only a small fraction has actually been pursued beyond the proof-of-principle stage. Notwithstanding this, radiolabeled probes for mapping of α v β 3 expression are widely considered to possess significant clinical potential for tumor imaging. The encouraging results of clinical applications of the first c(RGDfK)-based positron emission tomography (PET) radiotracer 18 F-galacto-RGD in cancer patients have triggered the development of numerous α v β 3-targeted radiopharmaceuticals for imaging of tumor angiogenesis [237–239]. A recent

review by Chen et al. [240] summarizes human PET images and additional data for eight different compounds [241–248], some of which are currently in clinical trials, highlighting their capability to visualize solid tumors. However, as has been repeatedly pointed out, the initially envisaged purpose of imaging $\alpha v\beta 3$, i.e., the in vivo quantification of the tumor angiogenesis and/or the patient stratification for antiangiogenic therapies, is somewhat thwarted by the finding that $\alpha v\beta 3$ expression does not necessarily correspond to angiogenic activity in (tumor) tissues [165,240,249]. Many tumor cell types display membranous or cytoplasmic $\alpha v\beta 3$ expression, but also substantial physiological expression is observed in many organs. As a result, the clinical value of the in vivo mapping of $\alpha v\beta 3$ remains to be defined [249].

In contrast to $\alpha v\beta 3$, which is not strictly required for tumor angiogenesis [183–185], $\alpha 5\beta 1$ suggests itself as a more attractive target, because of the evidence for a stricter link between angiogenesis and $\beta 1$ integrin expression [250]. Although $\alpha 5\beta 1$ is only weakly expressed in quiescent murine and human endothelial cells [251], it is upregulated on endothelial cells during vessel sprouting during the tumor angiogenic process [167]. Despite this promising perspective, only few imaging probes for $\alpha 5\beta 1$ have been described so far. The first in vivo imaging of $\alpha 5\beta 1$ expression was performed by using ^{68}Ga -labeled derivatives of a $\alpha 5\beta 1$ -selective peptidomimetic [252,253], followed by radiolabeled derivatives of a linear peptide derived by phage display [254] and of a cyclic peptide comprising the HisoDGR structural motif [255]. The best sensitivity and contrast for $\alpha 5\beta 1$ during PET imaging was hitherto achieved by using ^{68}Ga -Aqubepirin [256,257], a trimer of the aforementioned peptidomimetic [252] with an $\alpha 5\beta 1$ affinity (IC₅₀) of 80 pM. However, to the best of our knowledge, no $\alpha 5\beta 1$ imaging in humans has been reported, as yet.

Although in the context of molecular imaging, the ubiquity of c(RGDxK)-based probes has led to the common perception of an equivalence of integrin expression and angiogenesis, this distorted notion might soon be adjusted by the recent advent of imaging agents, targeting $\alpha v\beta 6$ [258].

6.2. Integrin $\alpha v\beta 6$

Several characteristics qualify $\alpha v\beta 6$ as a potential diagnostic and therapeutic target. First, it is not constitutively expressed under physiological conditions in differentiated adult epithelia but within a specific pathological context upon cellular alterations going along with EMT, i.e., wound healing, carcinogenesis, and tumor metastasis. Second, the use of targeting diagnostic and therapeutic tools should allow a clear demarcation between healthy and diseased tissue. As an EMT marker, $\alpha v\beta 6$ is specifically expressed at the epithelial-mesenchymal boundary and thus enables the specific targeting of invasive and migrating cells. Consequently, because of its high affinity and its *de novo* expression in cancer tissue, its involvement in cellular invasion and metastasis, and its expression in fibrotic tissue, $\alpha v\beta 6$ represents a promising cancer cell target [47,58,259]. To this end, increasing efforts are currently being undertaken to synthesize—besides ligands for targeting other integrins— $\alpha v\beta 6$ -specific peptidic and peptidomimetic integrin ligands/antagonists [260–264]. For the diagnostic imaging of human cancer cells, the peptide A20FMDV2, derived from the food-and-mouth disease virus sequence [262], the peptide H2009.1 [263], and the cyclic peptide S02 [264] are currently being explored for their use as radiolabeled integrin ligands. The idea of targeting RGD-binding integrins for tumor imaging has been followed extensively, however, mainly for $\alpha v\beta 3$ [265]. Nevertheless, $\alpha v\beta 6$ has been imaged in vivo by single-photon emission computed tomography (SPECT) [258–268] and PET [262,269–273].

To date, a series of various $\alpha v\beta 6$ -targeting tracers have been developed for diagnostic purposes, including linear 10- to 20-mer peptides and “stapled” cystine peptides [266–268]. Our group have developed and tested enzymatically stable cyclic peptides as novel $\alpha v\beta 6$ ligands, revealing so far the lowest molecular weight of all $\alpha v\beta 6$ -binding molecules with sub-nanomolar binding affinity [223]. Recent work has demonstrated that these novel ligands can serve as the basis for the synthesis of promising new PET-tracers [273]. ^{68}Ga -Avebehexin [271], a ^{68}Ga -labeled derivative of the metabolically stable cyclic nonapeptide cyclo-(FRGD₁LA₁Fp(NMe)K) [223], has shown the most advantageous tumor-to-background contrast because of excellent renal clearance (Figure 4). Especially for cancer

entities, which exhibit high $\alpha\beta6$ expression in more than 95% of cases, like in OSCC, its imaging encourages a novel and most promising tool for cancer diagnosis [47,262].




Figure 4. Imaging of various integrin subtypes.

Until now, only a single study on $\alpha\beta6$ targeting carried out in living human subjects has been reported. Using the ^{68}Ga - and ^{177}Lu -labeled compound SFITGv6, which comprises the binding sequence FRGDLMQL, Altmann et al. performed PET/CT scans of head and neck squamous cell carcinoma (HNSCC) and non-small-cell lung carcinoma (NSCLC) patients and found that the tracer accumulated specifically in tumors, but not in inflammatory lesions [274]. Finally, $\alpha\beta6$ -targeted imaging and therapy has been repeatedly pointed out to hold greatest promise for pancreatic carcinoma (PDAC) [51,73,267,269,275], evidence in humans is still lacking.

As examples, Figure 4 shows μPET images (maximal intensity projections) of SCID mice bearing subcutaneous tumor xenografts on the right shoulders (positions indicated by arrows). Left: M21 human melanoma with high $\alpha5\beta1$ expression, imaged using ^{68}Ga -Aquibepirin [256,257]. Right: H2009 human lung adenocarcinoma with high $\alpha\beta6$ expression, imaged using ^{68}Ga -Avebehexin [73,275].

7. Integrins in Cancer Therapy

Binding molecules with high affinity and selectivity against specific integrin subtypes can serve as key pharmacological tools for studying the biological functions of integrins. We suggest that a detailed analysis of the correlation of integrin subtype expression with cancer progression and the understanding of the underlying molecular mechanisms will help to generate molecular diagnostic and therapeutic tools to improve cancer patient care.

This knowledge will provide the basis for the clinical translation to diagnostic and therapeutic applications. Therefore, increasing efforts are currently being made to synthesize also meanwhile $\alpha\beta6$ -specific non-peptidic and peptidic integrin ligands with antagonizing/inhibiting effects [260–264].

The interruption of integrin-specific functions and signaling by specific integrin ligands has been considered as a promising potential therapeutic approach. This aspect will be exemplified here for $\alpha\beta6$: the US patent 7150871 describes the successful reduction of metastasis in $\alpha\beta6$ overexpressing lung cancer cells upon the specific binding of the $\alpha\beta6$ -specific function blocking monoclonal antibody 10D5 [47]. Moreover, in colon cancer cells, this antibody provokes increased cellular apoptosis, accompanied by enhanced ERK phosphorylation [276]. The binding of 10D5 to $\beta6$ has been suggested to disrupt binding to ERK and thus reduce metastasis. The major capsid protein VP1 of the O1 strain of the foot-and-mouth-disease virus is well known as a high affinity ligand for $\alpha\beta6$. A 17-mer peptide derived from the viral coat protein-1 VP1 has been used to create the humanized single

chain antibody scFv B63 that can be used to interfere with $\alpha v \beta 6$ -mediated cancer cell invasion [277]. As described above, small molecule peptidic and peptidomimetic integrin ligands/antagonists are also being explored for their efficacy in inhibiting specific integrin functions, tumor progression, and metastasis. Moreover, the inhibition of the binding of LAP-TGF- $\beta 1$ to $\alpha v \beta 6$ may represent a specific and context-dependent therapeutic approach for $\alpha v \beta 6$ -positive neoplasia.

8. Conclusions

The context-specific expression and selective functions of RGD-recognizing integrins open the possibility for targeting distinct physiological and pathological processes, including cell proliferation, differentiation, apoptosis, adhesion, and migration. All these processes are involved in cell invasion and angiogenesis. Selective ligands for the various integrin subtypes are the key for targeted diagnostic approaches, such as molecular imaging and drug-based therapeutic approaches. Potential clinical applications range from oncologic, to fibrotic and to inflammatory diseases management. Future studies in translationally orientated pre-clinical models and clinical trials will provide insights as to which application might be of clinical relevance. Current clinical trials are addressing the detection of integrin $\alpha v \beta 6$ in pancreatic cancer to evaluate the feasibility of [18 F]FP-R01-MG-F2 PET/CT imaging in patients afflicted with pancreatic cancer (NCT02683824) [278]. A further study will provide data considering the side effects of 18 F- $\alpha v \beta 6$ -binding-peptide and its feasibility for imaging in patients with primary tumors or cancer that has spread to the breast, colon, lung, or pancreas, in order to improve the detection of the cancer location within the body (NCT03164486) [279].

Acknowledgments: The authors declare no conflict of interest. The work had no specific funding. There are no financial disclosures from any authors.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

RGD	Arg-Gly-Asp
BM	Basement membrane
BMP	Bone morphogenetic protein
CAM	Cell adhesion molecule
CD	Crohn's disease
COPD	Chronic obstructive pulmonary disease
DC	Dendritic cells
ECM	Extracellular matrix
ERK	Extracellular signal regulated kinase
FA	Focal adhesion
TGF- β	Transforming growth factors- β
MAP	Mitogen-activated protein kinases
PI3K	Phosphoinositide kinase
GTP	Guanosine-5'-triphosphate
EGF-R	Epidermal growth factor receptor
VEGF	Vascular endothelial growth factor
PDGF-R	Platelet-derived growth factor
FAK	Focal adhesion kinase
CAS	Crk-associated substrate
LAP	Latency-associated peptide
MMP	Matrix metalloproteases
uPA	Urokinase-type plasminogen activator
HREs	Hypoxia response elements
HAX-1	Hematopoietic lineage cell-specific protein-1 (HS1) associated protein
HIF	Hypoxia-inducible factor
OSCC	Oral squamous cell carcinoma

BCC	Basal cell carcinomas
Shh	Sonic hedgehog
SPECT	Single-photon emission computed tomography
tTregs	T regulatory cells
SLC	Small latent complex
LLC	Large latent complex
LTBP	Latent TGF- β binding proteins
FMDV	Foot-and-mouth disease virus
HNSCC	Head and neck squamous cell carcinoma
NSCLC	Non-small-cell lung carcinoma
PDAC	Pancreatic carcinoma
SPECT	Single photon emission computed tomography
PET	Positron emission tomography

References

- Horwitz, A.R. The origins of the molecular era of adhesion research. *Nat. Rev. Mol. Cell Biol.* **2012**, *13*, 805–811. [[CrossRef](#)] [[PubMed](#)]
- Zhu, J.; Zhu, J.; Springer, T.A. Complete integrin headpiece opening in eight steps. *J. Cell Biol.* **2013**, *201*, 1053–1068. [[CrossRef](#)] [[PubMed](#)]
- Calderwood, D.A. Integrin activation. *J. Cell Sci.* **2004**, *117*, 657–666. [[CrossRef](#)] [[PubMed](#)]
- Mould, A.P.; Barton, S.J.; Askari, J.A.; Craig, S.E.; Humphries, M.J. Role of ADMIDAS cation-binding site in ligand recognition by integrin $\alpha 5\beta 1$. *J. Biol. Chem.* **2003**, *278*, 51622–51629. [[CrossRef](#)] [[PubMed](#)]
- Van Aghoven, J.F.; Xiong, J.-P.; Alonso, J.L.; Rui, X.; Adair, B.D.; Goodman, S.L.; Arnaout, M.A. Structural basis for pure antagonism of integrin $\alpha v\beta 3$ by a high-affinity form of fibronectin. *Nat. Struct. Mol. Biol.* **2014**, *21*, 384–388. [[CrossRef](#)] [[PubMed](#)]
- Rahmouni, S.; Lindner, A.; Rechenmacher, F.; Neubauer, S.; Sobahi, T.R.; Kessler, H.; Cavalcanti-Adam, E.A.; Spatz, J.P. Hydrogel micropillars with integrin selective peptidomimetic functionalized nanopatterned tops for the separate measurement of cell traction forces transmitted through $\alpha v\beta 3$ - or $\alpha 5\beta 1$ -integrins. *Adv. Mater.* **2013**, *25*, 5869–5874. [[CrossRef](#)] [[PubMed](#)]
- Dong, X.; Zhao, B.; Iacob, R.E.; Zhu, J.; Koksals, A.C.; Lu, C.; Engen, J.R.; Springer, T.A. Force interacts with macromolecular structure in activation of TGF- β . *Nature* **2017**, *542*, 55–59. [[CrossRef](#)] [[PubMed](#)]
- Pierschbacher, M.D.; Ruoslahti, E. Cell attachment activity of fibronectin can be duplicated by small synthetic fragments of the molecule. *Nature* **1984**, *309*, 30–33. [[CrossRef](#)] [[PubMed](#)]
- Suzuki, S.; Oldberg, A.; Hayman, E.G.; Pierschbacher, M.D.; Ruoslahti, E. Complete amino acid sequence of human vitronectin deduced from cDNA. Similarity of cell attachment sites in vitronectin and fibronectin. *EMBO J.* **1985**, *4*, 2519–2524. [[PubMed](#)]
- Plow, E.F.; Pierschbacher, M.D.; Ruoslahti, E.; Marguerie, G.A.; Ginsberg, M.H. The effect of Arg-Gly-Asp-containing peptides on fibrinogen and von Willebrand factor binding to platelets. *Proc. Natl. Acad. Sci. USA* **1985**, *82*, 8057–8061. [[CrossRef](#)] [[PubMed](#)]
- Oldberg, A.; Franzén, D.; Heinegård, D. Cloning and sequence analysis of rat bone sialoprotein (osteopontin) cDNA reveals an Arg-Gly-Asp cell-binding sequence. *Proc. Natl. Acad. Sci. USA* **1986**, *83*, 8819–8823. [[CrossRef](#)] [[PubMed](#)]
- Grant, D.S.; Tashiro, K.-I.; Segui-Real, B.; Yamada, Y.; Martin, G.R.; Kleinman, H.K. Two different laminin domains mediate the differentiation of human endothelial cells into capillary-like structures in vitro. *Cell* **1989**, *58*, 933–943. [[CrossRef](#)]
- Aota, S.; Nomizu, M.; Yamada, K.M. The short amino acid sequence Pro-His-Ser-Arg-Asn in human fibronectin enhances cell-adhesive function. *J. Biol. Chem.* **1994**, *269*, 24756–24761. [[PubMed](#)]
- Pierschbacher, M.D.; Ruoslahti, E. Arg-Gly-Asp: A versatile cell recognition signal. *Cell* **1986**, *44*, 517–518.
- Ruoslahti, E.; Pierschbacher, M.D. New perspectives in cell adhesion: RGD and integrins. *Science* **1987**, *238*, 491–497. [[CrossRef](#)] [[PubMed](#)]

16. Schittenhelm, J.; Klein, A.; Tatagiba, M.S.; Meyermann, R.; Fend, F.; Goodman, S.L.; Sipos, B. Comparing the expression of integrins $\alpha v\beta 3$, $\alpha v\beta 5$, $\alpha v\beta 6$, $\alpha v\beta 8$, fibronectin and fibrinogen in human brain metastases and their corresponding primary tumors. *Int. J. Clin. Exp. Pathol.* **2013**, *6*, 2719–2732. [[CrossRef](#)] [[PubMed](#)]
17. Huveneers, S.; Danen, E.H.J. Adhesion signaling-crosstalk between integrins, Src and Rho. *J. Cell Sci.* **2009**, *122*, 1059–1069. [[CrossRef](#)] [[PubMed](#)]
18. Geiger, B.; Spatz, J.; Bershadsky, A.D. Environmental sensing through focal adhesions. *Nat. Rev. Mol. Cell Biol.* **2009**, *10*, 21–33. [[CrossRef](#)] [[PubMed](#)]
19. Niu, J.; Li, Z. The roles of integrin $\alpha v\beta 6$ in cancer. *Cancer Lett.* **2017**, *403*, 128–137. [[CrossRef](#)] [[PubMed](#)]
20. Jin, H.; Varner, J. Integrins: Roles in cancer development and as treatment targets. *Br. J. Cancer* **2004**, *90*, 561–565. [[CrossRef](#)] [[PubMed](#)]
21. Desgrosellier, J.S.; Cheresh, D.A. Integrins in cancer: Biological implications and therapeutic opportunities. *Nat. Rev. Cancer* **2010**, *10*, 9–22. [[CrossRef](#)] [[PubMed](#)]
22. Sheldrake, H.M.; Patterson, L.H. Strategies to inhibit tumor associated integrin receptors: Rationale for dual and multi-antagonists. *J. Med. Chem.* **2014**, *57*, 6310–6315. [[CrossRef](#)] [[PubMed](#)]
23. Niland, S.; Eble, J.A. Integrin-mediated cell-matrix interaction in physiological and pathological blood vessel formation. *J. Oncol.* **2012**, 125278. [[CrossRef](#)] [[PubMed](#)]
24. Goodman, S.L.; Picard, M. Integrins as therapeutic targets. *Trends Pharmacol. Sci.* **2012**, *33*, 405–412. [[CrossRef](#)] [[PubMed](#)]
25. Goswami, S. Importance of integrin receptors in the field of pharmaceutical and medical science. *Adv. Biol. Chem.* **2013**, *3*, 224–252. [[CrossRef](#)]
26. Kiran Marelli, U.; Rechenmacher, F.; Ali Sobahi, T.R.; Mas-Moruno, C.; Kessler, H. Tumor targeting via integrin ligands. *Front. Pharmacol. Anti-Cancer Drugs* **2013**, *3*. [[CrossRef](#)]
27. Meyer, A.; Auernheimer, J.; Modlinger, A.; Kessler, H. Targeting RGD recognizing integrins: Drug development, biomaterial research, tumor imaging and targeting. *Curr. Pharm. Des.* **2006**, *12*, 2723–2747. [[CrossRef](#)] [[PubMed](#)]
28. Arndt, T.; Arndt, U.; Reuning, U.; Kessler, H. Integrins in angiogenesis: Implications for tumor therapy. In *Cancer Therapy. Molecular Targets in Tumor-Host Interactions*; Weber, G.F., Ed.; Horizon Bioscience: Norfolk, UK, 2005; pp. 93–141.
29. Xiong, J.P.; Stehle, T.; Diefenbach, B.; Zhang, R.; Dunker, R.; Scott, D.L.; Joachimiak, A.; Goodman, S.L.; Arnaout, M.A. Crystal structure of the extracellular segment of integrin $\alpha v\beta 3$. *Science* **2001**, *294*, 339–345. [[CrossRef](#)] [[PubMed](#)]
30. Xiong, J.P.; Stehle, T.; Zhang, R.; Joachimiak, A.; Frech, M.; Goodman, S.L.; Arnaout, M.A. Crystal structure of the extracellular segment of integrin $\alpha v\beta 3$ in complex with an Arg-Gly-Asp ligand. *Science* **2002**, *296*, 151–155. [[CrossRef](#)] [[PubMed](#)]
31. Müller, M.A.; Opfer, J.; Volkhardt, L.A.; Brunie, L.; Sinner, E.-K.; Boettiger, D.; Bochen, A.; Kessler, H.; Gottschalk, K.-E.; Reuning, U. The glycoporphin A transmembrane sequence within integrin $\alpha v\beta 3$ creates a non-signalling integrin with low basal affinity that is strongly adhesive under force. *J. Mol. Biol.* **2013**, *425*, 2988–3006. [[CrossRef](#)] [[PubMed](#)]
32. Gottschalk, K.-E.; Kessler, H. The structures of integrins and integrin-ligand complexes: Implications for drug design and signal transduction. *Angew. Chem. Int. Ed.* **2002**, *41*, 3767–3774. [[CrossRef](#)]
33. Gottschalk, K.-E.; Adams, P.D.; Brunger, A.T.; Kessler, H. Transmembrane signal transduction of the $\alpha IIb\beta 3$ integrin. *Protein Sci.* **2002**, *11*, 1800–1812. [[CrossRef](#)] [[PubMed](#)]
34. Moser, M.; Legate, K.R.; Zent, R.; Faessler, R. The tail of integrins, talin, and kindlins. *Science* **2009**, *324*, 895–899. [[CrossRef](#)] [[PubMed](#)]
35. Mas-Moruno, C.; Fraioli, R.; Rechenmacher, F.; Neubauer, S.; Kapp, T.G.; Kessler, H. $\alpha v\beta 3$ - or $\alpha 5\beta 1$ -integrin-selective peptidomimetics for surface coating. *Angew. Chem. Int. Ed.* **2016**, *55*, 7048–7067. [[CrossRef](#)] [[PubMed](#)]
36. Askari, J.A.; Buckley, P.A.; Mould, A.P.; Humphries, M.J. Linking integrin conformation to function. *J. Cell Sci.* **2009**, *122*, 165–170. [[CrossRef](#)] [[PubMed](#)]
37. Hoefling, M.; Kessler, H.; Gottschalk, K.-E. The transmembrane structure of integrin $\alpha IIb\beta 3$ —Significance to signal transduction. *Angew. Chem. Int. Ed.* **2009**, *48*, 6590–6593. [[CrossRef](#)] [[PubMed](#)]
38. Gottschalk, K.-E.; Kessler, H. A computational model of transmembrane integrin clustering. *Structure* **2004**, *12*, 1109–1116. [[CrossRef](#)] [[PubMed](#)]

39. Müller, M.A.; Brunie, L.; Bächer, A.S.; Kessler, H.; Gottschalk, K.-E.; Reuning, U. Cytoplasmic salt bridge formation in integrin $\alpha v \beta 3$ stabilizes its inactive state affecting integrin-mediated cell biological effects. *Cell Signal.* **2014**, *11*, 2493–2503. [[CrossRef](#)] [[PubMed](#)]
40. Schiller, H.B.; Hermann, M.R.; Polleux, J.; Vignaud, T.; Zanivan, S.; Friedel, C.C.; Sun, Z.Q.; Raducanu, A.; Gottschalk, K.E.; Thery, M.; et al. $\beta 1$ - and αv -class integrins cooperate to regulate myosin II during rigidity sensing of fibronectin-based microenvironments. *Nat. Cell Biol.* **2013**, *15*, 625–636. [[CrossRef](#)] [[PubMed](#)]
41. Stucci, S.; Tucci, M.; Passarelli, A.; Silvestris, F. $\alpha v \beta 3$ integrin: Pathogenetic role in osteotropic tumors. *Crit. Rev. Oncol. Hematol.* **2015**, *96*, 183–193. [[CrossRef](#)] [[PubMed](#)]
42. Janes, S.M.; Watt, F.M. Switch from $\alpha v \beta 1$ to $\alpha v \beta 6$ integrin expression protects squamous cell carcinomas from anoikis. *J. Cell Biol.* **2004**, *166*, 419–431. [[CrossRef](#)] [[PubMed](#)]
43. Takayama, S.; Ishii, S.; Ikeda, T.; Masamura, S.; Doi, M.; Kitajima, M. The relationship between bone metastasis from human breast cancer and integrin $\alpha v \beta 3$ expression. *Anticancer Res.* **2005**, *25*, 79–83. [[PubMed](#)]
44. Furger, K.A.; Allan, A.L.; Wilson, S.M.; Hota, C.; Vantuyghem, S.A.; Postenka, C.O.; Al-Katib, W.; Chambers, A.F.; Tuck, A.B. $\beta 3$ integrin expression increases breast carcinoma cell responsiveness to the malignancy-enhancing effects of osteopontin. *Mol. Cancer Res.* **2003**, *1*, 810–819. [[PubMed](#)]
45. Ju, J.A.; Godet, I.; Ye, I.C.; Byun, J.; Jayatilaka, H.; Lee, S.J.; Xiang, L.; Samanta, D.; Lee, M.H.; Wu, P.H.; et al. Hypoxia selectively enhances integrin $\alpha 5 \beta 1$ receptor expression in breast cancer to promote metastasis. *Mol. Cancer Res.* **2017**, *15*, 723–734. [[CrossRef](#)] [[PubMed](#)]
46. Bates, R.C.; Bellovin, D.I.; Brown, C.; Maynard, E.; Wu, B.; Kawakatsu, H.; Sheppard, D.; Oettgen, P.; Mercurio, A.M. Transcriptional activation of integrin beta6 during the epithelial-mesenchymal transition defines a novel prognostic indicator of aggressive colon carcinoma. *J. Clin. Investig.* **2005**, *115*, 339–347. [[CrossRef](#)] [[PubMed](#)]
47. Bandyopadhyay, A.; Raghavan, S. Defining the role of integrin $\alpha v \beta 6$ in cancer. *Curr. Drug Targets* **2009**, *10*, 645–652. [[CrossRef](#)] [[PubMed](#)]
48. Kawashima, A.; Tsugawa, S.; Boku, A.; Kobayashi, M.; Minamoto, T.; Nakanishi, I.; Oda, Y. Expression of αv integrin family in gastric carcinomas: Increased $\alpha v \beta 6$ is associated with lymph node metastasis. *Pathol. Res. Pract.* **2003**, *199*, 57–64. [[CrossRef](#)] [[PubMed](#)]
49. Ahmed, N.; Pansino, F.; Clyde, R.; Murthi, P.; Quinn, M.A.; Rice, G.E.; Agrez, M.V.; Mok, S.; Baker, M.S. Overexpression of alpha(v)beta6 integrin in serous epithelial ovarian cancer regulates extracellular matrix degradation via the plasminogen activation cascade. *Carcinogenesis* **2002**, *23*, 237–244. [[CrossRef](#)] [[PubMed](#)]
50. Ramos, D.M.; But, M.; Regezi, J.; Schmidt, B.L.; Atakilit, A.; Dang, D.; Ellis, D.; Jordan, R.; Li, X. Expression of integrin beta 6 enhances invasive behavior in oral squamous cell carcinoma. *Matrix Biol. J. Int. Soc. Matrix Biol.* **2002**, *21*, 297–307. [[CrossRef](#)]
51. Sipos, B.; Hahn, D.; Carceller, A.; Piulats, J.; Hedderich, J.; Kalthoff, H.; Goodman, S.L.; Kosmahl, M.; Kloppel, G. Immunohistochemical screening for beta6-integrin subunit expression in adenocarcinomas using a novel monoclonal antibody reveals strong up-regulation in pancreatic ductal adenocarcinomas in vivo and in vitro. *Histopathology* **2004**, *45*, 226–236. [[CrossRef](#)] [[PubMed](#)]
52. Patsenker, E.; Wilkens, L.; Banz, V.; Osterreicher, C.H.; Weimann, R.; Eisele, S.; Keogh, A.; Stroka, D.; Zimmermann, A.; Stickel, F. The $\alpha v \beta 6$ integrin is a highly specific immunohistochemical marker for cholangiocarcinoma. *J. Hepatol.* **2010**, *52*, 362–369. [[CrossRef](#)] [[PubMed](#)]
53. Zhang, Z.Y.; Xu, K.S.; Wang, J.S.; Yang, G.Y.; Wang, W.; Wang, J.Y.; Niu, W.B.; Liu, E.Y.; Mi, Y.T.; Niu, J. Integrin $\alpha v \beta 6$ acts as a prognostic indicator in gastric carcinoma. *Clin. Oncol. R. Coll. Radiol.* **2008**, *20*, 61–66. [[CrossRef](#)] [[PubMed](#)]
54. Arihiro, K.; Kaneko, M.; Fujii, S.; Inai, K.; Yokosaki, Y. Significance of $\alpha 9 \beta 1$ and $\alpha v \beta 6$ integrin expression in breast carcinoma. *Breast Cancer* **2000**, *7*, 19–26. [[CrossRef](#)] [[PubMed](#)]
55. Ahmed, N.; Riley, C.; Rice, G.E.; Quinn, M.A.; Baker, M.S. $\alpha v \beta 6$ integrin—A marker for the malignant potential of epithelial ovarian cancer. *J. Histochem. Cytochem.* **2002**, *50*, 1371–1380. [[CrossRef](#)] [[PubMed](#)]
56. Bates, R.C. Colorectal cancer progression: Integrin $\alpha v \beta 6$ and the epithelial-mesenchymal transition (EMT). *Cell Cycle* **2005**, *4*, 1350–1352. [[CrossRef](#)] [[PubMed](#)]
57. Bates, R.C.; Mercurio, A.M. The epithelial-mesenchymal transition (EMT) and colorectal cancer progression. *Cancer Biol. Ther.* **2005**, *4*, 365–370. [[CrossRef](#)] [[PubMed](#)]

58. Ramos, D.M.; Dang, D.; Sadler, S. The role of the integrin $\alpha\beta6$ in regulating the epithelial to mesenchymal transition in oral cancer. *Anticancer Res.* **2009**, *29*, 125–130. [[PubMed](#)]
59. Munshi, H.G.; Stack, M.S. Reciprocal interactions between adhesion receptor signaling and MMP regulation. *Cancer Metastasis Rev.* **2006**, *25*, 45–56. [[CrossRef](#)] [[PubMed](#)]
60. Böger, C.; Warneke, V.S.; Behrens, H.M.; Kalthoff, H.; Goodman, S.L.; Becker, T.; Röcken, C. Integrins $\alpha\beta3$ and $\alpha\beta5$ as prognostic, diagnostic, and therapeutic targets in gastric cancer. *Gastric Cancer* **2015**, *18*, 784–795. [[CrossRef](#)] [[PubMed](#)]
61. Schnell, O.; Krebs, B.; Wagner, E.; Romagna, A.; Beer, A.J.; Grau, S.J.; Thon, N.; Goetz, C.; Kretschmar, H.A.; Tonn, J.C.; et al. Expression of integrin $\alpha\beta3$ in gliomas correlates with tumor grade and is not restricted to tumor vasculature. *Brain Pathol.* **2008**, *18*, 378–386. [[CrossRef](#)] [[PubMed](#)]
62. Berghoff, A.S.; Kovanda, A.K.; Melchardt, T.; Bartsch, R.; Hainfellner, J.A.; Sipos, B.; Schittenhelm, J.; Zielinski, C.C.; Widhalm, G.; Dieckmann, K.; et al. $\alpha\beta3$, $\alpha\beta5$ and $\alpha\beta6$ integrins in brain metastases of lung cancer. *Clin. Exp. Metastasis* **2014**, *31*, 841–851. [[CrossRef](#)] [[PubMed](#)]
63. Böger, C.; Kalthoff, H.; Goodman, S.L.; Behrens, H.M.; Röcken, C. Integrins and their ligands are expressed in non-small cell lung cancer but not correlated with parameters of disease progression. *Virchows Arch.* **2014**, *464*, 69–78. [[CrossRef](#)] [[PubMed](#)]
64. Fabricius, E.M.; Wildner, G.P.; Kruse-Boitschenko, U.; Hoffmeister, B.; Goodman, S.L.; Raguse, J.D. Immunohistochemical analysis of integrins $\alpha\beta3$, $\alpha\beta5$ and $\alpha5\beta1$, and their ligands, fibrinogen, fibronectin, osteopontin and vitronectin, in frozen sections of human oral head and neck squamous cell carcinomas. *Exp. Ther. Med.* **2011**, *2*, 9–19. [[CrossRef](#)] [[PubMed](#)]
65. Hosotani, R.; Kawaguchi, M.; Masui, T.; Koshihara, T.; Ida, J.; Fujimoto, K.; Wada, M.; Doi, R.; Imamura, M. Expression of integrin $\alpha\beta3$ in pancreatic carcinoma: Relation to MMP-2 activation and lymph node metastasis. *Pancreas* **2002**, *25*, e30–e35. [[CrossRef](#)] [[PubMed](#)]
66. Heß, K.; Böger, C.; Behrens, H.M.; Röcken, C. Correlation between the expression of integrins in prostate cancer and clinical outcome in 1284 patients. *Ann. Diagn. Pathol.* **2014**, *18*, 343–350. [[CrossRef](#)] [[PubMed](#)]
67. Marsh, D.; Dickinson, S.; Neill, G.W.; Marshall, J.F.; Hart, I.R.; Thomas, G.J. $\alpha\beta6$ integrin promotes the invasion of morphoeic basal cell carcinoma through stromal modulation. *Cancer Res.* **2008**, *68*, 3295–3303. [[CrossRef](#)] [[PubMed](#)]
68. Moore, K.M.; Thomas, G.J.; Duffy, S.W.; Warwick, J.; Gabe, R.; Chou, P.; Ellis, I.O.; Green, A.R.; Haider, S.; Brouillette, K.; et al. Therapeutic targeting of integrin $\alpha\beta6$ in breast cancer. *J. Natl. Cancer Inst.* **2014**, *106*. [[CrossRef](#)] [[PubMed](#)]
69. Yang, G.Y.; Guo, S.; Dong, C.Y.; Wang, X.Q.; Hu, B.Y.; Liu, Y.F.; Chen, Y.W.; Niu, J.; Dong, J.H. Integrin $\alpha\beta6$ sustains and promotes tumor invasive growth in colon cancer progression. *World J. Gastroenterol.* **2015**, *21*, 7457–7467. [[CrossRef](#)] [[PubMed](#)]
70. Hecht, J.L.; Dolinski, B.M.; Gardner, H.A.; Violette, S.M.; Weinreb, P.H. Overexpression of the $\alpha\beta6$ integrin in endometrial cancer. *Appl. Immunohistochem. Mol. Morphol.* **2008**, *16*, 543–547. [[CrossRef](#)] [[PubMed](#)]
71. Elayadi, A.N.; Samli, K.N.; Prudkin, L.; Liu, Y.H.; Bian, A.; Xie, X.J.; Wistuba, I.I.; Roth, J.A.; McGuire, M.J.; Brown, K.C. A peptide selected by biopanning identifies the integrin $\alpha\beta6$ as a prognostic biomarker for nonsmall cell lung cancer. *Cancer Res.* **2007**, *67*, 5889–5895. [[CrossRef](#)] [[PubMed](#)]
72. Impola, U.; Uitto, V.J.; Hietanen, J.; Hakkinen, L.; Zhang, L.; Larjava, H.; Isaka, K.; Saarialho-Kere, U. Differential expression of matrilysin-1 (MMP-7), 92 kd gelatinase (MMP-9), and metalloelastase (MMP-12) in oral verrucous and squamous cell cancer. *J. Pathol.* **2004**, *202*, 14–22. [[CrossRef](#)] [[PubMed](#)]
73. Steiger, K.; Schlitter, A.M.; Weichert, W.; Esposito, I.; Wester, H.J.; Notni, J. Perspective of $\alpha\beta6$ -integrin imaging for clinical management of pancreatic carcinoma and its precursor lesions. *Mol. Imaging* **2017**, in press. [[CrossRef](#)] [[PubMed](#)]
74. Sawada, K.; Mitra, A.K.; Radjabi, A.R.; Bhaskar, V.; Kistner, E.O.; Tretiakova, M.; Jagadeeswaran, S.; Montag, A.; Becker, A.; Kenny, H.A.; et al. Loss of E-cadherin promotes ovarian cancer metastasis via $\alpha5$ -integrin, which is a therapeutic target. *Cancer Res.* **2008**, *68*, 2329–2339. [[CrossRef](#)] [[PubMed](#)]
75. Daley, W.P.; Peters, S.B.; Larsen, M. Extracellular matrix dynamics in development and regenerative medicine. *J. Cell Sci.* **2008**, *121*, 255–264. [[CrossRef](#)] [[PubMed](#)]
76. Jordan, C.T.; Guzman, M.L.; Noble, M. Cancer stem cells. *N. Engl. J. Med.* **2006**, *355*, 1253–1261. [[CrossRef](#)] [[PubMed](#)]

77. Taipale, J.; Beachy, P.A. The Hedgehog and Wnt signalling pathways in cancer. *Nature* **2001**, *411*, 349–354. [[CrossRef](#)] [[PubMed](#)]
78. Hurt, E.M.; Chan, K.; Serrat, M.A.; Thomas, S.B.; Veenstra, T.D.; Farrar, W.L. Identification of vitronectin as an extrinsic inducer of cancer stem cell differentiation and tumor formation. *Stem Cells* **2010**, *28*, 390–398. [[CrossRef](#)] [[PubMed](#)]
79. Seguin, L.; Kato, S.; Franovic, A.; Camargo, M.F.; Lesperance, J.; Elliott, K.C.; Yebra, M.; Mielgo, A.; Lowy, A.M.; Husain, H.; et al. An integrin β 3-KRAS-RalB complex drives tumour stemness and resistance to EGFR inhibition. *Nat. Cell Biol.* **2014**, *16*, 457–468. [[CrossRef](#)] [[PubMed](#)]
80. Vellon, L.; Menendez, J.A.; Liu, H.; Lupu, R. Upregulation of α v β 3 integrin expression is a novel molecular response to chemotherapy-induced cell damage in a heregulin-dependent manner. *Differentiation* **2007**, *75*, 819–830. [[CrossRef](#)] [[PubMed](#)]
81. Sheldrake, H.M.; Patterson, L.H. Function and antagonism of β 3 integrins in the development of cancer therapy. *Curr. Cancer Drug Targets* **2009**, *9*, 519–540. [[CrossRef](#)] [[PubMed](#)]
82. Gianni, T.; Massaro, R.; Campadelli-Fiume, G. Dissociation of HSV gL from gH by α v β 6- or α v β 8-integrin promotes gH activation and virus entry. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E3901–E3910. [[CrossRef](#)] [[PubMed](#)]
83. Chesnokova, L.S.; Nishimura, S.L.; Hutt-Fletcher, L.M. Fusion of epithelial cells by Epstein-Barr Virus proteins is triggered by binding of viral glycoproteins gHgL to integrins α v β 6 or α v β 8. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 20464–20469. [[CrossRef](#)] [[PubMed](#)]
84. Hutt-Fletcher, L.M.; Chesnokova, L.S. Integrins as triggers of Epstein-barr virus fusion and epithelial cell infection. *Virulence* **2010**, *1*, 395–398. [[CrossRef](#)] [[PubMed](#)]
85. Bear, J.E.; Haugh, J.M. Directed migration of mesenchymal cells: Where signaling and the cytoskeleton meet. *Curr. Opin. Cell Biol.* **2014**, *30*, 74–82. [[CrossRef](#)] [[PubMed](#)]
86. Parsons, J.T.; Horwitz, A.R.; Schwartz, M.A. Cell adhesion: Integrating cytoskeletal dynamics and cellular tension. *Nat. Rev. Mol. Cell Biol.* **2010**, *11*, 633–643. [[CrossRef](#)] [[PubMed](#)]
87. Missirlis, D.; Haraszti, T.; Scheele, C.; Wiegand, T.; Diaz, C.; Neubauer, S.; Rechenmacher, F.; Kessler, H.; Spatz, J.P. Substrate engagement of integrins α 5 β 1 and α v β 3 is necessary, but not sufficient, for high directional persistence in migration on fibronectin. *Sci. Rep.* **2016**, *6*, 23258. [[CrossRef](#)] [[PubMed](#)]
88. Ganguly, K.K.; Pal, S.; Moulik, S.; Chatterjee, A. Integrins and metastasis. *Cell Adhes. Migr.* **2013**, *7*, 251–261. [[CrossRef](#)] [[PubMed](#)]
89. Lawson, C.D.; Burridge, K. The on-off relationship of Rho and Rac during integrin-mediated adhesion and cell migration. *Small GTPases* **2014**, *5*, e27958. [[CrossRef](#)] [[PubMed](#)]
90. Cao, Z.; Livas, T.; Kyprianou, N. Anoikis and EMT: Lethal “liaisons” during cancer progression. *Crit. Rev. Oncol.* **2016**, *21*, 155–168. [[CrossRef](#)] [[PubMed](#)]
91. Kalluri, R.; Weinberg, R.A. The basics of epithelial-mesenchymal transition. *J. Clin. Investig.* **2009**, *119*, 1420–1428. [[CrossRef](#)] [[PubMed](#)]
92. Bartsch, J.E.; Staren, E.D.; Appert, H.E. Adhesion and migration of extracellular matrix-stimulated breast cancer. *J. Surg. Res.* **2003**, *110*, 287–294. [[CrossRef](#)]
93. Pytela, R.; Pierschbacher, M.D.; Ruoslahti, E. Identification and isolation of a 140 kd cell surface glycoprotein with properties expected of a fibronectin receptor. *Cell* **1985**, *40*, 191–198. [[CrossRef](#)]
94. Stefansson, S.; Lawrence, D.A. The serpin PAI-1 inhibits cell migration by blocking integrin α v β 3 binding to vitronectin. *Nature* **1996**, *383*, 441–443. [[CrossRef](#)] [[PubMed](#)]
95. Lian, J.; Guoping, C.; Shapiro, S.S.; Tran, L.P.; Beacham, D.A. Glycoprotein Ib α can mediate endothelial cell migration on von Willebrand factor-containing substrata. *Exp. Cell Res.* **1999**, *252*, 114–122. [[CrossRef](#)] [[PubMed](#)]
96. Hamidi, H.; Pietilä, M.; Ivaska, J. The complexity of integrins in cancer and new scopes for therapeutic targeting. *Br. J. Cancer* **2016**, *115*, 1017–1023. [[CrossRef](#)] [[PubMed](#)]
97. Gehler, S.; Ponik, S.M.; Riching, K.M.; Keely, P.J. Bi-directional signaling: Extracellular matrix and integrin regulation of breast tumor progression. *Crit. Rev. Eukaryot. Gene Expr.* **2013**, *23*, 139–157. [[CrossRef](#)] [[PubMed](#)]
98. Upheber, S.; Karle, A.; Miller, J.; Schlaugk, S.; Gross, E.; Reuning, U. Alternative splicing of KAI1 abrogates its tumor-suppressive effects on integrin α v β 3-mediated ovarian cancer biology. *Cell Signal.* **2015**, *27*, 652–662. [[CrossRef](#)] [[PubMed](#)]

99. Hapke, S.; Kessler, H.; Lubber, B.; Bengel, A.; Hutzler, P.; Höfler, H.; Schmitt, M.; Reuning, U. Ovarian cancer cell proliferation and motility is induced by engagement of integrin $\alpha v \beta 3$ /vitronectin interaction. *Biol. Chem.* **2003**, *384*, 1073–1083. [[CrossRef](#)] [[PubMed](#)]
100. Löffek, S.; Franzke, C.W.; Helfrich, I. Tension in Cancer. *Int. J. Mol. Sci.* **2016**, *17*, 1910. [[CrossRef](#)] [[PubMed](#)]
101. Morgan, M.R.; Byron, A.; Humphries, M.J.; Bass, M.D. Giving off mixed signals—distinct functions of $\alpha 5 \beta 1$ and $\alpha v \beta 3$ integrins in regulating cell behaviour. *IUBMB Life* **2009**, *61*, 731–738. [[CrossRef](#)] [[PubMed](#)]
102. Roca-Cusachs, P.; Iskratsch, T.; Sheetz, M.P. Finding the weakest link: Exploring integrin-mediated mechanical molecular pathways. *J. Cell Sci.* **2012**, *125*, 3025–3038. [[CrossRef](#)] [[PubMed](#)]
103. McCabe, N.P.; De, S.; Vasanji, A.; Brainard, J.; Byzova, T.V. Prostate cancer specific integrin $\alpha v \beta 3$ modulates bone metastatic growth and tissue remodeling. *Oncogene* **2007**, *26*, 6238–6243. [[CrossRef](#)] [[PubMed](#)]
104. Mierke, C.T.; Frey, B.; Fellner, M.; Herrmann, M.; Fabry, B. Integrin $\alpha 5 \beta 1$ facilitates cancer cell invasion through enhanced contractile forces. *J. Cell Sci.* **2011**, *124*, 369–383. [[CrossRef](#)] [[PubMed](#)]
105. Paul, N.R.; Allen, J.L.; Chapman, A.; Morlan-Mairal, M.; Zindy, E.; Jacquemet, G.; Fernandez del Ama, L.; Ferizovic, N.; Green, D.M.; Howe, J.D.; et al. $\alpha 5 \beta 1$ integrin recycling promotes Arp2/3-independent cancer cell invasion via the formin FHOD3. *J. Cell Biol.* **2015**, *210*, 1013–1031. [[CrossRef](#)] [[PubMed](#)]
106. Breuss, J.M.; Gillett, N.; Lu, L.; Sheppard, D.; Pytela, R. Restricted distribution of integrin beta 6 mRNA in primate epithelial tissues. *J. Histochem. Cytochem.* **1993**, *41*, 1521–1527. [[CrossRef](#)] [[PubMed](#)]
107. Niu, G.; Chen, X. Why integrin as a primary target for imaging and therapy. *Theranostics* **2011**, *1*, 30–47. [[CrossRef](#)] [[PubMed](#)]
108. Breuss, J.M.; Gallo, J.; DeLisser, H.M.; Klimanskaya, I.V.; Folkesson, H.G.; Pittet, J.F.; Nishimura, S.L.; Aldape, K.; Landers, D.V.; Carpenter, W. Expression of the beta 6 integrin subunit in development, neoplasia and tissue repair suggests a role in epithelial remodeling. *J. Cell Sci.* **1995**, *108*, 2241–2251. [[PubMed](#)]
109. Huang, R.; Zong, X. Aberrant cancer metabolism in epithelial-mesenchymal transition and cancer metastasis: Mechanisms in cancer progression. *Crit. Rev. Oncol. Hematol.* **2017**, *115*, 13–22. [[CrossRef](#)] [[PubMed](#)]
110. Jiang, X.; Teng, M.; Guo, X.; Zhang, D.; Zhang, Q.; Zhang, J.; Huang, Y. Switch from $\alpha v \beta 5$ to $\alpha v \beta 6$ integrin is required for CD9-regulated keratinocyte migration and MMP-9 activation. *FEBS Lett.* **2014**, *588*, 4044–4052. [[CrossRef](#)] [[PubMed](#)]
111. Guo, W.; Giancotti, F.G. Integrin signalling during tumour progression. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 816–826. [[CrossRef](#)] [[PubMed](#)]
112. Balzac, F.; Avolio, M.; Degani, S.; Kaverina, I.; Torti, M.; Silengo, L.; Small, J.V.; Retta, S.F. E-cadherin endocytosis regulates the activity of Rap1: A traffic light GTPase at the crossroads between cadherin and integrin function. *J. Cell Sci.* **2005**, *118*, 4765–4783. [[CrossRef](#)] [[PubMed](#)]
113. Oloumi, A.; McPhee, T.; Dedhar, S. Regulation of E-cadherin expression and beta-catenin/Tcf transcriptional activity by the integrin-linked kinase. *Biochim. Biophys. Acta* **2004**, *1691*, 1–15. [[CrossRef](#)] [[PubMed](#)]
114. Yang, G.Y.; Xu, K.S.; Pan, Z.Q.; Zhang, Z.Y.; Mi, Y.T.; Wang, J.S.; Chen, R.; Niu, J. Integrin $\alpha v \beta 6$ mediates the potential for colon cancer cells to colonize in and metastasize to the liver. *Cancer Sci.* **2008**, *99*, 879–887. [[CrossRef](#)] [[PubMed](#)]
115. Saldanha, R.G.; Molloy, M.P.; Bdeir, K.; Cines, D.B.; Song, X.; Uitto, P.M.; Weinreb, P.H.; Violette, S.M.; Baker, M.S. Proteomic identification of lynchpin urokinase plasminogen activator receptor protein interactions associated with epithelial cancer malignancy. *J. Proteome Res.* **2007**, *6*, 1016–1028. [[CrossRef](#)] [[PubMed](#)]
116. Li, X.; Yang, Y.; Hu, Y.; Dang, D.; Regezi, J.; Schmidt, B.L.; Atakilit, A.; Chen, B.; Ellis, D.; Ramos, D.M. $\alpha v \beta 6$ -fyn signaling promotes oral cancer progression. *J. Biol. Chem.* **2003**, *278*, 41646–41653. [[CrossRef](#)] [[PubMed](#)]
117. Al-Hazmi, N.; Thomas, G.J.; Speight, P.M.; Whawell, S.A. The 120 kDa cell-binding fragment of fibronectin up-regulates migration of $\alpha v \beta 6$ -expressing cells by increasing matrix metalloproteinase-2 and -9 secretion. *Eur. J. Oral Sci.* **2007**, *115*, 454–458. [[CrossRef](#)] [[PubMed](#)]
118. Morgan, M.R.; Thomas, G.J.; Russell, A.; Hart, I.R.; Marshall, J.F. The integrin cytoplasmic-tail motif ekqkvdldtc is sufficient to promote tumor cell invasion mediated by matrix metalloproteinase (MMP)-2 or MMP-9. *J. Biol. Chem.* **2004**, *279*, 26533–26539. [[CrossRef](#)] [[PubMed](#)]
119. Thomas, G.J.; Hart, I.R.; Speight, P.M.; Marshall, J.F. Binding of TGF- $\beta 1$ latency-associated peptide (LAP) to $\alpha v \beta 6$ integrin modulates behaviour of squamous carcinoma cells. *Br. J. Cancer* **2002**, *87*, 859–867. [[CrossRef](#)] [[PubMed](#)]

120. Thomas, G.J.; Lewis, M.P.; Hart, I.R.; Marshall, J.F.; Speight, P.M. $\alpha v \beta 6$ integrin promotes invasion of squamous carcinoma cells through up-regulation of matrix metalloproteinase-9. *Int. J. Cancer* **2001**, *92*, 641–650. [[CrossRef](#)]
121. Niu, J.; Dorahy, D.J.; Gu, X.; Scott, R.J.; Draganic, B.; Ahmed, N.; Agrez, M.V. Integrin expression in colon cancer cells is regulated by the cytoplasmic domain of the $\beta 6$ integrin subunit. *Int. J. Cancer* **2002**, *99*, 529–537. [[CrossRef](#)] [[PubMed](#)]
122. Ramsay, A.G.; Keppler, M.D.; Jazayeri, M.; Thomas, G.J.; Parsons, M.; Violette, S.; Weinreb, P.; Hart, I.R.; Marshall, J.F. HS1-associated protein X-1 regulates carcinoma cell migration and invasion via clathrin-mediated endocytosis of integrin $\alpha v \beta 6$. *Cancer Res.* **2007**, *67*, 5275–5284. [[CrossRef](#)] [[PubMed](#)]
123. Lanzetti, L.; Di Fiore, P.P. Endocytosis and Cancer: An ‘Insider’ Network with Dangerous Liaisons. *Traffic* **2008**, *9*, 2011–2021. [[CrossRef](#)] [[PubMed](#)]
124. Nishimura, S.L.; Sheppard, D.; Pytela, R. Integrin $\alpha v \beta 8$. Interaction with vitronectin and functional divergence of the $\beta 8$ cytoplasmic domain. *J. Biol. Chem.* **1994**, *269*, 28708–28715. [[PubMed](#)]
125. Worthington, J.J.; Klementowicz, J.E.; Travis, M.A. TGF- β : A sleeping giant awoken by integrins. *Trends Biochem. Sci.* **2011**, *36*, 47–54. [[CrossRef](#)] [[PubMed](#)]
126. Aluwihare, P.; Mu, Z.; Zhao, Z.; Yu, D.; Weinreb, P.H.; Horan, G.S.; Violette, S.M.; Munger, J.S. Mice that lack activity of $\alpha v \beta 6$ - and $\alpha v \beta 8$ -integrins reproduce the abnormalities of TGF- $\beta 1$ - and TGF- $\beta 3$ -null mice. *J. Cell Sci.* **2009**, *122*, 227–232. [[CrossRef](#)] [[PubMed](#)]
127. Ozawa, A.; Sato, Y.; Imabayashi, T.; Uemura, T.; Takagi, J.; Sekiguchi, K. Molecular basis of the ligand binding specificity of $\alpha v \beta 8$ integrin. *J. Biol. Chem.* **2016**, *291*, 11551–11565. [[CrossRef](#)] [[PubMed](#)]
128. Mu, D.; Cambier, S.; Fjellbirkeland, L.; Baron, J.L.; Munger, J.S.; Kawakatsu, H.; Sheppard, D.; Broaddus, V.C.; Nishimura, S.L. The integrin $\alpha v \beta 8$ mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF- $\beta 1$. *J. Cell Biol.* **2002**, *157*, 493–507. [[CrossRef](#)] [[PubMed](#)]
129. McCarty, J.H.; Cook, A.A.; Hynes, R.O. An interaction between $\alpha v \beta 8$ integrin and band 4.1b via a highly conserved region of the band 4.1 c-terminal domain. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 13479–13483. [[CrossRef](#)] [[PubMed](#)]
130. Cheerathodi, M.; Avci, N.G.; Guerrero, P.A.; Tang, L.K.; Popp, J.; Morales, J.E.; Chen, Z.; Carnero, A.; Lang, F.F.; Ballif, B.A.; et al. The cytoskeletal adapter protein spinophilin regulates invadopodia dynamics and tumor cell invasion in glioblastoma. *Mol. Cancer Res.* **2016**, *14*, 1277–1287. [[CrossRef](#)] [[PubMed](#)]
131. Neurohr, C.; Nishimura, S.L.; Sheppard, D. Activation of transforming growth factor-beta by the integrin $\alpha v \beta 8$ delays epithelial wound closure. *Am. J. Respir. Cell Mol. Biol.* **2006**, *35*, 252–259. [[CrossRef](#)] [[PubMed](#)]
132. Milner, R.; Huang, X.; Wu, J.; Nishimura, S.; Pytela, R.; Sheppard, D.; Ffrench-Constant, C. Distinct roles for astrocyte $\alpha v \beta 5$ and $\alpha v \beta 8$ integrins in adhesion and migration. *J. Cell Sci.* **1999**, *112*, 4271–4279. [[PubMed](#)]
133. Cambier, S.; Gline, S.; Mu, D.; Collins, R.; Araya, J.; Dolganov, G.; Einheber, S.; Boudreau, N.; Nishimura, S.L. Integrin $\alpha v \beta 8$ -mediated activation of transforming growth factor- β by perivascular astrocytes: An angiogenic control switch. *Am. J. Pathol.* **2005**, *166*, 1883–1894. [[CrossRef](#)]
134. Hanahan, D.; Weinberg, R.A. The hallmarks of cancer. *Cell* **2000**, *100*, 57–70. [[CrossRef](#)]
135. Cruet-Hennequart, S.; Maubant, S.; Luis, J.; Gauduchon, P.; Staedel, C.; Dedhar, S. αv integrins regulate cell proliferation through integrin-linked kinase (ILK) in ovarian cancer cells. *Oncogene* **2003**, *22*, 1688–1702. [[CrossRef](#)] [[PubMed](#)]
136. Wei, Q.; Pohl, T.L.; Seckinger, A.; Spatz, J.P.; Cavalcanti-Adam, E.A. Regulation of integrin and growth factor signaling in biomaterials for osteodifferentiation. *Beilstein J. Org. Chem.* **2015**, *11*, 773–783. [[CrossRef](#)] [[PubMed](#)]
137. Streuli, C.H.; Akhtar, N. Signal co-operation between integrins and other receptor systems. *Biochem. J.* **2009**, *418*, 491–506. [[CrossRef](#)] [[PubMed](#)]
138. Hannigan, G.; Troussard, A.A.; Dedhar, S. Integrin-linked kinase. A cancer therapeutic target unique among its ILK. *Nat. Rev. Cancer* **2005**, *5*, 51–63. [[CrossRef](#)] [[PubMed](#)]
139. Stupack, D.G.; Cheresch, D.A. Get a ligand, get a life: Integrins, signaling and cell survival. *J. Cell Sci.* **2002**, *115*, 3729–3738. [[CrossRef](#)] [[PubMed](#)]
140. Lössner, D.; Abou-Ajram, C.; Bengel, A.; Reuning, U. Integrin $\alpha v \beta 3$ mediates upregulation of epidermal growth-factor receptor expression and activity in human ovarian cancer cells. *Int. J. Biochem. Cell. Biol.* **2008**, *40*, 2746–2761. [[CrossRef](#)] [[PubMed](#)]

141. Schwartz, M.A.; Assoian, R.K. Integrins and cell proliferation: Regulation of cyclin-dependent kinases via cytoplasmic signaling pathways. *J. Cell Sci.* **2001**, *114*, 2553–2560. [[PubMed](#)]
142. Munger, J.S.; Huang, X.; Kawakatsu, H.; Griffiths, M.J.; Dalton, S.L.; Wu, J.; Pittet, J.F.; Kaminski, N.; Garat, C.; Matthay, M.A.; et al. The integrin $\alpha\beta6$ binds and activates latent TGF- β 1: A mechanism for regulating pulmonary inflammation and fibrosis. *Cell* **1999**, *96*, 319–328. [[CrossRef](#)]
143. Liang, D.; Xu, W.; Zhang, Q.; Tao, B.B. Study on the effect of Integrin $\alpha\beta6$ on proliferation and apoptosis of cervical cancer cells. *Eur. Rev. Med. Pharmacol. Sci.* **2017**, *21*, 2811–2815. [[PubMed](#)]
144. Fjellbirkeland, L.; Cambier, S.; Broaddus, V.C.; Hill, A.; Brunetta, P.; Dolganov, G.; Jablons, D.; Nishimura, S.L. Integrin $\alpha\beta8$ -mediated activation of transforming growth factor-beta inhibits human airway epithelial proliferation in intact bronchial tissue. *Am. J. Pathol.* **2003**, *163*, 533–542. [[CrossRef](#)]
145. Cambier, S.; Mu, D.Z.; O'Connell, D.; Boylen, K.; Travis, W.; Liu, W.H.; Broaddus, V.C.; Nishimura, S.L. A role for the integrin $\alpha\beta8$ in the negative regulation of epithelial cell growth. *Cancer Res.* **2000**, *60*, 7084–7093. [[PubMed](#)]
146. Giacotti, F.G.; Ruoslahti, E. Transduction-integrin signaling. *Science* **1999**, *285*, 1028–1032. [[CrossRef](#)]
147. Hayashido, Y.; Kitano, H.; Sakaue, T.; Fujii, T.; Suematsu, M.; Sakurai, S.; Okamoto, T. Overexpression of integrin $\alpha\nu$ facilitates proliferation and invasion of oral squamous cell carcinoma cells via MEK/ERK signaling pathway that is activated by interaction of integrin $\alpha\beta8$ with type collagen. *Int. J. Oncol.* **2014**, *45*, 1875–1882. [[CrossRef](#)] [[PubMed](#)]
148. Hashimoto, M.; Yanagisawa, H.; Minagawa, S.; Sen, D.; Ma, R.; Murray, L.A.; Tsui, P.; Lou, J.; Marks, J.D.; Baron, J.L.; et al. TGF- β -dependent dendritic cell chemokinesis in murine models of airway disease. *J. Immunol.* **2015**, *195*, 1182–1190. [[CrossRef](#)] [[PubMed](#)]
149. Brand, O.J.; Somanath, S.; Moermans, C.; Yanagisawa, H.; Hashimoto, M.; Cambier, S.; Markovics, J.; Bondesson, A.J.; Hill, A.; Jablons, D.; et al. Transforming growth factor- β and interleukin-1 β signaling pathways converge on the chemokine CCL20 promoter. *J. Biol. Chem.* **2015**, *290*, 14717–14728. [[CrossRef](#)] [[PubMed](#)]
150. Kitamura, H.; Cambier, S.; Somanath, S.; Barker, T.; Minagawa, S.; Markovics, J.; Goodsell, A.; Publicover, J.; Reichardt, L.; Jablons, D.; et al. Mouse and human lung fibroblasts regulate dendritic cell trafficking, airway inflammation, and fibrosis through integrin $\alpha\beta8$ -mediated activation of TGF- β . *J. Clin. Investig.* **2011**, *121*, 2863–2875. [[CrossRef](#)] [[PubMed](#)]
151. Worthington, J.J.; Kelly, A.; Smedley, C.; Bauche, D.; Campbell, S.; Marie, J.C.; Travis, M.A. Integrin $\alpha\beta8$ -mediated TGF- β activation by effector regulatory T cells is essential for suppression of T-cell-mediated inflammation. *Immunity* **2015**, *42*, 903–915. [[CrossRef](#)] [[PubMed](#)]
152. Edwards, J.P.; Thornton, A.M.; Shevach, E.M. Release of active TGF- β 1 from the latent TGF- β 1/GARP complex on T regulatory cells is mediated by integrin β 8. *J. Immunol.* **2014**, *193*, 2843–2849. [[CrossRef](#)] [[PubMed](#)]
153. Cheresh, D.A.; Stupack, D.G. Integrin-mediated death: An explanation of the integrin-knockout phenotype? *Nat. Med.* **2002**, *8*, 193–194. [[CrossRef](#)] [[PubMed](#)]
154. Frisch, S.M.; Ruoslahti, E. Integrins and anoikis. *Curr. Opin. Cell Biol.* **1997**, *9*, 701–770. [[CrossRef](#)]
155. Buchheit, C.L.; Weigel, K.J.; Schafer, Z.T. Cancer cell survival during detachment from the ECM: Multiple barriers to tumour progression. *Nat. Rev. Cancer* **2014**, *14*, 632–641. [[CrossRef](#)] [[PubMed](#)]
156. Teitz, T.; Wei, T.; Valentine, M.B.; Vanin, E.F.; Grenet, J.; Valentine, V.A.; Behm, F.G.; Look, A.T.; Lahti, J.M.; Kidd, V.J. Caspase 8 is deleted or silenced preferentially in childhood neuroblastomas with amplification of MYCN. *Nat. Med.* **2000**, *6*, 529–535. [[CrossRef](#)] [[PubMed](#)]
157. Alanko, J.; Mai, A.; Jacquemet, G.; Schauer, K.; Kaukonen, R.; Saari, M.; Goud, B.; Ivaska, J. Integrin endosomal signalling suppresses anoikis. *Nat. Cell Biol.* **2015**, *17*, 1412–1421. [[CrossRef](#)] [[PubMed](#)]
158. Plantefaber, L.C.; Hynes, R.O. Changes in integrin receptors on oncogenically transformed cells. *Cell* **1989**, *56*, 281–290. [[CrossRef](#)]
159. Salmeron-Sanchez, M.; Dalby, M.J. Synergistic growth factor microenvironments. *Chem. Commun. Camb.* **2016**, *52*, 13327–13336. [[CrossRef](#)] [[PubMed](#)]
160. Reginato, M.J.; Mills, K.R.; Paulus, J.K.; Lynch, D.K.; Sgroi, D.C.; Debnath, J.; Muthuswamy, S.K.; Brugge, J.S. Integrins and EGFR coordinately regulate the pro-apoptotic protein BIM to prevent anoikis. *Nat. Cell Biol.* **2003**, *5*, 733–740. [[CrossRef](#)] [[PubMed](#)]

161. Ray, A.-M.; Schaffner, F.; Janouskova, H.; Noulet, F.; Rognan, D.; Lelong-Rebel, I.; Choulier, L.; Blandin, A.-F.; Lehmann, M.; Martin, S.; et al. Single cell tracking assay reveals an opposite effect of selective small non-peptidic $\alpha 5\beta 1$ or $\alpha v\beta 3/\beta 5$ integrins antagonists in U87MG glioma cells. *Biochem. Biophys. Acta* **2014**, *1840*, 2978–2987. [[CrossRef](#)] [[PubMed](#)]
162. Renner, G.; Janouskova, H.; Noulet, F.; Guerin, E.; Bär, S.; Nuesch, J.; Rechenmacher, F.; Neubauer, S.; Kessler, H.; Choulier, L.; et al. Integrin $\alpha 5\beta 1$ and p53 convergent pathways in the control of anti-apoptotic proteins PEA-15 and survivin in high grade glioma. *Cell Death Differ.* **2016**, *23*, 640–653. [[CrossRef](#)] [[PubMed](#)]
163. Paoli, P.; Giannoni, E.; Chiarugi, P. Anoikis molecular pathways and its role in cancer progression. *Biochim. Biophys. Acta* **2013**, *1833*, 3481–3498. [[CrossRef](#)] [[PubMed](#)]
164. Demircioglu, F.; Hodivala-Dilke, K. $\alpha v\beta 3$ Integrin and tumour blood vessels-learning from the past to shape the future. *Curr. Opin. Cell Biol.* **2016**, *42*, 121–127. [[CrossRef](#)] [[PubMed](#)]
165. Avraamides, C.J.; Garmy-Susini, B.; Varnier, J.A. Integrins in angiogenesis and lymphangiogenesis. *Nat. Rev. Cancer* **2008**, *8*, 604–617. [[CrossRef](#)] [[PubMed](#)]
166. Bergers, G.; Benjamin, L.E. Tumorigenesis and the angiogenic switch. *Nat. Rev. Cancer* **2003**, *3*, 401–410. [[CrossRef](#)] [[PubMed](#)]
167. Kim, S.; Bell, K.; Mousa, S.A.; Varnier, J.A. Regulation of angiogenesis in vivo by ligation of integrin $\alpha 5\beta 1$ with the central cell-binding domain of fibronectin. *Am. J. Pathol.* **2000**, *156*, 1345–1362. [[CrossRef](#)]
168. Stefanidakis, M.; Koivunen, E. Cell-surface association between matrix metalloproteinases and integrins: Role of the complexes in leukocyte migration and cancer progression. *Blood* **2006**, *108*, 1441–1450. [[CrossRef](#)] [[PubMed](#)]
169. Mengele, K.; Napieralski, R.; Magdolen, V.; Reuning, U.; Gkazepis, A.; Sweep, F.; Brunner, N.; Foekens, J.; Harbeck, N.; Schmitt, M. Characteristics of the level-of-evidence-1 disease forecast cancer biomarkers uPA and its inhibitor PAI-1. *Expert Rev. Mol. Diagn.* **2010**, *10*, 947–962. [[CrossRef](#)] [[PubMed](#)]
170. Wang, G.L.; Jiang, B.H.; Rue, E.A.; Semenza, G.L. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-pas heterodimer regulated by cellular O_2 tension. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 5510–5514. [[CrossRef](#)] [[PubMed](#)]
171. Krock, B.L.; Skuli, N.; Simon, M.C. Hypoxia-induced angiogenesis: Good and evil. *Genes Cancer* **2011**, *2*, 1117–1133. [[CrossRef](#)] [[PubMed](#)]
172. Cowden Dahl, K.D.; Robertson, S.E.; Weaver, V.M.; Simon, M.C. Hypoxia-inducible factor regulates $\alpha v\beta 3$ integrin cell surface expression. *Mol. Biol. Cell* **2005**, *16*, 1901–1912. [[CrossRef](#)] [[PubMed](#)]
173. Keely, S.; Glover, L.E.; MacManus, C.F.; Campbell, E.L.; Scully, M.M.; Furuta, G.T.; Colgan, S.P. Selective induction of integrin $\beta 1$ by hypoxia-inducible factor: Implications for wound healing. *FASEB J.* **2009**, *23*, 1338–1346. [[CrossRef](#)] [[PubMed](#)]
174. Brooks, D.L.; Schwab, L.P.; Krutilina, R.; Parke, D.N.; Sethuraman, A.; Hoogewijs, D.; Schorg, A.; Gotwald, L.; Fan, M.; Wenger, R.H.; et al. Itga6 is directly regulated by hypoxia-inducible factors and enriches for cancer stem cell activity and invasion in metastatic breast cancer models. *Mol. Cancer* **2016**, *15*, 26. [[CrossRef](#)] [[PubMed](#)]
175. Chou, C.C.; Chuang, H.C.; Salunke, S.B.; Kulp, S.K.; Chen, C.S. A novel HIF-1 alpha-integrin-linked kinase regulatory loop that facilitates hypoxia-induced HIF-1alpha expression and epithelial-mesenchymal transition in cancer cells. *Oncotarget* **2015**, *6*, 8271–8285. [[CrossRef](#)] [[PubMed](#)]
176. Liu, W.; Shen, S.M.; Zhao, X.Y.; Chen, G.Q. Targeted genes and interacting proteins of hypoxia inducible factor-1. *Int. J. Biochem. Mol. Biol.* **2012**, *3*, 165–178. [[PubMed](#)]
177. Han, Z.B.; Ren, H.; Zhao, H.; Chi, Y.; Chen, K.; Zhou, B.; Liu, Y.J.; Zhang, L.; Xu, B.; Liu, B.; et al. Hypoxia-inducible factor (HIF)-1 alpha directly enhances the transcriptional activity of stem cell factor (scf) in response to hypoxia and epidermal growth factor (EGF). *Carcinogenesis* **2008**, *29*, 1853–1861. [[CrossRef](#)] [[PubMed](#)]
178. Kelly, B.D.; Hackett, S.F.; Hirota, K.; Oshima, Y.; Cai, Z.; Berg-Dixon, S.; Rowan, A.; Yan, Z.; Campochiaro, P.A.; Semenza, G.L. Cell type-specific regulation of angiogenic growth factor gene expression and induction of angiogenesis in nonischemic tissue by a constitutively active form of hypoxia-inducible factor 1. *Circ. Res.* **2003**, *93*, 1074–1081. [[CrossRef](#)] [[PubMed](#)]
179. Nikitenko, L.L.; Smith, D.M.; Bicknell, R.; Rees, M.C. Transcriptional regulation of the crlr gene in human microvascular endothelial cells by hypoxia. *FASEB J.* **2003**, *17*, 1499–1501. [[CrossRef](#)] [[PubMed](#)]

180. Weis, S.M.; Cheresh, D.A. α V integrins in angiogenesis and cancer. *Cold Spring Harb. Perspect. Med.* **2011**, *1*, a006478. [[CrossRef](#)] [[PubMed](#)]
181. Raguse, J.D.; Gath, H.J.; Bier, J.; Riess, H.; Oettle, H. Cilengitide (emd 121974) arrests the growth of a heavily pretreated highly vascularised head and neck tumour. *Oral Oncol.* **2004**, *40*, 228–230. [[CrossRef](#)] [[PubMed](#)]
182. Reynolds, A.R.; Hart, I.R.; Watson, A.R.; Welti, J.C.; Silva, R.G.; Robinson, S.D.; Da Violante, G.; Gourlaourou, V.; Salih, M.; Jones, M.C.; et al. Stimulation of tumor growth and angiogenesis by low concentrations of RGD-mimetic integrin inhibitors. *Nat. Med.* **2009**, *15*, 392–400. [[CrossRef](#)] [[PubMed](#)]
183. Bader, B.L.; Rayburn, H.; Crowley, D.; Hynes, R.O. Extensive vasculogenesis, angiogenesis, and organogenesis precede lethality in mice lacking all α v integrins. *Cell* **1998**, *95*, 507–519. [[CrossRef](#)]
184. Reynolds, L.E.; Wyder, L.; Lively, J.C.; Taverna, D.; Robinson, S.D.; Huang, X.; Sheppard, D.; Hynes, R.O.; Hodivala-Dilke, K.M. Enhanced pathological angiogenesis in mice lacking β 3 integrin or β 3 and β 5 integrins. *Nat. Med.* **2002**, *8*, 27–34. [[CrossRef](#)] [[PubMed](#)]
185. Van der Flier, A.; Badu-Nkansah, K.; Whittaker, C.A.; Crowley, D.; Bronson, R.T.; Lacy-Hulbert, A.; Hynes, R.O. Endothelial α 5 and α v integrins cooperate in remodeling of the vasculature during development. *Development* **2010**, *137*, 2439–2449. [[CrossRef](#)] [[PubMed](#)]
186. Wong, P.P.; Demircioglu, F.; Ghazaly, E.; Alrawashdeh, W.; Stratford, M.R.L.; Scudamore, C.L.; Cereser, B.; Crnogorac-Jurcevic, T.; McDonald, S.; Elia, G.; et al. Dual-action combination therapy enhances angiogenesis while reducing tumor growth and spread. *Cancer Cell* **2015**, *27*, 123–137. [[CrossRef](#)] [[PubMed](#)]
187. Kostourou, V.; Lechertier, T.; Reynolds, L.E.; Lees, D.M.; Baker, M.; Jones, D.T.; Tavora, B.; Ramjaun, A.R.; Birdsey, G.M.; Robinson, S.D.; et al. FAK-heterozygous mice display enhanced tumour angiogenesis. *Nat. Commun.* **2013**, *4*, 2020. [[CrossRef](#)] [[PubMed](#)]
188. Wong, P.P.; Bodrug, N.; Hodivala-Dilke, K.M. Exploring Novel Methods for Modulating Tumor Blood Vessels in Cancer Treatment. *Curr. Biol.* **2016**, *26*, R1161–R1166. [[CrossRef](#)] [[PubMed](#)]
189. Arnold, T.D.; Niaudet, C.; Pang, M.F.; Siegenthaler, J.; Gaengel, K.; Jung, B.; Ferrero, G.M.; Mukoyama, Y.S.; Fuxe, J.; Akhurst, R.; et al. Excessive vascular sprouting underlies cerebral hemorrhage in mice lacking α v β 8-TGF β signaling in the brain. *Development* **2014**, *141*, 4489–4499. [[CrossRef](#)] [[PubMed](#)]
190. Liu, J.; Zeng, L.; Kennedy, R.M.; Gruenig, N.M.; Childs, S.J. β Pix plays a dual role in cerebral vascular stability and angiogenesis, and interacts with integrin α v β 8. *Dev. Biol.* **2012**, *363*, 95–105. [[CrossRef](#)] [[PubMed](#)]
191. Hirota, S.; Liu, Q.; Lee, H.S.; Hossain, M.G.; Lacy-Hulbert, A.; McCarty, J.H. The astrocyte-expressed integrin α v β 8 governs blood vessel sprouting in the developing retina. *Development* **2011**, *138*, 5157–5166. [[CrossRef](#)] [[PubMed](#)]
192. Tchaicha, J.H.; Mobley, A.K.; Hossain, M.G.; Aldape, K.D.; McCarty, J.H. A mosaic mouse model of astrocytoma identifies α v β 8 integrin as a negative regulator of tumor angiogenesis. *Oncogene* **2010**, *29*, 4460–4472. [[CrossRef](#)] [[PubMed](#)]
193. Su, H.; Kim, H.; Pawlikowska, L.; Kitamura, H.; Shen, F.; Cambier, S.; Markovics, J.; Lawton, M.T.; Sidney, S.; Bollen, A.W.; et al. Reduced expression of integrin α v β 8 is associated with brain arteriovenous malformation pathogenesis. *Am. J. Pathol.* **2010**, *176*, 1018–1027. [[CrossRef](#)] [[PubMed](#)]
194. Smith, B.N.; Bhowmick, N.A. Role of EMT in metastasis and therapy resistance. *J. Clin. Med.* **2016**, *5*, 17. [[CrossRef](#)] [[PubMed](#)]
195. Khan, Z.; Marshall, J.F. The role of integrins in TGF- β -activation in the tumour stroma. *Cell Tissue Res.* **2016**, *365*, 657–673. [[CrossRef](#)] [[PubMed](#)]
196. Araki, K.; Shimura, T.; Suzuki, H.; Tsutsumi, S.; Wada, W.; Yajima, T.; Kobayahi, T.; Kubo, N.; Kuwano, H. E/N-cadherin switch mediates cancer progression via TGF- β -induced epithelial-to-mesenchymal transition in extrahepatic cholangiocarcinoma. *Br. J. Cancer* **2011**, *105*, 1885–1893. [[CrossRef](#)] [[PubMed](#)]
197. Dubois, C.M.; Laprise, M.H.; Blanchette, F.; Gentry, L.E.; Leduc, R. Processing of transforming growth factor β 1 precursor by human furin convertase. *J. Biol. Chem.* **1995**, *270*, 10618–10624. [[CrossRef](#)] [[PubMed](#)]
198. Lawrence, D.A.; Pircher, R.; Kryceve-Martinerie, C.; Jullien, P. Normal embryo fibroblasts release transforming growth factors in a latent form. *J. Cell Physiol.* **1984**, *121*, 184–188. [[CrossRef](#)] [[PubMed](#)]
199. Wipff, P.J.; Hinz, B. Integrins and the activation of latent transforming growth factor beta1—An intimate relationship. *Eur. J. Cell Biol.* **2008**, *87*, 601–615. [[CrossRef](#)] [[PubMed](#)]
200. Shi, M.; Zhu, J.; Wang, R.; Chen, X.; Mi, L.; Walz, T.; Springer, T.A. Latent TGF- β structure and activation. *Nature* **2011**, *474*, 343–349. [[CrossRef](#)] [[PubMed](#)]
201. Ramirez, F.; Pereira, L. The fibrillins. *Int. J. Biochem. Cell Biol.* **1999**, *31*, 255–259. [[CrossRef](#)]

202. Ahamed, J.; Burg, N.; Yoshinaga, K.; Janczak, C.A.; Rifkin, D.B.; Collier, B.S. In vitro and in vivo evidence for shear-induced activation of latent transforming growth factor- β 1. *Blood* **2008**, *112*, 3650–3660. [[CrossRef](#)] [[PubMed](#)]
203. Annes, J.P.; Chen, Y.; Munger, J.S.; Rifkin, D.B. Integrin α v β 6-mediated activation of latent TGF- β requires the latent TGF- β binding protein-1. *J. Cell Biol.* **2004**, *165*, 723–734. [[CrossRef](#)] [[PubMed](#)]
204. Munger, J.S.; Harpel, J.G.; Giancotti, F.G.; Rifkin, D.B. Interactions between growth factors and integrins: Latent forms of transforming growth factor- β are ligands for the integrin α v β 1. *Mol. Biol. Cell* **1998**, *9*, 2627–2638. [[CrossRef](#)] [[PubMed](#)]
205. Robertson, I.B.; Rifkin, D.B. Regulation of the bioavailability of TGF- β and TGF- β -related proteins. *Cold Spring Harb. Perspect. Biol.* **2016**, *8*. [[CrossRef](#)] [[PubMed](#)]
206. Eke, I.; Zscheppang, K.; Dickreuter, E.; Hickmann, L.; Mazzeo, E.; Unger, K.; Krause, M.; Cordes, N. Simultaneous β 1 integrin-EGFR targeting and radiosensitization of human head and neck cancer. *J. Natl. Cancer Inst.* **2015**, *107*, dju419. [[CrossRef](#)] [[PubMed](#)]
207. Tatler, A.L.; Goodwin, A.T.; Gbolahan, O.; Saini, G.; Porte, J.; John, A.E.; Clifford, R.L.; Violette, S.M.; Weinreb, P.H.; Parfrey, H.; et al. Amplification of TGF- β induced ITGB6 gene transcription may promote pulmonary fibrosis. *PLoS ONE* **2016**, *11*, e0158047. [[CrossRef](#)] [[PubMed](#)]
208. Yeh, Y.Y.; Chiao, C.C.; Kuo, W.Y.; Hsiao, Y.C.; Chen, Y.J.; Wei, Y.Y.; Lai, T.H.; Fong, Y.C.; Tang, C.H. TGF- β 1 increases motility and α v β 3 integrin up-regulation via PI3K, Akt and NF- κ B-dependent pathway in human chondrosarcoma cells. *Biochem. Pharmacol.* **2008**, *75*, 1292–1301. [[CrossRef](#)] [[PubMed](#)]
209. Wendt, M.K.; Smith, J.A.; Schiemann, W.P. Transforming growth factor- β -induced epithelial-mesenchymal transition facilitates epidermal growth factor-dependent breast cancer progression. *Oncogene* **2010**, *29*, 6485–6498. [[CrossRef](#)] [[PubMed](#)]
210. Pechkovsky, D.V.; Scaffidi, A.K.; Hackett, T.L.; Ballard, J.; Shaheen, F.; Thompson, P.J.; Thannickal, V.J.; Knight, D.A. Transforming growth factor β 1 induces α v β 3 integrin expression in human lung fibroblasts via a β 3 integrin-, c-SRC-, and p38 MAPK-dependent pathway. *J. Biol. Chem.* **2008**, *283*, 12898–12908. [[CrossRef](#)] [[PubMed](#)]
211. Kracklauer, M.P.; Schmidt, C.; Sclabas, G.M. TGF- β 1 signaling via α v β 6 integrin. *Mol. Cancer* **2003**, *2*, 28. [[CrossRef](#)] [[PubMed](#)]
212. Mamuya, F.A.; Duncan, M.K. α v integrins and TGF- β -induced EMT: A circle of regulation. *J. Cell. Mol. Med.* **2012**, *16*, 445–455. [[CrossRef](#)] [[PubMed](#)]
213. Heckmann, D.; Kessler, H. Design and chemical synthesis of integrin ligands. *Methods Enzymol.* **2007**, *426*, 463–503. [[CrossRef](#)] [[PubMed](#)]
214. Kapp, T.G.; Rechenmacher, F.; Neubauer, S.; Maltsev, O.V.; Cavalcanti-Adam, A.E.; Zarka, R.; Reuning, U.; Notni, J.; Wester, H.-J.; Mas-Moruno, C.; et al. A comprehensive evaluation of the activity and selectivity profile of ligands for RGD-binding integrins. *Sci. Rep.* **2017**, *7*, 39805. [[CrossRef](#)] [[PubMed](#)]
215. Dechantsreiter, M.A.; Planker, E.; Mathä, B.; Lohof, E.; Hölzemann, G.; Jonczyk, A.; Goodman, S.L.; Kessler, H. N-Methylated cyclic RGD Peptides as highly active and selective α v β 3 integrin antagonists. *J. Med. Chem.* **1999**, *42*, 3033–3040. [[CrossRef](#)] [[PubMed](#)]
216. Mas-Moruno, C.; Rechenmacher, F.; Kessler, H. Cilengitide: The first anti-angiogenic small molecule drug candidate. Design, synthesis and clinical evaluation. *Anticancer Agents Med. Chem.* **2010**, *10*, 753–768. [[CrossRef](#)] [[PubMed](#)]
217. Reardon, D.A.; Neyns, B.; Weller, M.; Tonn, J.C.; Nabors, L.B.; Stupp, R. Cilengitide: An RGD pentapeptide α v β 3 and α v β 5 integrin inhibitor in development for glioblastoma and other malignancies. *Future Oncol.* **2011**, *7*, 339–354. [[CrossRef](#)] [[PubMed](#)]
218. Nabors, L.B.; Fink, K.L.; Mikkelsen, T.; Grujicic, D.; Tarnawski, D.H.N.; Mazurkiewicz, M.; Salacz, M.; Ashby, L.; Zagonel, V.; Depenni, R.; et al. Two Cilengitide regimens in combination with standard treatment for patients with newly diagnosed glioblastoma and unmethylated MGMT gene promoter: Results of the open-label, controlled, randomized phase II CORE study. *Neuro-Oncology* **2015**, *17*, 708–717. [[CrossRef](#)] [[PubMed](#)]
219. Mason, W.P. End of the road: Confounding results of the CORE trial terminate the arduous journey of Cilengitide for glioblastoma. *Neuro-Oncology* **2015**, *17*, 634–635. [[CrossRef](#)] [[PubMed](#)]
220. Heckmann, D.; Meyer, A.; Laufer, B.; Zahn, G.; Stragies, R.; Kessler, H. Rational design of highly active and selective ligands for the α 5 β 1 integrin receptor. *ChemBioChem* **2008**, *9*, 1397–1407. [[CrossRef](#)] [[PubMed](#)]

221. Neubauer, S.; Rechenmacher, F.; Brimiouille, R.; Di Leva, F.S.; Bochen, A.; Sobahi, T.R.; Schottelius, M.; Novellino, E.; Mas-Moruno, C.; Marinelli, L.; et al. Pharmacophoric modifications lead to superpotent $\alpha v \beta 3$ integrin ligands with suppressed $\alpha 5 \beta 1$ activity. *J. Med. Chem.* **2014**, *57*, 3410–3417. [[CrossRef](#)] [[PubMed](#)]
222. Stragies, R.; Osterkamp, F.; Zischinsky, G.; Vossmeier, D.; Kalkhof, H.; Reimer, U.; Zahn, G. Design and Synthesis of a new class of selective integrin $\alpha 5 \beta 1$ antagonists. *J. Med. Chem.* **2007**, *50*, 3786–3794. [[CrossRef](#)] [[PubMed](#)]
223. Maltsev, O.V.; Marelli, U.K.; Kapp, T.G.; Di Leva, F.S.; Di Maro, S.; Nieberler, M.; Reuning, U.; Schwaiger, M.; Novellino, E.; Marinelli, L.; et al. Stable peptides instead of stapled peptides: Highly potent $\alpha v \beta 6$ -selective integrin ligands. *Angew. Chem. Int. Ed.* **2016**, *55*, 1535–1539. [[CrossRef](#)] [[PubMed](#)]
224. Indrevoll, B.; Kindberg, G.M.; Solbakken, M.; Bjurgert, E.; Johansen, J.H.; Karlsen, H.; Mendizabal, M.; Cuthbertson, A. NC-100717: A versatile RGD peptide scaffold for angiogenesis imaging. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 6190–6193. [[CrossRef](#)] [[PubMed](#)]
225. Hynes, R.O. Integrins: Versatility, modulation, and signaling in cell adhesion. *Cell* **1992**, *69*, 11–25. [[CrossRef](#)]
226. Aumailley, M.; Gurrath, M.; Müller, G.; Calvete, J.; Timpl, R.; Kessler, H. Arg-Gly-Asp constrained within cyclic pentapeptides. Strong and selective inhibitors of cell adhesion to vitronectin and laminin fragment P1. *FEBS Lett.* **1991**, *291*, 50–54. [[CrossRef](#)]
227. Haubner, R.; Finsinger, D.; Kessler, H. Stereoisomeric Peptide Libraries and Peptidomimetics for designing selective inhibitors of the $\alpha v \beta 3$ integrin for a new cancer therapy. *Angew. Chem. Int. Ed.* **1997**, *36*, 1374–1389. [[CrossRef](#)]
228. Gurrath, M.; Müller, G.; Kessler, H.; Aumailley, M.; Timpl, R. Conformation/activity studies of rationally designed potent anti-adhesive RGD peptides. *Eur. J. Biochem.* **1992**, *210*, 911–921. [[CrossRef](#)] [[PubMed](#)]
229. Chatterjee, J.; Rechenmacher, F.; Kessler, H. N-Methylation of peptides and proteins: An important element for modulating biological functions. *Angew. Chem. Int. Ed.* **2013**, *52*, 254–269. [[CrossRef](#)] [[PubMed](#)]
230. Collier, B.S.; Shattil, S.J. The GPIIb/IIIa (integrin $\alpha IIb \beta 3$) odyssey: A technology-driven saga of a receptor with twists, turns, and even a bend. *Blood* **2008**, *112*. [[CrossRef](#)] [[PubMed](#)]
231. Zhu, J.; Luo, B.-H.; Xiao, T.; Zhang, C.; Nishida, N.; Springer, T.A. Structure of a complete integrin ectodomain in a physiologic resting state and activation and deactivation by applied forces. *Mol. Cell* **2008**, *32*, 849–861. [[CrossRef](#)] [[PubMed](#)]
232. Nagae, M.; Re, S.; Mihara, E.; Nogi, T.; Sugita, Y.; Tagaki, J. Crystal structure of $\alpha 5 \beta 1$ integrin ectodomain: Atomic details of the fibronectin receptor. *J. Cell Biol.* **2012**, *187*, 131–140. [[CrossRef](#)] [[PubMed](#)]
233. Pallarola, D.; Platzman, I.; Bochen, A.; Cavalcanti-Adam, E.A.; Axmann, M.; Kessler, H.; Geiger, B.; Spatz, J.P. Focal adhesion stabilization by enhanced integrin-cRGD binding affinity. *BioNanoMaterials* **2017**, *18*, 1–2. [[CrossRef](#)]
234. Ley, K.; Rivera-Nieves, J.; Sandborn, W.J.; Shattil, S. Integrin-based therapeutics: Biological basis, clinical use and new drugs. *Nat. Rev. Drug Discov.* **2016**, *15*, 173–183. [[CrossRef](#)] [[PubMed](#)]
235. Hovlid, M.L.; Steinmetz, N.F.; Laufer, B.; Lau, J.L.; Kuzelka, J.; Wang QHyypä, T.; Nemerow, G.R.; Kessler, H.; Manchester, M.; Finn, M.G. Guiding plant virus particles to integrin-displaying cells. *Nanoscale* **2012**, *4*, 3698–3705. [[CrossRef](#)] [[PubMed](#)]
236. Shroff, K.; Kokkoli, E. PEGylated liposomal doxorubicin targeted to $\alpha 5 \beta 1$ -expressing MDA-MB-231 breast cancer cells. *Langmuir* **2012**, *28*, 4729–4736. [[CrossRef](#)] [[PubMed](#)]
237. Beer, A.J.; Haubner, R.; Goebel, M.; Luderschmidt, S.; Spilker, M.E.; Wester, H.J.; Weber, W.A.; Schwaiger, M. Biodistribution and pharmacokinetics of the $\alpha v \beta 3$ -selective tracer ^{18}F -galacto-RGD in cancer patients. *J. Nucl. Med.* **2005**, *46*, 133–141.
238. Beer, A.J.; Haubner, R.; Sarbia, M.; Goebel, M.; Luderschmidt, S.; Grosu, A.L.; Schnell, O.; Niemeyer, M.; Kessler, H.; Wester, H.J.; et al. Positron emission tomography using ^{18}F -Galacto-RGD identifies the level of integrin $\alpha v \beta 3$ expression in man. *Clin. Cancer Res.* **2006**, *12*, 3942–3949. [[CrossRef](#)] [[PubMed](#)]
239. Haubner, R.; Weber, W.A.; Beer, A.J.; Vabuliene, E.; Reim, D.; Sarbia, M.; Becker, K.F.; Goebel, M.; Hein, R.; Wester, H.J.; et al. Noninvasive visualization of the activated $\alpha v \beta 3$ integrin in cancer patients by positron emission tomography and ^{18}F -Galacto-RGD. *PLoS Med.* **2005**, *2*, e70. [[CrossRef](#)] [[PubMed](#)]
240. Chen, H.; Niu, G.; Wu, H.; Chen, X. Clinical application of radiolabeled RGD peptides for PET imaging of integrin $\alpha v \beta 3$. *Theranostics* **2016**, *6*, 78–92. [[CrossRef](#)] [[PubMed](#)]

241. Beer, A.J.; Haubner, R.; Wolf, I.; Goebel, M.; Luderschmidt, S.; Niemeyer, M.; Grosu, A.-L.; Martinez, M.-J.; Wester, H.J.; Weber, W.A.; et al. PET-based human dosimetry of ^{18}F -galacto-RGD, a new radiotracer for imaging $\alpha v\beta 3$ expression. *J. Nucl. Med.* **2006**, *47*, 763–769. [[PubMed](#)]
242. McParland, B.J.; Miller, M.P.; Spinks, T.J.; Kenny, L.M.; Osman, S.; Khela, M.K.; Aboagye, E.; Coombes, R.C.; Hui, A.M.; Cohen, P.S. The biodistribution and radiation dosimetry of the Arg-Gly-Asp peptide ^{18}F -AH111585 in healthy volunteers. *J. Nucl. Med.* **2008**, *49*, 1664–1667. [[CrossRef](#)] [[PubMed](#)]
243. Doss, M.; Kolb, H.C.; Zhang, J.J.; Belanger, M.J.; Stubbs, J.B.; Stabin, M.G.; Hostetler, E.D.; Alpaugh, R.K.; von Mehren, M.; Walsh, J.C.; et al. Biodistribution and radiation dosimetry of the integrin marker ^{18}F -RGD-K5 determined from whole-body PET/CT in monkeys and humans. *J. Nucl. Med.* **2012**, *53*, 787–795. [[CrossRef](#)] [[PubMed](#)]
244. Kim, J.H.; Lee, J.S.; Kang, K.W.; Lee, H.Y.; Han, S.W.; Kim, T.Y.; Lee, Y.S.; Jeong, J.M.; Lee, D.S. Whole-body distribution and radiation dosimetry of ^{68}Ga -NOTA-RGD, a positron emission tomography agent for angiogenesis imaging. *Cancer Biother. Radiopharm.* **2012**, *27*, 65–71. [[CrossRef](#)] [[PubMed](#)]
245. Wan, W.; Guo, N.; Pan, D.; Yu, C.; Weng, Y.; Luo, S.; Ding, H.; Xu, Y.; Wang, L.; Lang, L.; et al. First experience of ^{18}F -Alfatide in lung cancer patients using a new lyophilized kit for rapid radiofluorination. *J. Nucl. Med.* **2013**, *54*, 691–698. [[CrossRef](#)] [[PubMed](#)]
246. Mitra, E.S.; Goris, M.L.; Iagaru, A.H.; Kardan, A.; Burton, L.; Berganos, R.; Chang, E.; Lui, S.; Shen, B.; Chin, F.T.; et al. Pilot pharmacokinetic and dosimetric studies of ^{18}F -FPPRGD2: A PET radiopharmaceutical agent for imaging $\alpha v\beta 3$ integrin levels. *Radiology* **2011**, *260*, 182–191. [[CrossRef](#)] [[PubMed](#)]
247. Yu, C.; Pan, D.; Mi, B.; Xu, Y.; Lang, L.; Niu, G.; Yang, M.; Wan, W.; Chen, X. ^{18}F -Alfatide II PET/CT in healthy human volunteers and patients with brain metastases. *Eur. J. Nucl. Med. Mol. Imaging* **2015**, *42*, 2021–2028. [[CrossRef](#)] [[PubMed](#)]
248. Zheng, K.; Liang, N.; Zhang, J.; Lang, L.; Zhang, W.; Li, S.; Zhao, J.; Niu, G.; Li, F.; Zhu, Z.; et al. ^{68}Ga -NOTA-PRGD2 PET/CT for integrin imaging in patients with lung cancer. *J. Nucl. Med.* **2015**, *56*, 1823–1827. [[CrossRef](#)] [[PubMed](#)]
249. Atkinson, S.J.; Ellison, T.S.; Steri, V.; Gould, E.; Robinson, S.D. Redefining the role(s) of endothelial $\alpha v\beta 3$ -integrin in angiogenesis. *Biochem. Soc. Trans.* **2014**, *42*, 1590–1595. [[CrossRef](#)] [[PubMed](#)]
250. Fässler, R.; Meyer, M. Consequences of lack of $\beta 1$ integrin gene expression in mice. *Genes Dev.* **1995**, *9*, 1896–1908. [[CrossRef](#)] [[PubMed](#)]
251. Tanjore, H.; Zeisberg, E.M.; Gerami-Naini, B.; Kalluri, R. $\beta 1$ integrin expression on endothelial cells is required for angiogenesis but not for vasculogenesis. *Dev. Dyn.* **2007**, *237*, 75–82. [[CrossRef](#)] [[PubMed](#)]
252. Neubauer, S.; Rechenmacher, F.; Beer, A.J.; Curnis, F.; Pohle, K.; D'Alessandria, C.; Wester, H.J.; Reuning, U.; Corti, A.; Schwaiger, M.; et al. Selective imaging of the angiogenic relevant integrins $\alpha 5\beta 1$ and $\alpha v\beta 3$. *Angew. Chem. Int. Ed.* **2013**, *52*, 11656–11659. [[CrossRef](#)] [[PubMed](#)]
253. D'Alessandria, C.; Pohle, K.; Rechenmacher, S.; Neubauer, S.; Notni, J.; Wester, H.J.; Schwaiger, M.; Kessler, H.; Beer, A.J. In vivo biokinetic and metabolic characterization of the ^{68}Ga -labeled $\alpha 5\beta 1$ -selective peptidomimetic FR366. *Eur. J. Nucl. Med. Mol. Imaging* **2016**, *43*, 953–963. [[CrossRef](#)]
254. Haubner, R.; Maschauer, S.; Einsiedel, J.; Eder, I.E.; Rangger, C.; Gmeiner, P.; Virgolini, I.; Prante, O. H-CRRETAWAC-OH, a lead structure for the development of radiotracer targeting integrin $\alpha 5\beta 1$? *Biomed. Res. Int.* **2014**, *2014*, 243185. [[CrossRef](#)] [[PubMed](#)]
255. Zhao, H.T.; Gao, H.N.; Zhai, L.P.; Liu, X.J.; Jia, B.; Shi, J.Y.; Wang, F. Tc-99m-HisoDGR as a Potential SPECT Probe for orthotopic glioma detection via targeting of integrin $\alpha 5\beta 1$. *Bioconjug. Chem.* **2016**, *27*, 1259–1266. [[CrossRef](#)] [[PubMed](#)]
256. Notni, J.; Steiger, K.; Hoffmann, F.; Reich, D.; Kapp, F.G.; Rechenmacher, T.; Neubauer, S.; Kessler, H.; Wester, H.J. Complementary, selective PET-Imaging of integrin subtypes $\alpha 5\beta 1$ and $\alpha v\beta 3$ using Ga-68-aquibepirin and Ga-68-avebetrin. *J. Nucl. Med.* **2016**, *57*, 460–466. [[CrossRef](#)] [[PubMed](#)]
257. Notni, J.; Steiger, K.; Hoffmann, F.; Reich, D.; Schwaiger, M.; Kessler, H.; Wester, H.J. Variation of specific activities of Ga-68-aquibepirin and Ga-68-avebetrin enables selective PET-imaging of different expression levels of integrins $\alpha 5\beta 1$ and $\alpha v\beta 3$. *J. Nucl. Med.* **2016**, *57*, 1618–1624. [[CrossRef](#)] [[PubMed](#)]
258. Liu, H.; Wu, Y.; Wang, F.; Liu, Z. Molecular imaging of integrin $\alpha v\beta 6$ expression in living subjects. *Am. J. Nucl. Med. Mol. Imaging* **2014**, *4*, 333–345. [[PubMed](#)]

259. Zhang, Y.; Sun, Y.; Yang, F.; Guo, J.; He, J.; Wu, Q.; Cao, W.; Lv, L.; Zheng, H.; Zhang, Z. Induction of partial protection against foot and mouth disease virus in guinea pigs by neutralization with the integrin $\alpha\beta 6$ -1 subunit. *Viruses* **2013**, *5*, 1114–1130. [[CrossRef](#)] [[PubMed](#)]
260. Goodman, S.L.; Hölzemann, G.; Sulyok, G.A.; Kessler, H. Nanomolar small molecule inhibitors for $\alpha\beta 6$, $\alpha\beta 5$, and $\alpha\beta 3$ integrins. *J. Med. Chem.* **2002**, *45*, 1045–1051. [[CrossRef](#)] [[PubMed](#)]
261. Kraft, S.; Diefenbach, B.; Mehta, R.; Jonczyk, A.; Luckenbach, G.A.; Goodman, S.L. Definition of an unexpected ligand recognition motif for $\alpha\beta 6$ integrin. *J. Biol. Chem.* **1999**, *274*, 1979–1985. [[CrossRef](#)] [[PubMed](#)]
262. Hausner, S.H.; DiCara, D.; Marik, J.; Marshall, J.F.; Sutcliffe, J.L. Use of a peptide derived from foot-and-mouth disease virus for the noninvasive imaging of human cancer: Generation and evaluation of 4- ^{18}F fluorobenzoyl a20fmdv2 for in vivo imaging of integrin $\alpha\beta 6$ expression with positron emission tomography. *Cancer Res.* **2007**, *67*, 7833–7840. [[CrossRef](#)] [[PubMed](#)]
263. Li, S.; McGuire, M.J.; Lin, M.; Liu, Y.H.; Oyama, T.; Sun, X.; Brown, K.C. Synthesis and characterization of a high-affinity $\alpha\beta 6$ -specific ligand for in vitro and in vivo applications. *Mol. Cancer Ther.* **2009**, *8*, 1239–1249. [[CrossRef](#)] [[PubMed](#)]
264. Kimura, R.H.; Teed, R.; Hackel, B.J.; Pysz, M.A.; Chuang, C.Z.; Sathirachinda, A.; Willmann, J.K.; Gambhir, S.S. Pharmacokinetically stabilized cystine knot peptides that bind $\alpha\beta 6$ integrin with single-digit nanomolar affinities for detection of pancreatic cancer. *Clin. Cancer Res.* **2012**, *18*, 839–849. [[CrossRef](#)] [[PubMed](#)]
265. Ahmedah, H.T.; Patterson, L.H.; Shnyder, S.D.; Sheldrake, H.M. RGD-binding integrins in head and neck cancers. *Cancers* **2017**, *9*, 56. [[CrossRef](#)] [[PubMed](#)]
266. John, A.E.; Luckett, J.C.; Tatler, A.L.; Awais, R.O.; Desai, A.; Habgood, A.; Ludbrook, S.; Blanchard, A.D.; Perkins, A.C.; Jenkins, R.G.; et al. Preclinical SPECT/CT imaging of $\alpha\beta 6$ integrins for molecular stratification of idiopathic pulmonary fibrosis. *J. Nucl. Med.* **2013**, *54*, 2146–2152. [[CrossRef](#)] [[PubMed](#)]
267. Liu, Z.; Liu, H.; Ma, T.; Sun, X.; Shi, J.; Jia, B.; Sun, Y.; Zhan, J.; Zhang, H.; Zhu, Z.; et al. Integrin $\alpha\beta 6$ -targeted SPECT imaging for pancreatic cancer detection. *J. Nucl. Med.* **2014**, *55*, 989–994. [[CrossRef](#)] [[PubMed](#)]
268. Zhu, X.; Li, J.; Hong, Y.; Kimura, R.H.; Ma, X.; Liu, H.; Qin, C.; Hu, X.; Hayes, T.R.; Benny, P.; et al. $^{99\text{m}}\text{Tc}$ -labeled cystine knot peptide targeting integrin $\alpha\beta 6$ for tumor SPECT imaging. *Mol. Pharm.* **2014**, *11*, 1208–1217. [[CrossRef](#)] [[PubMed](#)]
269. Hausner, S.H.; Abbey, C.K.; Bold, R.J.; Gagnon, M.K.; Marik, J.; Marshall, J.F.; Stanecki, C.E.; Sutcliffe, J.L. Targeted in vivo imaging of integrin $\alpha\beta 6$ with an improved radiotracer and its relevance in a pancreatic tumor model. *Cancer Res.* **2009**, *69*, 5843–5850. [[CrossRef](#)] [[PubMed](#)]
270. Singh, A.N.; McGuire, M.J.; Li, S.; Hao, G.; Kumar, A.; Sun, X.; Brown, K.C. Dimerization of a phage-display selected peptide for imaging of $\alpha\beta 6$ -integrin: Two approaches to the multivalent effect. *Theranostics* **2014**, *4*, 745–760. [[CrossRef](#)] [[PubMed](#)]
271. Hausner, S.H.; Bauer, N.; Sutcliffe, J.L. In vitro and in vivo evaluation of the effects of aluminum ^{18}F fluoride radiolabeling on an integrin $\alpha\beta 6$ -specific peptide. *Nucl. Med. Biol.* **2014**, *41*, 43–50. [[CrossRef](#)] [[PubMed](#)]
272. Hausner, S.H.; Bauer, N.; Hu, L.Y.; Knight, L.M.; Sutcliffe, J.L. The effect of bi-terminal PEGylation of an integrin $\alpha\beta 6$ -targeted ^{18}F peptide on pharmacokinetics and tumor uptake. *J. Nucl. Med.* **2015**, *56*, 784–790. [[CrossRef](#)] [[PubMed](#)]
273. Hausner, S.H.; Carpenter, R.D.; Bauer, N.; Sutcliffe, J.L. Evaluation of an integrin $\alpha\beta 6$ -specific peptide labeled with ^{18}F fluorine by copper-free, strain-promoted click chemistry. *Nucl. Med. Biol.* **2013**, *40*, 233–239. [[CrossRef](#)] [[PubMed](#)]
274. Altmann, A.; Sauter, M.; Roesch, S.; Mier, W.; Warta, R.; Debus, J.; Dyckhoff, G.; Herold-Mende, C.; Haberkorn, U. Identification of a novel ITG $\alpha\beta 6$ -binding peptide using protein separation and phage display. *Clin. Cancer Res.* **2017**. [[CrossRef](#)] [[PubMed](#)]
275. Notni, J.; Reich, D.; Maltsev, O.V.; Kapp, T.G.; Steiger, K.; Hoffmann, F.; Esposito, I.; Weichert, W.; Kessler, H.; Wester, H.J. In Vivo pet imaging of the “cancer integrin” $\alpha\beta 6$ using gallium-68 labelled cyclic RGD nonapeptides. *J. Nucl. Med.* **2016**, *58*, 671–677. [[CrossRef](#)] [[PubMed](#)]
276. Zhang, Z.-Y.; Xu, K.; He, Q.-S.; Niu, W.-B.; Wang, J.-Y.; Mi, Y.-T.; Wang, J.-S.; Wang, G.-Q.; Yang, G.-Y.; Niu, J. Signaling and regulatory mechanisms of integrin $\alpha\beta 6$ on the apoptosis of colon cancer cells. *Cancer Lett.* **2008**, *266*, 209–215. [[CrossRef](#)] [[PubMed](#)]

277. Kogelberg, H.; Tolner, B.; Thomas, G.J.; Di Cara, D.; Minogue, S.; Ramesh, B.; Sodha, S.; Marsh, D.; Lowdell, M.W.; Meyer, T.; et al. Engineering a single-chain fv antibody to $\alpha v \beta 6$ integrin using the specificity-determining loop of a foot-and-mouth disease virus. *J. Mol. Biol.* **2008**, *382*, 385–401. [[CrossRef](#)] [[PubMed](#)]
278. Integrin Alpha-v-Beta and [^{18}F]-R01-MG-F2 PET/CT in Measuring Response in Patients with Pancreatic Cancer and Healthy Volunteers. Available online: <https://clinicaltrials.gov/ct2/show/NCT02683824> (accessed on 7 August 2017).
279. First-in-Human Positron Emission Tomography Study Using the ^{18}F - $\alpha v \beta 6$ -Binding Peptide. Available online: <https://clinicaltrials.gov/ct2/show/NCT03164486> (accessed on 7 August 2017).



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).