



Review Article

A review of ovary torsion

Ci Huang^a, Mun-Kun Hong^{a,b}, Dah-Ching Ding^{a,b*}

^aDepartment of Obstetrics and Gynecology, Buddhist Tzu Chi General Hospital and Tzu Chi University, Hualien, Taiwan,
^bInstitute of Medical Sciences, Tzu Chi University, Hualien, Taiwan

Received : 29-Mar-2017
Revised : 07-Apr-2017
Accepted : 02-May-2017

ABSTRACT

Ovarian torsion is a rare but emergency condition in women. Early diagnosis is necessary to preserve the function of the ovaries and tubes and prevent severe morbidity. Ovarian torsion refers to complete or partial rotation of the adnexal supporting organ with ischemia. It can affect females of all ages. Ovarian torsion occurs in around 2%–15% of patients who have surgical treatment of adnexal masses. The main risk in ovarian torsion is an ovarian mass. The most common symptom of ovarian torsion is acute onset of pelvic pain, followed by nausea and vomiting. Pelvic ultrasonography can provide information on ovarian cysts. Once ovarian torsion is suspected, surgery or detorsion is the mainstay of diagnosis and treatment.

KEYWORDS: *Abdominal pain, Ovarian cyst, Ovarian torsion, Pelvic pain, Ultrasound*

INTRODUCTION

Ovarian torsion, which affects females of all ages, is a gynecological emergency [1-3]. It refers to a complete or partial rotation of the adnexal supporting organ, resulting in ischemic changes in the ovary. Torsions more commonly involve both the ovary and fallopian tube, and there are fewer cases of isolated torsion involving either one (one in 1.5 million women) [4-6]. Torsion involving paratubal or paraovarian cysts has also been found [2,7,8]. Early diagnosis and surgery are essential to protect ovarian and tubal function and prevent severe morbidity [9,10].

INFUNDIBULOPELVIC AND UTERO-OVARIAN LIGAMENTS

The infundibulopelvic ligaments suspend the movable ovary, allowing the ovary to position laterally or posteriorly to the uterus. The ovarian vessels travel along the infundibulopelvic ligaments which attach to the pelvic sidewall. Because adnexal tissue is not fixed, a big leading point, such as tumorous growth, can induce twisting. The other side of the ovary is connected to the uterus by the utero-ovarian (UO) ligament. The UO ligament is composed of muscular and fibrous tissue. The function of the UO ligament is to connect the ovary to the uterus and support it, and it also supplies blood from the uterine artery to the ovary [11].

PATHOGENESIS

Ovarian torsion occurs when an ovarian cyst or mass presents and rotates both the infundibulopelvic ligament and the UO ligament. The cyst or mass is usually a benign lesion over 5 cm in diameter [12-15]. Torsion can also occur in

normal ovaries, however, particularly in premenarchal girls who have elongated infundibulopelvic ligaments [16-18]. However, the occurrence of ovarian torsion may decrease thereafter because the ligament shortens when premenarchal girls mature to puberty.

EPIDEMIOLOGY

A 10-year review of 128 patients with adnexal torsion states that 2.7% of emergency surgery cases involved ovarian torsion [19]. Another 10-year study showed that 15% of 135 patients with surgically treated adnexal masses had torsion [20]. Totally, around 2%–15% of patients who had surgical treatment of adnexal masses had ovarian torsion. Most ovarian torsion occurs in the reproductive age group, and it is less common in premenarchal girls and postmenopausal women (17.2% of cases) [21].

Risk factors

More than 80% patients with ovarian torsion had ovarian masses of 5 cm or larger, indicating that the primary risk in ovarian torsion is an ovarian mass [12,15,22]. The sizes of ovarian masses are correlated with the risk of the torsion. Ovarian torsion has been reported to occur with masses from 1 to 30 cm (mean 9.5 cm) [15], but it can happen with any size mass. Ovulation induction for treatment of infertility may cause multiple large ovarian follicular cysts; the large cysts carry an increased risk of torsion [23].

**Address for correspondence:*

Dr. Dah-Ching Ding,
Department of Obstetrics and Gynecology, Buddhist Tzu Chi General Hospital, 707, Section 3, Chung-Yang Road, Hualien, Taiwan.
E-mail: dah1003@yahoo.com.tw

Access this article online

Quick Response Code:



Website: www.tcmjmed.com

DOI: 10.4103/tcmj.tcmj_55_17

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Huang C, Hong MK, Ding DC. A review of ovary torsion. Tzu Chi Med J 2017;29:143-7.

Benign and malignant tumors in ovarian torsion

Ovarian torsion is more likely to occur with a benign tumor than in a malignancy. The incidence of ovarian torsion with ovarian malignancy was <2% in reported case series [13,22,24,25].

Ovarian torsion in the premenarchal population

Compared with older women, premenarchal girls with ovarian torsion are more commonly found to have a normal ovary [26]. More than 50% of patients under 15 years old with torsion have normal ovaries [24,27]. Torsion occurred more in patients with normal adnexa (7/11) than those with abnormal adnexa (4/46) [14].

Ovarian torsion in pregnancy

About 10%–22% of ovarian torsion occurred in pregnancy [15,22,24,28]. The incidence is higher at 10–17 weeks of gestation with ovarian masses larger than 4 cm [29,30]. Pregnant women with adnexal masses 4 cm or greater had a 1%–6% lower incidence of torsion compared with nonpregnant women [31,32].

Summary of epidemiology and risk factors

The incidence of ovarian torsion ranges from 2% to 15% in patients who have surgical treatment of adnexal masses. Ovarian tumors larger than 5 cm carry a risk of ovarian torsion. About 10%–22% of ovarian torsion occurs in pregnant women.

CLINICAL PRESENTATION

Ovarian torsion due to an adnexal mass causes various symptoms and signs on clinical presentation. The most common symptom is acute onset of lower abdominal pain, followed by nausea and vomiting [15,22,24,33-35]. Some patients experience waves of nausea with or without vomiting [15,22,24,33]. The abdominal pain is usually off and on with a sudden onset. Most reported patients presented for evaluation 1 or more days up as late as 210 days after pain onset [15,22,36,37]. Premenarchal patients tended to mention diffuse pain because it was difficult for them to localize the pain [38]. The uncomfortable symptoms and signs were considered to be caused by the adnexal torsion. Ovarian torsion without infective disease resulting in a low-grade fever has been found in some patients [15,22,24,33].

EVALUATION AND DIAGNOSIS

On clinical presentation, the first approach to a patient is a medical history and physical examination. The medical history should include any recent diagnosis of an adnexal mass, recurrent abdominal pain, and low-grade fever. The physical examination should include a search for a pelvic mass or pain. Laboratory evaluation should include serum human chorionic gonadotropin, a hematocrit, white blood cell count, and electrolyte panel.

There is no serum marker for a diagnosis of adnexal torsion. Several serum markers can hint at an adnexal tumor type. Serum human chorionic gonadotropin can reveal pregnancy or an ovarian germ cell tumor. CA-125 may indicate a malignant ovary tumor or endometrioma. Some studies have found an association between an increased level of serum interleukin-6

and ovarian torsion [39,40], although further research such as oxidative stress during ovarian torsion is needed [41].

Imaging studies are the most important when evaluating a pelvic mass [42]. Ultrasonography is the first-line diagnostic assessment. A torsed ovary may be rounded and enlarged compared with the contralateral ovary, because of edema or vascular and lymph engorgement [43,44]. An ultrasound can easily distinguish an ovarian mass by its components, location, density, Doppler flow, and size. There can be decreased or absent Doppler flow in the vessels of a torsed ovary [45-47]. One prospective study reported that Doppler flow has high sensitivity and specificity [48]; another retrospective study showed low sensitivity and high specificity in the diagnosis of ovarian torsion [26]. It is not the gold standard for diagnosis, but it is a good tool. Two other studies suggested that a whirlpool sign is highly sensitive for ovary torsion [49,50]. The whirlpool sign shows a twisted vascular pedicle, and a Doppler sonogram reveals circular vessels within the mass. However, further study on the diagnosis of ovarian torsion is necessary to determine the usefulness of this sign in ovary torsion.

Magnetic resonance imaging (MRI) is expensive but helpful in diagnosing ovarian torsion if findings on ultrasound are equivocal [51-57]. MRI can demonstrate the components of a mass in more detail than an ultrasound. Computed tomography (CT), however, is not typically used in ovary torsion because of radiation and density, but patients with acute abdominal or pelvic pain need to undergo CT to exclude diagnoses such as appendicitis, diverticulitis, and others.

Finally, direct visualization is needed for a definitive diagnosis of ovary torsion. Hence, the diagnosis needs to be surgical proven for early rescue of ovary function.

MANAGEMENT

The gold standard to treat ovary torsion is surgery, and this is also the only way to confirm the torsion. There are two surgical methods, laparoscopy and laparotomy. A laparoscopic approach has become a popular procedure. However, if cancer of the ovary or fallopian tube is suspected, a laparotomy should be done [58,59]. While performing the surgery, it is necessary to assess ovarian viability and preserve its function. The only way to determine the viability of a torsed ovary during surgery is by gross visual inspection. In the conventional view of point, dark and enlarged ovaries may have vascular and lymphatic congestion, and may seem nonviable. However, multiple studies have suggested that even those black or blue-like ovaries may retain ovarian function following detorsion [60-67]. Postoperative follow-up with ultrasound showed over 80% of patients had normal follicular development after detorsion [55,63,66,67]. Animal study showed that there may not be total occlusion of the artery in ovarian torsion even with venous and lymphatic congestion [68]. In recent years, the mainstay of the treatment for ovarian torsion has been surgical evaluation and preserving ovarian function. There are many ways to perform the surgery and detorsion and ovarian conservation are almost always recommended now rather than salpingo-oophorectomy [69]. An ovarian

cystectomy is often performed for a benign ovarian mass. If malignancy is highly suspected, a salpingo-oophorectomy is needed. According to many observational studies, detorsion is associated with preserved ovarian function [60-64]. The earlier the approach to torsion, the higher is the chance to preserve function. An animal study showed that necrosis might develop after occlusion of ovarian vessels for 36 h or longer [68]. After the symptoms have developed, ovarian conservation reportedly decreases with time [27,65]. No evidence suggests that detorsion increases adverse events postoperatively [70]. Management in pregnant women is similar to that in nonpregnant patients, and laparoscopic surgery is safe for torsion in pregnant women [28,71-73]. Neonates with ovarian torsion often present with irritability and the condition can be treated with laparoscopic surgery [74-76].

PREVENTION OF RECURRENCE

There is a risk of recurrence after detorsion, but the incidence and causes are unknown [14,17,77-79]. According to recent research, several methods can be used to decrease the risk of recurrence. One method is suppression of ovarian cysts by oral contraceptives [80-83]. Another method is an oophorectomy [72,84,85]. However, both approaches lack long-term follow-up and systematic study.

CONCLUSION

Although the diagnosis of ovarian torsion is difficult and challenging, careful analysis of presenting symptoms (such as sudden onset of lower abdominal pain) is very critical. Pelvic ultrasonography can provide information on ovarian cysts. Once ovarian torsion is suspected, surgery is the mainstay of diagnosis and treatment. Ovarian cystectomy, oophorectomy, or conservative treatment with detorsion can be the treatment of choice.

Acknowledgment

The authors would like to thank Dr. Jon-Son Kuo for English editing.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- McWilliams GD, Hill MJ, Dietrich CS 3rd. Gynecologic emergencies. *Surg Clin North Am* 2008;88:265-83, vi.
- Muolokwu E, Sanchez J, Bercau JL, Sangi-Haghpeykar H, Banszek T, Brandt ML, et al. The incidence and surgical management of paratubal cysts in a pediatric and adolescent population. *J Pediatr Surg* 2011;46:2161-3.
- Huchon C, Fauconnier A. Adnexal torsion: A literature review. *Eur J Obstet Gynecol Reprod Biol* 2010;150:8-12.
- Ding DC, Hsu S, Kao SP. Isolated torsion of the hydrosalpinx in a postmenopausal woman. *JLS* 2007;11:252-4.
- Antoniou N, Varras M, Akrivis C, Kitsiou E, Stefanaki S, Salamalekis E, et al. Isolated torsion of the fallopian tube: A case report and review of the literature. *Clin Exp Obstet Gynecol* 2004;31:235-8.
- van der Zanden M, Nap A, van Kints M. Isolated torsion of the fallopian tube: A case report and review of the literature. *Eur J Pediatr* 2011;170:1329-32.
- Schrager J, Robles G, Platz T. Isolated fallopian tube torsion: A rare entity in a premenarcheal female. *Am Surg* 2012;78:118-9.
- Said MR, Bamigboye V. Twisted paraovarian cyst in a young girl. *J Obstet Gynaecol* 2008;28:549-50.
- Argenta PA, Yeagley TJ, Ott G, Sondheimer SJ. Torsion of the uterine adnexa. Pathologic correlations and current management trends. *J Reprod Med* 2000;45:831-6.
- Robertson JJ, Long B, Koyfman A. Myths in the evaluation and management of ovarian torsion. *J Emerg Med* 2017;52:449-56.
- Anne A, Dalley A. Grant's Atlas of Anatomy. 13th ed. Philadelphia: Lippincott Williams & Wilkins; 2012.
- Varras M, Tsikini A, Polyzos D, Samara CH, Hadjopoulos G, Akrivis CH, et al. Uterine adnexal torsion: Pathologic and gray-scale ultrasonographic findings. *Clin Exp Obstet Gynecol* 2004;31:34-8.
- Oltmann SC, Fischer A, Barber R, Huang R, Hicks B, Garcia N. Cannot exclude torsion – A 15-year review. *J Pediatr Surg* 2009;44:1212-6.
- Pansky M, Smorgick N, Herman A, Schneider D, Halperin R. Torsion of normal adnexa in postmenarchal women and risk of recurrence. *Obstet Gynecol* 2007;109:355-9.
- Houry D, Abbott JT. Ovarian torsion: A fifteen-year review. *Ann Emerg Med* 2001;38:156-9.
- Celik A, Ergün O, Aldemir H, Ozcan C, Ozok G, Erdener A, et al. Long-term results of conservative management of adnexal torsion in children. *J Pediatr Surg* 2005;40:704-8.
- Germain M, Rarick T, Robins E. Management of intermittent ovarian torsion by laparoscopic oophorectomy. *Obstet Gynecol* 1996;88:715-7.
- Buss JG, Lee RA. Sequential torsion of the uterine adnexa. *Mayo Clin Proc* 1987;62:623-5.
- Hibbard LT. Adnexal torsion. *Am J Obstet Gynecol* 1985;152:456-61.
- Bouguizane S, Bibi H, Farhat Y, Dhifallah S, Darraji F, Hidar S, et al. Adnexal torsion: A report of 135 cases. *J Gynecol Obstet Biol Reprod (Paris)* 2003;32:535-40.
- Oelsner G, Shashar D. Adnexal torsion. *Clin Obstet Gynecol* 2006;49:459-63.
- White M, Stella J. Ovarian torsion: 10-year perspective. *Emerg Med Australas* 2005;17:231-7.
- Gorkemli H, Camus M, Clasen K. Adnexal torsion after gonadotrophin ovulation induction for IVF or ICSI and its conservative treatment. *Arch Gynecol Obstet* 2002;267:4-6.
- Tsafrir Z, Azem F, Hasson J, Solomon E, Almog B, Nagar H, et al. Risk factors, symptoms, and treatment of ovarian torsion in children: The twelve-year experience of one center. *J Minim Invasive Gynecol* 2012;19:29-33.
- Rotoli JM. Abdominal pain in the post-menopausal female: Is ovarian torsion in the differential? *J Emerg Med* 2017;52:749-52.
- Bar-On S, Mashiach R, Stockheim D, Soriano D, Goldenberg M, Schiff E, et al. Emergency laparoscopy for suspected ovarian torsion: Are we too hasty to operate? *Fertil Steril* 2010;93:2012-5.
- Anders JF, Powell EC. Urgency of evaluation and outcome of acute ovarian torsion in pediatric patients. *Arch Pediatr Adolesc Med* 2005;159:532-5.
- Ding DC, Chang YH. Laparoendoscopic single-site surgical cystectomy of a twisted ovarian dermoid cyst during early pregnancy: A case report and literature review. *Gynecol Minim Invasive Ther* 2016;5:173-7.
- Johnson TR Jr., Woodruff JD. Surgical emergencies of the uterine adnexae during pregnancy. *Int J Gynaecol Obstet* 1986;24:331-5.
- Yen CF, Lin SL, Murk W, Wang CJ, Lee CL, Soong YK, et al. Risk analysis of torsion and malignancy for adnexal masses during pregnancy. *Fertil Steril* 2009;91:1895-902.
- Bromley B, Benacerraf B. Adnexal masses during pregnancy: Accuracy of

- sonographic diagnosis and outcome. *J Ultrasound Med* 1997;16:447-52.
32. Schmeler KM, Mayo-Smith WW, Peipert JF, Weitzen S, Manuel MD, Gordinier ME, et al. Adnexal masses in pregnancy: Surgery compared with observation. *Obstet Gynecol* 2005;105:1098-103.
 33. Huchon C, Panel P, Kayem G, Schmitz T, Nguyen T, Fauconnier A, et al. Does this woman have adnexal torsion? *Hum Reprod* 2012;27:2359-64.
 34. Karaman E, Beger B, Çetin O, Melek M, Karaman Y. Ovarian torsion in the normal ovary: A diagnostic challenge in postmenarchal adolescent girls in the emergency department. *Med Sci Monit* 2017;23:1312-6.
 35. Rey-Bellet Gasser C, Gehri M, Joseph JM, Pauchard JY. Is it ovarian torsion? A systematic literature review and evaluation of prediction signs. *Pediatr Emerg Care* 2016;32:256-61.
 36. Kirkham YA, Lacy JA, Kives S, Allen L. Characteristics and management of adnexal masses in a canadian pediatric and adolescent population. *J Obstet Gynaecol Can* 2011;33:935-43.
 37. Sasso RA. Intermittent partial adnexal torsion after electrosurgical tubal ligation. *J Am Assoc Gynecol Laparosc* 1996;3:427-30.
 38. Ashwal E, Hirsch L, Krissi H, Eitan R, Less S, Wiznitzer A, et al. Characteristics and management of ovarian torsion in premenarchal compared with postmenarchal patients. *Obstet Gynecol* 2015;126:514-20.
 39. Cohen SB, Wattiez A, Stockheim D, Seidman DS, Lidor AL, Mashiach S, et al. The accuracy of serum interleukin-6 and tumour necrosis factor as markers for ovarian torsion. *Hum Reprod* 2001;16:2195-7.
 40. Daponte A, Pournaras S, Hadjichristodoulou C, Lialios G, Kallitsaris A, Maniatis AN, et al. Novel serum inflammatory markers in patients with adnexal mass who had surgery for ovarian torsion. *Fertil Steril* 2006;85:1469-72.
 41. Laganà AS, Sofò V, Salmeri FM, Palmara VI, Triolo O, Terzić MM, et al. Oxidative stress during ovarian torsion in pediatric and adolescent patients: Changing the perspective of the disease. *Int J Fertil Steril* 2016;9:416-23.
 42. Dahmouh H, Anupindi SA, Pawel BR, Chauvin NA. Multimodality imaging findings of massive ovarian edema in children. *Pediatr Radiol* 2017;47:576-83.
 43. Anthony EY, Caserta MP, Singh J, Chen MY. Adnexal masses in female pediatric patients. *AJR Am J Roentgenol* 2012;198:W426-31.
 44. Wilkinson C, Sanderson A. Adnexal torsion – A multimodality imaging review. *Clin Radiol* 2012;67:476-83.
 45. Lee EJ, Kwon HC, Joo HJ, Suh JH, Fleischer AC. Diagnosis of ovarian torsion with color doppler sonography: Depiction of twisted vascular pedicle. *J Ultrasound Med* 1998;17:83-9.
 46. Albayram F, Hamper UM. Ovarian and adnexal torsion: Spectrum of sonographic findings with pathologic correlation. *J Ultrasound Med* 2001;20:1083-9.
 47. Servaes S, Zurakowski D, Laufer MR, Feins N, Chow JS. Sonographic findings of ovarian torsion in children. *Pediatr Radiol* 2007;37:446-51.
 48. Nizar K, Deutsch M, Filmer S, Weizman B, Beloosesky R, Weiner Z. Doppler studies of the ovarian venous blood flow in the diagnosis of adnexal torsion. *J Clin Ultrasound* 2009;37:436-9.
 49. Valsky DV, Esh-Broder E, Cohen SM, Lipschuetz M, Yagel S. Added value of the gray-scale whirlpool sign in the diagnosis of adnexal torsion. *Ultrasound Obstet Gynecol* 2010;36:630-4.
 50. Vijayaraghavan SB, Senthil S. Isolated torsion of the fallopian tube: The sonographic whirlpool sign. *J Ultrasound Med* 2009;28:657-62.
 51. Born C, Wirth S, Stäbler A, Reiser M. Diagnosis of adnexal torsion in the third trimester of pregnancy: A case report. *Abdom Imaging* 2004;29:123-7.
 52. Haque TL, Togashi K, Kobayashi H, Fujii S, Konishi J. Adnexal torsion: MR imaging findings of viable ovary. *Eur Radiol* 2000;10:1954-7.
 53. Kawakami K, Murata K, Kawaguchi N, Furukawa A, Morita R, Tenzaki T, et al. Hemorrhagic infarction of the diseased ovary: A common MR finding in two cases. *Magn Reson Imaging* 1993;11:595-7.
 54. Schlaff WD, Lund KJ, McAleese KA, Hurst BS. Diagnosing ovarian torsion with computed tomography. A case report. *J Reprod Med* 1998;43:827-30.
 55. Kimura I, Togashi K, Kawakami S, Takakura K, Mori T, Konishi J. Ovarian torsion: CT and MR imaging appearances. *Radiology* 1994;190:337-41.
 56. Hiller N, Appelbaum L, Simanovsky N, Lev-Sagi A, Aharoni D, Sella T, et al. CT features of adnexal torsion. *AJR Am J Roentgenol* 2007;189:124-9.
 57. Naffaa L, Deshmukh T, Tumu S, Johnson C, Boyd KP, Meyers AB, et al. Imaging of acute pelvic pain in girls: Ovarian torsion and beyond. *Curr Probl Diagn Radiol* 2017;46:317-29.
 58. Oelsner G, Cohen SB, Soriano D, Admon D, Mashiach S, Carp H, et al. Minimal surgery for the twisted ischaemic adnexa can preserve ovarian function. *Hum Reprod* 2003;18:2599-602.
 59. Tsafir Z, Hasson J, Levin I, Solomon E, Lessing JB, Azem F, et al. Adnexal torsion: Cystectomy and ovarian fixation are equally important in preventing recurrence. *Eur J Obstet Gynecol Reprod Biol* 2012;162:203-5.
 60. Harkins G. Ovarian torsion treated with untwisting: Second look 36 hours after untwisting. *J Minim Invasive Gynecol* 2007;14:270.
 61. Mashiach S, Bider D, Moran O, Goldenberg M, Ben-Rafael Z. Adnexal torsion of hyperstimulated ovaries in pregnancies after gonadotropin therapy. *Fertil Steril* 1990;53:76-80.
 62. Bider D, Mashiach S, Dulitzky M, Kokia E, Lipitz S, Ben-Rafael Z, et al. Clinical, surgical and pathologic findings of adnexal torsion in pregnant and nonpregnant women. *Surg Gynecol Obstet* 1991;173:363-6.
 63. Oelsner G, Bider D, Goldenberg M, Admon D, Mashiach S. Long-term follow-up of the twisted ischemic adnexa managed by detorsion. *Fertil Steril* 1993;60:976-9.
 64. Dolgin SE, Lublin M, Shlasko E. Maximizing ovarian salvage when treating idiopathic adnexal torsion. *J Pediatr Surg* 2000;35:624-6.
 65. Aziz D, Davis V, Allen L, Langer JC. Ovarian torsion in children: Is oophorectomy necessary? *J Pediatr Surg* 2004;39:750-3.
 66. Wang JH, Wu DH, Jin H, Wu YZ. Predominant etiology of adnexal torsion and ovarian outcome after detorsion in premenarchal girls. *Eur J Pediatr Surg* 2010;20:298-301.
 67. Shalev J, Goldenberg M, Oelsner G, Ben-Rafael Z, Bider D, Blankstein J, et al. Treatment of twisted ischemic adnexa by simple detorsion. *N Engl J Med* 1989;321:546.
 68. Taskin O, Birincioglu M, Aydin A, Buhur A, Burak F, Yilmaz I, et al. The effects of twisted ischaemic adnexa managed by detorsion on ovarian viability and histology: An ischaemia-reperfusion rodent model. *Hum Reprod* 1998;13:2823-7.
 69. Hubner N, Langer JC, Kives S, Allen LM. Evolution in the management of pediatric and adolescent ovarian torsion as a result of quality improvement measures. *J Pediatr Adolesc Gynecol* 2017;30:132-37.
 70. Zweizig S, Perron J, Grubb D, Mishell DR Jr. Conservative management of adnexal torsion. *Am J Obstet Gynecol* 1993;168(6 Pt 1):1791-5.
 71. Mathevet P, Nessah K, Dargent D, Mellier G. Laparoscopic management of adnexal masses in pregnancy: A case series. *Eur J Obstet Gynecol Reprod Biol* 2003;108:217-22.
 72. Djavadian D, Braendle W, Jaenicke F. Laparoscopic oophoropexy for the treatment of recurrent torsion of the adnexa in pregnancy: Case report and review. *Fertil Steril* 2004;82:933-6.
 73. Bisharah M, Tulandi T. Laparoscopic surgery in pregnancy. *Clin Obstet Gynecol* 2003;46:92-7.
 74. Crombleholme TM, Craigo SD, Garmel S, D'Alton ME. Fetal ovarian cyst decompression to prevent torsion. *J Pediatr Surg* 1997;32:1447-9.
 75. Alrabeeah A, Galliani CA, Giacomantonio M, Heifetz SA, Lau H. Neonatal ovarian torsion: Report of three cases and review of the literature. *Pediatr Pathol* 1988;8:143-9.
 76. Bryant AE, Laufer MR. Fetal ovarian cysts: Incidence, diagnosis and management. *J Reprod Med* 2004;49:329-37.

77. Crouch NS, Gyampoh B, Cutner AS, Creighton SM. Ovarian torsion: To pex or not to pex? Case report and review of the literature. *J Pediatr Adolesc Gynecol* 2003;16:381-4.
78. Ozcan C, Celik A, Ozok G, Erdener A, Balik E. Adnexal torsion in children may have a catastrophic sequel: Asynchronous bilateral torsion. *J Pediatr Surg* 2002;37:1617-20.
79. Grunewald B, Keating J, Brown S. Asynchronous ovarian torsion – The case for prophylactic oophorectomy. *Postgrad Med J* 1993;69:318-9.
80. Functional ovarian cysts and oral contraceptives. negative association confirmed surgically. A cooperative study. *JAMA* 1974;228:68-9.
81. Caillouette JC, Koehler AL. Phasic contraceptive pills and functional ovarian cysts. *Am J Obstet Gynecol* 1987;156:1538-42.
82. Grimes DA, Godwin AJ, Rubin A, Smith JA, Laccarra M. Ovulation and follicular development associated with three low-dose oral contraceptives: A randomized controlled trial. *Obstet Gynecol* 1994;83:29-34.
83. Mishell DR Jr. Noncontraceptive benefits of oral contraceptives. *J Reprod Med* 1993;38 (12 Suppl):1021S-9S.
84. Kaleli B, Aktan E, Gezer S, Kirkali G. Reperfusion injury after detorsion of unilateral ovarian torsion in rabbits. *Eur J Obstet Gynecol Reprod Biol* 2003;110:99-101.
85. Dolgin SE. Acute ovarian torsion in children. *Am J Surg* 2002;183:95-6.