



Published in final edited form as:

Addiction. 2017 November ; 112(11): 1931–1941. doi:10.1111/add.13896.

Are Alcohol-Related Disparities between Sexual Minority and Heterosexual Youth Decreasing?

Jessica N. Fish, Ph.D.¹, Ryan J. Watson, Ph.D.², Carolyn M. Porta, Ph.D.,MPH,RN³, Stephen T. Russell, Ph.D.⁴, and Elizabeth M. Saewyc, Ph.D., RN, FSAHM, FCAHS, FAAN⁵

²Department of Human Development and Family Studies, University of Connecticut, 348 Mansfield Road, U-1058, Storrs, CT 06269

³School of Nursing, University of Minnesota, School of Nursing, Mail Stop 1331 5-140 WDH, 308 Harvard St. SE, Minneapolis, MN 55455

⁴Human Development and Family Sciences, University of Texas at Austin, 108 E. Dean Keeton St., Stop A2702, Austin, TX, 78712

⁵School of Nursing, University of British Columbia, Vancouver, BC, Canada

Abstract

Background and Aims—Although sexual-orientation-related alcohol use disparities are well established, researchers have not identified whether disparities are diminishing as societal attitudes towards lesbian/gay and bisexual (LGB) people become more accepting. We examined changes in four alcohol-related disparities between heterosexual and LGB youth from 1998–2013 by (1) estimating the prevalence of these behaviors; (2) estimating disparities in alcohol-related outcomes between heterosexual and LGB youth within each wave year; and (3) testing whether the degree of difference in alcohol-related disparities between heterosexual and LGB youth has changed.

Design—Logistic regression models and year-by-sexual-orientation interactions with repeated, cross-sectional, provincially-representative data.

Setting—British Columbia, Canada.

Participants—Students (ages 12–19) from the 1998 (n = 22,858), 2003 (n = 29,323), 2008 (n = 25,254), and 2013 (n = 21,938) British Columbia Adolescent Health Survey (total N = 99,373, 48.7% male, M @age = 14.84).

Measurements—We modeled age-adjusted differences in lifetime alcohol use, age of onset, past 30-day drinking, and past 30-day heavy episodic drinking between heterosexual and three subgroups of sexual minority youth (i.e., mostly heterosexual, bisexual, and lesbian/gay).

Findings—Generally, alcohol use declined for all youth, although less so among LGB youth (average aOR = .58 and aOR = .53 for heterosexual males and females and aOR = .71 and aOR = .57 for sexual minority males and females, respectively). Within-year comparisons demonstrated

¹Corresponding Author, Population Research Center, University of Texas at Austin, 305 E. 23rd St., Stop G1800, Austin, TX 78712, jessica.fish@utexas.edu, PH: 512-471-4270, F: 512-471-4886.

The authors report no conflicts of interest.

elevated rates of alcohol use among LGB compared with heterosexual youth for each of the four survey years, especially among females. Findings indicate few changes over time; however, results show an increase in risky alcohol use from 1998 to 2013 among mostly heterosexual (aOR = 1.58 for lifetime alcohol use, aOR = 1.58 for 30-day alcohol use, and aOR = 1.34 for 30-day heavy episodic drinking), and bisexual (aOR = 1.95 for lifetime alcohol use) females.

Conclusion—Despite the general decline in the prevalence of alcohol use among young people in Canada since 1998, lesbian/gay and bisexual youth in Canada continue to show elevated rates of alcohol use compared with heterosexual youth.

Introduction

Although rates have declined in recent years [1–3], adolescent alcohol use remains a major public health concern. Youth are particularly vulnerable to alcohol-related morbidity and mortality [4,5] and patterns of heavy drinking established during adolescence can persist into adulthood [6,7], leading to long-term health consequences [8]. These concerns related to adolescent alcohol use highlight the importance of identifying those youth most at risk to help inform prevention and intervention strategies [9,10].

Sexual minority (i.e., lesbian/gay and bisexual [LGB]) youth represent an at-risk group for alcohol use and misuse during adolescence [11–13]. Over a decade of research demonstrates disproportionately high rates of alcohol use among LGB compared to heterosexual youth across multiple indicators of risk including: lifetime alcohol use [14], early initiation [14,15], frequency of use [16,17], the participation in and frequency of heavy episodic drinking (HED) [14,18], and problems related to drinking [19]. Comparative studies also highlight the propensity for LGB adolescents to persist and accelerate alcohol use during the transition to adulthood [17,20–22], a factor which may explain the higher rates of alcohol abuse and dependence among sexual minority adults [23,24].

Prior studies clearly identify sexual minority disparities in alcohol use, yet at the same time there has been unprecedented, growing social acceptance for LGB people in North America [25,26]: Only 44% of US adults supported the legal recognition of marriage in 1996 compared to 68% in 2016 [27]. Laws and policies protecting the rights of LGB youth and adults have also been instituted in Canada [28] and in various states across the US [29,30], although specific anti-discrimination laws that enumerate sexual orientation have yet to receive federal US recognition. Considering the link between anti-LGB attitudes and increased alcohol use among sexual minority youth [31–33], we hypothesize that these trends in LGB acceptance could lead to a decrease in alcohol-related disparities between sexual minority and heterosexual youth. Although researchers have started to document trends in LGB health disparities over time [34], studies have yet to examine whether trends in alcohol-related disparities are changing as a function of broader social acceptance and the implementation of LGB-specific protections [29].

Sexual Identity, Gender, and Alcohol Use

Although rates of alcohol use among sexual minority youth, in general, are elevated compared to heterosexual peers, specific groups among sexual minority youth display

differential risk. For example, across studies, disparities in alcohol use among sexual minorities are larger and more consistent for females, whereas evidence of disparities is less consistent for males [11,12,14]. One meta-analytic review demonstrated that sexual minority females had five times greater odds of substance use compared to heterosexual females: Sexual minority male adolescents were 3.5 times more likely to report substance use than their heterosexual counterparts [35].

Bisexual-identified and attracted youth also appear to be at greater risk for alcohol and substance use behavior than their gay, lesbian, and heterosexual peers [12,14,19]. For example, Marshal and colleagues [35] found that bisexual youth had 4.5 times the odds of engaging in substance use behavior than heterosexuals, and the size of the effect was 7 times greater than that of gay/lesbian youth. Emerging studies also demonstrate risk for alcohol use among youth who question their identities or attractions [31] along with youth and adults who report “mostly heterosexual” identities [21,36,37]. These within-group differences among sexual minority youth are important to consider in studies investigating health disparities as they have important implications for focused prevention and intervention strategies.

The Current Study

We used a provincially-representative sample of Canadian adolescents to examine trends in alcohol use and trends of alcohol-related disparities between heterosexual and three sexual minority youth subgroups: mostly heterosexual, bisexual, and gay/lesbian. Our overall goal was threefold: (1) track trends in the prevalence of lifetime alcohol use, age of onset, past 30-day alcohol use, and past 30-day heavy episodic drinking (HED) by sexual orientation from 1998 to 2013; (2) estimate alcohol-related disparities between heterosexual and sexual minority subgroups for each survey year; and (3) test whether sexual-orientation-related disparities in alcohol use youth have increased, decreased, or remained stable over the past 15 years. Previous findings highlight gender [11,14] differences in alcohol use; in our data source, we found the proportion of sexual minority youth increases over time for girls, but not for boys. Therefore, disaggregating analyses by gender, in order to adhere to the new Sex and Gender Equity in Research (SAGER) international guidelines [38] is a salient consideration. The SAGER guidelines for researchers and editors are strongly promoted by the Canadian Institutes of Health Research’s Institute of Gender and Health; we have conducted and reported analyses separately by gender.

Methods

Design

Using data from a repeated, cross-sectional survey of middle- and high-school students in British Columbia, Canada, we first estimated the prevalence of lifetime alcohol use, age of onset, past 30-day use, and past 30-day HED among heterosexual and sexual minority students from 1998 to 2013. Second, we estimated differences between heterosexual and sexual minority youth for these four alcohol-related behaviors for each year of data collection: 1998, 2003, 2008, and 2013. Third, we tested whether the degree to which heterosexual and sexual minority youth differ in lifetime alcohol use, age of onset, past 30-

day use, and past 30-day HED has increased, decreased, or remained stable from 1998, 2003, and 2008 to 2013 (our reference category year).

Data and Sample

Data are from the 1998, 2003, 2008, and 2013 British Columbia Adolescent Health Survey (BCAHS), a cross-sectional survey developed to capture the health and risk behavior of youth in British Columbia. Conducted by the McCreary Centre Society, the BCAHS is a provincially-based cluster-stratified random classroom survey of Canadian public school students in grades 7 to 12 across BC. Sample estimates were stratified by grade and health service delivery area in the five health authorities in the province and data collected from participating school districts to be both regionally and provincially representative. Depending on the year, between 78–95% of eligible schools districts participated in the BCAHS, and the sample sizes for each area were selected to ensure standard errors of 3.5% within grade and region strata (original unweighted $N = 115,573$). Data were weighted to account for non-response and differential probability of sampling and scaled to replicate provincial enrollment (for more information see [39,40]).

To assess trends in alcohol use behaviors, we pooled data from the 1998, 2003, 2008, and 2013 surveys from school districts that participated in at least 3 of the 4 surveys. Students who did not provide a valid response to the sexual orientation measure were excluded bringing the total unweighted sample to $N = 99,373$ adolescents across 15 years, or 86% of the original samples. Table 1 displays prevalence of sexual orientation by survey year and average age across sexual orientation within survey year.

Measures

Sexual Orientation—A single item was used to measure sexual orientation and asked: “People have different feelings about themselves when it comes to questions of being attracted to other people. Which of the following best describes your feelings?” Response options in 1998–2008 were *100% heterosexual (attracted to persons of the opposite sex)*; *Mostly heterosexual = 2*; *Bisexual (attracted to both males and females)*; *Mostly homosexual*; *100% homosexual (gay/lesbian; attracted to persons of the same sex)*; and *Not Sure*. The 2013 survey response options included: *Completely heterosexual*; *Mostly heterosexual*; *Bisexual*; *Mostly homosexual*; *Completely homosexual*; *Questioning*; and *I don’t have attractions*. After extensive psychometric evaluation of these measures across survey collection years (see [41]), we merged those reporting “not sure” (1998–2008), “questioning” (2013), and “I don’t have attractions” (2013) with heterosexual identified youth. Those who reported being “mostly homosexual” were combined with individuals who reported “100% homosexual” or “completely homosexual” identities, depending on the year.

Sex—Participants stated whether they were *male* or *female*.

Prior to the section on alcohol use, the survey included an instruction box that defined a drink as “1 drink = 1 bottle of beer (12 oz.), 1 glass of wine (5 oz.), or 1 shot of hard liquor (1.5 oz.).”

Lifetime drinking—Youth reported lifetime alcohol use by responding *no* = 0 or *yes* = 1 to “Have you ever had alcohol in your lifetime?”.

Age of Onset—Participants were asked, “How old were you when you had your first drink of alcohol other than a few sips?” Item was recoded to reflect those who had a drink *12 or younger* = 1 and those had their first drink *after the age of 12* = 0.

Past 30-day drinking—Youth reported past 30-day drinking by stating how many days they had used alcohol in the previous month. The item was recoded to reflect *0 days/no use* = 0 and *1 or more days of alcohol use* = 1.

Past 30-day Heavy Episodic Drinking (HED)—Occurrences of HED were measured with a single item that asked participants to report, “During the past 30 days, on how many days did you have 5 or more drinks of alcohol within a couple of hours?” Responses were recoded to reflect *no HED* = 0 and *1 or more day of HED* = 1.

Age—Age was calculated based on participants’ reports of their age in years.

Analytic Approach

SPSS Complex Samples 22 was used to conduct all analyses to apply survey weights and account for the BCAHS complex sampling design. We used crosstabs to examine whether the prevalence of alcohol use behaviors changed across years of analysis (e.g., from 1998, 2003, 2008, 2013) within sexual orientation groups. This preliminary investigation provided valuable information on group-specific changes or stability in trends that inform the interpretation of later comparative analyses. Next, we tested whether sexual minority youth differed from heterosexual youth in the prevalence of each alcohol use behavior within each data collection year, adjusted for age.

Finally, we used logistic regression with year-by-orientation interactions terms to test whether the disparity between heterosexual and sexual minority youth changed (i.e., widened, declined, or maintained) since 1998. Comparing absolute measures of inequalities, such as the prevalence of substance use over time, can be misleading because age differences across samples can potentially explain the changes in trends [42]. Within each orientation group, students in 1998 had significantly younger mean ages than the other years, and within each year, lesbian, gay, and bisexual youth were somewhat older on average than heterosexual adolescents. Therefore, it is more appropriate to compare age-adjusted odds ratios. However, odds ratios cannot be directly compared across different samples [43], and cannot provide the scope of a disparity; to test whether trends changed over time, controlling for age, we computed an interaction term between survey year and sexual orientation in a logistic regression framework. The resulting year-by-sexual orientation interaction term calculates a ratio of odds ratios (*ORs*) which compare the age-adjusted *OR* of an alcohol use behavior (e.g., 30-day HED) for a particular subgroup (e.g., gay males) vs. the referent group (heterosexual males) in a given year (e.g., 2008) to the odds of those with the same identity in 2013 (the reference year) compared to their heterosexual peers. We chose 2013 as the reference year for each of our models to be able to consistently refer to the declining changes in substance use from a historical perspective; as a result, the estimated interaction

would need to be interpreted as the inverse of typical *ORs* (i.e., an interaction *OR* above 1 would indicate narrowing disparity). For ease of interpretation, we inverted estimated interaction *ORs* to reflect changes from the past to present. Therefore, an estimated *OR* for the interaction term above 1 indicates a widening disparity from earlier years to the reference year, and below 1 indicates a declining disparity for that behavior from earlier years for specific sexual orientation identities to the reference year. Essentially, this approach of producing age-adjusted year-by-sexual orientation interaction terms estimates the change in the disparity over time (and thus, the degree of substance use disparities between groups). More detailed explanations of this approach are available elsewhere [34].

Results

Prevalence and Trends by Sexual Orientation

Rates of alcohol use across all four behaviors generally declined within all sexual orientation groups over time with some notable exceptions (see Table 2). Unlike heterosexual and bisexual youth, mostly heterosexual and gay male youth did not experience a reduction in lifetime alcohol use from 2003 to 2013. Although the prevalence of early onset declined precipitously across survey years for heterosexual youth, sexual minority youth showed less consistent patterns in early onset from 2003 and 2013. Generally, early onset decreased for mostly heterosexual, bisexual, and gay males from 1998 to 2013, although comparison within mostly heterosexual, bisexual, and gay boys showed no difference in early onset between 2003 and 2013. Past 30-day alcohol use among heterosexual males declined across all survey years, however, the same reduction was not evident for mostly heterosexual, bisexual, and gay males. Unlike heterosexual, mostly heterosexual, and bisexual females, lesbians did not display a decrease in past 30-day alcohol use across the 15 years of data collection. Heterosexual males were the only group with decreasing HED for all year comparisons. HED decreased among mostly heterosexual and gay boys from 1998 to 2013, but bisexual males had stable rates of HED across all years. There were significant decreases for all female sexual orientation groups in the prevalence of HED when comparing 2013 to previous years.

Disparities in Alcohol Use between Sexual Minority and Heterosexual Youth across Years

Age-adjusted odds ratios (*ORs*) documenting disparities between sexual minority and heterosexual male youth by year are presented in Table 3. Mostly heterosexual and bisexual, but not gay male, youth reported higher odds of lifetime alcohol use in 2008 only. Compared to heterosexuals, bisexual boys were more likely to start drinking prior to the age of 13 in 1998 and 2008 and gay boys in 2008 and 2013. Mostly heterosexual boys were more likely than heterosexual boys to report drinking before the age of 12 in 2013. Bisexual and gay males had greater odds of past 30-day alcohol use than heterosexual peers in 2008 and mostly heterosexual males reported lower risk for past 30-day heavy episode drinking compared to heterosexual males in 2003 and 2013.

Female disparities in alcohol use behaviors, also adjusted for age, are presented in Table 4. Mostly heterosexual females had greater odds of lifetime alcohol use, early age of onset, and past 30-day alcohol use across all survey years except for past 30-day alcohol use in 1998.

Compared to heterosexual females, bisexual females had greater odds of lifetime alcohol use and past 30-day alcohol use in 2003, 2008, and 2013 and earlier age of onset for all survey years. Lesbian youth had higher odds of having their first drink prior to the age of 13 than heterosexual females in 2003, 2008, and 2013 and higher the odds of reporting lifetime alcohol use in 2013. Mostly heterosexual and bisexual females had greater odds of HED in 2003, 2008, and 2013, compared to heterosexual youth. Only lesbians had greater odds than heterosexual female youth for HED in 2008.

Trends in Alcohol Use Disparities over Time

Last, we tested whether the difference in alcohol use behaviors by sexual orientation have decreased, widened, or remained stable from 1998 to 2013 by modeling interactions terms of sexual orientation by survey year while adjusting for age (see Table 5). For males, the significant interaction between gay boys and the year 2003 indicates that the gap between the age of onset between heterosexual and gay males has widened from 2003 to 2013. All other differences between heterosexual and sexual minority youth were stable across years.

There were a number of changes in risk across outcomes for females. Disparities between heterosexual and mostly heterosexual females from 1998 to 2013 widened for lifetime alcohol use, past 30-day alcohol use, and HED. Bisexual females' risk for lifetime alcohol use increased from 1998 to 2013, but decreased for age of onset during that same time.

Discussion

This study is the first to our knowledge to examine trends in alcohol use disparities between heterosexual and sexual minority adolescents. Using a provincially-representative sample of British Columbia adolescents between 1998 and 2013 we found that, despite growing acceptance in societal attitudes towards LGB populations in North America, disparities in alcohol use among sexual minority youth have generally maintained their degree of difference, and in some cases, widened, as compared with their non-sexual minority counterparts.

Findings regarding within-sexual orientation prevalence of lifetime alcohol use, age of onset, past 30-day alcohol use, and past 30-day heavy episodic drinking (HED) are positive, and consistent with other North American data: adolescent alcohol use is declining [1,2]. Also similar to previous studies and meta-analyses [11,12,35], sexual minority girls demonstrated a greater number of disparities in alcohol use behavior across the 15-year time span than were observed in the comparisons between heterosexual and sexual minority boys. Furthermore, as with previous research, bisexual and mostly heterosexual youth in our study displayed greater risk for alcohol use behaviors across years when compared to their heterosexual peers, especially among females, while gay/lesbian youth did not differ from heterosexual youth in 23 of the 32 comparisons.

Despite the general decline in the prevalence of alcohol use behaviors since 1998, our novel assessment of the stability of alcohol use behaviors across sexual identity groups suggests that the difference between these groups has remained fairly consistent, with a few exceptions. Since 1998 mostly heterosexual females demonstrated increased disparity

compared to heterosexual females in lifetime alcohol use, past 30-day use, and past 30-day HED, and bisexual females showed a widening gap in lifetime alcohol use vs. heterosexual peers, although their disparity in risk of early onset narrowed from 1998 to 2013. Among gay males, our results showed an increase in disparity in early age of onset compared to heterosexual peers from 2003 to 2013.

Findings from our study support a need for further research to explain an unanswered question: If attitudes towards LGB people are improving, then why do disparities in alcohol use among sexual minority youth remain? Though we cannot make direct inferences from these trend data, one possible explanation might reside in the clash between increasing societal acceptance and developmental trends in peer social regulation [44]: Youth are disclosing their LGB identities at younger ages [44,45], due in part to greater acceptance of LGB identities in the broader society, yet early adolescence is a time during which peers begin to socially regulate gender and sexuality [46,47] and are more likely to report prejudicial attitudes and homophobic behavior [48,49]. Therefore, although overall attitudes are improving in society, youth may remain susceptible to developmentally situated and prejudiced experiences in their peer groups that lead to substance use. Interestingly, this trend might be reflected in our findings on the prevalence of heterosexuality among adolescent girls, which declined from 1998 (90.3%) to 2013 (85.6%). If secular trends in attitudes towards sexual minorities are contributing to more youth reporting lesbian, bisexual and unsure identities, sexual orientation comparisons in modern cohorts might provide more accurate samples of sexual minority girls, compared to earlier years.

The increase in alcohol use among mostly heterosexual and bisexual females is also counter to expectations given the changing social climate. Why does this group appear to be particularly at risk? Studies increasingly demonstrate the disproportionate risk of bisexual and mostly heterosexual youth and adults across a host of outcomes relative to heterosexual and lesbian/gay peers [16,35–37], often linked to prejudice coming from both heterosexual and gay/lesbian communities [50–52]. Although broader attitudes do not appear to mitigate risk for alcohol use among sexual minority youth, these changing attitudes may differentially favor lesbian/gay identities as opposed to those who identify as mostly heterosexual and bisexual.

Given persistent disparities in alcohol use among sexual minority adolescent over time, despite the shifting culture milieu, there is a clear need for targeted programs and policies for sexual minority youth. Because studies demonstrate the role of minority stress in sexual minority youth alcohol use [31], programs and policies that specifically address anti-LGB discrimination and homophobic bullying in schools help to reduce sexual minority youth experiences of prejudice [53] in ways that may subsequently reduce their alcohol use. Similarly, programs using prevention curricula should consider curricula that specifically address relevant experiences of prejudice and discrimination: For example, one of the only published interventions of a randomized clinical trial with young adult gay men showed that incorporating conversations about minority stress into therapy protocols helped reduce substance use and improved mental health [54].

Like all studies, there are limitations. Notably, we position our perspective of trends in alcohol use within the shifting social context, yet youth may have a different experience and understanding of changes. We were unable to capture that perspective with these data. We also use a single item measure of sexual orientation; however, reports of sexual identity, attraction, and behavior do not consistently align within persons and studies that simultaneously investigate alcohol use outcomes across different indicators of sexual orientation demonstrate differential risk [36]. Therefore, estimates of risk may vary depending on sexual orientation measurement. These data are also geographically limited, and reflect a single Canadian province; findings may not reflect the experience of youth in other countries, considering the large variability in local, regional, and country sociocultural and legal environments related to LGB people. Future studies should consider implementing this approach to other repeating population-based data sources to help broaden our understanding of changes in sexual minority youth experiences. Data were also collected using Statistics Canada's conventional measure for ethnicity, precluding the use of an ethnicity variable that provides mutually exclusive categories. Therefore, we do not include ethnicity in our adjustment of analytic models. Finally, the BCAHS data were collected in schools, and therefore do not represent the experience of youth who were not enrolled. Given the disproportionately high number of sexual minority youth who experience homelessness [55] and school pushout [56] and the link between these experiences and substance use [57], such limitations need to be acknowledged.

Despite these limitations, there are several strengths of the current study. First, the use of a large population-based survey of adolescents in schools provided us with an opportunity to disaggregate trends based on those with mostly heterosexual, bisexual, and gay/lesbian sexual orientations. Second, because odds ratios are not directly comparable across groups [58], but a widely used statistic, our novel statistical approach offers new opportunities for substance use researchers to examine how differences in risk may be changing over time. Notably, the applications of this approach with these data is a testament to the importance of sexual (and gender) minority identity markers in large, school-based samples—without these data, we are unable to document if youth experiences change over time and the (potential) effects of social movements and policy [59,60].

Our findings provide valuable information regarding the stability and widening of disparities in alcohol use among sexual minority youth from 1998 to 2013, even in the face of growing social acceptance of LGB people in North America. This juxtaposition may be partially explained by developmental vulnerabilities in peer interaction and cognitive processes that warrant the implementation of enumerated policies and targeted intervention programs for this population. Future studies are needed to better explore how contexts influence youth wellbeing and in what ways these changing attitudes, policies, and programs might alleviate alcohol use disparities among sexual minority youth.

Acknowledgments

Funding Acknowledgements: This study was funded by grants #CPP 86374 and #MOP 119472 from the Canadian Institutes of Health Research. The authors acknowledge the McCreary Centre Society (<http://www.mcs.bc.ca>) for access to the British Columbia Adolescent Health Survey. In addition, the work by the first author was funded by the National Institute on Alcohol Abuse and Alcoholism grant number F32AA023138. Fish and Russell also

received support from grant R24HD042849, awarded to the Population Research Center at The University of Texas at Austin by the Eunice Kennedy Shriver National Institute of Child Health and Human Development. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Centers for Disease Control. Trends in the prevalence of alcohol use national YRBS: 1991–2015. 2016. Available from: http://www.cdc.gov/healthyouth/data/yrbs/pdf/trends/2015_us_alcohol_trend_yrbs.pdf
2. Miech, RA., Johnston, LD., O'Malley, PM., Bachman, JG., Schulenberg, JE. Monitoring the future national survey results on drug use, 1975–2014: Volume I, Secondary school students. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2015.
3. World Health Organization. Global status report on alcohol and health 2014 [Internet]. Geneva, Switzerland: 2014. Available from: http://apps.who.int/iris/bitstream/10665/112736/1/9789240692763_eng.pdf?ua=1
4. Brown SA, McGue M, Maggs J, Schulenberg J, Hingson R, Swartzwelder S, et al. A developmental perspective on alcohol and youths 16 to 20 years of age. *Pediatrics*. 2008; 121:S290–S310. [PubMed: 18381495]
5. Kann L. Youth risk behavior surveillance—United States, 2015. *MMWR*. 2016; 65:1–174.
6. Schulenberg JE, Maggs JL. Destiny matters: distal developmental influences on adult alcohol use and abuse. *Addiction*. 2008; 103:1–6.
7. Pitkänen T, Kokko K, Lyyra A-L, Pulkkinen L. A developmental approach to alcohol drinking behaviour in adulthood: a follow-up study from age 8 to age 42. *Addiction*. 2008; 103:48–68. [PubMed: 18426540]
8. Sloan F, Grossman D, Platt A. Heavy episodic drinking in early adulthood and outcomes in midlife. *J Stud Alcohol Drugs*. 2011; 72:459–70. [PubMed: 21513683]
9. U.S. Department of Health and Human Services. The surgeon general's call to action to prevent and reduce underage drinking. Rockville, MD: Office of the Surgeon General; 2007.
10. Healthy People 2020 [Internet]. Washington, DC: 2016. [cited 2016 Jun 17]. Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/substance-abuse>
11. Plöderl M, Tremblay P. Mental health of sexual minorities: a systematic review. *Int Rev Psychiatry*. 2015; 27:367–85. [PubMed: 26552495]
12. Talley AE, Gilbert PA, Mitchell J, Goldbach J, Marshall BDL, Kaysen D. Addressing gaps on risk and resilience factors for alcohol use outcomes in sexual and gender minority populations. *Drug Alcohol Rev*. 2016 advanced online publication.
13. Bos H, van Beusekom G, Sandfort T. Drinking motives, alcohol use, and sexual attraction in youth. *J Sex Res*. 2015; 0:1–4.
14. Talley AE, Hughes TL, Aranda F, Birkett M, Marshal MP. Exploring alcohol-use behaviors among heterosexual and sexual minority adolescents: Intersections with sex, age, and race/ethnicity. *Am J Public Health*. 2014; 104:295–303. [PubMed: 24328614]
15. Corliss HL, Rosario M, Wypij D, Fisher LB, Austin S. Sexual orientation disparities in longitudinal alcohol use patterns among adolescents: findings from the growing up today study. *Arch Pediatr Adolesc Med*. 2008; 162:1071–8. [PubMed: 18981356]
16. Fish JN, Pasley K. Sexual (minority) trajectories, mental health, and alcohol use: a longitudinal study of youth as they transition to adulthood. *J Youth Adolesc*. 2015; 44:1508–27. [PubMed: 25956289]
17. Talley AE, Sher KJ, Littlefield AK. Sexual orientation and substance use trajectories in emerging adulthood. *Addiction*. 2010; 105:1235–1245. [PubMed: 20491728]
18. Baiocco R, D'Alessio M, Laghi F. Binge drinking among gay, and lesbian youths: the role of internalized sexual stigma, self-disclosure, and individuals' sense of connectedness to the gay community. *Addict Behav*. 2010; 35:896–9. [PubMed: 20584573]

19. Russell ST, Driscoll AK, Truong N. Adolescent same-sex romantic attractions and relationships: implications for substance use and abuse. *Am J Public Health*. 2002; 92:198–202. [PubMed: 11818291]
20. Hatzenbuehler ML, Corbin WR, Fromme K. Trajectories and determinants of alcohol use among LGB young adults and their heterosexual peers: results from a prospective study. *Dev Psychol*. 2008; 44:81. [PubMed: 18194007]
21. Marshal MP, Friedman MS, Stall R, Thompson AL. Individual trajectories of substance use in lesbian, gay and bisexual youth and heterosexual youth. *Addiction*. 2009; 104:974–981. [PubMed: 19344440]
22. Needham BL. Sexual attraction and trajectories of mental health and substance use during the transition from adolescence to adulthood. *J Youth Adolesc*. 2012; 41:179–190. [PubMed: 22076077]
23. McCabe SE, Hughes TL, Bostwick WB, West BT, Boyd CJ. Sexual orientation, substance use behaviors and substance dependence in the United States. *Addiction*. 2009; 104:1333–45. [PubMed: 19438839]
24. Goldberg S, Strutz KL, Herring AA, Halpern CT. Risk of substance abuse and dependence among young adult sexual minority groups using a multidimensional measure of sexual orientation. *Public Health Rep*. 2013; 128:144–52. [PubMed: 23633729]
25. Pew Research Center. Changing attitudes on gay marriage [Internet]. 2016. [cited 2016 Jun 20]; Available from: <http://www.pewforum.org/2016/05/12/changing-attitudes-on-gay-marriage/>
26. Pew Research Center. The global divide on homosexuality [Internet]. 2013. [cited 2016 Jun 20]; Available from: <http://www.pewglobal.org/2013/06/04/the-global-divide-on-homosexuality/>
27. Gallup. Gay and lesbian rights [Internet]. p. 2016 [cited 2016 Jun 20]; Available from: <http://www.gallup.com/poll/1651/Gay-Lesbian-Rights.aspx>
28. Hurley, M. Sexual orientation and legal rights: a chronological overview (PRB 04-13E). Ottawa: Library of Parliament; 2005.
29. Russell ST, Kosciw J, Horn S, Saewyc E. Safe schools policy for LGBTQ students. *Social Policy Report*. 2010; 24:1–24.
30. Gay, Lesbian, and Straight Education Network. Enumerated anti-bullying laws by state [Internet]. GLSEN2016 [cited 2016 Jun 20]; Available from: <http://www.glsen.org/article/state-maps>
31. Birkett M, Espelage DL, Koenig B. LGB and questioning students in schools: the moderating effects of homophobic bullying and school climate on negative outcomes. *J Youth Adolesc*. 2009; 38:989–1000. [PubMed: 19636741]
32. Pachankis JE, Hatzenbuehler ML, Starks TJ. The influence of structural stigma and rejection sensitivity on young sexual minority men's daily tobacco and alcohol use. *Soc Sci Med*. 2014; 103:67–75. [PubMed: 24507912]
33. Goldbach JT, Tanner-Smith EE, Bagwell M, Dunlap S. Minority stress and substance use in sexual minority adolescents: a meta-analysis. *Prev Sci*. 2014; 15:350–363. [PubMed: 23605479]
34. Homma Y, Saewyc E, Zumbo BD. Is it getting better? an analytical method to test trends in health disparities, with tobacco use among sexual minority vs. heterosexual youth as an example. *Int J Equity Health*. 2016; 15:1. [PubMed: 26728322]
35. Marshal MP, Friedman MS, Stall R, King KM, Miles J, Gold MA, et al. Sexual orientation and adolescent substance use: a meta-analysis and methodological review. *Addiction*. 2008; 103:546–56. [PubMed: 18339100]
36. McCabe SE, Hughes TL, Bostwick W, Boyd CJ. Assessment of difference in dimensions of sexual orientation: Implications for substance use research in a college-age population. *J Stud Alcohol*. 2005; 66:620. [PubMed: 16331847]
37. Hughes TL, Wilsnack SC, Kristjanson AF. Substance use and related problems among U.S women who identify as mostly heterosexual. *BMC Public Health*. 2015; 15:803. [PubMed: 26289792]
38. Heidari S, Babor TF, De Castro P, Tort S, Curno M. Sex and gender equity in research: rationale for the SAGER guidelines and recommended use. *Res Integr Peer Rev*. 2016; 1:2.
39. Saewyc, E., Green, R. Survey methodology for the 2008 BC adolescent health survey IV. Vancouver, BC, Canada: McCreary Centre Society; 2009.

40. Saewyc EM, Taylor D, Homma Y, Ogilvie G. Trends in sexual health and risk behaviours among adolescent students in British Columbia. *Can J Hum Sex.* 2008; 17:1–13. [PubMed: 23115485]
41. Saewyc EM, Bauer GR, Skay CL, Bearinger LH, Resnick MD, Reis E, et al. Measuring sexual orientation in adolescent health surveys: evaluation of eight school-based surveys. *J Adolesc Health.* 2004; 35:345.e1–345.e15.
42. Asada Y. On the choice of absolute or relative inequality measures. *Milbank Q.* 2010; 88:616–22. [PubMed: 21166871]
43. Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ.* 2003; 326:219. [PubMed: 12543843]
44. Russell ST, Fish JN. Mental health in lesbian, gay, bisexual, and transgender (LGBT) youth. *Annu Rev Clin Psychol.* 2016; 12:465–87. [PubMed: 26772206]
45. Floyd FJ, Bakeman R. Coming-out across the life course: implications of age and historical context. *Arch Sex Behav.* 2006; 35:287–96. [PubMed: 16804747]
46. Mulvey KL, Killen M. Challenging gender stereotypes: resistance and exclusion. *Child Dev.* 2015; 86:681–94. [PubMed: 25382748]
47. Pascoe, CJ. *Dude, you're a fag: masculinity and sexuality in high school.* Univ of California Press; 2011.
48. Poteat VP, Anderson CJ. Developmental changes in sexual prejudice from early to late adolescence: the effects of gender, race, and ideology on different patterns of change. *Dev Psychol.* 2012; 48:1403–15. [PubMed: 22250998]
49. Robinson JP, Espelage DL, Rivers I. Developmental trends in peer victimization and emotional distress in LGB and heterosexual youth. *Pediatrics.* 2013; 131:423–430. [PubMed: 23382442]
50. Balsam KF, Mohr JJ. Adaptation to sexual orientation stigma: a comparison of bisexual and lesbian/gay adults. *J Couns Psychol.* 2007; 54:306.
51. Bostwick WB, Boyd CJ, Hughes TL, West BT, McCabe SE. Discrimination and mental health among lesbian, gay, and bisexual adults in the United States. *Am J Orthopsychiatry.* 2014; 84:35–45. [PubMed: 24826824]
52. Israel T, Mohr JJ. Attitudes toward bisexual women and men. *J Bisexuality.* 2004; 4:117–34.
53. Marx RA, Kettrey HH. Gay-straight alliances are associated with lower levels of school-based victimization of LGBTQ+ youth: a systematic review and meta-analysis. *J Youth Adolesc.* 2016; 45:1269–82. [PubMed: 27221632]
54. Pachankis JE, Hatzenbuehler ML, Rendina HJ, Safren SA, Parsons JT. LGB-affirmative cognitive-behavioral therapy for young adult gay and bisexual men: a randomized controlled trial of a transdiagnostic minority stress approach. 2015; 83:875–89.
55. Durso, LE., Gates, GJ. *Serving our youth: findings from a national survey of services providers working with lesbian, gay, bisexual and transgender youth who are homeless or at risk of becoming homeless.* Los Angeles, CA: The Williams Institute; 2012.
56. Snapp SD, Hoenig JM, Fields A, Russell ST. Messy, butch, and queer LGBTQ youth and the school-to-prison pipeline. *J Adolesc Res.* 2015; 30:57–82.
57. Rosario M, Schrimshaw EW, Hunter J. Homelessness among lesbian, gay, and bisexual youth: implications for subsequent internalizing and externalizing symptoms. *J Youth Adolesc.* 2011; 41:544–60. [PubMed: 21656284]
58. Allison PD. Comparing logit and probit coefficients across groups. *Sociol Methods Res.* 1999; 28:186–208.
59. Arredondo, M., Gray, C., Russell, S., Skiba, R., Snapp, S. *Documenting disparities for LGBT students: expanding the collection and reporting of data on sexual orientation and gender identity.* Bloomington, IN: The Equity Project at Indiana University; 2016.
60. Snapp SD, Russell ST, Arredondo M, Skiba R. A right to disclose: LGBTQ youth representation in data, science, and policy. *Adv Child Dev Behav.* 2016; 50:135–59. [PubMed: 26956072]

Table 1
Samples By Sexual Orientation and Survey Year in the British Columbia Adolescent Health Survey

	1998		2003		2008		2013	
	<i>n</i> (%) _w	\bar{M}_{age} [95% CI]	<i>n</i> (%) _w	\bar{M}_{age} [95% CI]	<i>n</i> (%) _w	\bar{M}_{age} [95% CI]	<i>n</i> (%) _w	\bar{M}_{age} [95% CI]
Males								
Heterosexual	10,223 (93.8)	14.82 [14.70, 14.95]	13,880 (95.3)	15.04 [14.98, 15.09]	11,573 (94.1)	15.00 [14.96, 15.04]	10,024 (93.2)	14.98 [14.94, 15.02]
Mostly Hetero.	418 (4.2)	15.24 [15.07, 15.41]	432 (3.2)	15.39 [15.25, 15.53]	459 (3.9)	15.55 [15.47, 15.64]	463 (4.4)	15.54 [15.46, 15.62]
Bisexual	127 (1.1)	14.74 [14.55, 14.93]	122 (0.9)	15.52 [15.34, 15.69]	141 (1.2)	15.55 [15.41, 15.70]	153 (1.4)	15.59 [15.49, 15.68]
Gay	84 (0.9)	15.25 [15.00, 15.49]	89 (0.6)	15.86 [15.73, 15.99]	104 (0.9)	15.76 [15.57, 15.95]	118 (1.1)	15.73 [15.61, 15.85]
Females								
Heterosexual	10,829 (90.3)	14.79 [14.68, 14.89]	13,013 (87.4)	14.87 [14.80, 14.93]	11,268 (86.7)	14.98 [14.94, 15.03]	9,601 (85.6)	14.89 [14.85, 14.93]
Mostly Hetero.	919 (7.6)	15.59 [15.44, 15.73]	1,304 (9.2)	15.58 [15.49, 15.66]	1,200 (9.6)	15.55 [15.48, 15.62]	1,019 (9.5)	15.62 [15.55, 15.69]
Bisexual	217 (1.9)	15.48 [15.28, 15.68]	428 (3.0)	15.61 [15.49, 15.73]	444 (3.2)	15.44 [15.32, 15.56]	468 (4.0)	15.55 [15.46, 15.64]
Lesbian	41 (0.3)	14.75 [14.63, 14.86]	55 (0.3)	14.93 [14.80, 15.06]	65 (0.5)	15.58 [15.41, 15.76]	92 (0.9)	15.48 [15.37, 15.58]

Note. Percentages are weighted.

Table 2
Trends in Prevalence of Lifetime Use, Age of Onset, Past 30-day Use, and HED Across Years, within Sexual Orientation Groups

	Prevalence Comparison				Trend Comparison			
	1998	2003	2008	2013	1998 ^a	2003 ^a	2008 ^a	2013 ^a
	%	%	%	%	<i>aOR</i> ^b	[95% CI]	<i>aOR</i> ^b	[95% CI]
<u>Lifetime Use</u>								
Male								
Heterosexual	65.8	58.6	54.8	45.2	.34	[-.31, .38]	.44	[-.39, .48]
Mostly Hetero	71.6	60.0	66.5	56.8	.46	[-.32, .67]	.71	[-.44, 1.13]
Bisexual	60.9	67.5	72.3	57.4	.16	[.08, .33]	.30	[.13, .70]
Gay	73.2	65.4	63.7	57.4	.48	[-.24, .94]	1.54	[.87, 2.72]
Female								
Heterosexual	61.5	55.9	54.1	44.2	.36	[-.32, .40]	.48	[-.43, .54]
Mostly Hetero	74.2	76.5	72.3	66.6	.37	[-.28, .49]	.51	[-.40, .66]
Bisexual	71.7	82.5	79.4	68.6	.20	[-.13, .31]	.64	[.45, .91]
Lesbian	67.6	58.2	65.4	64.6	.43	[-.24, .77]	.27	[-.16, .46]
<u>Age of Onset</u>								
Male								
Heterosexual	51.1	42.5	36.7	23.8	.41	[-.38, .45]	.57	[-.51, .62]
Mostly Hetero	45.5	34.6	37.3	28.1	.39	[-.29, .54]	.80	[.55, 1.15]
Bisexual	67.6	48.2	48.6	21.3	.64	[-.39, 1.05]	.80	[.48, 1.33]
Gay	59.8	25.8	46.0	36.3	.58	[-.37, .90]	.92	[.56, 1.51]
Female								
Heterosexual	40.4	33.1	28.3	17.5	.41	[-.38, .45]	.56	[-.51, .62]
Mostly Hetero	38.2	32.7	33.9	20.0	.81	[-.65, 1.02]	.67	[.54, .83]
Bisexual	63.3	41.6	53.5	31.4	.65	[.45, .94]	.52	[.38, .72]
Lesbian	54.5	60.9	55.4	30.1	.39	[-.20, .77]	.60	[.36, .98]
<u>Past 30-Day Use</u>								
Male								
Heterosexual	43.9	39.5	36.4	28.0	.76	[-.67, .85]	.76	[-.67, .86]
Mostly Hetero	52.2	38.2	44.4	34.4	.56	[-.37, .84]	.95	[.59, 1.53]
Bisexual	42.7	43.0	52.4	38.2	.67	[-.40, 1.13]	.91	[.44, 1.91]

Table 3

Male Sexual Orientation Disparities in Lifetime Alcohol Use, Age of Onset, Past 30-day Alcohol Use, and Past 30-day Heavy Episodic Drinking, within Year

	1998		2003		2008		2013	
	<i>aOR^b</i>	[95% CI]	<i>aOR^b</i>	[95% CI]	<i>aOR^b</i>	[95% CI]	<i>aOR^b</i>	[95% CI]
Lifetime Alcohol Use ^a								
Mostly Hetero	1.09	[.83, 1.44]	.89	[.64, 1.25]	1.33	[1.06, 1.68]	1.24	[.99, 1.55]
Bisexual	.82	[.48, 1.39]	1.21	[.74, 1.97]	1.79	[1.10, 2.92]	1.26	[.85, 1.87]
Gay	1.22	[.65, 2.26]	.90	[.50, 1.61]	1.06	.63, 1.78]	1.19	[.78, 1.80]
Age of Onset ^a								
Mostly Hetero	1.01	[.75, 1.36]	.83	[.54, 1.27]	1.21	[.91, 1.61]	1.35	[1.01, 1.81]
Bisexual	1.93	[1.08, 3.44]	1.40	[.71, 2.77]	1.86	[1.16, 3.00]	.95	[.54, 1.67]
Gay	1.76	[.96, 3.26]	.66	[.36, 1.18]	1.85	[1.05, 3.25]	2.38	[1.39, 4.05]
Past 30-Day Alcohol Use ^a								
Mostly Hetero	1.20	[.94, 1.52]	.79	[.57, 1.09]	1.09	[.87, 1.37]	1.07	[.85, 1.34]
Bisexual	.98	[.59, 1.65]	.96	[.59, 1.55]	1.60	[1.05, 2.44]	1.28	[.87, 1.86]
Gay	1.36	[.73, 2.55]	.94	[.48, 1.83]	1.68	[1.00, 2.85]	1.56	[1.03, 2.35]
Past 30-Day HED ^a								
Mostly Hetero	1.06	[.80, 1.41]	.56	[.37, .86]	.92	[.72, 1.18]	.79	[.61, 1.02]
Bisexual	1.23	[.68, 2.22]	1.07	[.64, 1.78]	1.45	[.91, 2.31]	1.41	[.95, 2.08]
Gay	1.33	[.69, 2.58]	.66	[.32, 1.36]	1.39	[.80, 2.41]	1.42	[.88, 2.30]

Note. Data were weighted. Odds ratios in bold indicate $p < .05$. Mostly Hetero = Mostly Heterosexual.

^a Heterosexual is reference group

^b Adjusted for age

Table 4

Female Sexual Orientation Disparities in Lifetime Alcohol Use, Age of Onset, Past 30-day Alcohol Use, and Past 30-day Heavy Episodic Drinking, within Year

	1998		2003		2008		2013	
	<i>aOR^b</i>	95% CI	<i>aOR^b</i>	95% CI	<i>aOR^b</i>	95% CI	<i>aOR^b</i>	95% CI
Lifetime Alcohol Use ^a								
Mostly Hetero	1.34	[1.09, 1.65]	2.05	[1.68, 2.50]	1.85	[1.57, 2.19]	1.96	[1.64, 2.34]
Bisexual	1.23	[.77, 1.96]	2.99	[1.94, 4.60]	3.07	[2.33, 4.07]	2.35	[1.80, 3.07]
Lesbian	1.48	[.69, 3.14]	1.07	[.52, 2.22]	1.25	[.64, 2.47]	1.90	[1.09, 3.33]
Age of Onset ^a								
Mostly Hetero	1.38	[1.07, 1.78]	1.27	[1.07, 1.52]	1.54	[1.27, 1.86]	1.35	[1.10, 1.66]
Bisexual	4.46	[2.75, 7.22]	1.79	[1.35, 2.37]	3.38	[2.55, 4.48]	2.37	[1.80, 3.11]
Lesbian	1.75	[.68, 4.50]	4.11	[1.33, 12.68]	4.53	[1.74, 11.80]	2.19	[1.13, 4.25]
Past 30-Day Alcohol Use ^a								
Mostly Hetero	1.18	[.99, 1.42]	1.70	[1.43, 2.02]	1.75	[1.52, 2.03]	1.78	[1.52, 2.09]
Bisexual	1.43	[.95, 2.15]	2.08	[1.48, 2.93]	2.32	[1.80, 3.00]	1.63	[1.30, 2.06]
Gay	1.76	[.82, 3.75]	1.26	[.61, 2.61]	1.44	[.76, 2.71]	1.04	[.60, 1.80]
Past 30-Day HED ^a								
Mostly Hetero	1.19	[.99, 1.44]	1.49	[1.23, 1.80]	1.52	[1.31, 1.78]	1.56	[1.32, 1.83]
Bisexual	1.40	[.92, 2.14]	1.99	[1.37, 2.87]	2.00	[1.58, 2.53]	1.58	[1.24, 2.01]
Lesbian	1.88	[.88, 4.02]	1.09	[.56, 2.12]	1.98	[1.06, 3.71]	1.01	[.58, 1.77]

Note. Data were weighted. Odds ratios in bold indicate $p < .05$. Mostly Hetero = Mostly Heterosexual.

^a Heterosexual is reference group

^b Adjusted for age

Table 5

Trends in Disparities in Lifetime Alcohol Use, Age of Onset, Past 30-Day Alcohol Use, and Past 30-Day HED [HED]: Interactions Between Sexual Orientation and Year^a

	Lifetime Alcohol Use		Age of Onset		30-Day Alcohol Use		30-Day HED	
	aOR ^b	[95% CI]	aOR ^b	[95% CI]	aOR ^b	[95% CI]	aOR ^b	[95% CI]
Males								
Mostly Hetero by Yr1998	1.19	[.83, 1.69]	1.36	[.90, 2.06]	.69	[-42, 1.13]	.77	[-53, 1.12]
Mostly Hetero by Yr2003	1.45	[.97, 2.18]	1.62	[.96, 2.72]	1.26	[.70, 2.29]	1.45	[.88, 2.37]
Mostly Hetero by Yr2008	.96	[.70, 1.31]	1.12	[.75, 1.68]	.85	[.53, 1.38]	.88	[.61, 1.25]
Bisexual by Yr1998	1.60	[.83, 3.09]	.49	[.22, 1.10]	.77	[-31, 1.90]	1.16	[-57, 2.37]
Bisexual by Yr2003	1.09	[.59, 2.02]	.67	[.28, 1.64]	1.19	[.39, 3.67]	1.35	[.71, 2.55]
Bisexual by Yr2008	.73	[.39, 1.34]	.51	[.25, 1.07]	1.47	[.63, 3.44]	.98	[.54, 1.78]
Gay by Yr1998	1.03	[.48, 2.18]	1.36	[.61, 3.05]	.78	[-26, 2.34]	1.09	[-48, 2.49]
Gay by Yr2003	1.41	[.70, 2.86]	3.56	[1.60, 7.90]	1.80	[.58, 5.57]	2.23	[.95, 5.25]
Gay by Yr2008	1.17	[.61, 2.25]	1.30	[.60, 2.81]	.79	[-30, 2.07]	1.04	[-51, 2.12]
Females								
Mostly Hetero by Yr1998	1.52	[1.16, 1.99]	.99	[.72, 1.38]	1.58	[1.25, 2.01]	1.34	[1.05, 1.72]
Mostly Hetero by Yr2003	.98	[.76, 1.27]	1.05	[.80, 1.38]	1.09	[.86, 1.37]	1.08	[.85, 1.39]
Mostly Hetero by Yr2008	1.08	[.85, 1.37]	.89	[.68, 1.18]	1.04	[.84, 1.28]	1.05	[.84, 1.30]
Bisexual by Yr1998	1.95	[1.14, 3.33]	.54	[.31, .94]	1.20	[.75, 1.92]	1.16	[.71, 1.88]
Bisexual by Yr2003	.79	[.48, 1.30]	1.31	[.88, 1.95]	.81	[.54, 1.22]	.82	[.53, 1.27]
Bisexual by Yr2008	.77	[.53, 1.11]	.72	[.48, 1.06]	.72	[-52, 1.01]	.81	[-58, 1.13]
Lesbian by Yr1998	1.29	[.51, 3.26]	1.26	[.40, 3.96]	.60	[-24, 1.54]	.55	[-21, 1.41]
Lesbian by Yr2003	1.80	[.73, 4.45]	.52	[.14, 1.95]	.86	[-35, 2.11]	.95	[-40, 2.27]
Lesbian by Yr2008	1.54	[.65, 3.63]	.50	[.16, 1.57]	.74	[-33, 1.67]	.52	[-23, 1.20]

Note. Data were weighted. Odds ratios in bold indicate $p < .05$; Mostly Hetero = Mostly heterosexual

^a Heterosexual by Yr2013 is reference group

^b Adjusted model included sexual orientation, survey year, and age along with orientation-by-year interactions.