

■ Erratum

Correction: The Effect of Lifestyle Changes on Blood Pressure Control among Hypertensive Patients

Myung Hwa Yang¹, Seo Young Kang¹, Jung Ah Lee¹, Young Sik Kim^{1*}, Eun Ju Sung², Ka-Young Lee³, Jun-Su Kim³, Han Jin Oh⁴, Hee Chul Kang⁵, Sang Yeoup Lee⁶

¹Department of Family Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

²Department of Family Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Korea

³Department of Family Medicine, Inje University Busan Paik Hospital, Inje University College of Medicine, Busan, Korea

⁴Department of Family Medicine, Vievisnamuh Hospital, Seoul, Korea

⁵Department of Family Medicine, Yonsei University College of Medicine, Seoul, Korea

⁶Department of Family Medicine, Pusan National University Yangsan Hospital, Pusan National University School of Medicine, Yangsan, Korea

<https://doi.org/10.4082/kjfm.2017.38.4.173>

Korean J Fam Med 2017;38:173-180

There was an error in the article, “The effect of lifestyle changes on blood pressure control among hypertensive patients¹⁾”. The authors found an error after the publication of this work¹⁾ and asked the editorial board to correct it. So we would like to replace Table 3 of the original paper with a revised version. A revised version of Table 3 in the original article is given below.

We sincerely apologized to the readers.

REFERENCE

1. Yang MH, Kang SY, Lee JA, Kim YS, Sung EJ, Lee KY, et al. The effect of lifestyle changes on blood pressure control among hypertensive patients. Korean J Fam Med 2017;38:173-80.

*Corresponding Author: Young Sik Kim Tel: +82-2-3010-3811, Fax: +82-2-3010-0578, E-mail: youngkim@amc.seoul.kr

<Original version Table 3>

Table 3. Changes in the performance index between patients with successful and failed blood pressure control

Variable	Total	Failure	Success	Univariate OR		Adjusted OR*	
				(95% CI)	P-value	(95% CI)	P-value
Reduced alcohol use among drinkers	1,139						
Yes	88 (7.7)	22 (25.0)	66 (75.0)	1 (reference)		1 (reference)	
No	243 (21.3)	43 (17.7)	200 (82.3)	0.645 (0.360–1.157)	0.141	0.727 (0.398–1.330)	0.301
Unknown	808 (70.9)						
Reduction in frequency of alcohol consumption	1,139						
No change	201 (17.6)	38 (18.9)	163 (81.1)	1 (reference)		1 (reference)	
<1/wk	24 (2.1)	2 (8.3)	22 (91.7)	0.390 (0.088–1.730)	0.215	0.421 (0.093–1.895)	0.260
1–2/wk	14 (1.2)	2 (14.3)	12 (85.7)	0.715 (0.154–3.328)	0.669	0.569 (0.119–2.713)	0.479
>3/wk	4 (0.4)	1 (25.0)	3 (75.0)	1.430 (0.145–14.127)	0.760	1.500 (0.145–150.494)	0.734
Unknown†	896 (78.7)						
Increased body mass index	1,062						
Yes	250	59 (23.6)	191 (76.4)	1 (reference)		1 (reference)	
No	812	101 (12.4)	711 (87.6)	2.175 (1.519–3.113)	<0.001	2.010 (1.388–2.911)	<0.001
>0.5 kg/m ²	96 (9.0)	25 (26.0)	71 (74.0)	1 (reference)		1 (reference)	
≤0.5 kg/m ²	966 (91.0)	135 (14.0)	831 (86.0)	2.044 (1.420–2.942)	<0.001	2.189 (1.309–3.659)	0.003
Increased physical activity	965						
Yes	882	128 (14.5)	754 (85.5)	1 (reference)		1 (reference)	
No	83	21 (25.3)	62 (74.7)	1.995 (1.175–3.387)	0.011	2.377 (1.351–4.183)	0.003
Reduced salt intake	1,096						
Yes	408	52 (12.7)	356 (87.3)	1 (reference)		1 (reference)	
No	688	121 (17.6)	567 (82.4)	1.461 (1.029–2.075)	0.034	1.324 (0.914–1.917)	0.138

Values are presented as number or number (%).

OR, odds ratio; CI, confidence interval.

*Multivariate OR calculated by adjusting for age, sex, and comorbidity. †Persons with no classification.

<Revised version Table 3>

Table 3. Changes in the performance index between patients with successful and failed blood pressure control

Variable	Total	Failure	Success	Univariate OR		Adjusted OR*	
				(95% CI)	P-value	(95% CI)	P-value
Reduced alcohol use among drinkers	1,139						
Yes	88 (7.7)	22 (25.0)	66 (75.0)	1 (reference)		1 (reference)	
No	243 (21.3)	43 (17.7)	200 (82.3)	0.645 (0.360–1.157)	0.141	0.727 (0.398–1.330)	0.301
Unknown	808 (70.9)						
Reduction in frequency of alcohol consumption	1,139						
No change	201 (17.6)	38 (18.9)	163 (81.1)	1 (reference)		1 (reference)	
<1/wk	24 (2.1)	2 (8.3)	22 (91.7)	0.390 (0.088–1.730)	0.215	0.421 (0.093–1.895)	0.260
1–2/wk	14 (1.2)	2 (14.3)	12 (85.7)	0.715 (0.154–3.328)	0.669	0.569 (0.119–2.713)	0.479
>3/wk	4 (0.4)	1 (25.0)	3 (75.0)	1.430 (0.145–14.127)	0.760	1.500 (0.145–15.494)	0.734
Unknown†	896 (78.7)						
Increased body mass index	1,062						
No	812	101 (12.4)	711 (87.6)	1 (reference)		1 (reference)	
Yes	250	59 (23.6)	191 (76.4)	2.175 (1.519–3.113)	<0.001	2.010 (1.388–2.911)	<0.001
>0.5 kg/m ²	966 (91.0)	135 (14.0)	831 (86.0)	1 (reference)		1 (reference)	
≤0.5 kg/m ²	96 (9.0)	25 (26.0)	71 (74.0)	2.044 (1.420–2.942)	<0.001	2.189 (1.309–3.659)	0.003
Increased physical activity	965						
Yes	882	128 (14.5)	754 (85.5)	1 (reference)		1 (reference)	
No	83	21 (25.3)	62 (74.7)	1.995 (1.175–3.387)	0.011	2.377 (1.351–4.183)	0.003
Reduced salt intake	1,096						
Yes	408	52 (12.7)	356 (87.3)	1 (reference)		1 (reference)	
No	688	121 (17.6)	567 (82.4)	1.461 (1.029–2.075)	0.034	1.324 (0.914–1.917)	0.138

Values are presented as number or number (%).

OR, odds ratio; CI, confidence interval.

*Multivariate OR calculated by adjusting for age, sex, and comorbidity. †Persons with no classification.