



HHS Public Access

Author manuscript

Nature. Author manuscript; available in PMC 2017 October 15.

Published in final edited form as:

Nature. 2016 November 24; 539(7630): 498–499. doi:10.1038/nature20475.

Clamping down on copy errors

NEIL M. KAD and

School of Biosciences, University of Kent, Canterbury CT2 7NJ, UK

BENNETT VAN HOUTEN

Department of Pharmacology and Chemical Biology, University of Pittsburgh, University of Pittsburgh Cancer Institute, Hillman Cancer Center, Pittsburgh, Pennsylvania 15213, USA

Abstract

Repair enzymes must communicate across hundreds of nucleotides to undo errors made during DNA replication. Imaging reveals that the enzymes do this by forming a series of ring-like clamps that diffuse along the DNA.

As we write up results for publication, any spelling errors are quickly identified by our computer's built-in spellchecker. However, this is not an invention of the modern world; nature has been using DNA spellcheckers for millions of years to avoid genetic errors that arise during DNA replication, which it corrects through a process called mismatch repair. In the bacterium *Escherichia coli*, repair requires communication between enzymes across long stretches of DNA, and how this occurs has been hotly debated for decades^{1–3}. On page 583, Liu *et al.*⁴ help to solve this mystery by using state-of-the-art techniques to analyse mismatch repair at the single-molecule level.

DNA exists as a double-stranded duplex, connected across the strands by complementary base-pairing — guanine (G) with cytosine (C) and adenine (A) with thymine (T). During DNA replication, the duplex unwinds, and each strand is copied using new nucleotides to create a freshly synthesized daughter strand. Errors occur when incorrect bases are incorporated into the daughter strand, creating a mismatch that requires subsequent repair to prevent mutations from arising. In *E. coli*, DNA with the base sequence GATC is normally tagged with a methyl group, but the newly synthesized DNA is temporarily unmethylated. These 'hemi-methylated' regions provide a means for mismatch-repair enzymes to distinguish between parent and daughter DNA.

Hemi-methylated sequences are also the sites at which the endonuclease enzyme MutH generates a single-strand break in the erroneous DNA strand during mismatch repair. Because of the rarity of GATC sequences, MutH-generated breaks can occur hundreds of bases from the site of the mismatch. Following breakage, the damaged strand is excised by an exonuclease enzyme that works from the break to the error, and the DNA is resynthesized to incorporate the correct base. A key aspect of this process is the coordination of mismatch detection with loading of the DNA-excision machinery, to ensure that excision occurs towards, rather than away from, the mismatch.

For the past decade, it has been thought^{5,6} that the protein MutS, which recognizes DNA mismatches, acts as a sliding clamp — a ring around DNA that can efficiently move up or down the two strands, enabling communication between the mismatch and the hemi-methylated sites. A third protein, MutL, activates the cutting activity of MutH. But how MutS links mismatch detection to the activation of MutH has been unclear.

To study initiation of mismatch repair, Liu *et al.* added fluorescent tags to MutH, MutS and MutL and analysed the movements of the individual proteins along single DNA molecules. They found that MutL is loaded onto DNA through interactions with a MutS molecule that has already located a mismatch. This observation builds on a structural study⁷ showing that the DNA duplex is positioned in the open channel of the MutS ring following mismatch verification, and that MutL binding sites on MutS are revealed during this process, permitting formation of the MutS–MutL complex.

Next, the authors confirmed another previous observation⁶ — that the MutS–MutL complex forms a single sliding clamp. Extending these observations, they showed that MutL can detach from MutS to act as a clamp on its own. In this role, MutL undertakes short excursions away from MutS. Surprisingly, some of these excursions pass behind MutS, an observation that Liu *et al.* took to imply that MutL might pass through the MutS channel. Therefore, MutL can be thought of as acting like a yo-yo, separately scouting the DNA. But scouting for what?

The answer comes from the group's demonstration that MutL can recruit MutH to the DNA, thereby setting the stage for the excision phase of mismatch repair. The MutL–MutH complex behaves similarly to MutL alone, detaching from MutS and diffusing along DNA more quickly than when the three proteins are bound together — presumably in search of hemi-methylated sites. The authors propose that the MutS–MutL–MutH complex is the dominant species, because the sliding-clamp nature of MutS means that this complex is tethered to the DNA. However, efficient searching is also possible with MutL–MutH, which Liu and colleagues posit hops along the DNA.

Together, these observations might have solved the long-standing mystery of how the mismatch and excision sites interact, superseding previous models in which the DNA is looped through the MutS clamp or in which the clamp moves in a targeted manner along the duplex^{1,8}. The model that emerges from these data suggests that MutS acts as a guiding clamp, sending out scouting clamps to search for hemi-methylated GATC sequences in a highly energy-efficient manner (Fig. 1).

In 2015, Paul Modrich received the Nobel Prize in Chemistry for his seminal contributions to our understanding of mismatch repair¹. However, many aspects of the process remain unresolved. One is that mismatch repair in eukaryotic cells (those that have a nucleus) involves more-complex protein–protein and protein–DNA interactions that have yet to be fully described. For instance, the eukaryotic equivalent of MutL has endonuclease activity, and eukaryotes use different mechanisms for excision depending on whether the break point is 3' or 5' of the mismatch^{1,3}. Given this complexity, a better understanding of eukaryotic mismatch repair will surely present new mechanistic surprises. The use of single-molecule

approaches to enable analysis of the eukaryotic process, already under way in several laboratories, will be central to elucidating these systems.

The principles outlined by Liu *et al.* might apply to enzymes involved in other types of DNA repair. A recent report⁹ revealed that the yeast protein Rad4, which is involved in nucleotide-excision repair, does not directly bind damaged sites, but instead diffuses for up to 1 kilobase around the damage, providing a dynamic platform for the recruitment of other repair proteins. Furthermore, the protein Mfd, which dislodges RNA polymerase enzymes that become stalled while transcribing DNA, then scouts ahead for DNA damage on the transcribed strand¹⁰.

The next chapter of the mismatch-repair story will surely see experiments that follow the entire process of repair, enabling observation of every protein in real time in a single assay. With single-molecule fluorescence techniques developing rapidly, such experiments are in sight, and should provide a molecular understanding of biology *in vitro* that can be adapted to systems *in vivo*.

References

1. Modrich P. *Angew Chem Int Edn.* 2016; 55:8490–8501.
2. Fishel R. *J Biol Chem.* 2015; 290:26395–26403. [PubMed: 26354434]
3. Kunkel TA, Erie DA. *Annu Rev Genet.* 2015; 49:291–313. [PubMed: 26436461]
4. Liu J, et al. *Nature.* 2016; 539:583–587. [PubMed: 27851738]
5. Jeong C, et al. *Nature Struct Mol Biol.* 2011; 18:379–385. [PubMed: 21278758]
6. Gorman J, et al. *Proc Natl Acad Sci USA.* 2012; 109:E3074–E3083. [PubMed: 23012240]
7. Groothuizen FS, et al. *eLife.* 2015; 4:e06744. [PubMed: 26163658]
8. Iyer RR, Pluciennik A, Burdett V, Modrich PL. *Chem Rev.* 2006; 106:302–323. [PubMed: 16464007]
9. Kong M, et al. *Mol Cell.* 2016; 64:376–387. [PubMed: 27720644]
10. Haines NM, Kim YIT, Smith AJ, Savery NJ. *Proc Natl Acad Sci USA.* 2014; 111:4037–4042. [PubMed: 24554077]




Figure 1. Scouting for errors

a, Following DNA replication, the protein MutS clamps around DNA at mismatches — sites of incorrect pairing between the two complementary DNA strands. **b**, Liu *et al.*⁴ report that, once a mismatch is detected, MutS recruits the protein MutL, and the two proteins slowly slide together up and down DNA. **c**, MutL in turn recruits the enzyme MutH, and the three move as one complex. **d**, MutL and MutH can detach from MutS to move rapidly along DNA in search of hemi-methylated sites (H) — sequences at which the parent strand is tagged with a methyl group but the mismatched daughter strand is not. The daughter strand is subsequently cleaved by MutH, initiating DNA excision and repair.