



HHS Public Access

Author manuscript

J Neurosci Res. Author manuscript; available in PMC 2017 October 16.

Published in final edited form as:

J Neurosci Res. 2010 November 01; 88(14): 2991–3001. doi:10.1002/jnr.22444.

The Decade of the Dendritic NMDA Spike

Srdjan D. Antic, Wen-Liang Zhou, Anna R. Moore, Shaina M. Short, and Katerina D. Ikonomu

Department of Neuroscience, Univ. Connecticut Health Center, Farmington, CT 06030, USA

Abstract

In the field of cortical cellular physiology, much effort has been invested in understanding thick apical dendrites of pyramidal neurons and the regenerative sodium and calcium spikes that take place in the apical trunk. Here we focus on thin dendrites of pyramidal cells (basal, oblique, and tuft dendrites), and we discuss one relatively novel form of an electrical signal (“NMDA spike”) that is specific for these branches. Basal, oblique, and apical tuft dendrites receive a high density of glutamatergic synaptic contacts. Synchronous activation of 10–50 neighboring glutamatergic synapses triggers a local dendritic regenerative potential, NMDA spike/plateau, which is characterized by significant local amplitude (40–50 mV) and an extraordinary duration (up to several hundred milliseconds). The NMDA plateau potential, when it is initiated in an apical tuft dendrite, is able to maintain a good portion of that tuft in a sustained depolarized state. However, if NMDA-dominated plateau potentials originate in proximal segments of basal dendrites, they regularly bring the neuronal cell body into a sustained depolarized state, which resembles a cortical up state. At each dendritic initiation site (basal, oblique, and tuft) an NMDA spike creates favorable conditions for causal interactions of active synaptic inputs, including the spatial or temporal binding of information, as well as processes of short-term and long-term synaptic modifications (e.g., long-term potentiation or long-term depression). Because of their strong amplitudes and durations, local dendritic NMDA spikes make up the cellular substrate for multisite independent subunit computations that enrich the computational power and repertoire of cortical pyramidal cells. We propose that NMDA spikes are likely to play significant roles in cortical information processing in awake animals (spatiotemporal binding, working memory) and during slow-wave sleep (neuronal up states, consolidation of memories).

Keywords

synaptic integration; plateau potential; summation; regeneration; up state; neocortex; basal; oblique; glutamate

The previous decade 2000 – 2009 started with a paper describing N-methyl-D-aspartate (NMDA) spikes in basal dendrites of cortical layer 5 pyramidal neurons (Schiller et al., 2000), and ended with a paper describing NMDA spikes in the apical tuft dendrites of the same neuron type (Larkum et al., 2009). The study conducted by Schiller et al., (2000) has pioneered this area of research in two fundamental ways. First, it has provided direct

experimental evidence that an isolated NMDA conductance can support regenerative membrane potential (spike) in a dendritic branch of a living central nervous system (CNS) neuron. Second, it has expanded the focus of dendritic physiology from the apical dendrite to other branches of cortical pyramidal neurons (e.g. basal, oblique and tuft) – here termed “non-apical” dendrites.

For didactic purposes, one can divide the last twenty years of cortical dendritic physiology into two nicely rounded calendar decades (Table 1). In the first decade, 1990 – 1999, experimenters systematically explored the physiological properties of the apical dendrite in pyramidal neurons. They found that apical dendrites express voltage-gated sodium, voltage-gated potassium and voltage-gated calcium channels, which allowed apical dendrites to actively propagate action potentials (AP) back into the dendritic tree (Stuart and Sakmann, 1994; Magee and Johnston, 1995; Spruston et al., 1995; Waters et al., 2005) and participate in the process known as spike timing dependent synaptic plasticity (STDP) - reviewed in (Lisman and Spruston, 2005; Sjöstrom et al., 2008). Aside from this role in AP backpropagation, upon adequate stimulation the active apical membrane conductances are also involved in producing regenerative membrane potentials, also known as “dendritic spikes” (Table 1, *Local Regenerative Potentials*)(Herreras, 1990; Kim and Connors, 1993; Schiller et al., 1997; Kamondi et al., 1998). Among the spikes that initiate in the apical dendrite the two most impressive entities were local spikes that failed to invade the soma (Golding et al., 2002), and calcium spikes which could be triggered either by coincident arrival of backpropagating AP and a distal excitatory postsynaptic potential in brain slice preparations (Larkum et al., 1999), or by whisker deflection in living animals (Helmchen et al., 1999).

Local (dendrite-restricted) spikes that fail to invade the cell body brought a lot of excitement in the field, because they were able to mediate long-term potentiation of synaptic inputs in the absence of neuronal action potential firing (Golding et al., 2002). Before 2002 one could argue that local dendritic spikes were just side effects of the lateral diffusion of voltage-gated channels from the soma into the apical trunk; a trifling epiphenomenon, with no specific physiological relevance. The 2002 paper by Golding et al. provided a solid factual demonstration of the utility of a purely dendritic regenerative event, and the critical role such an event might play in cellular neurobiology (see also (Remy and Spruston, 2007)).

Calcium action potentials in the dendritic initiation zone near the main apical bifurcation, described by Larkum et al., 1999 (Table 1, *Calcium Spike*)(Fig. 1B₂), constitute a mechanism by which the most distal synaptic inputs in neocortical pyramidal cells (tuft inputs) could impact the neuronal output (axonal action potential initiation zone). Besides a powerful boost for layer 1 and layer 2 synaptic inputs, the dendritic calcium action potentials also serve as reliable coincidence detectors. Recall that in layer 5 pyramidal neurons these spikes initiate near the primary bifurcation of the apical trunk (Fig. 1B₂, *Initiation Site*) in response to synchronous strong input into both deep and superficial cortical layers (Larkum et al., 1999, 2001).

It can be safely said that this 10-year long (1990 – 1999), and quite prolific, experimental quest on the apical dendrite function was propelled by two technical achievements that took

place in the early nineties: (1) The infrared differential interference contrast (IR-DIC) visualization of the apical trunk (infrared video microscopy)(Dodt, 1993); and (2) The dendritic patch-electrode recordings (Stuart and Sakmann, 1994). Ten years ago, these two methods worked well on thick apical dendrites, but not so well in thin submicron diameter dendritic branches. The thin branches (basal, oblique and tuft dendrites) were poorly visible in standard IR-DIC setup, which rendered their visually-guided patching quite difficult and impractical for systematic analysis. This may explain a relatively low yield in the number of studies addressing the membrane physiology of non-apical dendrites in that period. Only occasionally, a small section of an elaborate paper would explore the physiology of a thin dendrite as a side project (Regehr and Tank, 1992; Yuste et al., 1994; Schiller et al., 1995; Cossart et al., 2000).

Schiller et al. (2000) turned the searchlight upon small-diameter branches by introducing one novel experimental approach, suitable for the analysis of the thin-dendrite function, and by showing that some exciting physiological phenomena reside not in the apical trunk, but rather in non-apical branches, in this case, basal dendrites of cortical pyramidal neurons. The novel experimental approach was not based on the improvement in recording of dendritic physiological signals per se, but rather on the improvement in the dendritic stimulation technique. Fueled by advances in laser scanning technology and confocal microscopy, a new stimulation technique emerged that allowed a precise delivery of glutamate ions on one thin dendrite of choice. This technique, also known as glutamate uncaging (Callaway and Katz, 1993; Pettit et al., 1997; Wei et al., 2001), provided researchers with the ability to precisely control four critical parameters of glutamatergic stimulation: **(1)** A specific location (in any given segment along the thin dendritic branch); **(2)** An exact area of the membrane receiving the stimulus in the shape of a glutamate cloud of variable diameter (several micrometers usually); **(3)** Stimulus intensity (by adjusting the intensity of the laser beam); and **(4)** Stimulus duration (by setting the time for which a laser beam is turned on). Once it has been established that glutamatergic stimuli were restrictively targeting one thin dendritic branch, the somatic whole-cell recordings could then be used to detect membrane potential changes that arise in the targeted thin dendrite and propagate to the cell body. In addition to whole-cell recordings Schiller et al., (2000) also used dendritic calcium imaging to monitor the glutamate evoked dendritic calcium influx. This trinity of experimental techniques [**(1)** Glutamate stimulation (either by laser uncaging or microiontophoresis); **(2)** Optical imaging (either calcium-sensitive dye or voltage-sensitive dye); and **(3)** Somatic whole-cell recordings] would become the “Holy Trinity” of thin-dendrite physiology in the decade to come (Oakley et al., 2001b; Wei et al., 2001; Cai et al., 2004; Carter and Sabatini, 2004; Milojkovic et al., 2004; Milojkovic et al., 2005a; Noguchi et al., 2005; Losonczy and Magee, 2006; Cai et al., 2007; Milojkovic et al., 2007; Sobczyk and Svoboda, 2007; Losonczy et al., 2008; Major et al., 2008; Suzuki et al., 2008; Remy et al., 2009). The availability of experimental methods was not the only propellant for the ensuing interest in thin dendrites. A genuine scientific curiosity was also at play. In a particular case of thin dendrites – understanding their properties is an important step towards understanding operation of cortical neurons and networks. Basal, oblique and tuft branches receive the overwhelming majority of excitatory glutamatergic synaptic inputs impinging on any given cortical pyramidal cell. It has been estimated that as much as 85% of the total number of excitatory

glutamatergic drive, which impinges on an individual layer 5 pyramidal neuron, actually arrives on its basal, oblique and tuft dendrites, while just a tiny fraction, less than 15%, is being received by the soma and the apical trunk below main bifurcation (Larkman, 1991). Recall that dendritic spines are putative sites of long-term potentiation and long-term depression (Emptage et al., 1999; Luscher et al., 2000; Matsuzaki et al., 2004), reviewed in (Yuste and Bonhoeffer, 2001; Nimchinsky et al., 2002). Being the principle bearers of dendritic spines in pyramidal neurons, the basal oblique and tuft dendrites are most likely the key elements of cellular mechanisms for learning and memory (Yang et al., 2009). Moreover, if thin dendrites were involved in the processes of cortical computations (synaptic integration) and cortical learning (synaptic plasticity) then it would be prudent to know more about their intrinsic physiological properties. This was, in our opinion, the second catalyst for the wave of thin-dendrite physiology exploration in the past decade (2000 – 2009).

Arguably, the most exciting feature of the thin-dendrite membrane is its ability to initiate a local regenerative membrane potential (spike), whose major ionic contributor is not a sodium or calcium voltage-gated channel, but rather a ligand-gated receptor-channel, NMDAr (Schiller et al., 2000; Rhodes, 2006; Major et al., 2008; Larkum et al., 2009). When glutamate-binding sites of an NMDA receptor-channel become occupied with two glutamate ions then the voltage sensitivity of the NMDAr current exhibits a region of negative slope conductance due to the relief of magnesium block (Mayer et al., 1984; Nowak et al., 1984). In other words, in conditions of abundant glutamate supply the current-voltage (I–V) relationship of an NMDAr current has a similar waveform as the I–V relationship of the voltage-gated sodium channel, and therefore, during strong glutamatergic release the NMDA receptor-channels fire a regenerative NMDA spike, just like sodium channels upon adequate depolarization fire an action potential. Much earlier research had established that NMDAr current has a region of negative slope conductance, which is a biophysical substrate of regenerative property (Mayer et al., 1984; Nowak et al., 1984), but in 2000 it was not clear if CNS neurons could actually support NMDA spikes. At that time there were three obstacles to solving the “NMDA spike question”: **(1)** It was not clear if any neuronal compartment possessed an adequate NMDA channel density (conductance density) to support NMDA spike initiation. **(2)** What type and what intensity of glutamatergic stimulation would be needed to efficiently activate those conductances?; and **(3)** What kind of experimental design one should employ to demonstrate the existence of NMDA spikes in real neurons? The first problem **(1)** was solved by shifting the experimental focus from the apical trunk to smaller diameter branches (basal dendrites), where surface to volume ratio is high, and where NMDAr-bearing dendritic spines reside in great numbers. The second problem **(2)** was solved by using the power of a ultra-violet laser beam to raise the local concentration of glutamate to millimolar values. The third problem **(3)** was solved by combining an exogenous glutamate pulse with bath application of drugs that blocked both voltage-gated sodium and calcium channels (Tetrodotoxin (TTX) and Cd^{2+} , respectively). It is important to note that synaptic stimulation is not possible in the presence of TTX and Cd^{2+} , because sodium and calcium channels are necessary for triggering the synaptic release of glutamate. Schiller et al., 2000 arrived at an excellent experimental design to solve the “dendritic NMDA spike” question by combining three aforementioned approaches: (1) glutamatergic stimulation, (2) whole-cell recording and (3) dendritic calcium imaging. Glutamatergic

stimulations were not only instrumental for finding the adequate (threshold) stimulus intensity; but also, for the graded control of the stimulus intensity between sub and suprathreshold levels, and for co-application with voltage-channel antagonists (TTX and Cd^{2+}). Whole-cell recordings were critical for precise detection of nonlinearity in the family of membrane potential changes (hallmark of spike generation – see Schiller et al., (2000); their figure 1b). Finally, dendritic calcium imaging unequivocally proved that only one segment of one dendritic branch had been receiving the glutamate input (Schiller et al., 2000, their figure 1d).

Physiological Properties of Dendritic NMDA Spikes

Glutamate threshold

The glutamate concentration needed to induce an NMDA spike can be deduced from glutamate uncaging experiments. A typical concentration of caged glutamate in the bath solution is 1 mM (Schiller et al., 2000). Under these conditions the maximum concentration of uncaged (free) glutamate in the center of the laser beam (assuming 100% success rate of UV photolysis) cannot exceed 1 mM. Since glutamate uncaging is a very reliable stimulus for dendritic NMDA spikes (Schiller et al., 2000), we conclude that glutamate concentrations ≥ 1 mM are sufficient to generate this event.

Ionic composition

Pure NMDA spikes (Fig. 1A, NMDA Spike) initiate in thin dendrites when two major voltage-gated conductances (Na^+ and Ca^{2+}) are blocked with drugs (TTX and Cd^{2+}) and when the concentration of glutamate reaches a certain threshold. In normal physiological saline (no drugs) suprathreshold glutamatergic stimulations generate NMDA spikes mixed with strong activation of dendritic voltage-gated Na^+ and Ca^{2+} channels. The resulting plateau potential (Fig. 1A, *Plateau Potential*) is a complex spike comprised of several complementary conductances that activate in a relatively strict temporal order. According to model predictions (Schiller et al., 2000) the plateau potential seems to involve a “spike-chain” mechanism: the AMPA response evokes a fast local sodium spikelet, which elicits a slower calcium-mediated regenerative response, which in turn initiates the full blown NMDA spike. These model predictions were based on whole-cell somatic recordings (Schiller et al., 2000). Voltage-sensitive dye recordings performed in basal dendrites that were stimulated either synaptically, or by exogenous glutamate application, revealed the actual waveform of the plateau potential at dendritic loci (Milojkovic et al., 2004). This dendritic membrane potential transient is characterized by a fast onset, initial spikelet (Fig. 1A, *Na⁺ Spikelet*), a prominent plateau phase and an abrupt collapse at the end of the plateau phase (Fig. 1A, *Plateau potential*). The initial sodium spikelet was not a robust and widespread phenomenon. On the contrary, in the greater majority of basal dendrites the initial sodium spikelet did not accompany glutamate-evoked plateau potentials (Milojkovic et al., 2005b; Acker and Antic, 2009). Although suprathreshold glutamate stimulations invariably evoked plateau potentials in every basal dendrite tested so far (several hundred), only a fraction (10 – 25%) of basal branches belonging to neocortical layer 5 pyramidal neurons possess the adequate density of sodium channels to support local sodium regenerative potentials (Milojkovic et al., 2005b; Nevian et al., 2007; Acker and Antic,

2009). Thus the activation of dendritic voltage-gated sodium channels is not the determining factor for the initiation of glutamate-dependent dendritic spikes. Sodium channels assist in spike initiation, but are not necessary for it (Schiller et al., 2000; Oakley et al., 2001b; Wei et al., 2001; Milojkovic et al., 2005a; Rhodes, 2006; Milojkovic et al., 2007; Major et al., 2008; Larkum et al., 2009).

Amplitude

Voltage-sensitive dye recordings have determined that the amplitude of a glutamate-evoked dendritic plateau potential during its plateau phase is approximately 2/3 of the amplitude of a backpropagating action potential in the same dendritic site (Milojkovic et al., 2004; their figure 4). Previously, computer simulation based on realistic neuronal morphology and multi-site voltage-sensitive dye measurements predicted that in basal dendritic segments 150 μm from the soma, the amplitude of the back-propagating action potential (AP) is at least 60 mV (Antic, 2003). Based on these two pieces of data Milojkovic et al (2004) concluded that the amplitude of the glutamate-evoked plateau at 150 μm from the soma was at least 40 mV. Direct patch electrode recordings from basal dendrites showed that the peak-to-peak amplitude of the dendritic glutamate-evoked regenerative event (spike) is in the range between 40 and 50 mV (Nevian et al., 2007). Therefore, in terms of the peak amplitude, the glutamate-evoked NMDA-dependent regenerative potentials are quite comparable and could stand shoulder to shoulder with previously described dendritic calcium spikes (Fig. 1B₂, *Ca²⁺ Spike*) and backpropagating sodium action potentials that sweep through basal, oblique and apical tuft branches of cortical pyramidal neurons (not shown in this figure)(Antic, 2003; Canepari et al., 2007; Nevian et al., 2007; Zhou et al., 2007; Zhou et al., 2008; Acker and Antic, 2009; Holthoff et al., 2010).

Duration

The duration of a pure NMDA spike obtained with near-threshold stimulation intensity is in the order of 50 – 100 ms (Fig. 1B₁)(Schiller et al., 2000; Polsky et al., 2004; Rhodes, 2006; Major et al., 2008). In normal physiological saline (saline without blockers of dendritic conductances), and upon suprathreshold stimuli, the durations of dendritic glutamate-evoked plateau potentials are somewhat greater than the duration of a pure NMDA spike (Fig. 1A), presumably due to the activation of voltage-gated calcium channels. One interesting aspect of dendritic plateau potentials is that their durations depend on the intensity of glutamatergic stimulation. This is in striking contrast to sodium action potentials (Fig. 1B₃, *AP*), which are largely insensitive to stimulus properties.

In a typical experiment designed to study properties of NMDA-dependent dendritic spikes, a series of consecutive glutamate pulses of gradually increasing intensities is applied on a dendritic segment (Fig. 2A), and both the dendritic and somatic voltage transients are recorded simultaneously (Fig. 2B). The dendritic voltage imaging performed in the presence of sodium channel blocker TTX, have shown that the local dendritic voltage response to suprathreshold glutamate levels was quickly saturated. The amplitude of the dendritic spike did not increase with further increase in stimulus intensity (Fig. 2B, Intensities 1 – 4). This could be explained by the fact that the maximal amplitude of the NMDA-dependent spike is determined by the glutamate reversal potential, which is around 0 mV (Jonas and Sakmann,

1992). We think that dendritic plateau potentials approach the glutamate reversal potential and thus saturate (Fig. 2B, Intensities 1 – 4).

While the peak amplitude of the dendritic membrane potential transient remained the same throughout the wide range of suprathreshold stimulus intensities, the duration of the dendritic plateau potential, on the other hand, grew linearly with the stimulus intensity (Fig. 2B, sweeps 1–4). Identical results were obtained in normal physiological saline, where glutamate-evoked sustained depolarizations are accompanied by action potential generation in the cell body (Fig. 2D). Here again, a gradual increase in glutamate stimulation intensity causes a gradual increase in the duration of the somatic slow component measured at half-amplitude (Fig. 2E). That means that despite amplitude saturation of their voltage response the thin dendritic branches have not lost the ability to code the intensity of glutamatergic excitatory input. In the suprathreshold range of input intensities (Fig. 2B, Intensities 1 – 4) the coding mechanism is based not on the amplitude of the slow depolarization (Fig. 2E, amp.), but rather on its duration (Fig. 2E), see also Nevian et al., (2007); their figure 7B,C. An *in vitro* study has indicated that when strong excitatory inputs converge in middle segments of basilar branches then all aspects of the ensuing dendritic plateau-potentials (timing, amplitude, duration) are mirrored in the cell body (Milojkovic et al., 2005a). Consequently, the dendritic input-output function becomes essentially the neuronal input-output profile. A flexible duration of dendritic plateau potentials (Fig. 2B) is the basis for linear correlation between the intensity of glutamatergic stimulation received in the basilar dendrite and the number of somatic action potentials per stimulation event (not shown here but see Milojkovic et al., 2005a; their figure 2G).

Propagation of dendritic NMDA spikes

Calcium imaging and computational modeling suggested that NMDA spikes are highly localized events that envelop just one small dendritic segment, 10 – 40 μm in length (Schiller et al., 2000; Rhodes, 2006; Larkum and Nevian, 2008; Major et al., 2008). Simultaneous multi-site recordings have determined that glutamate-evoked dendritic plateau potentials propagate from their initiation site (Fig. 3, *Glut. Input*) in two directions, distally towards the end of the dendritic branch (Fig. 3, grey arrows) and proximally towards the cell body (Fig. 3, white arrows). The degree of amplitude attenuation is significantly smaller in the direction from the initiation site (distally) towards the dendritic tip (Milojkovic et al., 2007; their figure 4A). This is because the sealed end of the basal dendrite has high impedance and it can be easily charged by a spreading plateau potential (Fig. 3, *Dend. Tip*). In contrast, in the orthograde (proximal) direction (Fig. 3, white arrows) the rate of amplitude attenuation is much faster. Due to an impedance mismatch between the thin basal dendrite and large soma the plateau potentials are drastically reduced in amplitude as they approach the cell body (Milojkovic et al., 2004; their figure 7). Nevertheless, suprathreshold glutamatergic stimulations delivered in the middle and proximal segments of a single basal dendrite (Fig. 3, rectangle) often produced 10 – 20 mV sustained depolarizations of the cell body (Figs. 2 and 3). The timing and duration of the somatic depolarization phase (Fig. 3, *Soma*) are in strict relationship with the time course of the dendritic plateau potential at the initiation site (Fig. 3, *Glut. Input*). Several milliseconds after the onset of a dendritic plateau potential the somatic membrane shifts into the sustained depolarized state (Fig. 2C). The cell body

remains in a depolarized state for the whole duration of the dendritic plateau phase, and it returns to resting immediately after the collapse of the dendritic plateau potential (Fig. 2C) (Milojkovic et al., 2005a). In other words, the onset of the somatic plateau depolarization is a consequence of the onset of the dendritic plateau spike, and, by the same token, the breakdown of the somatic depolarized state occurs as a consequence of the collapse of the glutamate-evoked dendritic plateau potential (Fig. 2C). Upon strong glutamatergic stimulations one can often detect a sharp inflection on the rising phase of the somatic plateau potential (Fig. 2C, arrow), which is quite prominent in the first derivative of the voltage waveform (Acker and Antic, 2009). It is now firmly established that the early inflection on the somatic waveform is caused by a sodium spikelet firing at the dendritic initiation site (Ariav et al., 2003; Milojkovic et al., 2005b; Nevian et al., 2007; Larkum and Nevian, 2008; Losonczy et al., 2008; Remy et al., 2009).

Functional Significance of NMDA Spikes

Neuronal UP state

During sleep, and sometimes even in awake state, cortical pyramidal neurons alternate between intervals of strong activity (UP states) and intervals of almost complete silence (Down states) (Cowan and Wilson, 1994; Timofeev et al., 2000; Petersen et al., 2003). The UP states are based solely on patterned synaptic (glutamatergic) excitation (Steriade et al., 1993; Cowan and Wilson, 1994; Lampl et al., 1999; Shu et al., 2003). Because UP states spread consistently and rapidly throughout the entire cortex at a typical frequency of ~1 Hz, and seem to engage every cortical neuron in the way (Volgushev et al., 2006), they are poised to generate synchronous activity of afferent axon terminals impinging onto an individual cell. The density of glutamatergic afferents is highest in middle segments of thin branches (Larkman, 1991; Benavides-Piccione et al., 2002). Therefore, a cortical UP state may trigger a sudden surge in the concentration of extracellular glutamate around thin dendrites of cortical pyramidal neurons. Transient increases in both exogenous and synaptic glutamate in basal and oblique dendrites have been shown to elicit local regenerative dendritic potentials (Schiller et al., 2000; Oakley et al., 2001a; Wei et al., 2001; Polsky et al., 2004; Gordon et al., 2006; Nevian et al., 2007) and therefore dendritic NMDA-dependent spikes and plateau potentials may occur during cortical UP states (discussed in (Antic et al., 2007)).

In acute brain slice preparations glutamate-evoked dendritic plateau potentials cause sustained depolarizations of the pyramidal cell somata (Fig. 2B, D). These sustained somatic depolarizations resemble cortical UP states in respect to: (1) amplitude of the slow component; (2) duration of the depolarized state; (3) irregular firing of accompanying action potentials; (4) fast onset; and (5) abrupt collapse at the end of the plateau depolarization (Fig. 2H) (Milojkovic et al., 2004); their figure 1). This is not to say that NMDA-dependent plateau potentials are causing cortical UP states. Quite the opposite, we are suggesting that cortical UP states, via a massive increase in extracellular glutamate concentration, create favorable conditions for initiation of dendritic plateau potentials. The all-or-none nature of dendritic plateau potentials, combined with neuronal “restorative” (repolarizing) conductances (see (Wilson, 2008)), may explain the constant and uniform voltage

waveforms (amplitude and duration) regularly observed in individual cortical pyramidal neurons *in vivo* during the consecutive cortical UP states (Cowan and Wilson, 1994; Branchereau et al., 1996; Lewis and O'Donnell, 2000; Timofeev et al., 2000; Petersen et al., 2003). So far, there has not been an adequate explanation for the observed uniformity of amplitudes in consecutive cortical UP states. A perfectly balanced excitatory/inhibitory activity is one possibility (Shu et al., 2003; Waters and Helmchen, 2006). Here we propose that saturation of the dendritic voltage response during the NMDA-dependent plateau potential (Fig. 2B–F) may also produce consecutive UP states of uniform amplitudes (Fig. 2H). In summary, through their potential role in cortical UP states, the NMDA-dependent dendritic plateau potentials could be of importance for arousal of attention, proper cognition and working memory in awake (Funahashi et al., 1989; Major and Tank, 2004; Durstewitz and Seamans, 2006; Haider and McCormick, 2009), as well as for memory consolidation in asleep mammals (Stickgold et al., 2000; Sirota et al., 2003; Rasch et al., 2007), reviewed in (Diekelmann and Born, 2010).

Independent multisite subcellular integration

Calcium spikes that initiate in the apical trunk (Fig. 1B₂) and sodium action potentials that initiate in the axon initial segment (Fig. 1B₃) both propagate several hundred micrometers to invade relatively large sections of the axo-dendritic arbor, acting as global broadcasting signals (Stuart et al., 1997). In contrast to calcium spikes or action potentials, the NMDA spikes in thin dendritic branches (basal, oblique and tuft) can be truly local and spatially restricted phenomena involving only one sister branch in the entire dendritic tree (Schiller et al., 2000; Oakley et al., 2001b; Wei et al., 2001; Holthoff et al., 2004; Milojkovic et al., 2004; Milojkovic et al., 2005b; Major et al., 2008; Larkum et al., 2009). By being both localized and regenerative, NMDA spikes could contribute uniquely to processes of synaptic integration and synaptic plasticity (Gordon et al., 2006), reviewed in (Holthoff et al., 2006). Glutamatergic inputs clustered over approximately 20 – 50 micrometers of dendritic length can elicit local NMDA spike/plateau potentials in terminal dendrites of cortical pyramidal neurons, inspiring the notion that a single terminal dendrite can function as a decision-making computational subunit (Major et al., 2008). This notion has been explored in biophysically detailed compartmental models, and those numerical studies concluded that active dendrites of a single neuron can perform the multiple independent subunit computations and enrich the computational power and repertoire of cortical pyramidal cells (Archie and Mel, 2000; Poirazi et al., 2003; Gollo et al., 2009). Multiple-subunit computations combined with long durations of the dendritic glutamate-evoked regenerative potentials (Milojkovic et al., 2004; Milojkovic et al., 2005a) amply support the neuronal persistent activity (Goldman et al., 2003; Major and Tank, 2004; Morita, 2008).

Early in the “decade of the NMDA spike” it has been proposed that functionally related synaptic contacts are spatially clustered on particular dendritic branches of cortical pyramidal neurons. Namely, in response to every-day life experience, synaptic afferents that carry similar informational content would tend to aggregate in a restricted part of the dendritic tree, presumably on the same dendritic branch (Poirazi and Mel, 2001). Clustering of synaptic inputs in space (and time) improves the chances for reaching the dendritic threshold for firing a regenerative (amplified) response and provides the opportunity for

Author Manuscript
Author Manuscript
Author Manuscript
Author Manuscript

faster and more frequent cooperation among synaptic contacts involved in the same computational task. This intriguing concept, if true, may have profound implications for cortical information processing. Instead of thousands of synaptic inputs, the pyramidal cell needs only a 'correct' set of <50 active synaptic contacts (Gasparini et al., 2004; Larkum et al., 2009) to trigger a regenerative dendritic response (e.g. NMDA/plateau potential). Inside the dendritic segment that is experiencing a clustered glutamatergic input sufficient to trigger an NMDA spike, there is a massive calcium influx (calcium plateau, (Milojkovic et al., 2007; Major et al., 2008; Takahashi and Magee, 2009)), which could serve a multitude of calcium-dependent cellular processes, including the short-term and long-term synaptic plasticities (Nimchinsky et al., 2002; Zucker and Regehr, 2002; Holthoff et al., 2006; Lau et al., 2009). Owing to its prolonged plateau phase the glutamate-evoked dendritic NMDA/plateau potentials may enable dendrites to retain information for hundreds of milliseconds, which potentially extends the time window for binding of fragmented information, such as multiple modalities of a sensory stimulus (Nakamura et al., 1992; Wilson et al., 1993). Each individual feature of a perceptual event, regardless of whether the event was based on external (sensory perception) or internal presentation (recall of memory), is coded by an assembly of cortical neurons. The neural assemblies, defined in terms of time-resolved correlated neural firing patterns, are the basic functional units in cortical information processing (Singer, 2001; Wolters and Raffone, 2008). During the DOWN state cortical pyramidal neurons are electrically quiescent, as documented in several species and several cortical areas, including the visual, somatosensory and prefrontal cortices (Cowan and Wilson, 1994; Lampl et al., 1999; Lewis and O'Donnell, 2000; Timofeev et al., 2000; Petersen et al., 2003; Volgushev et al., 2006). Dendritic NMDA spikes can bring cortical pyramidal neurons to a sustained depolarized state - UP state (Fig. 2H). A transition from a DOWN to UP state dramatically increases the AP firing probability and therefore provides a basis for synchronization with an active assembly of neurons (group of neurons already in an UP state performing a computational task). Furthermore, the glutamate-evoked plateau depolarization is a special form of a "high-conductance state", which causes the shortening of the membrane time constant, which, in turn, favors finer temporal discrimination of distant synaptic inputs (Bernander et al., 1991). Computer simulations predict that cortical neurons in these high-conductance states can resolve higher frequency inputs and therefore efficiently participate in temporal binding (Shadlen and Movshon, 1999; Shelley et al., 2002; Destexhe et al., 2003). In summary, two features of dendritic glutamate-evoked plateau potentials may serve in the temporal binding processes of the CNS: (i) Maintenance of neuronal sustained depolarization - UP state (Fig. 2C); and (ii) Shortening of the neuronal membrane time constant.

Author Manuscript

The impact of individual synapses on the generation of the neuronal output (axonal AP) is weak due to the biophysical constraints imposed by the morphology and membrane properties of the dendritic tree (Gulledge et al., 2005; Spruston, 2008). That is, distal synaptic inputs are severely attenuated as they travel from their site of origin (e.g. terminal tuft branch) through the entire length of the apical dendrite, then through soma, to finally reach the AP initiation zone in the initial segment of axon. Spatio-temporal clustering of glutamatergic inputs and the ensuing generation of dendritic spikes can dramatically increase the impact of distal synapses on the AP initiation process and neuronal AP burst

firing (Fig. 2D). Neurons equipped with dendrites capable of firing NMDA spikes (active dendrites) can exhibit a greater specificity of spiking responses, and perform a greater number of transformations of synaptic input into AP output, that would otherwise require more than one neuron with passive dendrites (discussed in (Larkum and Nevian, 2008)). Apart from the “classic” synaptic integration process (summation of electrical events) the dendritic NMDA spikes may also serve to couple synaptic inputs to intracellular signaling cascades. For example, strong activation of NMDA receptor channels in oblique dendrites synergistically enhances metabotropic glutamate receptor-mediated regenerative Ca^{2+} release from internal stores (Nakamura et al., 2002). The large amplitude and long duration of the intracellular Ca^{2+} increases due to Ca^{2+} release from stores is thought to affect downstream signaling mechanisms in pyramidal neurons (Ross et al., 2005). The benefits of synaptic clustering are thus multiple, and they all seem to depend on the ability of dendrites to generate an NMDA spike.

Concluding remarks

Thin dendrites of cortical pyramidal neurons (basal, oblique and tuft dendrites) are crowded with glutamatergic synaptic contacts. Each glutamatergic contact has its own anatomical substrate in the form of a dendritic spine. Synchronous activation of 10 – 50 neighboring glutamatergic inputs (spines) triggers a local regenerative potential, NMDA spike/plateau, which is characterized with significant local amplitude (40–50 mV) and duration (up to several hundred milliseconds). Depending on the path distance from the cell body these glutamate-evoked NMDA-dependent plateau potentials can either bring the cell body into a sustained depolarization (e.g. proximal segments of basal dendrites), or maintain an isolated dendritic segment in a sustained depolarized state (e.g. the distal part of a long basal branch, or an apical tuft dendrite). Regardless of the thin-dendrite subtype (basal, oblique or tuft), at each dendritic initiation site an NMDA spike creates favorable conditions for causal interactions of active synaptic inputs, including the spatial or temporal binding of information, as well as, processes of short-term and long-term synaptic modifications (e.g. LTP or LTD). Local dendritic NMDA spikes comprise the cellular substrate for multisite independent subunit computations that enrich the computational power and repertoire of cortical pyramidal cells, and are likely to play significant roles in cortical information processing in awake animals (spatio-temporal binding; expansion of the dynamic range; working memory) and during sleep (neuronal UP states, consolidation of memories).

Acknowledgments

This work was supported by NIH grant MH063503 and NARSAD Young Investigator Award.

References

- Acker CD, Antic SD. Quantitative assessment of the distributions of membrane conductances involved in action potential backpropagation along Basal dendrites. *J Neurophysiol.* 2009; 101:1524–1541. [PubMed: 19118105]
- Antic SD. Action potentials in basal and oblique dendrites of rat neocortical pyramidal neurons. *J Physiol.* 2003; 550:35–50. [PubMed: 12730348]
- Antic, SD., Acker, CD., Zhou, WL., Moore, AR., Milojkovic, BA. The Role of Dendrites in the Maintenance of the UP State. In: Timofeev, I., editor. *Mechanisms of spontaneous active states in*

- the neocortex, Mechanisms of spontaneous active states in the neocortex Edition. Research Signpost; Kerala, India: 2007. p. 45-72.
- Archie KA, Mel BW. A model for intradendritic computation of binocular disparity. *Nature Neuroscience*. 2000; 3:54–63. [PubMed: 10607395]
- Ariav G, Polsky A, Schiller J. Submillisecond precision of the input-output transformation function mediated by fast sodium dendritic spikes in basal dendrites of CA1 pyramidal neurons. *J Neurosci*. 2003; 23:7750–7758. [PubMed: 12944503]
- Benavides-Piccione R, Ballesteros-Yanez I, DeFelipe J, Yuste R. Cortical area and species differences in dendritic spine morphology. *J Neurocytol*. 2002; 31:337–346. [PubMed: 12815251]
- Bernander O, Douglas RJ, Martin KA, Koch C. Synaptic background activity influences spatiotemporal integration in single pyramidal cells. *Proc Natl Acad Sci U S A*. 1991; 88:11569–11573. [PubMed: 1763072]
- Branchereau P, Van Bockstaele EJ, Chan J, Pickel VM. Pyramidal neurons in rat prefrontal cortex show a complex synaptic response to single electrical stimulation of the locus coeruleus region: evidence for antidromic activation and GABAergic inhibition using in vivo intracellular recording and electron microscopy. *Synapse*. 1996; 22:313–331. [PubMed: 8867026]
- Cai X, Liang CW, Muralidharan S, Kao JP, Tang CM, Thompson SM. Unique roles of SK and Kv4.2 potassium channels in dendritic integration. *Neuron*. 2004; 44:351–364. [PubMed: 15473972]
- Cai X, Wei DS, Gallagher SE, Bagal A, Mei YA, Kao JP, Thompson SM, Tang CM. Hyperexcitability of distal dendrites in hippocampal pyramidal cells after chronic partial deafferentation. *J Neurosci*. 2007; 27:59–68. [PubMed: 17202472]
- Callaway EM, Katz LC. Photostimulation using caged glutamate reveals functional circuitry in living brain slices. *Proc Natl Acad Sci U S A*. 1993; 90:7661–7665. [PubMed: 7689225]
- Canepari M, Djuricic M, Zecevic D. Dendritic signals from the rat hippocampal CA1 pyramidal neurons during coincident pre- and post-synaptic activity: a combined voltage- and calcium-imaging study. *J Physiol*. 2007; 580:463–484. [PubMed: 17272348]
- Carter AG, Sabatini BL. State-dependent calcium signaling in dendritic spines of striatal medium spiny neurons. *Neuron*. 2004; 44:483–493. [PubMed: 15504328]
- Cossart R, Hirsch JC, Cannon RC, Dinoncourt C, Wheal HV, Ben-Ari Y, Esclapez M, Bernard C. Distribution of spontaneous currents along the somato-dendritic axis of rat hippocampal CA1 pyramidal neurons. *Neuroscience*. 2000; 99:593–603. [PubMed: 10974423]
- Cowan RL, Wilson CJ. Spontaneous firing patterns and axonal projections of single corticostriatal neurons in the rat medial agranular cortex. *Journal of Neurophysiology*. 1994; 71:17–32. [PubMed: 8158226]
- Destexhe A, Rudolph M, Pare D. The high-conductance state of neocortical neurons in vivo. *Nat Rev Neurosci*. 2003; 4:739–751. [PubMed: 12951566]
- Diekelmann S, Born J. The memory function of sleep. *Nat Rev Neurosci*. 2010; 11:114–126. [PubMed: 20046194]
- Dodt HU. Infrared-interference videomicroscopy of living brain slices. *Adv Exp Med Biol*. 1993; 333:245–249. [PubMed: 8362664]
- Durstewitz D, Seamans JK. Beyond bistability: biophysics and temporal dynamics of working memory. *Neuroscience*. 2006; 139:119–133. [PubMed: 16326020]
- Emptage N, Bliss TV, Fine A. Single synaptic events evoke NMDA receptor-mediated release of calcium from internal stores in hippocampal dendritic spines. *Neuron*. 1999; 22:115–124. [PubMed: 10027294]
- Funahashi S, Bruce CJ, Goldman-Rakic PS. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *Journal of Neurophysiology*. 1989; 61:331–349. [PubMed: 2918358]
- Gasparini S, Migliore M, Magee JC. On the initiation and propagation of dendritic spikes in CA1 pyramidal neurons. *J Neurosci*. 2004; 24:11046–11056. [PubMed: 15590921]
- Golding NL, Staff NP, Spruston N. Dendritic spikes as a mechanism for cooperative long-term potentiation. *Nature*. 2002; 418:326–331. [PubMed: 12124625]

- Goldman MS, Levine JH, Major G, Tank DW, Seung HS. Robust Persistent Neural Activity in a Model Integrator with Multiple Hysteretic Dendrites per Neuron. *Cereb Cortex*. 2003; 13:1185–1195. [PubMed: 14576210]
- Gollo LL, Kinouchi O, Copelli M. Active dendrites enhance neuronal dynamic range. *PLoS Comput Biol*. 2009; 5:e1000402. [PubMed: 19521531]
- Gordon U, Polsky A, Schiller J. Plasticity compartments in basal dendrites of neocortical pyramidal neurons. *J Neurosci*. 2006; 26:12717–12726. [PubMed: 17151275]
- Gulledge AT, Kampa BM, Stuart GJ. Synaptic integration in dendritic trees. *J Neurobiol*. 2005; 64:75–90. [PubMed: 15884003]
- Haider B, McCormick DA. Rapid neocortical dynamics: cellular and network mechanisms. *Neuron*. 2009; 62:171–189. [PubMed: 19409263]
- Helmchen F, Svoboda K, Denk W, Tank DW. In vivo dendritic calcium dynamics in deep-layer cortical pyramidal neurons. *Nature Neuroscience*. 1999; 2:989–996. [PubMed: 10526338]
- Herreras O. Propagating dendritic action potential mediates synaptic transmission in CA1 pyramidal cells in situ. *J Neurophysiol*. 1990; 64:1429–1441. [PubMed: 2178183]
- Holthoff K, Kovalchuk Y, Konnerth A. Dendritic spikes and activity-dependent synaptic plasticity. *Cell Tissue Res*. 2006; 326:369–377. [PubMed: 16816965]
- Holthoff K, Kovalchuk Y, Yuste R, Konnerth A. Single-shock LTD by local dendritic spikes in pyramidal neurons of mouse visual cortex. *J Physiol*. 2004; 560:27–36. [PubMed: 15319420]
- Holthoff KP, Zecevic DP, Konnerth A. Rapid time-course of action potentials in spines and remote dendrites of mouse visual cortex neurons. *J Physiol*. 2010
- Jonas P, Sakmann B. Glutamate receptor channels in isolated patches from CA1 and CA3 pyramidal cells of rat hippocampal slices. *J Physiol*. 1992; 455:143–171. [PubMed: 1282929]
- Kamondi A, Acsady L, Buzsaki G. Center for M, Behavioral Neuroscience RTSUoNJJNJUSA. Dendritic spikes are enhanced by cooperative network activity in the intact hippocampus. *Journal of Neuroscience*. 1998; 18:3919–3928. [PubMed: 9570819]
- Kim HG, Connors BW. Apical dendrites of the neocortex: correlation between sodium- and calcium-dependent spiking and pyramidal cell morphology. *J Neurosci*. 1993; 13:5301–5311. [PubMed: 8254376]
- Lampl I, Reichova I, Ferster D. Synchronous membrane potential fluctuations in neurons of the cat visual cortex. *Neuron*. 1999; 22:361–374. [PubMed: 10069341]
- Larkman AU. Dendritic morphology of pyramidal neurones of the visual cortex of the rat: III. Spine distributions. *Journal of Comparative Neurology*. 1991; 306:332–343. [PubMed: 1711059]
- Larkum ME, Nevian T. Synaptic clustering by dendritic signalling mechanisms. *Curr Opin Neurobiol*. 2008; 18:321–331. [PubMed: 18804167]
- Larkum ME, Zhu JJ, Sakmann B. A new cellular mechanism for coupling inputs arriving at different cortical layers. *Nature*. 1999; 398:338–341. [PubMed: 10192334]
- Larkum ME, Zhu JJ, Sakmann B. Dendritic mechanisms underlying the coupling of the dendritic with the axonal action potential initiation zone of adult rat layer 5 pyramidal neurons. *Journal of Physiology*. 2001; 533:447–466. [PubMed: 11389204]
- Larkum ME, Nevian T, Sandler M, Polsky A, Schiller J. Synaptic integration in tuft dendrites of layer 5 pyramidal neurons: a new unifying principle. *Science*. 2009; 325:756–760. [PubMed: 19661433]
- Lau CG, Takeuchi K, Rodenas-Ruano A, Takayasu Y, Murphy J, Bennett MV, Zukin RS. Regulation of NMDA receptor Ca²⁺ signalling and synaptic plasticity. *Biochem Soc Trans*. 2009; 37:1369–1374. [PubMed: 19909278]
- Lewis BL, O'Donnell P. Ventral tegmental area afferents to the prefrontal cortex maintain membrane potential 'up' states in pyramidal neurons via D(1) dopamine receptors. *Cerebral Cortex*. 2000; 10:1168–1175. [PubMed: 11073866]
- Lisman J, Spruston N. Postsynaptic depolarization requirements for LTP and LTD: a critique of spike timing-dependent plasticity. *Nat Neurosci*. 2005; 8:839–841. [PubMed: 16136666]
- Losonczy A, Magee JC. Integrative properties of radial oblique dendrites in hippocampal CA1 pyramidal neurons. *Neuron*. 2006; 50:291–307. [PubMed: 16630839]

- Losonczy A, Makara JK, Magee JC. Compartmentalized dendritic plasticity and input feature storage in neurons. *Nature*. 2008; 452:436–441. [PubMed: 18368112]
- Luscher C, Nicoll RA, Malenka RC, Muller D. Synaptic plasticity and dynamic modulation of the postsynaptic membrane. *Nat Neurosci*. 2000; 3:545–550. [PubMed: 10816309]
- Magee JC, Johnston D. Characterization of single voltage-gated Na⁺ and Ca²⁺ channels in apical dendrites of rat CA1 pyramidal neurons. *Journal of Physiology*. 1995; 487:67–90. [PubMed: 7473260]
- Major G, Tank D. Persistent neural activity: prevalence and mechanisms. *Curr Opin Neurobiol*. 2004; 14:675–684. [PubMed: 15582368]
- Major G, Polsky A, Denk W, Schiller J, Tank DW. Spatiotemporally graded NMDA spike/plateau potentials in basal dendrites of neocortical pyramidal neurons. *J Neurophysiol*. 2008; 99:2584–2601. [PubMed: 18337370]
- Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature*. 2004; 429:761–766. [PubMed: 15190253]
- Mayer ML, Westbrook GL, Guthrie PB. Voltage-dependent block by Mg²⁺ of NMDA responses in spinal cord neurones. *Nature*. 1984; 309:261–263. [PubMed: 6325946]
- Milojkovic BA, Radojicic MS, Antic SD. A strict correlation between dendritic and somatic plateau depolarizations in the rat prefrontal cortex pyramidal neurons. *J Neurosci*. 2005a; 25:3940–3951. [PubMed: 15829646]
- Milojkovic BA, Zhou WL, Antic SD. Voltage and Calcium Transients in Basal Dendrites of the Rat Prefrontal Cortex. *J Physiol*. 2007; 585:447–468. [PubMed: 17932150]
- Milojkovic BA, Radojicic MS, Goldman-Rakic PS, Antic SD. Burst generation in rat pyramidal neurones by regenerative potentials elicited in a restricted part of the basilar dendritic tree. *J Physiol*. 2004; 558:193–211. [PubMed: 15155788]
- Milojkovic BA, Wuskell JP, Loew LM, Antic SD. Initiation of sodium spikelets in basal dendrites of neocortical pyramidal neurons. *J Membr Biol*. 2005b; 208:155–169. [PubMed: 16645744]
- Morita K. Possible role of dendritic compartmentalization in the spatial working memory circuit. *J Neurosci*. 2008; 28:7699–7724. [PubMed: 18650346]
- Nakamura K, Mikami A, Kubota K. Oscillatory neuronal activity related to visual short-term memory in monkey temporal pole. *Neuroreport*. 1992; 3:117–120. [PubMed: 1611028]
- Nakamura T, Lasser-Ross N, Nakamura K, Ross WN. Spatial segregation and interaction of calcium signalling mechanisms in rat hippocampal CA1 pyramidal neurons. *J Physiol*. 2002; 543:465–480. [PubMed: 12205182]
- Nevian T, Larkum ME, Polsky A, Schiller J. Properties of basal dendrites of layer 5 pyramidal neurons: a direct patch-clamp recording study. *Nat Neurosci*. 2007; 10:206–214. [PubMed: 17206140]
- Nimchinsky EA, Sabatini BL, Svoboda K. Structure and function of dendritic spines. *Annu Rev Physiol*. 2002; 64:313–353. [PubMed: 11826272]
- Noguchi J, Matsuzaki M, Ellis-Davies GC, Kasai H. Spine-neck geometry determines NMDA receptor-dependent Ca²⁺ signaling in dendrites. *Neuron*. 2005; 46:609–622. [PubMed: 15944129]
- Nowak L, Bregestovski P, Ascher P, Herbert A, Prochiantz A. Magnesium gates glutamate-activated channels in mouse central neurones. *Nature*. 1984; 307:462–465. [PubMed: 6320006]
- Oakley JC, Schwindt PC, Crill WE. Dendritic calcium spikes in layer 5 pyramidal neurons amplify and limit transmission of ligand-gated dendritic current to soma. *Journal of Neurophysiology*. 2001a; 86:514–527. [PubMed: 11431529]
- Oakley JC, Schwindt PC, Crill WE. Initiation and propagation of regenerative Ca(2+)-dependent potentials in dendrites of layer 5 pyramidal neurons. *Journal of Neurophysiology*. 2001b; 86:503–513. [PubMed: 11431528]
- Petersen CC, Hahn TT, Mehta M, Grinvald A, Sakmann B. Interaction of sensory responses with spontaneous depolarization in layer 2/3 barrel cortex. *Proc Natl Acad Sci U S A*. 2003; 100:13638–13643. [PubMed: 14595013]
- Pettit DL, Wang SS, Gee KR, Augustine GJ. Chemical two-photon uncaging: a novel approach to mapping glutamate receptors. *Neuron*. 1997; 19:465–471. [PubMed: 9331338]

- Poirazi P, Mel BW. Impact of active dendrites and structural plasticity on the memory capacity of neural tissue. *Neuron*. 2001; 29:779–796. [PubMed: 11301036]
- Poirazi P, Brannon T, Mel BW. Pyramidal neuron as two-layer neural network. *Neuron*. 2003; 37:989–999. [PubMed: 12670427]
- Polsky A, Mel BW, Schiller J. Computational subunits in thin dendrites of pyramidal cells. *Nat Neurosci*. 2004; 7:621–627. [PubMed: 15156147]
- Rasch B, Buchel C, Gais S, Born J. Odor cues during slow-wave sleep prompt declarative memory consolidation. *Science*. 2007; 315:1426–1429. [PubMed: 17347444]
- Regehr WG, Tank DW. Calcium concentration dynamics produced by synaptic activation of CA1 hippocampal pyramidal cells. *J Neurosci*. 1992; 12:4202–4223. [PubMed: 1359030]
- Remy S, Spruston N. Dendritic spikes induce single-burst long-term potentiation. *Proc Natl Acad Sci U S A*. 2007; 104:17192–17197. [PubMed: 17940015]
- Remy S, Csicsvari J, Beck H. Activity-dependent control of neuronal output by local and global dendritic spike attenuation. *Neuron*. 2009; 61:906–916. [PubMed: 19323999]
- Rhodes P. The properties and implications of NMDA spikes in neocortical pyramidal cells. *J Neurosci*. 2006; 26:6704–6715. [PubMed: 16793878]
- Ross WN, Nakamura T, Watanabe S, Larkum M, Lasser-Ross N. Synaptically activated Ca^{2+} release from internal stores in CNS neurons. *Cell Mol Neurobiol*. 2005; 25:283–295. [PubMed: 16047542]
- Schiller J, Helmchen F, Sakmann B. Spatial profile of dendritic calcium transients evoked by action potentials in rat neocortical pyramidal neurones. *Journal of Physiology*. 1995; 487:583–600. [PubMed: 8544123]
- Schiller J, Schiller Y, Stuart G, Sakmann B. Calcium action potentials restricted to distal apical dendrites of rat neocortical pyramidal neurons. *Journal of Physiology*. 1997; 505:605–616. [PubMed: 9457639]
- Schiller J, Major G, Koester HJ, Schiller Y. NMDA spikes in basal dendrites of cortical pyramidal neurons. *Nature*. 2000; 404:285–289. [PubMed: 10749211]
- Shadlen MN, Movshon JA. Synchrony unbound: a critical evaluation of the temporal binding hypothesis. *Neuron*. 1999; 24:67–77. 111–125. [PubMed: 10677027]
- Shelley M, McLaughlin D, Shapley R, Wia laard J. States of high conductance in a large-scale model of the visual cortex. *J Comput Neurosci*. 2002; 13:93–109. [PubMed: 12215724]
- Shu YS, Hasenstaub A, McCormick DA. Turning on and off recurrent balanced cortical activity. *Nature*. 2003; 423:288–293. [PubMed: 12748642]
- Singer W. Consciousness and the binding problem. *Ann N Y Acad Sci*. 2001; 929:123–146. [PubMed: 11349422]
- Sirota A, Csicsvari J, Buhl D, Buzsaki G. Communication between neocortex and hippocampus during sleep in rodents. *Proc Natl Acad Sci U S A*. 2003; 100:2065–2069. [PubMed: 12576550]
- Sjostrom PJ, Rancz EA, Roth A, Hausser M. Dendritic excitability and synaptic plasticity. *Physiol Rev*. 2008; 88:769–840. [PubMed: 18391179]
- Sobczyk A, Svoboda K. Activity-Dependent Plasticity of the NMDA-Receptor Fractional Ca^{2+} Current. *Neuron*. 2007; 53:17–24. [PubMed: 17196527]
- Spruston N. Pyramidal neurons: dendritic structure and synaptic integration. *Nat Rev Neurosci*. 2008; 9:206–221. [PubMed: 18270515]
- Spruston N, Schiller Y, Stuart G, Sakmann B. Activity-dependent action potential invasion and calcium influx into hippocampal CA1 dendrites. *Science*. 1995; 268:297–300. [PubMed: 7716524]
- Steriade M, Nunez A, Amzica F. Intracellular analysis of relations between the slow (< 1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *Journal of Neuroscience*. 1993; 13:3266–3283. [PubMed: 8340807]
- Stickgold R, James L, Hobson JA. Visual discrimination learning requires sleep after training. *Nature Neuroscience*. 2000; 3:1237–1238. [PubMed: 11100141]
- Stuart G, Spruston N, Sakmann B, Hausser M. Action potential initiation and backpropagation in neurons of the mammalian CNS. *Trends in Neurosciences*. 1997; 20:125–131. [PubMed: 9061867]

- Stuart GJ, Sakmann B. Active propagation of somatic action potentials into neocortical pyramidal cell dendrites. *Nature*. 1994; 367:69–72. [PubMed: 8107777]
- Suzuki T, Kodama S, Hoshino C, Izumi T, Miyakawa H. A plateau potential mediated by the activation of extrasynaptic NMDA receptors in rat hippocampal CA1 pyramidal neurons. *Eur J Neurosci*. 2008; 28:521–534. [PubMed: 18702724]
- Takahashi H, Magee JC. Pathway interactions and synaptic plasticity in the dendritic tuft regions of CA1 pyramidal neurons. *Neuron*. 2009; 62:102–111. [PubMed: 19376070]
- Timofeev I, Grenier F, Bazhenov M, Sejnowski TJ, Steriade M. Origin of slow cortical oscillations in deafferented cortical slabs. *Cerebral Cortex*. 2000; 10:1185–1199. [PubMed: 11073868]
- Volgushev M, Chauvette S, Mukovski M, Timofeev I. Precise long-range synchronization of activity and silence in neocortical neurons during slow-wave sleep. *J Neurosci*. 2006; 26:5665–5672. [PubMed: 16723523]
- Waters J, Helmchen F. Background synaptic activity is sparse in neocortex. *J Neurosci*. 2006; 26:8267–8277. [PubMed: 16899721]
- Waters J, Schaefer A, Sakmann B. Backpropagating action potentials in neurones: measurement, mechanisms and potential functions. *Prog Biophys Mol Biol*. 2005; 87:145–170. [PubMed: 15471594]
- Wei DS, Mei YA, Bagal A, Kao JP, Thompson SM, Tang CM. Compartmentalized and binary behavior of terminal dendrites in hippocampal pyramidal neurons. *Science*. 2001; 293:2272–2275. [PubMed: 11567143]
- Wilson CJ. Up and down states. *Scholarpedia*. 2008; 3:1410. revision #68992.
- Wilson FA, Scalaidhe SP, Goldman-Rakic PS. Dissociation of object and spatial processing domains in primate prefrontal cortex. *Science*. 1993; 260:1955–1958. [PubMed: 8316836]
- Wolters G, Raffone A. Coherence and recurrency: maintenance, control and integration in working memory. *Cogn Process*. 2008; 9:1–17. [PubMed: 17901994]
- Yang G, Pan F, Gan WB. Stably maintained dendritic spines are associated with lifelong memories. *Nature*. 2009; 462:920–924. [PubMed: 19946265]
- Yuste R, Bonhoeffer T. Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annual Review of Neuroscience*. 2001; 24:1071–1089.
- Yuste R, Gutnick MJ, Saar D, Delaney KR, Tank DW. Ca²⁺ accumulations in dendrites of neocortical pyramidal neurons: an apical band and evidence for two functional compartments. *Neuron*. 1994; 13:23–43. [PubMed: 8043278]
- Zhou WL, Yan P, Wuskell JP, Loew LM, Antic SD. Intracellular long wavelength voltage-sensitive dyes for studying the dynamics of action potentials in axons and thin dendrites. *Journal of Neuroscience Methods*. 2007; 164:225–239. [PubMed: 17560661]
- Zhou WL, Yan P, Wuskell JP, Loew LM, Antic SD. Dynamics of action potential backpropagation in basal dendrites of prefrontal cortical pyramidal neurons. *Eur J Neurosci*. 2008; 27:1–14. [PubMed: 18093178]
- Zucker RS, Regehr WG. Short-term synaptic plasticity. *Annu Rev Physiol*. 2002; 64:355–405. [PubMed: 11826273]




Figure 1. Each Class of Neurites (Basal, Oblique, Tuft, Apical Trunk, and Axon) Supports a Characteristic Regenerative Potential (Spike)

(A) In thin dendrites of pyramidal neurons (basal, oblique and tuft) weak glutamatergic inputs produce EPSP-like depolarizations (“*Subthreshold EPSP*”). Stronger (suprathreshold) inputs regularly trigger dendritic “*Plateau Potentials*”. Plateau potentials are characterized by a rapid onset, initial spikelet (Na⁺ Spikelet), plateau phase, and an abrupt collapse at the end of the plateau phase. A glutamate-evoked plateau potential is the product of several dendritic conductances. The major ionic contributor is regenerative NMDA receptor current. Block of sodium and calcium channels with TTX and Cd²⁺, respectively, reveals a pure “*NMDA Spike*” in the dendrite (Schiller et al., 2000). (B) The waveforms of three regenerative potentials (spikes) are shown side by side on the same scale (B₁-B₃). The neuronal compartment which serves as the spike initiation site is colored black below each spike waveform (Initiation Sites). The principle compartments of cortical pyramidal cells are: *Soma, Axon, Apical Trunk, Basal, Oblique and Tuft dendrites*. (B1)

NMDA spikes last approximately 50–100 ms and they can be initiated in the apical tuft, apical oblique and basal dendrites. **(B2)** Calcium spikes have slightly greater amplitudes and durations than NMDA spikes. Calcium spikes are often found in the calcium spike initiation zone on the apical trunk. **(B3)** Action potential (AP) initiates in the axon initial segment. Once initiated in the axon the AP propagates back into the dendritic tree (Stuart et al., 1997). The voltage waveforms of a backpropagating action potential at dendritic sites are not shown in this figure (see (Antic, 2003; Nevian et al., 2007; Zhou et al., 2008; Holthoff et al., 2010)). Dendritic-spike initiation sites (B₁-B₃) are based on Larkum et al., 2009.




Figure 2. Strict correlation between dendritic plateau potential and somatic sustained depolarization

(A) Recording configuration showing glutamate iontophoresis electrode positioned in the middle segment of a basal dendrite. (B) Voltage waveforms are recorded at the stimulation site optically (dend.) and in the cell body electrically (patch) as indicated in “A”. Intensity of glutamate current pulse was increased in 4 equal increments (1 – 4). In this and the following panels the timing of glutamate pulse is marked by a vertical tick below the recording trace. In the presence of TTX the somatic membrane potential waveform closely follows that in the dendrite. While the amplitudes of both dendritic and somatic responses are saturated, their durations increase gradually in consecutive sweeps. There is a positive correlation between the stimulus intensity and duration of the electrical event (Milojkovic et al., 2005a). (C) In normal physiological saline the glutamate-evoked dendritic plateau potential is often preceded by a fast sodium spikelet. At this same moment when the spikelet fired in the dendrite the cell body showed fast inflection (arrow)(Milojkovic et al., 2005b; Nevian et al., 2007; Remy et al., 2009). (D) Somatic voltage waveforms generated by a sequence of incrementally increasing glutamate stimuli delivered on a basal dendrite, as shown in “A”. (E) Slow component of the somatic depolarization can be described in terms of amplitude (amp.) and half-width (duration, dur.). The amplitude and duration of the slow component (somatic) are plotted against stimulus intensity in F and G, respectively. (F) The

amplitude of somatic slow depolarization exhibits an all-or-none highly non-linear behavior (Binary), while duration exhibits steady growth with stimulus intensity (Linear “G”). (H) Cortical UP state-like somatic depolarizations were triggered in acute brain slice by a sequence of three identical glutamate puffs (glut.) delivered on a single basilar branch (Milojkovic et al., 2004).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Two-Directional Propagation of Glutamate-evoked Dendritic Plateau Potentials

The dendritic plateau potential (Plateau Potential) propagates from its initiation site (*Glut. Input*) in two directions: Distally towards the end of the basal dendrite (*Dend. Tip*); and proximally towards the cell body (*Soma*). In the proximal direction (white arrows) the rate of amplitude decline is much faster than in the distal direction (grey arrows). Glutamate-evoked dendritic plateau potentials initiated in the mid section of a typical basal dendrite (rectangle) regularly cause 10 – 20 mV sustained (> 100 ms) depolarizations of the cell body (Milojkovic et al., 2004; Milojkovic et al., 2005a). Note that at the glutamate stimulation site (*Glut. Input*) the dendritic plateau potential is followed by a small post-plateau depolarization (*Post-plateau Potential*). Both, the plateau and post-plateau potential, support calcium influxes, which when combined constitute the glutamate-evoked dendritic “calcium plateau”, not shown, but see (Milojkovic et al., 2007).

Table 1

Two Decades of Dendritic Spikes in Cortical Pyramidal Neurons:

Decade	1990 – 1999 Decade of the thick apical dendrite.	2000 – 2009 Decade of <i>thin branches</i> .
Dendrite Class	Apical Trunk	Basal Oblique Tuft
Local Regenerative Potential	Calcium Spike	Sodium Spikelet NMDA Spike Plateau Potential

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript