

SHORT COMMUNICATION



Regulatory interplay between *LEAFY*, *APETALA1/CAULIFLOWER* and *TERMINAL FLOWER1*: New insights into an old relationship

Antonio Serrano-Mislata^{a,†}, Kevin Goslin^{b,c}, Beibei Zheng^b, Liina Rae^b, Frank Wellmer^{b,†}, Emmanuelle Graciet^{c,†}, and Francisco Madueño^{a,†}

^aInstituto de Biología Molecular y Celular de Plantas, CSIC-UPV, Valencia, Spain; ^bSmurfit Institute of Genetics, Trinity College Dublin, Dublin, Ireland; ^cDepartment of Biology, National University of Ireland Maynooth, Maynooth, Ireland

ABSTRACT

The gene regulatory network comprised of *LEAFY* (*LFY*), *APETALA1* (*AP1*), the *AP1* paralog *CAULIFLOWER* (*CAL*), and *TERMINAL FLOWER1* (*TFL1*) is a major determinant of the flowering process in *Arabidopsis thaliana*. *TFL1* activity in the shoot apical meristem provides inflorescence identity while the transcription factors *LFY* and *AP1/CAL* confer floral identity to emerging floral primordia. It has been thought that *LFY* and *AP1/CAL* control the onset of flowering in part by repressing *TFL1* expression in flowers. However, in the June issue of *Plant Physiology*, we reported that *LFY* and *AP1* act antagonistically in the regulation of several key flowering regulators, including *TFL1*. Specifically, *TFL1* transcription was suppressed by *AP1* but promoted by *LFY*. Here, we present additional evidence for the role of *LFY* as an activator of *TFL1* and propose that this regulatory activity is pivotal for the indeterminate growth of the SAM during the reproductive phase of development.

ARTICLE HISTORY

Received 1 August 2017
Accepted 16 August 2017

KEYWORDS

APETALA1; *Arabidopsis*;
CAULIFLOWER; flowering;
inflorescence development;
LEAFY; meristem identity;
TERMINAL FLOWER1

Angiosperms integrate a multitude of endogenous and environmental signals to determine a time for flowering that ensures reproductive success. Research conducted over the past 25 years, especially in *Arabidopsis thaliana* (*Arabidopsis*), has revealed many of the genes that orchestrate the flowering process.¹⁻³ These include inflorescence meristem identity genes, such as *TERMINAL FLOWER1* (*TFL1*), and floral meristem identity genes, including the transcription factor-coding genes *LEAFY* (*LFY*), *APETALA1* (*AP1*) and the *AP1*-paralog *CAULIFLOWER* (*CAL*). Because these genes together determine when and where flowers will be formed, it has been proposed that differences in their expression patterns or in the functions of the corresponding proteins can explain much of the diversity of inflorescence architectures observed among the angiosperms.⁴

The inflorescence of *Arabidopsis* is characterized by a main axis (shoot) with an indeterminate shoot apical meristem (SAM) that laterally produces flowers. A number of elegant studies showed that *TFL1* and *LFY/AP1* are essential for building the *Arabidopsis* inflorescence. *TFL1* is expressed in the central region of the SAM, providing inflorescence identity and allowing the indeterminate growth of the shoot apex, while *LFY* and *AP1/CAL* are expressed in the flanks of the SAM, providing floral identity to emerging primordia.⁵⁻⁷ In *tfl1* mutants, *LFY/AP1* expression expands into the SAM that, consequently, acquires floral identity and abruptly terminates with the formation of a flower-like structure.⁷⁻⁹ Conversely, in *lfy* and *ap1* mutants, flowers are substituted by shoot-like structures.^{7,10-11} An even more dramatic conversion of flowers into

inflorescence-like meristems is observed in *ap1 cal* double-mutant plants.¹² In recent years, several studies have provided a molecular basis for the antagonism between *TFL1* and *LFY/AP1*: both *LFY* and *AP1* proteins were shown to bind to essential *cis*-regulatory elements in the *TFL1* promoter.¹³⁻¹⁵

LFY and *AP1* regulate floral development in a partially redundant manner and share many target genes.¹³⁻¹⁷ However, it is not known whether *LFY* and *AP1* act together or provide independent inputs to these targets. We addressed this question by analyzing the transcriptional activity of *LFY* in the absence of *AP1/CAL* function.¹⁸ To this end, we used a *p35S:LFY-GR* line introgressed into an *ap1 cal* double-mutant background to determine the gene expression changes caused by *LFY* activation in the inflorescence. We found that *LFY* can regulate some of its known target genes independently of *AP1/CAL* activity. In contrast, other *LFY* targets, including the floral homeotic genes *APETALA3* and *AGAMOUS*, appear to require functional *AP1/CAL*. In agreement with the results of a previous meta-analysis of published data sets,^{18,19} we further found that *LFY* and *AP1/CAL* regulate certain targets antagonistically. These included regulators of floral initiation such as *FLOWERING LOCUS D*, *TEMPRANILLO1*, *APETALA2* and, notably, *TFL1*. *TFL1* was upregulated in response to *LFY-GR* activation but downregulated by *AP1-GR* in *ap1 cal* inflorescences. In agreement with the transcriptional response of *TFL1*, activation of *LFY-GR* in *ap1 cal* plants led to a significant inhibition of flower formation while activation of *AP1-GR*

caused an immediate and synchronized onset of flowering, as previously reported.¹⁷

These results, as well as a set of previous observations, led us to reconsider the nature of the relationship between LFY and TFL1. Firstly, the expression domains of *TFL1* and *LFY* overlap in the inflorescence-like meristems of *ap1 cal* plants.²⁰⁻²¹ Moreover, weak *LFY* expression has been detected in the stem of wild-type inflorescences,²² where *TFL1* is also expressed.²³ The reanalysis of published transcriptomics data sets further showed that activation of LFY-GR in seedlings leads to upregulation of *TFL1* expression.³ Taken together, these results imply that *TFL1* and *LFY* are not necessarily antagonists and that LFY may be able to activate *TFL1*, at least, in the absence of AP1/CAL activity. In line with this idea, LFY was shown to bind to a region approximately 2.8 kilobases (kb) downstream of *TFL1*, which is essential for the maintenance of *TFL1* expression in the inflorescence meristem and, consequently, for SAM indeterminacy.^{14,23} Although *LFY* itself does not appear to be expressed in the SAM, it has been demonstrated that

LFY protein is mobile and can travel to the inflorescence meristem.²⁴ Thus, in addition to its role in flower development, LFY might be needed for indeterminate growth of the SAM. This would not be the first described function of LFY in a shoot meristem, as it has been shown previously that LFY stimulates axillary meristem growth.²⁵

To test the putative role of LFY in SAM identity via activation of *TFL1*, we made use of a previously generated set of *pTFL1:GUS* reporter lines²³ and monitored *TFL1* expression in genetic backgrounds with modified LFY activity (Fig. 1). First, we tested whether the LFY binding sites located in the 3' region of the *TFL1* promoter¹⁴ were essential for the transcriptional response of *TFL1* to LFY. To this end, we analyzed the activity of two *pTFL1:GUS* reporters – one containing the LFY binding sites, the other one lacking them – in plants that express a fusion protein between LFY and the VP16 transcriptional activator under the control of a heat-shock inducible promoter (*pHS:LFY-VP16*).^{26,27} As described in our recent publication,¹⁸ the activity of the *pTFL1:GUS* reporter containing the full




Figure 1. Analysis of *pTFL1:GUS* reporter lines in genetic backgrounds with modified LFY activity. (A) Summary of the experiments performed to test the functional relevance of the LFY binding sites in the 3' region of the *TFL1* promoter. The activity of 2 *pTFL1:GUS* constructs was assayed in *pHS:LFY-VP16* seedlings: one with the full length *TFL1* promoter (2.2 kb of the 5' region plus 4.6 kb of the 3') and one with a truncated version of the promoter lacking the LFY binding region (2.2 kb of the 5' region plus 2.7 kb of the 3', Δ LFY). 'Ectopic GUS (+)' denotes staining in roots, cotyledons and developing leaves after a heat-shock treatment. Growth conditions, heat-shock treatments (incubation for 3 h at 37°C during 3 consecutive days) and GUS staining were conducted as described previously.¹⁸ (B-D) Representative images of *pHS:LFY-VP16* seedlings containing a *pTFL1:GUS* reporter and stained for GUS: (B) reporter activity of the full length *TFL1* promoter in a *pHS:LFY-VP16* seedling grown under control conditions without heat-shock, (C) reporter activity of the full length *TFL1* promoter in a *pHS:LFY-VP16* seedling after heat-shock, (D) reporter activity of the truncated version of the *TFL1* promoter (Δ LFY) in a *pHS:LFY-VP16* seedling after heat-shock. Asterisks point to the position of the SAM. (E-F) Reporter activity of the full length *TFL1* promoter in representative inflorescence apices of wild type (accession Landsberg erecta) (E), *lfy-26* (F) and *ap1-1* (G) plants. The inset in (F) shows the terminal carpelloid structure in *lfy-26* shoots. Arrows in the main panels point to the position of the SAM. GUS staining was conducted as described previously.²³

length promoter of *TFL1* was broadly activated in wild-type seedlings after the heat-shock treatment but restricted to the shoot apex in plants grown under normal conditions (Fig. 1A-C). In contrast, plants carrying a truncated version of *pTFL1:GUS* without the LFY binding sites (Δ LFY in Fig. 1) did not exhibit ectopic GUS activity after induction of LFY-VP16 expression (Fig. 1D). Therefore, the LFY binding sites in the 3' region of the *TFL1* promoter appear to be necessary for the response of *pTFL1:GUS* to LFY-VP16.

Next, we asked whether a loss of LFY function affects *TFL1* transcription in the SAM. To test this, we monitored the activity of *pTFL1:GUS* in the strong *lfy-26* allele.²⁷ Compared to the wild type, the intensity of the GUS signal significantly decreased in the center of *lfy-26* inflorescences (Fig. 1E-F). This result is in agreement with the absence of *TFL1* expression in the SAM of *lfy-7* mutant plants.²⁰ In contrast, *pTFL1:GUS* activity was not apparently affected in the inflorescence apex of *ap1-1* mutants (Fig. 1E, G), which also exhibit impaired floral meristem identity.¹⁰ As described previously,⁷ we observed that the inflorescences of *lfy-26* mutants terminated in carpelloid structures (Fig. 1F inset). This determinacy phenotype of *lfy-26* plants may be caused by the low levels of *TFL1* expression in the inflorescence apex we detected with the *pTFL1:GUS* reporter. Taken together, these results suggest that LFY promotes *TFL1* expression in the SAM to ensure indeterminate growth.

LFY may also activate *TFL1* in flowers, at least under conditions where AP1/CAL are non-functional. In fact, the results presented in our *Plant Physiology* paper suggest that LFY, AP1/CAL and *TFL1* may be part of an incoherent feedback loop²⁸ during early establishment flower development, where LFY activates both *TFL1* and the repressors of *TFL1*, AP1/CAL.¹⁸ This regulatory loop might ensure that flower formation commences only when AP1/CAL levels are sufficiently high to repress *TFL1* expression and to trigger the genetic program required for flower development. A characterization of the protein complexes that regulate inflorescence and floral development may be required to explain the antagonistic activities of LFY and AP1 in the control of *TFL1* and other flowering regulators.

Acknowledgments

A.S.-M. is funded by the European Union's Horizon 2020 Research and Innovation program under Marie Skłodowska-Curie grant agreement No. 746396. F.W. and E.G. are supported by grants from Science Foundation Ireland. F.M. is supported by MINECO and FEDER, grant No. BIO2015-64307-R. B.Z. is supported by the Irish Research Council.

References

- Blázquez MA, Ferrándiz F, Madueño F, Parcy F. How floral meristems are built. *Plant Mol Biol.* 2006;60:855–70. doi:10.1007/s11103-006-0013-z.
- Fornara F, de Montaigu A, Coupland G. SnapShot: Control of flowering in Arabidopsis. *Cell.* 2010;141:550. doi:10.1016/j.cell.2010.04.024.
- Denay G, Chahtane H, Tichtinsky G, Parcy F. A flower is born: an update on Arabidopsis floral meristem formation. *Curr Opin Plant Biol.* 2017;35:15–22. doi:10.1016/j.pbi.2016.09.003.
- Benlloch R, Berbel A, Serrano-Mislata A, Madueño F. Floral initiation and inflorescence architecture: a comparative view. *Ann Bot.* 2007;100:659–76. doi:10.1093/aob/mcm146.
- Bradley D, Ratcliffe O, Vincent C, Carpenter R, Coen E. Inflorescence commitment and architecture in Arabidopsis. *Science* 1997;275:80–3. doi:10.1126/science.275.5296.80.
- Mandel MA, Gustafson-Brown C, Savidge B, Yanofsky MF. Molecular characterization of the Arabidopsis floral homeotic gene APETALA1. *Nature* 1992;360:273–7. doi:10.1038/360273a0.
- Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM. LEAFY controls floral meristem identity in Arabidopsis. *Cell* 1992;69:843–59. doi:10.1016/0092-8674(92)90295-N.
- Bowman JL, Alvarez J, Weigel D, Meyerowitz EM, Smyth DR. Control of flower development in Arabidopsis thaliana by APETALA1 and interacting genes. *Development* 1993;119:721–43.
- Gustafson-Brown C, Savidge B, Yanofsky MF (1994). Regulation of the Arabidopsis floral homeotic gene APETALA1. *Cell.* 1994;76:131–43. doi:10.1016/0092-8674(94)90178-3.
- Irish VF, Sussex IM. Function of the apatela-1 gene during Arabidopsis floral development. *Plant Cell* 1990;2:741–53. doi:10.1105/tpc.2.8.741.
- Huala E, Sussex IM (1992). LEAFY interacts with floral homeotic genes to regulate Arabidopsis floral development. *Plant Cell* 1992;4:901–13. doi:10.1105/tpc.4.8.901.
- Kempin SA, Savidge B, Yanofsky MF. Molecular basis of the cauliflower phenotype in Arabidopsis. *Science.* 1995;267:522–5. doi:10.1126/science.7824951.
- Kaufmann K, Wellmer F, Muiño JM, Ferrier T, Wuest SE, Kumar V, Serrano-Mislata A, Madueño F, Krajewski P, Meyerowitz EM et al. Orchestration of floral initiation by APETALA1. *Science.* 2010;328:85–9. doi:10.1126/science.1185244.
- Moyroud E, Minguet EG, Ott F, Yant L, Posé D, Monniaux M, Blanchet S, Bastien O, Thévenon E, Weigel D et al. Prediction of regulatory interactions from genome sequences using a biophysical model for the Arabidopsis LEAFY transcription factor. *Plant Cell.* 2011;23:1293–306. doi:10.1105/tpc.111.083329.
- Winter CM, Austin RS, Blanvillain-Baufume S, Reback MA, Monniaux M, Wu MF, Sang Y, Yamaguchi A, Yamaguchi N, Parker JE et al. LEAFY target genes reveal floral regulatory logic, cis motifs, and a link to biotic stimulus response. *Dev Cell.* 2011;20:430–43. doi:10.1016/j.devcel.2011.03.019.
- William DA, Su Y, Smith MR, Lu M, Baldwin DA, Wagner D. Genomic identification of direct target genes of LEAFY. *Proc Natl Acad Sci USA.* 2004;101:1775–80. doi:10.1073/pnas.0307842100.
- Wellmer F, Alves-Ferreira M, Dubois A, Riechmann JL, Meyerowitz EM. Genome-wide analysis of gene expression during early Arabidopsis flower development. *PLoS Genet.* 2006;2:e117. doi:10.1371/journal.pgen.0020117.
- Goslin K, Zheng B, Serrano-Mislata A, Rae L, Ryan PT, Kwaśniewska K, Thomson B, O'Maoileidigh D, Madueño F, Wellmer F, Graciet E. Transcription factor Interplay between LEAFY and APETALA1/CAULIFLOWER during floral initiation. *Plant Physiol.* 2017;174:1097–109. doi:10.1104/pp.17.00098.
- Winter CM, Yamaguchi N, Wu MF, Wagner D (2015). Transcriptional programs regulated by both LEAFY and APETALA1 at the time of flower formation. *Physiol Plant.* 2015;155:55–73. doi:10.1111/ppl.12357.
- Ratcliffe OJ, Bradley DJ, Coen ES. Separation of shoot and floral identity in Arabidopsis. *Development.* 1999;126:1109–20.
- Ferrandiz C, Gu Q, Martienssen R, Yanofsky MF. Redundant regulation of meristem identity and plant architecture by FRUITFULL, APETALA1 and CAULIFLOWER. *Development.* 2000;127:725–34.
- Blázquez MA, Soowal LN, Lee I, Weigel D. LEAFY expression and flower initiation in Arabidopsis. *Development.* 1997;124:3835–44.
- Serrano-Mislata A, Fernández-Nohales P, Doménech MJ, Hanzawa Y, Bradley D, Madueño F. Separate elements of the TFL1 cis-regulatory region integrate pathways to control flowering time and shoot meristem identity. *Development.* 2016;143:3315–27. doi:10.1242/dev.135269.

24. Sessions A, Yanofsky MF, Weigel D. Cell-cell signaling and movement by the floral transcription factors *LEAFY* and *APETALA1*. *Science*. 2000;289:779–82. doi:10.1126/science.289.5480.779.
25. Chahtane H, Vachon G, Le Masson M, Thévenon E, Périgon S, Mihajlovic N, Kalinina A, Michard R, Moyroud E, Monniaux M et al. A variant of *LEAFY* reveals its capacity to stimulate meristem development by inducing *RAX1*. *Plant J*. 2013;74:678–89. doi:10.1111/tpj.12156.
26. Benlloch R, Kim MC, Sayou C, Thevenon E, Parcy F, Nilsson O. Integrating long-day flowering signals: a *LEAFY* binding site is essential for proper photoperiodic activation of *APETALA1*. *Plant J*. 2011;67:1094–102. doi:10.1111/j.1365-3113.2011.04660.x.
27. Lee I, Wolfe DS, Nilsson O, Weigel D. A *LEAFY* co-regulator encoded by *UNUSUAL FLORAL ORGANS*. *Curr Biol*. 1997;7:95–104. doi:10.1016/S0960-9822(06)00053-4.
28. Kim D, Kwon YK, Cho KH. The biphasic behavior of incoherent feed-forward loops in biomolecular regulatory networks. *BioEssays*. 2008;30:1204–11. doi:10.1002/bies.20839.