

REVIEW

Open Access



The protective role of estrogen and estrogen receptors in cardiovascular disease and the controversial use of estrogen therapy

Andrea Iorga², Christine M. Cunningham¹, Shayan Moazeni¹, Gregoire Ruffenach¹, Soban Umar¹ and Mansoureh Eghbali^{1*}

Abstract

Epidemiologic studies have previously suggested that premenopausal females have reduced incidence of cardiovascular disease (CVD) when compared to age-matched males, and the incidence and severity of CVD increases postmenopause. The lower incidence of cardiovascular disease in women during reproductive age is attributed at least in part to estrogen (E2). E2 binds to the traditional E2 receptors (ERs), estrogen receptor alpha (ER α), and estrogen receptor beta (ER β), as well as the more recently identified G-protein-coupled ER (GPR30), and can exert both genomic and non-genomic actions. This review summarizes the protective role of E2 and its receptors in the cardiovascular system and discusses its underlying mechanisms with an emphasis on oxidative stress, fibrosis, angiogenesis, and vascular function. This review also presents the sexual dimorphic role of ERs in modulating E2 action in cardiovascular disease. The controversies surrounding the clinical use of exogenous E2 as a therapeutic agent for cardiovascular disease in women due to the possible risks of thrombotic events, cancers, and arrhythmia are also discussed. Endogenous local E2 biosynthesis from the conversion of testosterone to E2 via aromatase enzyme offers a novel therapeutic paradigm. Targeting specific ERs in the cardiovascular system may result in novel and possibly safer therapeutic options for cardiovascular protection.

Keywords: Estrogen, Estrogen receptor alpha, Estrogen receptor beta, GPR30, Cardiovascular disease, Hormone replacement therapy, Fibrosis, Angiogenesis, Oxidative stress, Vasodilation

Background

While cardiovascular disease (CVD) is the leading cause of death among women, epidemiologic studies indicate that females prior to menopause are somewhat protected against the development of CVD when compared to men. Women have reduced incidence of CVD when compared to age-matched males and present with CVD 10 years later than men [1]. In response to aortic stenosis, males have significantly more maladaptive cardiac remodeling than women, and transcriptome characterization reveals that fibrosis and inflammation-related genes and pathways are upregulated in males but not in females [2].

Premenopausal women also withstand ischemia/reperfusion (I/R) injury during open heart surgery better than males [3]. Female protection against CVD is associated with sex hormone levels as the incidence and severity of CVD increases in women postmenopause [3], and the prevalence of coronary artery disease (CAD) is greater in young women who had an oophorectomy compared to those with intact ovaries [4]. Taken together, this data has sparked much investigation into the potential of female sex hormones in conferring cardioprotection. The results of years of research into female sex hormones indicate that the main circulating female hormone, estrogen (E2), is cardioprotective through a plethora of mechanisms that are highlighted throughout this review. While there is abundant data supporting E2 as a cardioprotective agent in experimental models of CVD, E2 replacement therapy in postmenopausal women has been very controversial [5].

* Correspondence: meghbali@ucla.edu

¹Department of Anesthesiology, Division of Molecular Medicine, David Geffen School of Medicine at University of California, Los Angeles, BH-160CHS, Los Angeles, CA 90095-7115, USA

Full list of author information is available at the end of the article




Furthermore, while postmenopausal women are at greater risk than premenopausal women of developing CVD, women still have significantly lower incidence of CVD compared to age-matched men well beyond menopause [6]. Discrepancies in the development and prognosis of CVD indicate that the cardioprotection conferred by E2 in females is complex and potentially not the only sex-biasing factor, as other sex-biasing factors such as the sex chromosome composition are also known to contribute to sex biases in CVD [6–8]. The purpose of this review is to summarize the role of E2, E2 receptors (ERs), and aromatase in experimental models of CVD as well as highlight the controversies of hormone replacement therapy in women. We also present the sexual dimorphic role of ERs in modulating E2 action in CVD.

Estrogen and estrogen receptors

Estradiol, also known as 17beta-estradiol or estrogen (E2), is the most abundant form of circulating estrogens and considered the main female hormone. Two other naturally occurring forms occur in lower abundance (estrone (E1) and estriol (E3)), while a third form (estretrol (E4)) is produced only during pregnancy [9].

E2 is predominantly synthesized and secreted by the ovaries in premenopausal women. Some E2 is also produced in other tissue types including adipose, brain, and bone tissues as well as the vascular endothelium and aortic smooth muscle cells [10]. While gonadal E2 acts largely as an endocrine factor affecting distal tissues, extragonadal production of E2 acts locally as a paracrine or intracrine factor in the tissue where it is synthesized [10]. This extragonadal E2 production plays an important role, as it remains the only source of endogenous E2 production in postmenopausal women and men [10].

E2 binds to the traditional ERs, ER α and ER β , as well as the newly identified G-protein-coupled ER (GPR30) [11] (Fig. 1). Binding to ER α and ER β can confer both genomic and rapid non-genomic action [12]. In the genomic pathway, E2 binding triggers intracellular localization of ER α and ER β , which dimerize and enter the nucleus. Once in the nucleus, they bind to E2 response elements (ERE), or activator protein-1 (Ap1) and specificity protein-1 (Sp1), on the promoter of E2-responsive genes to regulate transcription. E2 also binds to membrane-bound ER α and ER β receptors as well as GPR30 to rapidly activate nuclear transcription factors



via the MAPK pathway. The E2 pathway is summarized in Fig. 1.

While still considered controversial by some, all three E2 receptors have been identified as functional and present in adult cardiomyocytes [11, 13–15]. The role of each ER and the mechanism by which it confers cardioprotection is actively being investigated in various animal models, and the current findings will be summarized in the following sections of this review.

Estrogen, mitochondria, and oxidative stress

Heart failure (HF) and ischemic heart injury are linked to several bioenergetic abnormalities including decreased energy metabolism, increased apoptosis, production of reactive oxygen species (ROS), and dysfunctional calcium signaling. These abnormalities can largely be attributed to changes in mitochondrial homeostasis [16, 17]. Both traditional ERs and GPR30 are localized on the mitochondrial membrane in cardiac tissue, and activation of these receptors via E2 can trigger transcriptional changes in nuclear and mitochondrial genes influencing mitochondrial function, cell survival, and ultimately cardioprotection [18, 19]. While the beneficial role of E2 in cardioprotection has been demonstrated in both male and female rodents, there is growing evidence demonstrating that E2-induced cardioprotection via mitochondrial homeostasis and oxidative stress may exhibit sex differences.

Lagranha et al. found that adult female rats are protected against ex vivo I/R injury in the presence of E2 as they had smaller infarct size and increased cardiac contractility compared to female rats without gonadal E2 (post ovariectomy, OVX) [20]. Male rats with E2 treatment were also protected against I/R injury with smaller infarct and higher cardiac contractility than non-treated controls. Mitochondria from both females and E2-treated males had increased levels of PKC-dependent phosphorylation of aldehyde dehydrogenase 2 (ALDH2) resulting in increased ALDH activity. Activation of ALDH has previously been reported to protect the heart against ischemic damage [21]. Lagranha et al. also investigated if E2 conferred cardioprotection via activation of PI3K signaling pathway. Pretreatment of hearts with the PI3K inhibitor wortmannin diminished the cardioprotection in female rats following I/R injury as well as reduced p-ALDH2 [20]. This study also linked increased p-ALDH2 to decreased ROS production and found that cardiomyocytes from female rats had less ROS production than cardiomyocytes from male rats following I/R injury [20].

A study by Zhai et al. highlighted the protective role of ER α in male mice following global myocardial I/R by comparing male ER α knockout (ERKO) and wild-type mice [22]. Following ischemia, the hearts of ERKO mice

had lower functional recovery, higher incidence of fibrillation and/or tachycardia, and a significantly lower coronary flow rate leading to larger infarct size compared to wild-type mice. Transmission electron microscopy revealed that ERKO hearts had swollen and fragmented mitochondria with loss of matrix and ruptured cristae [22]. Functionally, the mitochondria isolated from ERKO hearts were found to have more severe respiratory dysfunction than wild-type controls [22]. This study demonstrates that ER α plays a cardioprotective role against I/R injury in male mice and that the abolishment of ER α promotes I/R injury via impaired calcium influx and mitochondrial dysfunction. The importance of ER α in mediating E2 cardioprotection has also been identified in females as well. In an in vivo model of I/R using adult female rabbits, acute pretreatment with E2 or ER α -specific agonist PPT, but not ER β -specific agonist DPN, significantly decreased infarct size suggesting that ER α plays a significant role in the acute cardioprotective action of E2 [12, 23].


E2 also mediates cardioprotection through GPR30 activation on the mitochondrial surface. Bopassa et al. found that during I/R injury, GPR30 activation by E2 in adult male mice results in cardioprotection by inhibiting the opening of the mitochondrial permeability transition pore (mPTP) via activation of the ERK pathway [11]. E2 binding to GPR30 activates adenylyl cyclase, mitogen-activated protein kinases (MAPK), and extracellular signal-regulated kinases, ERK1 and ERK2, resulting in mobilization of intracellular calcium [24]. The mPTP appears to play a pivotal role in apoptosis following I/R [25]. During ischemia, the mPTP remains closed and opens during the first few minutes of reperfusion due to oxidative stress, calcium overload, and ATP depletion [26]. Bopassa et al. found that in the presence of the GPR30 agonist G-1, mouse hearts had better functional recovery as well as a smaller infarct size. Furthermore, the presence of G-1 allowed for the mitochondria to uptake more calcium prior to the opening of the mPTP. All the protective effects of GPR30 activation were blocked by ERK inhibition indicating that GPR30 confers cardioprotection by inhibiting the opening of the mPTP via ERK activation [11].

In agreement with the beneficial role of E2 on the heart, studies have shown a detrimental effect of E2 depletion via OVX on cardioprotection. Pavón et al. found that following OVX, mitochondria from adult female rats have decreased Ca²⁺ retention capacity, which is restored when mitochondria are incubated with E2, thus indicating a protective role of E2 against ischemic injury [27]. A later study by Pavón et al. found that in the absence of E2, female rats also exhibit increased mitochondrial dysfunction: mitochondria were more susceptible to stress-induced dysfunction of complex I and exhibited

significant changes in the oxidative phosphorylation-related proteins including cytochrome c oxidase and ATP synthase [28]. Finally, Pavón et al. also reported that E2 is protective against damaging oxidative stress resulting from lipid peroxidation and free radical formation. They showed that levels of thiobarbituric acid reactive substances (TBARS) were increased following I/R injury in OVX females when compared to intact females. The expression of TBARS was also greatly reduced in intact females compared to intact males [27]. A study by Liu et al. suggests that higher levels of circulating E2 in female rodents may protect against cardiovascular stressors due to increased antioxidant expression [30]. Intact female mice were found to have increased cardiac levels of the antioxidant mitochondrial superoxide dismutase 2 (SOD2) when compared to male mice. Following OVX, SOD2 expression decreased to match that of males, and E2 therapy in OVX female mice resulted in increased SOD2 expression. Furthermore, OVX mice fed high-fat diet produced more aortic ROS than intact and E2-treated females fed high-fat diet. E2 stimulated the expression of SOD2 through both classical ERs [30].

A study by Wang et al. further highlighted the anti-oxidative effects of E2 in cardioprotection. The study found that activation of ER α by E2 in cardiomyocytes decreased levels of microRNA 22 (miR-22) through Sp1 coactivation [29]. Decreased miR-22 expression resulted in an upregulation of cystathionine γ -lyase (CSE) in cardiomyocytes. CSE is known to be upregulated in the myocardium of female rats and in rats treated with E2 and confers cardioprotection by increasing synthesis of the antioxidant hydrogen sulfide [30]. Additionally, female cardiomyocytes are also protected against oxidative stress via ER α -mediated activation of AKT/GSK-3 β and AKT/Bcl-2 and inhibition of caspase 3 [31].

In summary, E2 increases cardioprotection against I/R injury and high-fat diet by improving mitochondrial function and reducing ROS mainly through ER α and GPR30 (Fig. 2). E2-enhanced mitochondrial structure and function by activating the cardioprotective signaling pathways (PI3K/ERK1/2) and inhibiting mPTP opening. E2 reduces ROS production and protects against oxidative stress by increasing the production of powerful antioxidants SOD2 and hydrogen sulfide.



Estrogen and fibrosis

Myocardial extracellular matrix (ECM) proteins, matrix metalloproteinases (MMPs), growth factors, and cytokines are mainly produced by cardiac fibroblasts, all of which contribute to the myocardial structure maintenance as well as cardiac remodeling in diseased hearts [32]. Cardiac remodeling is manifested by changes in the size, shape, and function of the heart including cardiomyocyte hypertrophy, fibroblast proliferation, fibrosis, an accumulation of fibrillar collagens, and myocyte apoptosis [33]. Increased expression of fibrillar collagens leads to stiffening and electrical uncoupling of the cardiac muscle impeding both relaxation and contraction of the heart. Furthermore, increased levels of fibrosis cause a decrease in angiogenesis in the heart as well as an increase of oxygen diffusion distance, thus leading to myocyte hypoxia [34].

The normal ECM, mainly composed of type I and type III collagens, plays an important adaptive role in HF by preventing excessive dilatation when ventricular overload occurs. An imbalance in the synthesis and/or inhibition in the expression of ECM proteins results in fibrosis. In the pressure overloaded ventricles, cardiomyocyte hypertrophy is accompanied by increased collagen deposition between and around myocytes, resulting in interstitial and perivascular fibrosis [35]. The ECM is enzymatically digested by MMPs, and an imbalance of collagen deposition and the activity or expression of MMPs can lead to cardiac remodeling. In the normal heart, the majority of cells are fibroblasts [36], and external stressors cause fibroblasts to change their phenotype to myofibroblasts, which plays a pivotal role in inflammation and fibrosis [32].

A study by Lee and Eghbali-Webb first investigated the expression of the classic ERs in cardiac fibroblasts isolated from female rats. The study found that both ER α and ER β are expressed in fibroblasts, with the predominant receptor being ER β which is expressed in the cytosol and nucleus [37]. Lee and Eghbali-Webb also found that treatment of fibroblasts with E2 increases DNA synthesis, which was prevented in the presence of tamoxifen, a selective ER modulator with high inhibitory affinity for both ERs in the heart. This upregulation of DNA synthesis occurs through nuclear translocation of receptor protein and activation of MAPK [37]. These data suggest that E2 acts on the fibroblasts to regulate ECM remodeling and modulate myocardial function via ER and MAPK-dependent mechanisms. Paradoxically, another early study by Dubey et al. found that E2 inhibited fetal calf serum-induced fibroblast proliferation and collagen synthesis in both male and female rats [38]. Furthermore, the E2 metabolites 2-hydroxyestradiol and 2-methoxyestradiol were found to be even more potent than E2 in inhibiting fibroblast proliferation and collagen

synthesis, and these effects were enhanced in the presence of progesterone and 4-hydroxytamoxifen (a high-affinity ER ligand) [38]. Considering that the results were similar in both sexes, E2 inhibition of fibroblast proliferation is most likely not acting in a sex-specific manner [38].

Recent studies further support the anti-fibrotic effect of E2 in the heart. Pedram et al. found that ER β activation prevents cardiac fibrosis by blocking the effects of angiotensin II (AngII) and endothelin-1 (ET-1)-induced pro-fibrotic signaling in female mice [39]. AngII and ET-1 promote cardiac fibrotic deposition by driving the transition of fibroblasts to myofibroblasts and stimulating the synthesis of transforming growth factor- β 1 (TGF β 1), a potent inducer of cardiac fibrosis. Upregulation of TGF β 1 in the myofibroblast leads to production of vimentin, fibronectin, and collagens I and III. E2, as well as an ER β agonist, inhibit TGF β 1 through cAMP and protein kinase A. E2 inhibits fibrosis through ER β as E2 administration was shown to prevent AngII-induced cardiac fibrosis in female OVX wild-type (WT) mice but not in ER β knockout mice [39]. Therefore, this study demonstrated the strong anti-fibrotic effects of ER β activation via E2.

Recent work by our lab found that E2 treatment reverses fibrotic deposition induced by the transaortic constriction model of HF in male mice [40]. E2 treatment diminished the expression of pro-fibrotic genes that are increased in response to cardiac pressure overload including collagens I and III, TGF β 1, lysyl oxidase, and fibrosin. E2 treatment resulted in an upregulation of ER β transcript expression which could elucidate the mechanism by which E2 exerts its effects inhibiting fibrosis [40]. Our study further contributed to the growing body of evidence highlighting E2 as a powerful inhibitor of profibrotic genes that are detrimental to cardiac function.

To study the effect of ERs on fibrosis in the in vivo I/R injury model, mice with overexpression of either ER α or ER β were subjected to left anterior descending (LAD) coronary artery ligation and subsequent reperfusion. Mahmoodzadeh et al. found that cardiac fibrosis was attenuated in female mice overexpressing ER α after they were subjected to LAD ligation [41]. This suggests that ER α has an important role in protecting the hearts of female rodents against ischemic insults. Another study by Schuster et al. examined the role of ER β in regulating myocardial infarct size post-LAD [42]. They found that cardiomyocyte-specific ER β overexpression led to improved heart function and survival in both sexes. Interestingly, male mice with ER β overexpression had increased left ventricular (LV) volume and EF with reduced cardiac fibrosis when compared to female mice overexpressing ER β .

The more recently investigated GPR30 may also play a role in conferring cardioprotection via anti-fibrotic

mechanisms. Wang et al. found that GPR30 activation via G-1 prevented cardiac fibroblast proliferation and fibrosis both in vitro and in vivo. They demonstrated that cultured cardiac fibroblasts from male rats displayed GPR30 at the cell surface and treatment with G-1 reduced proliferation. In vivo, G-1 treatment attenuated the adverse interstitial collagen deposition and fibroblast proliferation resulting from high-salt diet or E2 depletion from OVX in female rats [43].

MMP-2, which exhibits increased expression and activity following aortic stenosis, myocardial infarction (MI), and LV hypertrophy, is believed to have an adverse effect on cardiac remodeling [44–47]. Furthermore, constitutive expression of MMP-2 causes systolic dysfunction and severe remodeling [48], while its inhibition attenuates cardiac remodeling and improves survival after MI or pressure overload [49–51]. Mahmoodzadeh et al. found that E2 significantly reduced MMP-2 gene expression in male and female adult rat and human fibroblasts via activation of the classic ERs. This effect was due to E2-dependent phosphorylation of the transcription factor Elk-1 via the MAPK signaling pathway [44]. Thus, E2-induced repression of MMP-2 may contribute to sex differences in fibrotic processes, although the authors did not note sex differences in MMP-2 expression.

Lastly, our group has shown that E2 and ER β activation reverse right ventricular failure caused by pulmonary hypertension (PH) in intact male rats [52]. Right ventricular (RV) failure was associated with upregulation of ECM-interacting cardiac fetal gene osteopontin. We found that the transcript levels of two novel ECM-degrading disintegrin-metalloproteinases (ADAM), ADAM15 and ADAM17, as well as osteopontin and phosphorylated AKT were all elevated in the failing RV and E2 therapy after the onset of PH reversed these effects likely through ER β activation [52].

In summary, E2 has been demonstrated to have a significant role in ECM remodeling by acting on fibroblasts via ER and MAPK-dependent mechanisms (Fig. 2). Recent data demonstrates that E2 protects against cardiac fibrosis and harmful ECM remodeling by altering fibroblast proliferation and inhibiting various pro-fibrotic genes.

Estrogen and angiogenesis

By definition, angiogenesis is the physiological process by which new blood vessels form from pre-existing vessels. E2 has previously been shown to be pro-angiogenic in various tissues including the uterus, breast, brain, and limbs [53–55]. Increased cardiac angiogenesis is necessary to maintain LV function during adaptive hypertrophy and compensated heart hypertrophy as the larger LV has an obvious need for more blood vessels [56, 57]. However, an imbalance between cardiac growth and neoangiogenesis over time can induce the progression of

compensated heart hypertrophy to HF [58]. It is therefore postulated that stimulation of cardiac neoangiogenesis could be vital in preventing the transition to HF.

Several agents have been shown to induce angiogenesis. The fibroblast growth factor (FGF) family, particularly FGF-1 and FGF-2, are potent angiogenic growth factors that promote endothelial proliferation and organization into tubular structures [59, 60]. Vascular endothelial growth factor (VEGF) is an important angiogenic factor and a critical determinant of capillary growth and density [61]. When bound to its receptor, VEGF starts a signaling cascade which stimulates nitric oxide synthase 3 (eNOS) and nitric oxide (NO) production, endothelial cell proliferation, migration of tube structures, and differentiation into mature blood vessels [61, 62]. We recently showed that VEGF transcript levels are significantly downregulated (~5-fold) in male mice with HF and exogenous E2 treatment after establishment of HF was able to restore VEGF expression in the LV [40]. Furthermore, we showed that E2 therapy in male mice significantly enhanced capillary density by ~4-fold compared to the HF group alone [40]. Stimulation of cardiac neoangiogenesis by E2 in HF is not only limited to the LV, as E2 also stimulates angiogenesis in RV failure secondary to PH in male rats [67]. Both studies found that E2 was no longer able to rescue HF in the presence of an angiogenesis inhibitor, which indicates that E2 protects against LV and RV HF in male rodents by stimulating angiogenesis [40, 63].

In agreement with our finding, E2 has also been shown to promote vascularization in the experimental model of myocardial ischemic injury. Iwakura et al. showed that E2 augments the incorporation of endothelial progenitor cells (EPCs) in the sites of cardiac ischemic vascularization, resulting in protection from ischemic injury [64]. Using OVX female mice pretreated with placebo or E2 prior to MI induced by LAD ligation, the group found that E2 induced a significant increase in circulating EPCs, which led to an increase in cardiac capillary density. The study also found a greater preponderance of EPCs in the E2-treated animals at the ischemic sites [64]. E2 failed to increase homing of EPCs to the injured myocardium in eNOS-null mice [68]. Therefore, E2 maintains the integrity of ischemic heart tissue by increasing the mobilization and incorporation of EPCs by eNOS. The same group then investigated the role of ER α and ER β in the E2-induced mobilization of EPCs, which favorably affected neovascularization post cardiac ischemic injury [65]. The study found that the E2-induced endothelial migration, tube formation, and adhesion were impaired when using EPCs derived from ER α or ER β KO female mice; however, the impairment was more severe when EPCs were derived from ER α KO mice. When bone marrow was transplanted from either ER α or ER β KO female mice into WT female mice, capillary density at the

ischemic border zone was significantly reduced when compared to bone marrow transplanted from WT female mice. ER α transcript levels were more abundant than ER β in EPCs, and VEGF expression was downregulated only in the ER α knockout EPCs both in vivo and in vitro when compared to WT [65]. Thus, both ER α and ER β contribute to E2-induced EPC mobilization and preservation of cardiac function in female mice following MI, but ER α plays a more predominant role in this cardioprotective process.

Interestingly, a study by Mahmoodzadeh et al. using intact male and female mice with a cardiomyocyte-specific overexpression of ER α found that ER α increases expression of angiogenesis and lymphangiogenesis markers (such as VEGF and Lyve-1) and neovascularization in the peri-infarct area following MI in a sex-specific manner. Only female mice overexpressing ER α exhibited 100% survival 14 days post-MI concomitant with reduced pathological cardiac remodeling when compared to male mice overexpressing ER α and wild-type mice of both sexes [41]. Thus, ER α overexpression increased cardiac angiogenesis and induced cardioprotection in females, but was not sufficient to confer cardioprotection in males which is likely a result of the increased circulating E2 in females.

In summary, stimulation of neoangiogenesis by E2 therapy in the heart plays an important role in preventing the transition from compensated hypertrophy to HF in both the LV and RV. E2 therapy in HF rodents stimulates neoangiogenesis by enhancing VEGF expression, which is normally suppressed during HF, and increasing capillary density in the heart (Fig. 2). Induction of neoangiogenesis via E2 therapy is dependent on eNOS activation, as eNOS-null mice do not exhibit the pro-angiogenic effects following E2 therapy. Moreover, ER α plays a more dominant role than ER β in E2-induced EPC activation, upregulation of VEGF transcripts, and preservation of cardiac function post MI, and overexpression of ER α in cardiomyocytes is protective against MI in female but not male mice.

Estrogen and vasodilation

Under healthy conditions, the vascular endothelium plays an important role in maintaining vascular tone and blood flow. When dysfunctional, the endothelium promotes inflammation, prothrombotic factors, and vasoconstriction, which increase the risk of developing CVD [66–69]. Reports show that E2, when bound to either classical ERs or GPR30, can help maintain endothelial homeostasis and vasodilation by activating the transcription of eNOS and subsequent upregulation of NO through genomic and non-genomic pathways [5, 70–72]. While the effect of E2 on the vascular endothelium is largely time and tissue dependent, studies show that

some of the cardioprotection conferred by E2 may be a result of increased vasodilation.

While there is ample evidence supporting NO signaling through ER activation, the mechanisms by which E2 exerts its powerful vasoactive effects are not straightforward. White et al. found that E2 relaxed porcine coronary arteries in an endothelium-independent and dose-dependent fashion; however, when arteries were pretreated with agents to uncouple NO production from eNOS, E2 elicited dose-dependent contractions of arteries [73]. After E2-induced coronary contraction had reached its maximum levels, an inhibitor of superoxide (O $_2^-$, tempol) was able to reverse this response by reducing O $_2^-$. Thus, the authors proposed that acute E2 administration produces both coronary vasodilation and vasoconstriction via NO and O $_2^-$, respectively [73].

In an attempt to investigate the role of ER α and ER β in NO stimulation and cardioprotection, several groups studied the effect of OVX, E2, and selective ER agonists on vasodilation and hypertension in spontaneously hypertensive rats. Treatment of spontaneously hypertensive OVX female rats with E2 or a selective ER α agonists resulted in improved endothelial function, reduced cardiac hypertrophy, and improved cardiac outcomes when compared to OVX rats without treatment [74, 75]. While Widder et al. found that treatment with E2 or selective ER α agonist resulted in increased eNOS expression in the aorta, there was no statistically significant attenuation of hypertension when compared to OVX rats in this study or in a later study by Pelzer et al. [74, 75]. A follow-up study by Jazbutyte et al. revealed that administration of a selective ER β agonist to spontaneously hypertensive OVX female rats attenuated hypertrophy, lowered peripheral artery resistance, and decreased systolic blood pressure [76]. Taken together, these studies indicate that E2 improves cardiovascular function in spontaneously hypertensive female rats via both classical ERs, although ER β elicits a stronger anti-hypertensive effect. In the AngII-induced cardiac hypertension model, however, Xue et al. found that AngII-induced hypertension was greater in OVX and ER α KO female mice than in intact WT female mice suggesting that E2 acts mainly through ER α to reduce AngII-induced hypertension [77].

In the context of I/R injury, Gabel et al. also concluded that ER β confers cardioprotection in females by modulating the expression of eNOS and other genes involved in fatty acid metabolism [78]. In this study, gonadally intact males treated with isoproterenol immediately prior to ex vivo ischemia exhibited significantly lower functional recovery following reperfusion than intact females. Transgenic female ER α -KO mice exhibited similar functional recovery to WT females, while female ER β -KO mice exhibited diminished functional recovery similar to

that of WT males. Notably, cardiac eNOS protein expression was found to be significantly lower in the ER β -KO mice when compared to female WT and ER α -KO mice, indicating that the ER β -mediated expression of eNOS may be responsible for sex differences in functional recovery following I/R injury [78]. In a separate study, female ER β -KO mice were also found to exhibit abnormal vascular function, hypertension, increased mortality, and aggravated heart failure following MI when compared to female WT controls [79]. Interestingly, Pelzer et al. did not find significant differences in LV eNOS expression between groups [79].

Similarly, Nikolic et al. found that 2-week pretreatment with DPN, an ER β agonist, is cardioprotective in I/R injury in OVX female mice [79]. Gene profiling in this experimental model revealed that treatment with DPN is associated with a significant increase in cardioprotective genes, including those encoding NO biosynthesis and anti-apoptotic proteins [80]. Lin et al. showed that chronic E2 or DPN treatment also leads to activation of protein S-nitrosylation and cardioprotection, which was blocked by eNOS inhibition [81]. Taken together, these data indicate that chronic E2 exposure protects the heart largely via ER β activation and subsequent NO signaling.

Studies suggest that classical ERs are not solely responsible for E2-induced vasodilation and that GPR30 activation can also impact vascular tone. Lindsey et al. found that activation of GPR30 via G-1 induced similar vasodilatory effects as E2 in isolated aortic rings from both intact and OVX female rats [82]. Pretreatment with GPR30 antagonist, G15, successfully blocked G-1- and E2-induced vasodilation. Similarly, removal of the endothelium or pretreatment with l-NAME, a constitutive eNOS inhibitor, partially attenuated vasorelaxation. These data suggest that GPR30 activation by E2 or selective agonist induces vasorelaxation in female rodent aortas through a partially endothelium and eNOS-dependent mechanism [82].

More recently, Fredette et al. elucidated the mechanism by which E2 and G-1 induce vasodilation in male human endothelial cells and GPR30-KO mice [83]. The study demonstrated that GPR30 stimulates endothelial NO production via activation of the c-Src/EGFR/PI3K/ERK1/2 signaling pathway. E2 and G-1 treatment induced NO release in cultured endothelial cells and extracted aortic rings from WT mice. In GPR30-KO mice, the vasodilatory effect of G-1 was completely abrogated while the effect of E2 was reduced by 50% [83].

The significant involvement of GPR30 in eliciting a vasodilatory response may contribute to some of the sex differences in CVD. A study by Lenhart et al. reported that female mice have more membrane-localized GPR30 when compared to male mice [84]. Furthermore, they found that GPR30 directly interacts with receptor active modifying protein 3 (RAMP3), a protein known to affect

the trafficking and activity of several GPCRs. After subjecting male and female WT and Ramp3-KO mice to GPR30 activation via G-1, the authors found the cardio-protection conferred by G-1 to be both Ramp3 and sex specific [84].

Sudhir et al. found that E2 induced a significant increase in coronary cross-sectional area, flow velocity, and volumetric blood flow in male and female canine coronary arteries that was not attenuated by pretreatment with l-NAME or the classic ER antagonist, ICI 162,780 [85]. The investigators thus conclude that acute E2-induced dilation of coronary arteries is endothelium-independent and is not mediated via the classic intracellular E2 receptors, but presumably via non-genomic mechanisms involving GPRs at the plasma membrane [85].

In addition to stimulating eNOS production, E2 has also been shown to enhance vasodilation through inhibition of the angiotensin pathway. Isolated RNA from different tissues (kidney cortex, kidney medulla, lung, and aorta) of OVX female rats treated with E2 for 21 days revealed a downregulation of angiotensin-converting enzyme (ACE) transcript levels [86]. Thus, downregulation of ACE with a consequent reduction in the circulating levels of the vasoconstrictor AngII could be another mechanism by which E2 confers its beneficial cardiovascular effects.

In summary, studies suggest that E2 plays a very dynamic role in modulating vasorelaxation, vasoconstriction, and endothelial function via largely eNOS-dependent mechanisms (Fig. 2). This mechanism may contribute to the cardioprotective effect of E2 as E2 reduced hypertrophy and increased cardiac function in the spontaneous hypertensive rat model and in models of I/R injury. Further studies using animal models of hypertension found that stimulation of classical ERs may also attenuate hypertension. Stimulation of GPR30 by E2 or GPR30-specific agonists also stimulated eNOS and NO production in a sex-specific manner. By reducing the expression of ACE, E2 treatment also results in lower levels of AngII and reduced vasoconstriction.

Aromatase and the cardiovascular system

While the main site of E2 production is the female ovary, peripheral tissues can also act as sites of local E2 production by converting androgens into E2 via the aromatase enzyme [87]. Aromatase is widely distributed in gonadal and extragonadal tissues including the bone, brain, adipose tissue, and blood vessels [88]. Local production of E2 in these tissues by aromatase plays an important role in the physiological functions of these tissues [89]. For example, the presence of aromatase in the coronary endothelium suggests that the local cardiac

conversion of androgens to E2 likely affects cardiac function and structural modeling [90].

Aromatase is encoded by the CYP19 gene, and genetic variations in the CYP19 gene result in alterations of circulating levels of sex hormones [91–93]. Ma et al. suggests that genetic variations in CYP19 might contribute to variations in the pathophysiology of E2-dependent diseases [94]. Wang et al. identified a single nucleotide polymorphism in the CYP19 gene to be associated with a decreased risk of coronary heart disease among a Chinese population in which both sexes were equally represented [95].

Aromatase activity has been confirmed in female rat arterial smooth muscle cells (SMCs) and in bovine coronary endothelial cells (ECs) *in vitro* [96, 97]. Harada et al. further demonstrated aromatase activity in human male and female arterial SMCs, but not in female ECs, which could potentially indicate autocrine/paracrine E2 activity [89]. It is speculated that E2 produced in vascular SMCs acts to promote cardiac contractility, vasodilation, and collagen synthesis, while also stimulating EC functions such as NO production and angiogenesis [89]. Within vasculature, E2 protects against atherosclerotic progression in experimental animal models of atherosclerosis in both males and females, and aromatase has been shown to play a pivotal role in promoting this vascular protection [98–100]. The ablation of aromatase has also been demonstrated to be associated with increased adiposity [101] and progressive insulin resistance [102], which are significant factors in the development of cardiovascular diseases including atherosclerosis and diabetic cardiomyopathy.

While E2 is largely considered to be protective in both male and female rodents in the context of I/R injury and HF, the role of aromatase in conferring cardioprotection is unclear. Bell et al. investigated the role of aromatase deficiency in I/R injury using aromatase knockout (ArKO) female mice and found that the recovery of left ventricular developed pressure as well calcium handling was substantially improved in ArKO versus wild-type (WT) mouse hearts [103]. Later on, the same group demonstrated that isolated cardiomyocytes from ArKO female mice exhibited greater basal calcium transient amplitude and shortening than in WT [104]. Isolated cardiomyocytes from ArKO female mice exposed to a high-calcium load also showed increased calcium transient and contractile amplitudes. The study concluded that the relative withdrawal of E2 in favor of testosterone may be positive inotropic via optimized calcium handling in response to stress and aromatase inhibition which could lead to cardioprotection [104]. Bell et al. also studied whether upregulation of tissue aromatase expression could improve ischemic resilience in male hearts by subjecting male mice in which aromatase was transgenically

upregulated (AROM+) to I/R injury [105]. As expected, male AROM+ mice exhibited markedly lower testosterone and higher E2 levels when compared to wild-type male mice. AROM+ male hearts had better functional recovery post ischemia, whereas female AROM+ hearts did not exhibit the same improvement in post-ischemic function [105]. Thus, these findings demonstrate that aromatase modulates cardiac function in a sex-specific manner [105].

A recent study by our group found that cardiac aromatase transcript levels are reduced in male mice in the TAC-induced model of LV HF [40]. Interestingly, these mice also exhibited decreased local cardiac E2 levels, but not circulating plasma E2. E2 therapy for 10 days after the onset of HF restored aromatase transcript levels in the heart similar to control levels and upregulated local cardiac E2 concentrations. We stipulated that the normalization of cardiac aromatase by E2 could either be attributed to improvement in cardiac function or increased E2 bioavailability in the heart [40]. As the aromatase gene has several half E2 response elements as well as Sp1 and Ap1 elements, it is possible that increased heart E2 concentrations reported in the study post-E2 therapy could have stimulated the transcription of aromatase.

Recent studies have also implicated the role of aromatase in regulating RV function in the Sugden hypoxia rat model, an experimental model of PH that is associated with an elevated RV systolic pressure and RV hypertrophy. Aromatase inhibition using anastrozole (an aromatase inhibitor) reversed adverse RV remodeling in female Sugden hypoxia rats [106]. Studies have also shown that metformin (a drug used to treat type 2 diabetes by controlling blood glucose levels) reversed elevated RV systolic pressure and hypertrophy in Sugden hypoxia female rats, independent of glycemia, without any effects on systemic blood pressure, cardiac output, or heart rate [107]. Interestingly, metformin was shown to inhibit aromatase without any adverse effects on the RV [108]. Taken together, these studies suggest that aromatase inhibition may reverse PH and that metformin could be used to treat PH by inhibiting aromatase activity.

It is well-known that the incidence of cardiovascular events is significantly lower in women when compared to age-matched men even following menopause [109]. In postmenopausal women, peripheral aromatase activity becomes the primary source of E2 synthesis, and studies indicate that aromatase activity may also increase with age. Plasma concentrations of cytokines known to stimulate aromatase activity were shown to increase over time in males and females [87]. Differences in aromatase activity between males and females may contribute to sex biases even after menopause. Aromatase inhibitors, which are used clinically for treatment of breast cancer,

have been shown to carry the potential risk of increased cardiovascular events. One clinical study found that treatment with aromatase inhibitors is associated with a higher risk of ischemic heart disease versus tamoxifen, a broad ER antagonist [110]. Another population-based observational study of a heterogeneous population of 74 women with early breast cancer who received adjuvant hormonal therapy and subsequently underwent cardiac angiography concluded that aromatase inhibitors significantly increased the hazard for CAD compared to tamoxifen [111]. These studies, however, are in direct conflict with recent genome-wide association studies that attribute this finding to the cardioprotective effects of tamoxifen versus the detrimental effects of aromatase inhibitors [112].

Interestingly, the concurrent use of aromatase inhibitors and adjuvant radiotherapy for early-stage left-sided breast cancer, in the absence of tamoxifen treatment, yielded a significant decrease in RV systolic function and LV diastolic function when compared to radiotherapy alone [113]. While more long-term studies are necessary, these findings indicate that aromatase inhibition may lead to reduced cardioprotection in females.

Aromatase inhibition may increase the risk of CVD in men as well. A recent intervention pilot study by De Smet et al. investigated the effect of aromatase inhibition on the cardiovascular system on young healthy men 20–40 years old [114]. The study employed 20 participants who were randomized to the aromatase inhibitor group (letrozole, 2.5 mg daily for 7 days) or letrozole plus an E2 patch (75 µg/day for 7 days). The letrozole plus E2 group, which yielded a similar sex steroid profile to that observed in obese men, demonstrated a statistically significant decrease in circumferential strain indicating subclinically impaired function of the LV subendocardial layers [114].

In summary, the aromatase enzyme converts androgens to E2 in many extragonadal tissues. While the bulk of experiments suggest that E2 is cardioprotective, the effect of increased aromatase activity remains controversial. In vasculature, aromatase activity seems to protect against the development of atherosclerosis, and aromatase null mice exhibit increased adiposity and insulin resistance which are risk factors in the development of CVD. Conversely, aromatase null mice also exhibit increased calcium handling and better functional recovery when subjected to I/R injury. In direct contradiction to this finding, aromatase overexpression resulted in better functional recovery following I/R in male mice but not in female mice. Interestingly, aromatase expression is downregulated in HF, yet inhibition of aromatase activity helps to reverse RV HF and hypertrophy in rodents. In humans, the use of aromatase inhibitor in women and men may increase the risk of developing CVD or CVD

risk factors. These findings, while contradictory, demonstrate that aromatase may play a role in mediating cardioprotection; however, more research is needed to investigate the nature of aromatase role.

Controversies regarding the use of estrogen for therapeutic purposes

While the use of E2 therapy in CVD has shown predominantly beneficial effects in animal models, the use of E2 in humans in the form of hormone replacement therapy has been very controversial. The following are the main controversial aspects of hormonal therapy in humans.

Efficacy of hormone replacement therapy

Two large-scale randomized, double-blind, placebo-controlled clinical trials, the Women's Health Initiative (WHI) and Heart and Estrogen/progestin Replacement Study (HERS), were designed to examine the efficacy of hormone replacement therapy (HRT) in reducing the risk of ischemic heart disease in postmenopausal women [115]. In WHI trials, postmenopausal women (50–79 years old) without a history of CVD were administered a daily dose of conjugated equine E2 (CEE, 0.625 mg) plus medroxyprogesterone acetate (MPA, 2.5 mg). Unfortunately, the WHI trial was discontinued prematurely due to adverse cardiac events and increased risk of breast cancer [116]. The HERS trial, conducted in postmenopausal women with pre-existing CVD (44–79 years old), found that daily use of CEE (0.625 mg) plus MPA (2.5 mg) had no effect on primary or secondary cardiovascular outcomes [117, 118].

The failure of both trials to support the protective role of HRT in reducing CVD could, in part, be due to the HRT being administered long after the onset of menopause in most of the women enrolled. A new hypothesis deemed the “Critical Window of Hormone Therapy” has recently received a lot of attention. The hypothesis supports the view that HRT could be effective if started early at the onset of menopause [119]. A more recent randomized study that administered HRT to women (42–58 years old) within 6–36 months following the onset of menopause supports the “Critical Window” hypothesis as certain risk factors for CVD were reduced [120]. Another randomized study in women (45–58 years old) also showed a significantly reduced risk of mortality, heart failure, or myocardial infarction if HRT was initiated early after menopause [121].

Another potential reason for the failure of the WHI and HERS trials could be that the hormone dosage and the combination of E2 and progestin employed in WHI and HERS may not have been optimal for preventing or reducing CVD risk. CEE administered alone at a lower dose (0.3 mg/day) was shown to decrease major coronary events in women [122] and coronary artery

atherosclerosis in adult female monkeys [123]. Progestins, however, have been shown to downregulate ERs and stimulate direct progestin receptor-mediated effects that oppose estrogenic action [116]. MPA, a progestin administered in conjunction with E2 in the WHI and HERS trial, may also negate any cardioprotective effects of E2 therapy. A randomized placebo-controlled crossover study showed, however, that oral administration of E2 for 3 months without progestin does not improve coronary flow reserve in postmenopausal women [124].

Currently, the Kronos Early Estrogen Prevention Study (KEEPS) is an ongoing trial testing the effect of HRT in atherosclerosis and coronary calcification in women within 3 years of menopause. KEEPS will study the effect of HRT with or without progestin as well as test different modes of E2 administration (CEE or transdermal E2) [125].

Finally, disparity between the efficacy of HRT in animal studies and the WHI and HERS clinical trials could be due to the downregulation of ERs in aged women. Li et al. found that the expression of ER β and GPR30, but not ER α , are significantly decreased in the arteries of female rats [126]. Since most effects of estrogen are mediated through its receptors, downregulation of ERs could silence the beneficial effects of E2. Further studies are needed to examine the expression of ERs in aged women and identify if the efficacy of HRT changes with age.

In summary, despite numerous animal studies demonstrating the beneficial cardioprotective effects of E2, the WHI and HERS clinical trials both failed to support the effectiveness of HRT in reducing CVD. However, this may be attributed to several factors including the initiation of HRT long after the start of menopause, the dose, and combination of E2 and progestin. The ongoing large randomized KEEPS trial, which administers E2 with and without progestin early after the onset of menopause, has the potential to shed valuable light on the efficacy of HRT in reducing CVD in early postmenopausal women.

Arrhythmias, sudden cardiac death, and estrogen

Sex differences exist in the underlying electrophysiology of the heart that may influence the risk of developing arrhythmia and sudden cardiac death. It is likely that these differences are modulated by sex hormones and that sex hormones also affect the presentation of these arrhythmias and the response to antiarrhythmic drugs and therapies [127]. In fact, women have a higher risk of acquired long QT syndrome from antiarrhythmic drugs than men. Thus, understanding the effects of hormones in mediating arrhythmic risk and response to treatment is important in order to provide effective and individualized treatment for patients.

In experimental models of CVD, the role of E2 in reducing ventricular arrhythmias has been controversial.

In support of antiarrhythmic effects of E2, Philp et al. examined the effect of E2 on ischemia-induced cardiac arrhythmias and found that administration of E2 before coronary artery ligation exhibited antiarrhythmic activity in a dose-dependent manner in female rats [128]. Interestingly, the antiarrhythmic activity of E2 in the myocardial ischemia setting was greater in female rats than in male rats [128], which could be due to the effects of E2 on the expression and function of ion channels that control cardiac cell excitation and repolarization [129]. Sykes et al. showed that inhibition of E2 synthesis with an aromatase inhibitor triggered cardiac arrhythmias in the developing zebrafish [130]. While these data suggest that E2 is antiarrhythmic, several studies demonstrate that E2 actually promotes ventricular arrhythmias. For example, female rats treated with E2 and subjected to ex vivo ischemia experienced an increase in sustained ventricular arrhythmia upon reperfusion [131]. The proarrhythmic action of E2 was due to alterations of calcium handling in cardiomyocytes mediated through ER β signaling [131]. Interestingly, while E2 increased the incidence and duration of ventricular arrhythmia, the infarct size post-I/R was significantly reduced which highlights the cardioprotective role of E2 in I/R injury [131]. In an in vivo model of drug-induced arrhythmia, E2 was also found to trigger ventricular arrhythmias in both male and female rabbits [132]. Odening et al. showed that E2 also increases ventricular arrhythmias and sudden cardiac death in OVX transgenic long QT rabbits [133].

In the Oregon Sudden Unexpected Death Study (catchment population approximately one million), Narayanan et al. compared cases of sudden cardiac arrest (SCA) with matched controls. A multivariate analysis on testosterone and E2 levels measured from blood samples drawn at the time of the SCA event revealed that higher testosterone levels only in men were associated with lower SCA events. On the other hand, higher estradiol levels in both women and men were associated with higher SCA events [134].

In summary, in the experimental settings of CVD, the role of E2 in ventricular arrhythmias is controversial as E2 was found to be both pro- and antiarrhythmic in females or both sexes depending on the study. In humans, increased plasma E2 levels are found to be associated with higher SCA events in both men and women. Since females are more prone to drug-induced arrhythmia than males [135], further studies are needed to examine whether the female heart has a unique sensitivity to E2 in the presence of certain drugs.

Estrogen and thrombosis risks

The use of E2 as a therapeutic agent for cardiovascular protection is controversial as the use of oral contraceptives containing E2 has been linked to an increased risk

of venous thrombosis which can lead to MI, stroke, and peripheral artery disease. While the intent of the WHI clinical trials was to investigate the role of E2 in postmenopausal women, the investigators did find that the estrogen component of HRT was also associated with increased risk of venous thrombosis. Canonico et al. reported a similar finding [136]. These studies suggest that E2 may have prothrombotic ability, and while controversial and not fully understood, E2 may achieve this by increasing procoagulant factors (factors VII, X, XII, and XIII) and decreasing anticoagulant factors (protein S and antithrombin) [137]. Thus, E2 supplementation via oral contraceptives or HRT enhances the risk of venous thrombosis especially in women with pre-existing coagulation abnormalities. Progestin also contributes to this risk as oral contraceptives or HRT containing third-generation progestogens (desogestrel or gestodene) are associated with a higher risk of venous thrombosis than those containing second-generation progestogens (such as levonorgestrel) [138].

A study by Nayak et al. found that tamoxifen, a well-established cancer drug that reduces the effects of E2, has an antithrombotic effect in mice and human blood samples. In this study, tamoxifen significantly inhibited platelet functions in human plasma while it prolonged tail bleeding time and prevented thrombus formation at injured arterial walls in mice [139]. This study further demonstrates that ER modulation impacts coagulation.

While most evidence suggests that E2 has procoagulatory and prothrombotic effects, a clinical trial assessing HRT on cardiovascular events in recently postmenopausal women over 10 years did not reveal an increased risk of thrombotic events. Women receiving HRT early after menopause had a significantly reduced risk of CVD, without any apparent increase in risk of cancer, venous thromboembolism, or stroke [121].

In summary, E2 treatment in the form of oral contraceptives and HRT have been linked to increased risk of venous thrombosis, possibly due to E2's modulatory role on procoagulant and anticoagulant factors. Research demonstrating that tamoxifen exhibits antithrombotic effects further suggests that ER modulation is involved in coagulation. A long-term clinical trial studying HRT in recently menopausal women, however, was not associated with increased risk of thrombotic events.

Conclusions

The protection against cardiovascular disease in women during reproductive age is believed to be related at least in part to E2 since endogenous levels of E2 and the expression of ERs differ considerably between sexes. E2 mediates its cardioprotective actions by increasing angiogenesis and vasodilation and decreasing ROS, oxidative stress, and fibrosis. Through these mechanisms,

E2 limits cardiac remodeling and attenuates heart hypertrophy (Fig. 2). Although the use of E2 as a therapeutic agent in humans has remained controversial, targeting specific ERs in the cardiovascular system may result in novel and possibly safer therapeutic options for the use of E2 for cardiovascular protection.

Abbreviations

ACE: Angiotensin-converting enzyme; ADAM: Disintegrin-metalloproteinase; ALDH2: Aldehyde dehydrogenase 2; AngII: Angiotensin II; Ap1: Activator protein-1; ArKO: Aromatase knockout; AROM+: Mice in which aromatase was transgenically upregulated; CAD: Coronary artery disease; CEE: Conjugated equine estrogens; CSE: Cystathionine γ -lyase; CVD: Cardiovascular disease; E2: 17 β -estradiol, estrogen; EC: Endothelial cells; ECM: Extracellular matrix; eNOS: Nitric oxide synthase 3; EPC: Endothelial progenitor cells; ER: Estrogen receptors; ERKO: Estrogen receptor α knockout; ER α : Estrogen receptor α ; ER β : Estrogen receptor β ; ET-1: Endothelin-1; FGF: Fibroblast growth factor; GPR30: G-protein-coupled estrogen receptor; HERS: Heart and Estrogen/progestin Replacement Study; HF: Heart failure; HRT: Hormone replacement therapy; I/R: Ischemia/reperfusion; KEEPS: Kronos Early Estrogen Prevention Study; LAD: Left anterior descending; LV: left ventricular; MI: Myocardial infarction; miR-22: MicroRNA 22; MMP: matrix metalloproteinases; MPA: Medroxyprogesterone acetate; mPTP: Mitochondrial permeability transition pore; NO: Nitric oxide; OVX: Ovariectomy; PH: Pulmonary hypertension; RAMP3: Receptor active modifying protein 3; ROS: Reactive oxygen species; RV: Right ventricular; SCA: Sudden cardiac arrest; SMC: Smooth muscle cells; SOD2: Superoxide dismutase 2; Sp1: Specificity protein-1; O $_2^{\cdot -}$: Superoxide; TBARS: Thiobarbituric acid reactive substances; TGF: Transforming growth factor- β 1; TGF β : Vascular endothelial growth factor; VEGFR-2: VEGF binding to its receptor-2; WT: Wild type; WHI: Women's Health Initiative

Acknowledgements

None

Funding

This work was supported by NIH and NRSA training grants T32-GM065823 and HD00722 (Iorga), AHA-17PRE33420159 (Cunningham), and NIH R01HL131182-01A1 and AHA16GRANT27760058 (Eghbali).

Availability of data and materials

Not applicable

Authors' contributions

AI, CC, SU, GR, and ME contributed to the conception and design of the study. AI, CC, and SM drafted the article. AI, CC, SM, SU, and ME revised critically the article for important intellectual content. GR generated the figures. All authors approved this version of the manuscript.

Ethics approval

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Anesthesiology, Division of Molecular Medicine, David Geffen School of Medicine at University of California, Los Angeles, BH-160CHS, Los Angeles, CA 90095-7115, USA. ²Present address: Department of Medicine, Division of Gastroenterology/Liver, Keck School of Medicine of the University of Southern California, Los Angeles, CA 90033, USA.

Received: 23 May 2017 Accepted: 4 October 2017

Published online: 24 October 2017

References

- Wake R, Yoshiyama M. Gender differences in ischemic heart disease. *Recent Patents Cardiovasc Drug Discov*. 2009;4:234–40.
- Kararigas G, Dworatzek E, Petrov G, Summer H, Schulze TM, Baczko I, et al. Sex-dependent regulation of fibrosis and inflammation in human left ventricular remodelling under pressure overload. *Eur J Heart Fail*. 2014;16:1160–7.
- Hayward CS, Kelly RP, Collins P. The roles of gender, the menopause and hormone replacement on cardiovascular function. *Cardiovasc Res*. 2000;46:28–49.
- Parker WH, Jacoby V, Shoupe D, Rocca W. Effect of bilateral oophorectomy on women's long-term health. *Womens Health Lond Engl*. 2009;5:565–76.
- Menazza S, Murphy E. The expanding complexity of estrogen receptor signaling in the cardiovascular system. *Circ Res*. 2016;118:994–1007.
- Arnold AP, Cassis LA, Eghbali M, Reue K, Sandberg K. Sex hormones and sex chromosomes cause sex differences in the development of cardiovascular diseases. *Arterioscler. Thromb. Vasc. Biol*. 2017;ATVBAHA.116.307301.
- Arnold AP, Reue K, Eghbali M, Vilain E, Chen X, Ghahramani N, et al. The importance of having two X chromosomes. *Philos Trans R Soc Lond Ser B Biol Sci*. 2016;37(5):746–56. <https://doi.org/10.1161/ATVBAHA.116.307301>.
- Li J, Chen X, McClusky R, Ruiz-Sundstrom M, Itoh Y, Umar S, et al. The number of X chromosomes influences protection from cardiac ischaemia/reperfusion injury in mice: one X is better than two. *Cardiovasc Res*. 2014;102:375–84.
- Pharmacodynamic effects of the fetal estrogen estrorel in... : menopause [Internet]. LWW. [cited 2017 Aug 24]. Available from: http://journals.lww.com/menopausejournal/Fulltext/2017/06000/Pharmacodynamic_effects_of_the_fetal_estrogen.13.aspx
- Simpson ER. Sources of estrogen and their importance. *J Steroid Biochem Mol Biol*. 2003;86:225–30.
- Bopassa JC, Eghbali M, Toro L, Stefani E. A novel estrogen receptor GPER inhibits mitochondria permeability transition pore opening and protects the heart against ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol*. 2010;298:H16–23.
- Deschamps AM, Murphy E, Sun J. Estrogen receptor activation and cardioprotection in ischemia reperfusion injury. *Trends Cardiovasc. Med*. 2010;20:73–8.
- Babiker FA, De Windt LJ, van Eickels M, Thijssen V, Bronsaeer RJP, Grohé C, et al. 17beta-estradiol antagonizes cardiomyocyte hypertrophy by autocrine/paracrine stimulation of a guanylyl cyclase A receptor-cyclic guanosine monophosphate-dependent protein kinase pathway. *Circulation*. 2004;109:269–76.
- Ropero AB, Eghbali M, Minosyan TY, Tang G, Toro L, Stefani E. Heart estrogen receptor alpha: distinct membrane and nuclear distribution patterns and regulation by estrogen. *J Mol Cell Cardiol*. 2006;41:496–510.
- Grohé C, Kahlert S, Löbbeck K, Stimpel M, Karas RH, Vetter H, et al. Cardiac myocytes and fibroblasts contain functional estrogen receptors. *FEBS Lett*. 1997;416:107–12. IPresented in part at the 67th Scientific Sessions of the American Heart Association, Dallas, 14–17 November 1994.
- Chen J-Q, Yager JD, Russo J. Regulation of mitochondrial respiratory chain structure and function by estrogens/estrogen receptors and potential physiological/pathophysiological implications. *Biochim Biophys Acta BBA - Mol Cell Res*. 2005;1746:1–17.
- Chen L, Knowlton AA. Mitochondria and heart failure: new insights into an energetic problem. *Minerva Cardioangiol*. 2010;58:213–29.
- Klinge CM. Estrogenic control of mitochondrial function and biogenesis. *J Cell Biochem*. 2008;105:1342–51.
- Klinge CM. Estrogens regulate life and death in mitochondria. *J Bioenerg Biomembr*. 2017;1–18.
- Lagranha CJ, Deschamps A, Aponte A, Steenbergen C, Murphy E. Sex differences in the phosphorylation of mitochondrial proteins result in reduced production of reactive oxygen species and cardioprotection in females. *Circ Res*. 2010;106:1681–91.
- Chen C-H, Budas GR, Churchill EN, Disatnik M-H, Hurlley TD, Mochly-Rosen D. Activation of aldehyde dehydrogenase-2 reduces ischemic damage to the heart. *Science*. 2008;321:1493–5.
- Zhai P, Eurell TE, Cooke PS, Lubahn DB, Gross DR. Myocardial ischemia-reperfusion injury in estrogen receptor-alpha knockout and wild-type mice. *Am J Physiol Heart Circ Physiol*. 2000;278:H1640–7.
- Booth EA, Obeid NR, Lucchesi BR. Activation of estrogen receptor-alpha protects the in vivo rabbit heart from ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol*. 2005;289:H2039–47.
- Filardo EJ. Epidermal growth factor receptor (EGFR) transactivation by estrogen via the G-protein-coupled receptor, GPR30: a novel signaling pathway with potential significance for breast cancer. *J Steroid Biochem Mol Biol*. 2002;80:231–8.
- Zoratti M, Szabó I. The mitochondrial permeability transition. *Biochim Biophys Acta*. 1995;1241:139–76.
- Lemasters JJ, Qian T, He L, Kim J-S, Elmore SP, Cascio WE, et al. Role of mitochondrial inner membrane permeabilization in necrotic cell death, apoptosis, and autophagy. *Antioxid Redox Signal*. 2002;4:769–81.
- Pavón N, Martínez-Abundis E, Hernández L, Gallardo-Pérez JC, Alvarez-Delgado C, Cerbón M, et al. Sexual hormones: effects on cardiac and mitochondrial activity after ischemia-reperfusion in adult rats. *Gender difference*. *J Steroid Biochem. Mol. Biol.* 2012;132:135–46.
- Pavón N, Cabrera-Orefice A, Gallardo-Pérez JC, Uribe-Alvarez C, Rivero-Segura NA, Vazquez-Martínez ER, et al. In female rat heart mitochondria, oophorectomy results in loss of oxidative phosphorylation. *J Endocrinol*. 2017;232:221–35.
- Wang L, Tang Z-P, Zhao W, Cong B-H, Lu J-Q, Tang X-L, et al. MiR-22/Sp-1 links estrogens with the up-regulation of cystathionine γ -lyase in myocardium, which contributes to estrogenic cardioprotection against oxidative stress. *Endocrinology*. 2015;156:2124–37.
- Zhu X, Tang Z, Cong B, Du J, Wang C, Wang L, et al. Estrogens increase cystathionine- γ -lyase expression and decrease inflammation and oxidative stress in the myocardium of ovariectomized rats. *Menopause N. Y. N.* 2013;20:1084–91.
- Wang F, He Q, Sun Y, Dai X, Yang X-P. Female adult mouse cardiomyocytes are protected against oxidative stress. *Hypertension*. Dallas Tex 1979. 2010;55:1172–8.
- Manabe I, Shindo T, Nagai R. Gene expression in fibroblasts and fibrosis: involvement in cardiac hypertrophy. *Circ Res*. 2002;91:1103–13.
- Swynghedauw B. Molecular mechanisms of myocardial remodeling. *Physiol Rev*. 1999;79:215–62.
- Sabbah HN, Sharov VG, Lesch M, Goldstein S. Progression of heart failure: a role for interstitial fibrosis. *Mol Cell Biochem*. 1995;147:29–34.
- Weber KT, Sun Y, Bhattacharya SK, Ahokas RA, Gerling IC. Myofibroblast-mediated mechanisms of pathological remodeling of the heart. *Nat Rev Cardiol*. 2013;10:15–26.
- Maisch B. Extracellular matrix and cardiac interstitium: restriction is not a restricted phenomenon. *Herz*. 1995;20:75–80.
- Lee HW, Eghbali-Webb M. Estrogen enhances proliferative capacity of cardiac fibroblasts by estrogen receptor- and mitogen-activated protein kinase-dependent pathways. *J Mol Cell Cardiol*. 1998;30:1359–68.
- Dubey RK, Gillespie DG, Jackson EK, Keller PJ. 17Beta-estradiol, its metabolites, and progesterone inhibit cardiac fibroblast growth. *Hypertension*. Dallas Tex 1979. 1998;31:522–8.
- Pedram A, Razandi M, Narayanan R, Levin ER. Estrogen receptor beta signals to inhibition of cardiac fibrosis. *Mol Cell Endocrinol*. 2016;434:57–68.
- Iorga A, Li J, Sharma S, Umar S, Bopassa JC, Nadadur RD, et al. Rescue of Pressure Overload-Induced Heart Failure by estrogen therapy. *J Am Heart Assoc*. 2016;5. 5(1): e002482. doi: <https://doi.org/10.1161/JAHA.115.002482>
- Mahmoodzadeh S, Leber J, Zhang X, Jaisser F, Messaoudi S, Morano I, et al. Cardiomyocyte-specific estrogen receptor alpha increases angiogenesis, lymphangiogenesis and reduces fibrosis in the female mouse heart post-myocardial infarction. *J Cell Sci Ther*. 2014;5:153.
- Schuster I, Mahmoodzadeh S, Dworatzek E, Jaisser F, Messaoudi S, Morano I, et al. Cardiomyocyte-specific overexpression of oestrogen receptor β improves survival and cardiac function after myocardial infarction in female and male mice. *Clin Sci*. 2016;130:365–76.
- Wang H, Zhao Z, Lin M, Groban L. Activation of GPR30 inhibits cardiac fibroblast proliferation. *Mol Cell Biochem*. 2015;405:135–48.
- Mahmoodzadeh S, Dworatzek E, Fritschka S, Pham TH, Regitz-Zagrosek V. 17Beta-estradiol inhibits matrix metalloproteinase-2 transcription via MAP kinase in fibroblasts. *Cardiovasc Res*. 2010;85:719–28.
- Peterson JT, Li H, Dillon L, Bryant JW. Evolution of matrix metalloprotease and tissue inhibitor expression during heart failure progression in the infarcted rat. *Cardiovasc Res*. 2000;46:307–15.
- Spinale FG, Coker ML, Thomas CV, Walker JD, Mukherjee R, Hebban L. Time-dependent changes in matrix metalloproteinase activity and expression during the progression of congestive heart failure: relation to ventricular and myocyte function. *Circ Res*. 1998;82:482–95.
- Thomas CV, Coker ML, Zellner JL, Handy JR, Crumbley AJ, Spinale FG. Increased matrix metalloproteinase activity and selective upregulation in LV myocardium from patients with end-stage dilated cardiomyopathy. *Circulation*. 1998;97:1708–15.

48. Bergman MR, Teerlink JR, Mahimkar R, Li L, Zhu B-Q, Nguyen A, et al. Cardiac matrix metalloproteinase-2 expression independently induces marked ventricular remodeling and systolic dysfunction. *Am J Physiol Heart Circ Physiol*. 2007;292:H1847–60.
49. Hayashidani S, Tsutsui H, Ikeuchi M, Shiomi T, Matsusaka H, Kubota T, et al. Targeted deletion of MMP-2 attenuates early LV rupture and late remodeling after experimental myocardial infarction. *Am J Physiol Heart Circ Physiol*. 2003;285:H1229–35.
50. Matsumura S, Iwanaga S, Mochizuki S, Okamoto H, Ogawa S, Okada Y. Targeted deletion or pharmacological inhibition of MMP-2 prevents cardiac rupture after myocardial infarction in mice. *J Clin Invest*. 2005;115:599–609.
51. Matsusaka H, Ide T, Matsushima S, Ikeuchi M, Kubota T, Sunagawa K, et al. Targeted deletion of matrix metalloproteinase 2 ameliorates myocardial remodeling in mice with chronic pressure overload. *Hypertension*. Dallas Tex 1979. 2006;47:711–7.
52. Nadadur RD, Umar S, Wong G, Eghbali M, Iorga A, Matori H, et al. Reverse right ventricular structural and extracellular matrix remodeling by estrogen in severe pulmonary hypertension. *J Appl Physiol*. Bethesda Md 1985. 2012;113:149–58.
53. Ardelt AA, McCullough LD, Korach KS, Wang MM, Munzenmaier DH, Hurn PD. Estradiol regulates angiotensin-1 mRNA expression through estrogen receptor-alpha in a rodent experimental stroke model. *Stroke*. 2005;36:337–41.
54. Elkin M, Orgel A, Kleinman HK. An angiogenic switch in breast cancer involves estrogen and soluble vascular endothelial growth factor receptor 1. *J Natl Cancer Inst*. 2004;96:875–8.
55. Rodger FE, Young FM, Fraser HM, Illingworth PJ. Endothelial cell proliferation follows the mid-cycle luteinizing hormone surge, but not human chorionic gonadotrophin rescue, in the human corpus luteum. *Hum Reprod Oxf Engl*. 1997;12:1723–9.
56. Sano M, Minamoto T, Toko H, Miyauchi H, Orimo M, Qin Y, et al. p53-induced inhibition of Hif-1 causes cardiac dysfunction during pressure overload. *Nature*. 2007;446:444–8.
57. Shiojima I, Sato K, Izumiya Y, Schiekofe S, Ito M, Liao R, et al. Disruption of coordinated cardiac hypertrophy and angiogenesis contributes to the transition to heart failure. *J Clin Invest*. 2005;115:2108–18.
58. Sutendra G, Dromparis P, Paulin R, Zervopoulos S, Haromy A, Nagendran J, et al. A metabolic remodeling in right ventricular hypertrophy is associated with decreased angiogenesis and a transition from a compensated to a decompensated state in pulmonary hypertension. *J Mol Med Berl Ger*. 2013;91:1315–27.
59. Khurana R, Simons M. Insights from angiogenesis trials using fibroblast growth factor for advanced arteriosclerotic disease. *Trends Cardiovasc Med*. 2003;13:116–22.
60. Stegmann TJ. FGF-1: a human growth factor in the induction of neoangiogenesis. *Expert Opin Investig Drugs*. 1998;7:2011–5.
61. Flamme I, Breier G, Risau W. Vascular endothelial growth factor (VEGF) and VEGF receptor 2 (flk-1) are expressed during vasculogenesis and vascular differentiation in the quail embryo. *Dev Biol*. 1995;169:699–712.
62. Kroll J, Waltenberger J. VEGF-A induces expression of eNOS and iNOS in endothelial cells via VEGF receptor-2 (KDR). *Biochem Biophys Res Commun*. 1998;252:743–6.
63. Umar S, Iorga A, Matori H, Nadadur RD, Li J, Maltese F, et al. Estrogen rescues preexisting severe pulmonary hypertension in rats. *Am J Respir Crit Care Med*. 2011;184:715–23.
64. Iwakura A, Shastry S, Luedemann C, Hamada H, Kawamoto A, Kishore R, et al. Estradiol enhances recovery after myocardial infarction by augmenting incorporation of bone marrow-derived endothelial progenitor cells into sites of ischemia-induced neovascularization via endothelial nitric oxide synthase-mediated activation of matrix metalloproteinase-9. *Circulation*. 2006;113:1605–14.
65. Hamada H, Kim MK, Iwakura A, Li M, Thorne T, Qin G, et al. Estrogen receptors alpha and beta mediate contribution of bone marrow-derived endothelial progenitor cells to functional recovery after myocardial infarction. *Circulation*. 2006;114:2261–70.
66. Hadi HA, Carr CS, Al SJ. Endothelial dysfunction: cardiovascular risk factors, therapy, and outcome. *Vasc Health Risk Manag*. 2005;1:183–98.
67. Gimbrone MA. Vascular endothelium: an integrator of pathophysiologic stimuli in atherosclerosis. *Am J Cardiol*. 1995;75:67B–70B.
68. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res*. 2000;87:840–4.
69. Khazaei M, Moien-Afshari F, Laher I. Vascular endothelial function in health and diseases. *Pathophysiol Off J Int Soc Pathophysiol*. 2008;15:49–67.
70. Chakrabarti S, Davidge ST. G-protein coupled receptor 30 (GPR30): a novel regulator of endothelial inflammation. *PLoS One*. 2012;7:e2357.
71. Zhou L, Chen H, Mao X, Qi H, Baker PN, Zhang H. G-protein-coupled receptor 30 mediates the effects of estrogen on endothelial cell tube formation in vitro. *Int J Mol Med*. 2017;39:1461–7.
72. Guo X, Razandi M, Pedram A, Kassab G, Levin ER. Estrogen induces vascular wall dilation: mediation through kinase signaling to nitric oxide and estrogen receptors alpha and beta. *J Biol Chem*. 2005;280:19704–10.
73. White RE, Han G, Dimitropoulou C, Zhu S, Miyake K, Fulton D, et al. Estrogen-induced contraction of coronary arteries is mediated by superoxide generated in vascular smooth muscle. *Am J Physiol Heart Circ Physiol*. 2005;289:H1468–75.
74. Pelzer T, Jazbutyte V, Hu K, Segerer S, Nahrendorf M, Nordbeck P, et al. The estrogen receptor-alpha agonist 16alpha-LE2 inhibits cardiac hypertrophy and improves hemodynamic function in estrogen-deficient spontaneously hypertensive rats. *Cardiovasc Res*. 2005;67:604–12.
75. Widder J, Pelzer T, von Poser-Klein C, Hu K, Jazbutyte V, Fritzsche K-H, et al. Improvement of endothelial dysfunction by selective estrogen receptor-alpha stimulation in ovariectomized SHR. *Hypertension*. Dallas Tex 1979. 2003;42:991–6.
76. Jazbutyte V, Arias-Loza PA, Hu K, Widder J, Govindaraj V, von Poser-Klein C, et al. Ligand-dependent activation of ER[beta] lowers blood pressure and attenuates cardiac hypertrophy in ovariectomized spontaneously hypertensive rats. *Cardiovasc Res*. 2008;77:774–81.
77. Xue B, Pamidimukkala J, Lubahn DB, Hay M. Estrogen receptor-alpha mediates estrogen protection from angiotensin II-induced hypertension in conscious female mice. *Am J Physiol Heart Circ Physiol*. 2007;292:H1770–6.
78. Gabel SA, Walker VR, London RE, Steenbergen C, Korach KS, Murphy E. Estrogen receptor beta mediates gender differences in ischemia/reperfusion injury. *J Mol Cell Cardiol*. 2005;38:289–97.
79. Pelzer T, Loza P-AA HK, Bayer B, Dienesch C, Calvillo L, et al. Increased mortality and aggravation of heart failure in estrogen receptor-beta knockout mice after myocardial infarction. *Circulation*. 2005;111:1492–8.
80. Nikolic I, Liu D, Bell JA, Collins J, Steenbergen C, Murphy E. Treatment with an estrogen receptor-beta-selective agonist is cardioprotective. *J Mol Cell Cardiol*. 2007;42:769–80.
81. Lin J, Steenbergen C, Murphy E, Sun J. Estrogen receptor-beta activation results in S-nitrosylation of proteins involved in cardioprotection. *Circulation*. 2009;120:245–54.
82. Lindsey SH, Carver KA, Prossnitz ER, Chappell MC. Vasodilation in response to the GPR30 agonist G-1 is not different from estradiol in the mRen2.Lewis female rat. *J Cardiovasc Pharmacol*. 2011;57:598–603.
83. Fredette NC, Meyer MR, Prossnitz ER. Role of GPER in estrogen-dependent nitric oxide formation and vasodilation. *J Steroid Biochem Mol Biol*. [Internet]. 2017; Available from: <http://www.sciencedirect.com/science/article/pii/S0960076017301309>
84. Lenhart PM, Broselid S, Barrick CJ, Leeb-Lundberg LMF, Caron KM. G-protein coupled receptor 30 interacts with receptor activity modifying protein 3 and confers sex-dependent cardioprotection. *J Mol Endocrinol*. 2013;51:191–202.
85. Sudhir K, Chou TM, Mullen WL, Hausmann D, Collins P, Yock PG, et al. Mechanisms of estrogen-induced vasodilation: in vivo studies in canine coronary conductance and resistance arteries. *J Am Coll Cardiol*. 1995;26:807–14.
86. Gallagher PE, Li P, Lenhart JR, Chappell MC, Brosnihan KB. Estrogen regulation of angiotensin-converting enzyme mRNA. *Hypertension*. Dallas Tex 1979. 1999;33:323–8.
87. Purohit A, Reed MJ. Regulation of estrogen synthesis in postmenopausal women. *Steroids*. 2002;67:979–83.
88. Bulun SE, Sebastian S, Takayama K, Suzuki T, Sasano H, Shozu M. The human CYP19 (aromatase P450) gene: update on physiologic roles and genomic organization of promoters. *J Steroid Biochem Mol Biol*. 2003;86:219–24.
89. Harada N, Sasano H, Murakami H, Ohkuma T, Nagura H, Takagi Y. Localized expression of aromatase in human vascular tissues. *Circ Res*. 1999;84:1285–91.
90. Jazbutyte V, Stumpner J, Redel A, Lorenzen JM, Roewer N, Thum T, et al. Aromatase inhibition attenuates desflurane-induced preconditioning against acute myocardial infarction in male mouse heart in vivo. *PLoS One*. 2012;7:e42032.
91. Wang H, Li Q, Wang T, Yang G, Wang Y, Zhang X, et al. A common polymorphism in the human aromatase gene alters the risk for polycystic ovary syndrome and modifies aromatase activity in vitro. *Mol Hum Reprod*. 2011;17:386–91.
92. Zhang X-L, Zhang C-W, Xu P, Liang F-J, Che Y-N, Xia Y-J, et al. SNP rs2470152 in CYP19 is correlated to aromatase activity in Chinese polycystic ovary syndrome patients. *Mol Med Rep*. 2012;5:245–9.
93. Koudu Y, Onouchi T, Hosoi T, Horiuchi T. Association of CYP19 gene polymorphism with vertebral fractures in Japanese postmenopausal women. *Biochem Genet*. 2012;50:389–96.

94. Ma CX, Adjei AA, Salavaggione OE, Coronel J, Pellemounter L, Wang L, et al. Human aromatase: gene resequencing and functional genomics. *Cancer Res.* 2005;65:11071–82.
95. Wang B, Fu Z-Y, Ma Y-T, Huang D, Liu F, Dong C-L, et al. Identification of a CYP19 gene single-nucleotide polymorphism associated with a reduced risk of coronary heart disease. *Genet Test Mol Biomark.* 2016;20:2–10.
96. Bayard F, Clamens S, Meggetto F, Blaes N, Delsol G, Faye JC. Estrogen synthesis, estrogen metabolism, and functional estrogen receptors in rat arterial smooth muscle cells in culture. *Endocrinology.* 1995;136:1523–9.
97. Bayard F, Clamens S, Delsol G, Blaes N, Maret A, Faye JC. Oestrogen synthesis, oestrogen metabolism and functional oestrogen receptors in bovine aortic endothelial cells. *CIBA Found Symp.* 1995;191:122–32. 138
98. Adams MR, Kaplan JR, Manuck SB, Koritnik DR, Parks JS, Wolfe MS, et al. Inhibition of coronary artery atherosclerosis by 17-beta estradiol in ovariectomized monkeys. Lack of an effect of added progesterone. *Arterioscler. Dallas Tex.* 1990;10:1051–7.
99. Chen SJ, Li H, Durand J, Oparil S, Chen YF. Estrogen reduces myointimal proliferation after balloon injury of rat carotid artery. *Circulation.* 1996;93:577–84.
100. Sullivan TR, Karas RH, Aronovitz M, Faller GT, Ziar JP, Smith JJ, et al. Estrogen inhibits the response-to-injury in a mouse carotid artery model. *J Clin Invest.* 1995;96:2482–8.
101. Jones ME, Thorburn AW, Britt KL, Hewitt KN, Wreford NG, Proietto J, et al. Aromatase-deficient (ArKO) mice have a phenotype of increased adiposity. *Proc Natl Acad Sci U S A.* 2000;97:12735–40.
102. Takeda K, Toda K, Saibara T, Nakagawa M, Saika K, Onishi T, et al. Progressive development of insulin resistance phenotype in male mice with complete aromatase (CYP19) deficiency. *J Endocrinol.* 2003;176:237–46.
103. Bell JR, Mellor KM, Wollermann AC, Ip WTK, Reichelt ME, Meachem SJ, et al. Aromatase deficiency confers paradoxical postischemic cardioprotection. *Endocrinology.* 2011;152:4937–47.
104. Bell JR, Bernasochi GB, Wollermann AC, Raaijmakers AJA, Boon WC, Simpson ER, et al. Myocardial and cardiomyocyte stress resilience is enhanced in aromatase-deficient female mouse hearts through CaMKII δ activation. *Endocrinology.* 2015;156:1429–40.
105. Bell JR, Bernasochi GB, Varma U, Boon WC, Ellem SJ, Risbridger GP, et al. Aromatase transgenic upregulation modulates basal cardiac performance and the response to ischemic stress in male mice. *Am. J. Physiol. - heart circ. Physiol.* 2014;306:H1265–74.
106. Mair KM, Wright AF, Duggan N, Rowlands DJ, Hussey MJ, Roberts S, et al. Sex-dependent influence of endogenous estrogen in pulmonary hypertension. *Am J Respir Crit Care Med.* 2014;190:456–67.
107. Hadad SM, Hardie DG, Appleyard V, Thompson AM. Effects of metformin on breast cancer cell proliferation, the AMPK pathway and the cell cycle. *Clin Transl Oncol Off Publ Fed Span Oncol Soc Natl Cancer Inst Mex.* 2014;16:746–52.
108. Dean A, Nilsen M, Loughlin L, Salt IP, MacLean MR. Metformin reverses development of pulmonary hypertension via Aromatase inhibition. *Hypertension. Dallas Tex* 1979. 2016;68:446–54.
109. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics—2016 update: a report from the American Heart Association. *Circulation.* 2016;133:e38–360.
110. Abdel-Qadir H, Amir E, Fischer HD, Fu L, Austin PC, Harvey PJ, et al. The risk of myocardial infarction with aromatase inhibitors relative to tamoxifen in post-menopausal women with early stage breast cancer. *Eur J Cancer Oxf. Engl.* 1990. 2016;68:11–21.
111. Seruga B, Zadnik V, Kuhar CG, Marinko T, Cufur T, Zakotnik B, et al. Association of aromatase inhibitors with coronary heart disease in women with early breast cancer. *Cancer Investig.* 2014;32:99–104.
112. Khosrow-Khavar F, Filion KB, Al-Qurashi S, Torabi N, Bouganim N, Suissa S, et al. Cardiotoxicity of aromatase inhibitors and tamoxifen in postmenopausal women with breast cancer: a systematic review and meta-analysis of randomized controlled trials. *Ann Oncol Off J Eur Soc Med Oncol.* 2017;28:487–96.
113. Skyttä T, Tuohinen S, Virtanen V, Raatikainen P, Kellokumpu-Lehtinen P-L. The concurrent use of aromatase inhibitors and radiotherapy induces echocardiographic changes in patients with breast cancer. *Anticancer Res.* 2015;35:1559–66.
114. De Smet M a. J, Lapauw B, de backer T. Sex steroids in relation to cardiac structure and function in men. *Andrologia* 2017;49.
115. Grady D, Herrington D, Bittner V, Blumenthal R, Davidson M, Hlatky M, et al. Cardiovascular disease outcomes during 6.8 years of hormone therapy: heart and estrogen/progestin replacement study follow-up (HERS II). *JAMA.* 2002;288:49–57.
116. Sarrel PM. Risks and benefits of hormone replacement therapy for the prevention of cardiovascular disease. *Cardiol. Rome Italy.* 1999;44(Suppl 1):515–8.
117. Grady D, Applegate W, Bush T, Furberg C, Riggs B, Hulley SB. Heart and estrogen/progestin replacement study (HERS): design, methods, and baseline characteristics. *Control Clin Trials.* 1998;19:314–35.
118. Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and estrogen/progestin replacement study (HERS) research group. *JAMA.* 1998;280:605–13.
119. Maki PM. Critical window hypothesis of hormone therapy and cognition: a scientific update on clinical studies. *Menopause N Y N.* 2013;20:695–709.
120. Harman SM, Black DM, Naftolin F, Brinton EA, Budoff MJ, Cedars MI, et al. Arterial imaging outcomes and cardiovascular risk factors in recently menopausal women: a randomized trial. *Ann Intern Med.* 2014;161:249–60.
121. Schierbeck LL, Rejnmark L, Tofteng CL, Stilgren L, Eiken P, Mosekilde L, et al. Effect of hormone replacement therapy on cardiovascular events in recently postmenopausal women: randomised trial. *BMJ.* 2012;345:e6409.
122. Grodstein F, Chen J, Pollen DA, Albert MS, Wilson RS, Folstein MF, et al. Postmenopausal hormone therapy and cognitive function in healthy older women. *J Am Geriatr Soc.* 2000;48:746–52.
123. Appt SE, Kaplan JR, Clarkson TB, Cline JM, Christian PJ, Hoyer PB. Destruction of primordial ovarian follicles in adult cynomolgus macaques after exposure to 4-vinylcyclohexene diepoxide: a nonhuman primate model of the menopausal transition. *Fertil Steril.* 2006;86:1210–6.
124. Schwitzer J, Kozerke S, Bremerich J, Baltes C, Attenhofer Jost C, Birkhäuser M, et al. Oral administration of 17beta-estradiol over 3 months without progestin co-administration does not improve coronary flow reserve in post-menopausal women: a randomized placebo-controlled cross-over CMR study. *J Cardiovasc Magn Reson Off J Soc Cardiovasc Magn Reson.* 2007;9:665–72.
125. Wharton W, Gleason CE, Miller VM, Asthana S. Rationale and design of the Kronos Early Estrogen Prevention Study (KEEPS) and the KEEPS cognitive and affective sub study (KEEPS Cog). *Brain Res.* 2013;1514:12–7.
126. Li T, Xiao X, Zhang J, Zhu Y, Hu Y, Zang J, et al. Age and sex differences in vascular responsiveness in healthy and trauma patients: contribution of estrogen receptor-mediated rho kinase and PKC pathways. *Am J Physiol - Heart Circ Physiol.* 2014;306:H1105–15.
127. Bailey MS, Curtis AB. The effects of hormones on arrhythmias in women. *Curr Womens Health Rep.* 2002;2:83–8.
128. Philp KL, Hussain M, Byrne NF, Diver MJ, Hart G, Coker SJ. Greater antiarrhythmic activity of acute 17beta-estradiol in female than male anaesthetized rats: correlation with Ca $^{2+}$ channel blockade. *Br J Pharmacol.* 2006;149:233–42.
129. Odening KE, Koren G. How do sex hormones modify arrhythmogenesis in long QT syndrome? Sex hormone effects on arrhythmogenic substrate and triggered activity. *Heart Rhythm.* 2014;11:2107–15.
130. Sykes BG, Van Steyn PM, Vignali JD, Winalski J, Lozier J, Bell WE, et al. The relationship between estrogen and nitric oxide in the prevention of cardiac and vascular anomalies in the developing Zebrafish (*Danio Rerio*). *Brain Sci.* 2016;6
131. Yan S, Song W, Chen Y, Hong K, Rubinstein J, Wang H-S. Low-dose bisphenol A and estrogen increase ventricular arrhythmias following ischemia-reperfusion in female rat hearts. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res Assoc.* 2013;56:75–80.
132. Philp KL, Hart G, Coker SJ. A gender-independent proarrhythmic action of 17beta-estradiol in anaesthetized rabbits. *Eur J Pharmacol.* 2007;575:113–21.
133. Odening KE, Choi B-R, Liu GX, Hartmann K, Ziv O, Chaves L, et al. Estradiol promotes sudden cardiac death in transgenic long QT type 2 rabbits while progesterone is protective. *Heart Rhythm.* 2012;9:823–32.
134. Narayanan K, Havmoeller R, Reinier K, Jerger K, Teodorescu C, Uy-Evanado A, et al. Sex hormone levels in patients with sudden cardiac arrest. *Heart Rhythm.* 2014;11:2267–72.
135. Darpo B, Karnad DR, Badiilini F, Florian J, Garnett CE, Kothari S, et al. Are women more susceptible than men to drug-induced QT prolongation? Concentration-QTc modelling in a phase 1 study with oral rac-sotalol. *Br J Clin Pharmacol.* 2014;77:522–31.
136. Canonico M, Oger E, Plu-Bureau G, Conard J, Meyer G, Lévesque H, et al. Hormone therapy and venous thromboembolism among postmenopausal women: impact of the route of estrogen administration and progestogens: the ESTHER study. *Circulation.* 2007;115:840–5.
137. Hormone therapy and the risk of venous thromboembolism [Internet]. [cited 2017 Aug 2]. Available from: <http://www.clevelandclinicmeded.com/>

medicalpubs/diseasemanagement/womens-health/risk-of-venous-thromboembolism/

138. Rosendaal FR, Van Hylckama VA, Tanis BC, Helmerhorst FM. Estrogens, progestogens and thrombosis. *J Thromb Haemost JTH*. 2003;1:1371–80.
139. Nayak MK, Singh SK, Roy A, Prakash V, Kumar A, Dash D. Anti-thrombotic effects of selective estrogen receptor modulator tamoxifen. *Thromb Haemost*. 2011;106:624–35.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

