



HHS Public Access

Author manuscript

J Immunol. Author manuscript; available in PMC 2017 November 15.

Published in final edited form as:

J Immunol. 2017 May 15; 198(10): 3775–3789. doi:10.4049/jimmunol.1700373.

A Consensus Definitive Classification of Scavenger Receptors and Their Roles in Health and Disease

Mercy R. PrabhuDas^{*}, Cynthia L. Baldwin^{†,‡}, Paul L. Bollyky[§], Dawn M. E. Bowdish[¶], Kurt Drickamer^{||}, Maria Febbraio[#], Joachim Herz^{**††‡‡§§¶¶}, Lester Kobzik^{|||}, Monty Krieger^{###}, John Loike^{***}, Benita McVicker^{†††}, Terry K. Means^{†††}, Soren K. Moestrup^{§§§}, Steven R. Post^{¶¶¶}, Tatsuya Sawamura^{||||}, Samuel Silverstein^{####}, Robert C. Speth^{****}, Janice C. Telfer[†], Geoffrey M. Thiele^{††††}, Xiang-Yang Wang^{††††}, Samuel D. Wright^{§§§§}, and Joseph El Khoury^{¶¶¶¶}

^{*}Division of Allergy, Immunology and Transplantation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD 20852

[†]Department of Veterinary and Animal Sciences, University of Massachusetts, Amherst, MA 01003

[‡]Program in Molecular and Cellular Biology, University of Massachusetts, Amherst, MA 01003

[§]Department of Medicine, Stanford University, Stanford, CA 94305

[¶]Department of Pathology and Molecular Medicine, McMaster Immunology Research Centre, M.G. DeGroot Institute of Infectious Disease Research, McMaster University, Hamilton, Ontario L8S 4K1, Canada

^{||}Department of Life Sciences, Imperial College, London SW7 2AZ, United Kingdom

[#]Department of Dentistry, Katz Group Centre for Pharmacy and Health Research, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta T6G 2E1, Canada

^{**}Department of Molecular Genetics, University of Texas Southwestern Medical Center, Dallas, TX 75390

^{††}Department of Neuroscience, University of Texas Southwestern Medical Center, Dallas, TX 75390

^{‡‡}Department of Neurology and Neurotherapeutics, University of Texas Southwestern Medical Center, Dallas, TX 75390

^{§§}Center for Translational Neurodegeneration Research, University of Texas Southwestern Medical Center, Dallas, TX 75390

Address correspondence and reprint requests to Mercy R. PrabhuDas or Joseph El Khoury, National Institute of Allergy and Infectious Diseases, 5601 Fishers Lane, #7B57, Rockville, MD 20852 (M.R.P.) or Infectious Disease Division, Center for Immunology and Inflammatory Diseases, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA 02129 (J.E.K.).
mprabhudas@niaid.nih.gov (M.R.P.) or jelkhoury@mgh.harvard.edu (J.E.K.).

ORCID: 0000-0002-6434-2136 (R.C.S.); 0000-0001-5688-8596 (G.M.T.); 0000-0003-2421-5367 (X.-Y.W.).

Disclosures

The authors have no financial conflicts of interest.

^{¶¶}Peter O'Donnell Brain Institute, University of Texas Southwestern Medical Center, Dallas, TX 75390

^{¶¶¶}Department of Environmental Health, Harvard School of Public Health, Boston, MA 02115

^{##}Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139

^{***}Department of Physiology and Cellular Biophysics, Columbia University, New York, NY 10032

^{†††}University of Nebraska Medical Center, Omaha VA Nebraska-Western Iowa Health Care System, Omaha, NE 68105

^{†††}Center for Immunology and Inflammatory Diseases, Division of Rheumatology, Allergy, and Immunology, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA 02129

^{§§§}Department of Biomedicine, University of Aarhus, 8000 Aarhus C, Denmark

^{¶¶¶}Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, AR 72205

^{¶¶¶}Department of Physiology, Research Institute, Shinshu University School of Medicine, Matsumoto, Nagano, Japan

^{###}Department of Physiology and Cellular Biophysics, Columbia University, New York, NY 10032

^{****}Department of Pharmaceutical Sciences, Nova Southeastern University, Fort Lauderdale, FL 33328

^{††††}Division of Rheumatology, Department of Medicine, College of Medicine, University of Nebraska Medical Center, Omaha, NE 68105

^{††††}Department of Human and Molecular Genetics, Institute of Molecular Medicine, Massey Cancer Center, Virginia Commonwealth University School of Medicine, Richmond, VA 23298

^{§§§§}Cardiovascular Therapeutics, CSL Behring, King of Prussia, PA 19406

^{¶¶¶¶}Infectious Disease Division, Center for Immunology and Inflammatory Diseases, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA 02129

Abstract

Scavenger receptors constitute a large family of proteins that are structurally diverse and participate in a wide range of biological functions. These receptors are expressed predominantly by myeloid cells and recognize a diverse variety of ligands including endogenous and modified host-derived molecules and microbial pathogens. There are currently eight classes of scavenger receptors, many of which have multiple names, leading to inconsistencies and confusion in the literature. To address this problem, a workshop was organized by the United States National Institute of Allergy and Infectious Diseases, National Institutes of Health, to help develop a clear definition of scavenger receptors and a standardized nomenclature based on that definition. Fifteen experts in the scavenger receptor field attended the workshop and, after extensive discussion, reached a consensus regarding the definition of scavenger receptors and a proposed scavenger receptor nomenclature. Scavenger receptors were defined as cell surface receptors that typically bind multiple ligands and promote the removal of nonself or altered-self targets. They often function by mechanisms that include endocytosis, phagocytosis, adhesion, and signaling that

ultimately lead to the elimination of degraded or harmful substances. Based on this definition, nomenclature and classification of these receptors into 10 classes were proposed. This classification was discussed at three national meetings and input from participants at these meetings was requested. The following manuscript is a consensus statement that combines the recommendations of the initial workshop and incorporates the input received from the participants at the three national meetings.

Scavenger receptor activity was first described by Drs. Brown and Goldstein (1) when they identified receptors in macrophages that endocytosed and degraded modified (acetylated), but not native, low-density lipoprotein (LDL). Drs. Brown and Goldstein showed that these receptors for acetylated LDL recognized a wide variety of polyanionic ligands (2). Purification and cloning of the corresponding receptor protein and cDNA (3, 4) was soon followed by the identification of other modified LDL receptors (LDLRs) with broad binding specificity. In an initial attempt to systematically categorize these receptors, M. Krieger proposed to subdivide them into “classes” (A, B, etc.) based on their sequences, and each class subdivided further into “types” based on additional variations in their sequences due to alternative splicing (5).

There are currently eight classes of scavenger receptors (classes A–H). In some cases, multiple names have been assigned to the same receptor (e.g., MSR1, SR-AI, CD204, and SCARA1). In addition, there are proteins exhibiting scavenger receptor activity that have been named based on other criteria and have not been included in a general scavenger receptor nomenclature. Some examples include RAGE, LRP1, LRP2, ASGP, CD163, SR-PSOX, and CXCL16. New scavenger receptors also continue to be discovered and identified (6, 7). Therefore, it is important to convey consistent understanding and minimize redundancy and miscommunication in the scavenger receptor field. Developing a standardized nomenclature has the benefit of decreasing redundancy and facilitating communication and collaboration among investigators within a field, as well as in different fields, by building on a common understanding. Efforts to standardize other scientific nomenclatures have occurred in areas such as chemokines and their receptors (8), TNF family of receptors involved in bone metabolism (9), complement peptide receptors (10), and glutamate receptors (11). Committees, such as the International Union of Basic and Clinical Pharmacology, formed specifically for these processes by scientific societies or investigators in the field have developed systematic nomenclatures, for example, <http://www.guidetopharmacology.org/nomenclature.jsp> (accessed April 10, 2016), within each field for adoption by the broader scientific community.

To address the lack of a unified nomenclature system for scavenger receptors and better organize the field, the United States National Institute of Allergy and Infectious Diseases (NIAID), at the National Institutes of Health (NIH), represented by M.R. PrabhuDas, initiated discussions with J. El Khoury and Dr. Siamon Gordon which led to organizing and convening a workshop in which 15 investigators from five countries proposed a consensus definition of scavenger receptors and a systematic, standard nomenclature. The workshop participants and their areas of expertise are listed in Table I.

The objectives of this workshop were to: 1) establish guidelines and recommendations for standardizing the nomenclature for scavenger receptors; 2) consider strategies for dealing with future discoveries of scavenger receptors; and 3) communicate the recommendations to the wider scientific community as a point of reference and for further discussion and final scavenger receptor nomenclature standardization. The discussion and nomenclature recommendations described herein only refer to mammalian scavenger receptors.

Scavenger receptor background and history

The original description of scavenger receptors was in the context of studies of lipoproteins, atherosclerosis, and familial hypercholesterolemia (1, 12). Drs. Michael Brown and Joseph Goldstein (1) had previously identified LDLRs and LDLR-mediated endocytosis and established that familial hypercholesterolemia was caused by loss-of-function mutations in the LDLRs, which led to increased plasma LDL, premature atherosclerosis, and heart disease (1). Analysis of familial hypercholesterolemia patients showed that plasma LDL could be cleared by an LDLR-independent pathway called the “scavenger pathway.” The earliest use of the term “scavenger receptor” to refer to the broad specificity receptor identified by Dr. Brown and Dr. Goldstein was in 1981 by Fogelman and colleagues (13).

The first receptors that carried the name “scavenger receptor” were purified in 1988 (14) and cloned in 1990 (15, 16) by the Krieger group and were found to be alternatively spliced gene products of a single gene which were initially named type I and type II macrophage scavenger receptors. When additional scavenger receptors were identified, the type I and II macrophage receptors were renamed scavenger receptor, class A, type I and type II (SR-AI and SR-AII) by the Krieger group. One distinct feature of the broad binding specificity of scavenger receptors was that they could recognize a large repertoire of ligands, ranging from bacteria and yeast to self (native proteins) and modified-self targets (2, 17–21). Besides playing a role in host defense, class A scavenger receptors have been shown to be involved in functions such as homeostasis, Ag presentation, and pathogenesis of neurodegenerative disorders (22, 23).

Based on their broad ligand-binding specificities and expression in macrophages, Krieger and colleagues (5) proposed that scavenger receptors could serve as pattern recognition receptors for innate immunity. Work done by groups led by Drs. S. Gordon, M. Krieger, K. Joiner, T. Kodama, and J. El Khoury provided direct evidence that the class A receptors are adhesion receptors, and that they recognize Gram-positive bacteria and β -amyloid, confirming the broad ligand specificity for these receptors (17, 24, 25). Additional members of the class A scavenger receptors such as MARCO (26) and SCARA5 (27) were subsequently identified and demonstrated a role in homeostasis, adhesion, and cancer progression (28). Other multiligand pattern recognition receptors were independently identified, including CD14 (29), mannose receptors (30), and CD36 (31).

A report that served as an initial step in presenting the proposed nomenclature to the research community was published in *The Journal of Immunology* (32). A Web site was set up to invite comments from fellow investigators in response to the proposed nomenclature. The Web site (<https://respond.niaid.nih.gov/ScavengerReceptors/>) was open for three months

for public comment. In addition, NIAID organized several follow-up sessions at the following national meetings: Experimental Biology 2014 in San Diego, CA (<http://experimentalbiology.org/2014/Home.aspx>), Immunology 2014 in Pittsburgh, PA (<http://www.immunology2014.org/>), and Federation of Clinical Immunology Societies (FOCIS 2014) in Chicago, IL (<http://www.focisnet.org/focis-2014>). The sessions were open to the attendees of these national meetings to provide the broader scientific community with the opportunity to provide their comments and recommendations on the proposed scavenger receptor nomenclature. We now incorporate the comments received at the three national meetings and present a unified, consensus, and definitive classification and nomenclature for scavenger receptors.

Scavenger receptor nomenclature recommendations

At the initial workshop convened by the NIH, participants developed a definition of scavenger receptors, incorporating the various facets and characteristics of these receptors. This definition was used as the benchmark against which proposed nomenclatures were considered. As mentioned above, scavenger receptors were originally defined as cell surface proteins that bound chemically modified lipoproteins with high affinity. Additional characteristics included the ability to bind multiple ligands (broad-binding specificity), mediate endocytosis, in some cases mediate signaling, and act as innate immune pattern recognition receptors.

Based on the discussion, the group agreed upon the following working definition for scavenger receptors: Scavenger receptors are cell surface receptors that typically bind multiple ligands and promote the removal of nonself or altered-self targets. They often function by mechanisms that include endocytosis, phagocytosis, adhesion, and signaling that ultimately lead to the elimination of degraded or harmful substances.

Fig. 1 illustrates the consensus nomenclature formula agreed upon at the initial meeting. As an example, a scavenger receptor belonging to class A will be designated as SR-A1, where SR stands for scavenger receptor and is followed by a hyphen, then a capital letter representing the class of scavenger receptor, then an Arabic numeral representing the type of molecule within the class. Alternate splice variants are designated by a dot and an Arabic numeral following the type of molecule within the class (e.g., SR-A1.1). The numbering is based on the order in which the molecules were identified.

In the three follow up meetings there were discussions regarding the definition. There were questions as to whether receptors that strictly bind only pathogens should be included. The examples given were for viral receptors for rhinovirus or severe acute respiratory syndrome (SARS) (ICAM-1 or ACE2, respectively). There was also discussion on whether binding to modified self is a requirement. If a pathogen evolved to bind to a host molecule (including lipids) and that host molecule had a completely different primary function(s), one would not consider such a binding molecule on the host to be a scavenger receptor. Another idea proposed included the suggestion that if one of the principal functions of the host molecule is to bind pathogens (multiple pathogens), it could be considered a scavenger receptor even if there is no evidence for modified self-binding.

Finally, there were also discussions on whether to include binding specificity and affinity in the definition to differentiate various receptors. However, it was determined that this may not be appropriate for all scavenger receptors because some scavenger receptors bind some of their ligands in the micromolar range, whereas other ligand receptors bind in the nanomolar range.

Following the three follow up sessions at Experimental Biology 2014, Immunology 2014, and FOCIS 2014, it was decided to keep the original definition agreed upon at the initial meeting.

Tables II and III provide the current and revised nomenclature for mouse and human scavenger receptors. The following sections provide a brief overview of the currently known scavenger receptor classes and the revised nomenclature for each class and member, as recommended by the workshop participants and approved in the three follow up sessions discussed above.

Class A scavenger receptors: SR-A

Class A scavenger receptors (SR-A) are expressed primarily on tissue macrophages and macrophage subtypes such as Kupffer cells and cortical and medullary thymic macrophages (33). They have also been observed on high endothelial venules (33) and on subpopulations of dendritic cells (34), binding to a variety of foreign and self ligands. Members in this class have a similar structure comprised of an N-terminal cytoplasmic tail, transmembrane domain, spacer region, α -helical coiled coil domain, collagenous domain, and a C-terminal scavenger receptor cysteine-rich (SRCR) domain. SRCR domains are ~ 100–110 aa long, with 6–8 conserved cysteines predicted to form 3–4 disulfide bonds that are present in varying numbers in a variety of host defense-related proteins, including different classes of scavenger receptors (35, 36). In some cases alternative mRNA splicing deletes the SRCR domain. SCARA1/MSR1, the prototypical SR-A molecule, was the first scavenger receptor to be cloned and was initially isolated from bovine lung mRNA. It has subsequently been identified in other species, including mice and humans. There are three isoforms, which are alternative splice variants of the same gene that have been identified, namely, SR-AI, SR-AII, and SR-AIII (15, 16, 37, 38). Other members of this class of scavenger receptors include macrophage receptor with collagenous structure (MARCO), scavenger receptor A 5 (SCARA5), and scavenger receptor with C-type lectin domain (SRCL-I/II), also designated collectin, from placenta receptor-I (28, 39, 40).

The scavenger receptor currently designated as SCARA1 or MSR1 will now be referred to as SR-A1. SR-A1 is predominantly found in macrophages, monocytes, mast cells, and dendritic cells in both mice and humans (41). A diverse array of ligands is recognized by members of this class. SR-A1 binds to β -amyloid (42), heat shock proteins (43), surface molecules of Gram-positive and Gram-negative bacteria (17, 44), hepatitis C virus (38), and modified LDL such as acetylated LDL (termed AcLDL) and oxidized LDL (OxLDL), but not native LDL (15, 16).

SR-A1.1, an alternatively spliced form of SR-A1 (currently referred to as SR-AII), is characterized by a shortened C terminus. SR-A1.2, another alternatively spliced form of SR-

A1 (currently referred to as SR-AIII), is characterized by a truncated C terminus and remains trapped in the endoplasmic reticulum. The collagenous region of SR-A1 and SR-A1.1 has been identified as a major site of polyanionic ligand binding (45). There will be no receptor designated as SR-A2 to avoid confusion with the current SR-AII (new designation is: SR-A1.1).

Although SR-A1 and SR-A6 (currently known as MARCO) are primarily expressed in macrophages, SR-A3 (also referred to as SCARA3, cellular stress response/CSR), SR-A4 (SCARA4, SRCL, collectin placenta 1/CL-P1), and SR-A5 (SCARA5) are expressed in a variety of other tissues and cell types, including the lung, placenta, intestine, heart, and epithelial cells (46). SR-A3 has been associated with protecting cells from the detrimental effects of reactive oxygen species (21). SR-A4 functions both as an endocytic receptor for lipoproteins and mediates the recognition, internalization, and degradation of oxidatively modified LDL by vascular endothelial cells (47). SR-A5 (currently known as SCARA5) expression is restricted to epithelial cells within the testis, airway, thymus, and the adrenal gland. SR-A5 has the capacity to bind bacteria and may play an important role in host defense (27). Under normal conditions, in the absence of inflammation, the expression of SR-A6 is restricted to macrophages in the lymph nodes and marginal zone of the spleen. Studies have shown that SR-A6 mediates the clearance of bacteria from the lungs (48) and bloodstream (49).

Consensus member nomenclature

SR-A1. The protein is currently known as MSR1, SR-AI, or SCARA1 [prototype]; the current gene name is SCARA1.

SR-A1.1. The protein is currently known as SR-AII. This is an alternatively spliced form of SR-A1. There will be no receptor designated as SR-A2. This exception was agreed upon during the meeting to avoid confusion with the current SR-AII, which will now be designated as SR-A1.1.

SR-A3. The protein is currently known as MSRL1 or APC7; the current gene name is SCARA3.

SR-A4. The protein is currently referred to as scavenger receptor C-type lectin (SRCL) or CL-P1; the current gene name is COLEC12.

SR-A5. The protein is currently known as TESR; the current gene name is SCARA5.

SR-A6. The gene name is MARCO; the alternative name is SCARA2. The workshop participants recommended that the gene name be changed to SCARA6 for consistency.

Class B scavenger receptors: SR-B

Class B scavenger receptors include receptors currently known as SR-BI, and its alternatively spliced isoform SR-BII, as well as CD36 and LIMP2. Receptors belonging to this class are characterized by two transmembrane domains flanking an extracellular loop, with both the amino and C termini located within the cytoplasm. The extracellular domain of these receptors is extensively N-linked glycosylated, and this modification provides

protection against proteases often found in inflammatory sites. CD36 is the prototype class B type scavenger receptor and was initially identified as a receptor for thrombospondin (50). In this capacity, it modulates angiogenesis and cell-to-cell interactions. It is one of the most widely studied scavenger receptors and is involved in multiple aspects of macrophage biology, including migration and signaling and inflammatory processes such as foam cell formation. CD36 also plays an important role in the host immune response to fungi and bacteria and binds erythrocytes infected with the malaria parasite, *Plasmodium falciparum* (51), and it also functions as a facilitator of long-chain fatty acid uptake (52). It is expressed in many cell types, including insulin-responsive cells, hematopoietic cells such as platelets, monocytes, and macrophages, endothelial cells, and specialized epithelial cells in the breast and eye. Endemann et al. (31) were the first to characterize CD36 as a receptor for OxLDL, thereby cementing its role as a scavenger receptor. CD36 also binds polyanionic ligands of both mammalian hosts (e.g., high-density lipoprotein [HDL]) (53) and pathogens (e.g., *Staphylococcus aureus* and *Candida albicans*) (20, 54), and plays an important role in the recognition and endocytic uptake of oxidized phospholipids, apoptotic cells, and amyloid proteins (20). Stewart and colleagues (55) have shown that CD36 (proposed name SR-B2) cooperates with TLR family members (TLR4 and TLR6) to activate the innate immune response to ligands that accumulate in Alzheimer's disease and atherosclerosis, inducing proinflammatory mediators such as IL-1 β and RANTES.

SR-B1 (also known as SCARB1, SR-BI, or CD36L1) was initially cloned as a modified LDLR that could also bind unmodified LDL (56) and, together with CD36 (SR-B2), was the first surface receptor shown to bind anionic phospholipids (57). It was subsequently shown to be the first HDL receptor to be identified and to mediate the selective transport of lipids, such as cholesteryl esters from HDL, from circulating lipoproteins into cells (58, 59). SR-B1 is most highly expressed by hepatocytes and steroidogenic cells, but is also found in cells within the arterial wall and macrophages, such as those in human and murine atherosclerotic lesions (60). SR-B1 has been shown to have a protective effect on atherosclerosis development; total body ablation of SR-B1 in apoE-knockout mice fed a normal chow diet led to severe coronary arterial atherosclerosis, myocardial infarction, and death (61). Ablation of SR-B1 in bone marrow cells of LDLR knockout mice also resulted in increased atherosclerosis, but not death, indicative of partial protection against atherosclerosis (62).

SR-B2, an alternatively spliced form of SR-B1 expressed at substantially lower levels than SR-B1, contains a distinctly different cytoplasmic tail from SR-B1. SR-B2 is expressed in adrenal glands, testes, and liver, and has been shown to mediate selective uptake of cholesteryl ester from HDL with ~4-fold lower efficiency than SR-B1 (63).

Consensus member nomenclature

SR-B1. The protein is currently also known as SR-BI or CD36L1; the gene name is SCARB1.

SR-B1.1. This is an alternatively spliced form of SR-B1, currently known as SR-BII.

SR-B2. The protein is currently known as CD36, SCARB3, or PAS4 [prototype].

Class C scavenger receptors

Class C scavenger receptors have been described in *Drosophila* (5), but there are currently no known mammalian class C scavenger receptors. This scavenger receptor class was not discussed because the workshop focused on mammalian scavenger receptors.

Class D scavenger receptors: SR-D

CD68 is the only member of the class D scavenger receptors. CD68, also known as macrophage mannose receptor in mouse and by other names in mice and humans (Tables II, III), is a type I transmembrane glycoprotein that belongs to the lysosome-associated membrane protein (or LAMP) family of molecules (64). There is a 300 aa extracellular domain that is rich in threonine and serine, which may serve as attachment sites for carbohydrates. It has a short cytoplasmic tail. Human CD68 shares 75% aa sequence identity with mouse CD68 in the extracellular domain. CD68 is found on monocytes and tissue-specific macrophages in the peritoneum, lungs, liver, spleen, Langerhans cells, and microglia, and serves as a scavenger receptor for OxLDL (65). CD68 is a differentiation marker of hematopoietic cells of the monocyte/macrophage lineage. It is mostly found in the late endosomal compartment in macrophages and dendritic cells, with limited expression on resting cells. CD68 plays a minor role in the binding and uptake of oxidized lipoproteins and apoptotic cells by macrophages (23).

Consensus member nomenclature

SR-D1. The protein is currently known as CD68 [prototype]; the gene name is CD68 molecule.

Class E scavenger receptors: SR-E

Class E scavenger receptors are type 2 transmembrane proteins with C-type lectin-like domains and demonstrated scavenger receptor activity; sequence homology alone is not sufficient to include a protein in this class of receptors. Based upon their structure, these proteins belong to a subfamily of the NK cell C-type lectin-like (CLEC) receptor family. The NK receptor gene complex encodes a large number of CLEC-like receptors that are expressed on NK cells and other leukocytes. Analysis of the NK receptor gene complex in five species revealed variations in the sizes of the gene clusters: human (29 genes), dog (22 genes), cattle (32 genes), rat (75 genes), and mouse (57 genes) (66). Among these gene clusters, the first two members of the CLEC family, LOX-1 and dectin-1, have been well characterized and shown to have scavenger activity.

Lectin-like oxidized LDLR (LOX-1) is expressed on macrophages and dendritic cells, and binds oxidized LDL and the acute phase protein C-reactive protein (or CRP) (67). It was originally cloned from a bovine aortic endothelial cDNA expression library screened for receptors for oxidized LDL (68). LOX-1 is expressed on vascular endothelial cells, platelets, smooth muscle cells, adipocytes, and macrophages. In addition to binding oxidized LDL, LOX-1 has been implicated in recognizing other ligands, including apoptotic cells, and Gram-positive and Gram-negative bacteria (23, 67). A splicing variant of human LOX-1 lacking a lectin-like domain, called LOXIN, can act as a dominant negative isoform forming

a heterodimeric complex with LOX-1 (69). There are no data that this splicing variant is present in mouse cells.

Dectin-1 is expressed predominantly on myeloid cells (macrophages, dendritic cells, and neutrophils), and can be regulated by cytokines and microbial stimuli. Human dectin-1 is alternatively spliced, resulting in two functional forms. This receptor recognizes various bacterial, fungal, and plant carbohydrates (β -1,3 and/or β 1-6 glucans) in addition to intact fungi and parasites (70). Dectin-1 mediates proinflammatory cytokine production in response to β -glucan particles, in conjunction with TLR2. It can also function as a phagocytic receptor through both Syk-dependent and Syk-independent pathways (71). Dectin-1 has been reported to act as a costimulatory molecule to activate both CD4 and CD8 T cells. Moreover, dectin-1 is expressed on macrophages and dendritic cells in the medullary regions of the thymus, suggesting a role in thymocyte development (72).

Some CLEC family members are known to recognize pathogen-associated molecules. However, most of the members of the CLEC family have not been shown to have scavenger receptor activity. At this time, only LOX-1 and dectin-1 can be classified as scavenger receptors using the definition proposed by the workshop participants.

Mannose receptor (CD206)

The human mannose receptor, CD206 (SR-E3), is a 175 kDa transmembrane glycoprotein characterized by eight C-type lectin carbohydrate recognition domains. Depending on the particle size and the cell type expressing the receptor, it may be involved in phagocytosis of mannose-coated particles, in endocytosis of mannosylated glycoproteins, or in receptor-mediated facilitated Ag presentation. Multiple roles of SR-E3 have been observed. As a pattern recognition receptor, SR-E3 recognizes glycans on the surface of pathogenic microorganisms, internalizes them via phagocytosis, and transports them to lysosomes for degradation. Cells isolated from skin, bone marrow, thymus, lymph node, and peripheral blood have demonstrated that SR-E3 is a differentiation marker of immature monocyte-derived dendritic cells. Studies have shown that inflammatory dendritic epidermal cells that are found in inflammatory skin diseases such as atopic dermatitis express the mannose receptor, SR-E3. These results suggest that SR-E3 expression in the epidermis may play an important role in capturing mannosylated Ags for subsequent Ag presentation (73). Other studies have shown that mannose receptors may also play a role in regulating the levels of molecules such as myeloperoxidase that are released in response to invading pathogenic microorganisms. SR-E3 may then remove these glycoproteins from the circulation to prevent damage to host tissues.

Asialoglycoprotein receptor 1 (ASGPR1)

The asialoglycoprotein receptor (ASGPR), also termed the hepatic binding protein or the Ashwell receptor, was discovered over four decades ago as a hepatocellular surface receptor that binds glycoproteins lacking terminal sialic acid residues (asialoglycoproteins) (74, 75). Subsequently, many studies have led to the detailed characterization of the ASGPR, describing its functional role in the binding, internalization and transport of a wide range of

glycoproteins that have galactose or N-acetylgalactosamine residues via the process of receptor-mediated endocytosis (75–78). More specifically, it has been demonstrated that the ASGPR can bind a variety of clinically important plasma proteins that include transferrin (79), Igs including IgA (80), apoptotic cells (81), fibronectin (82, 83), enzymes such as alkaline phosphatase (84, 85), and immune cells (86, 87). The role of ASGPRs as a major liver clearance/scavenger mechanism in disease states remains a topic of current research efforts.

Several laboratories have investigated the relationship between altered ASGPR function and specific mechanisms involved in the pathogenesis of liver disease. Studies examining translatable consequences of altered hepatic clearance/scavenging mechanisms revealed that liver damaging mechanisms such as aberrant apoptotic cell regulation and innate immune defenses were associated with impaired ASGPR function. Particularly, it was shown that ethanol-mediated impairments in ASGPR clearance of apoptotic cells could be related to enhancements of proinflammatory mechanisms during liver disease (88, 89). Additionally, it was found that ethanol-mediated alterations to ASGPRs led to defective hepatocyte killing of T cells, which could impact liver immunologic homeostasis and be related to the accumulation of lymphocytes, especially CD8⁺ T cells, that occurs during alcoholic hepatitis (90, 91). And more recently, it was determined that impaired ASGPR function due to alcohol resulted in altered uptake and trafficking of alkaline phosphatases, leading to defective detoxification of LPS and enhanced inflammatory liver injury (92, 93).

These findings indicate that proper functioning of the ASGPR may mediate universal protective mechanisms against liver injury by removing a range of desialylated proteins from the circulation (94), demonstrating the physiological importance of this hepatic scavenging receptor system. Moreover, it is clear that there is an association between altered ASGPR function and the progression of liver disease. Further characterization of ASGPR's contribution in liver disease mechanisms could bring forth a better understanding and the potential development of successful therapeutic options for liver disease patients.

Consensus member nomenclature

SR-E1. The protein is currently known as LOX-1; the gene name is OLR1: oxidized LDLR 1.

SR-E1.1. An alternatively spliced form of SR-E1, the protein is known as LOXIN (no known scavenger receptor activity but works as a dominant negative form).

SR-E2. The protein is currently known as dectin-1; the gene name is CLEC7A.

SR-E3. The protein is currently known as mannose receptor 1, CD206, or MRC1; the gene name is mannose receptor C type 1.

SR-E4. The protein is currently known as ASGPR1, CLEC4H1, or HL-1; the gene name is ASGR1.

Class F scavenger receptors: SR-F

There are currently three members belonging to the class F scavenger receptors. The first class F scavenger receptor (SCARF or SCARF1) was identified as an endothelial receptor

for modified LDL, and is characterized by the presence of an extracellular ligand-binding domain, with multiple extracellular epidermal growth factor-like repeats, a transmembrane domain, and a long cytoplasmic tail that includes a serine/proline-rich region followed by a glycine-rich region and is rich in positively charged residues (38). It was named scavenger receptor expressed by endothelial cells (SREC or SREC-I) (95). SCARF1/SREC-I is also expressed in macrophages, and functions to bind calreticulin and associated peptides, which are transported into the MHC class I pathway in macrophages and dendritic cells (23). SCARF1 is also a receptor for fungal pathogens and heat shock proteins (20, 38). Interestingly, a *Caenorhabditis elegans* ortholog of SCARF1, CED-1, plays a role in engulfing apoptotic cells and has been implicated in host defense (23). A recent report indicates that SCARF1, similar to CED-1, participates in the clearance of apoptotic cells (96).

SREC-II, also known as SCARF2, was isolated by Ishii et al. (97) in 2002. Although SREC-II, like SREC-I, was initially identified on endothelial cells, SREC-II is also expressed on macrophages. SREC-II can bind to SREC-I in trans (97), but a scavenger receptor function for SREC-II has not yet been reported.

Another member of the SCARF family, multiple EGF-like domains-10 (MEGF10), recently has been shown to be a receptor for the protein amyloid- β in the brain, and is thought to be involved in the pathogenesis of Alzheimer's disease (23). Recently, MEGF10 has been shown to be a receptor for the complement protein C1Q and consequently to mediate the clearance of apoptotic cells in the developing mouse cerebellum (98). Mutations in MEGF10 in humans result in early-onset myopathy, areflexia, respiratory distress, and dysphagia (EMARDD). These EMARDD mutations also have a defect in binding C1Q (98). Whether such a defect is related to the clinical presentation of EMARDD patients is not known. It is not clear yet whether other MEGF family members, such as MEGF8, 9, and 11, also have a scavenger receptor function, because no functional studies have been published to date on these molecules.

Consensus member nomenclature

SR-F1. The protein is currently known as SREC-I/SCARF1 [prototype]; the gene name is SCARF1.

SR-F2. The protein is currently known as MEGF10.

Class G scavenger receptors: SR-G

Currently, one receptor belongs to scavenger receptor class G, SR-PSOX/CXCL16. Shimaoka et al. (99) used expression cloning from a cDNA library of PMA-treated human monocytic THP-1 cells to identify scavenger receptors that bound OxLDL, and discovered this receptor, which they called SR-PSOX (scavenger receptor that binds phosphatidylserine and oxidized lipids). At around the same time, the groups of Dr. Jason Cyster and Dr. Michael Briskin discovered a new chemokine that was capable of activating CXCR6 (also known as BONZO, STRL33, or TYMSTR), resulting in migration of T cells in vitro (100–102). This chemokine was named CXCL16 and was found to be identical to SR-PSOX in its amino acid sequence.

Studies show that the SR-PSOX/CXCL16 gene is located on chromosome 17p13, and its protein exists in both membrane and soluble forms (100). It is a type I transmembrane glycoprotein with a CXC chemokine motif, mucin stalk, transmembrane, and extracellular/soluble domains. SR-PSOX/CXCL16 does not share structural similarity with other scavenger receptors. SR-PSOX/CXCL16 has the capacity to mediate adhesion and phagocytosis of bacteria by APCs (101). The membrane-bound form can function as an adhesion molecule for CXCR6-expressing T cells without requiring CXCR6-induced signal transduction and integrin activation.

This multifunctional transmembrane protein has a chemokine-like CXC motif and appears to be involved in several phases of an immune response from Ag recognition to migration of immune cells to formation of inflammatory foci (103). The soluble form of SR-PSOX/CXCL16, produced by cleavage from the membrane domain, functions as a chemoattractant for activated T cells and bone marrow plasma cells via its receptor CXCR6 (104). The membrane-bound form acts as a scavenger receptor for OxLDL, and facilitates phagocytosis of bacteria by APCs (104).

Consensus member nomenclature

SR-G1. The protein is currently known as SR-PSOX/CXCL16 [prototype]; the gene name is CXCL16.

Class H scavenger receptors: SR-H

Receptors in this class are transmembrane proteins that consist of fasciclin, EGF-like, and lamin-type EGF-like (FEEL) domains. The currently known proposed SR-H group members are FEEL-1/stabilin-1/CLEVER-1, and FEEL-2/stabilin-2/HARE. FEEL-1 is predominantly expressed in macrophages, mononuclear cells, hematopoietic stem cells, and endothelial cells, whereas FEEL-2 is found mainly in sinusoidal endothelial cells. FEEL-1 and FEEL-2 have been shown to be involved in lymphocyte adhesion, transmigration, angiogenesis, apoptotic cell clearance, and intracellular trafficking. FEEL-1 and FEEL-2 also have been implicated in the clearance of aged RBCs by macrophages (105, 106) in a phosphatidylserine-dependent manner. FEEL-1 has been shown to directly interact with and phagocytose phosphatidylserine-coated beads, but this action has yet to be confirmed in vivo. However, FEEL-1 and FEEL-2 double knockout mice have been shown to have impairment in the clearance of noxious blood factors (107).

Consensus member nomenclature

SR-H1. The protein is currently known as FEEL-1, stabilin-1, or CLEVER-1 [prototype]; the gene name is STAB1.

SR-H2. The protein is currently known as FEEL-2, or stabilin-2; the gene name is STAB2.

Class I scavenger receptors: SR-I

The CD163 family of molecules are type I transmembrane receptors characterized by multiple group B SRCR domains in their extracellular domain, with membrane proximal

lymph node, trachea, and lung. The bovine genome contains one *CD163* gene and one *CD163c- α /SCART1* gene, which are both expressed in $\gamma\delta$ T cells, as well as peripheral blood monocytes, mesenteric lymph node, lung, and intestinal epithelial lymphocytes (113).

In addition to the class I scavenger receptors expressed in humans (CD163A, CD163b, and CD163c- α /SCART1), other mammalian species also express a closely related molecule WC1 (T19) (113). In species examined thus far, WC1 is encoded in duplicated multigenic arrays. The bovine genome has 13 *WC1* genes (116, 117). Twelve of the bovine WC1 genes encode 11 SRCR domains; one encodes six SRCR domains in the extracellular domain. All WC1 molecules are type I transmembrane receptors with a signal peptide, SRCR domains in the extracellular domain organized into the b-c-d-e-d' cassette that has been conserved in the CD163 family, a transmembrane domain, and a cytoplasmic domain (117, 118). Table IV provides the nomenclature for the bovine scavenger receptors. In the bovine, WC1 is only expressed on $\gamma\delta$ T cells where it serves as a hybrid coreceptor and pattern recognition receptor in the $\gamma\delta$ T cell memory response to bacteria (119–121). Coreceptor signaling through WC1 and TCR is dependent on the phosphorylation of tyrosines and serines in the WC1 cytoplasmic domain, and is downregulated by endocytosis via a WC1 dileucine motif (119–121). Individual WC1 SRCR domains from selected WC1 molecules bind to the bacteria *Leptospira* spp., *Borrelia burgdorferi*, and *Mycobacteria* spp., suggesting that the natural selection pressure necessary to maintain multiple WC1 open reading frames is derived from the specificity for diverse pathogens of individual WC1 molecules (122).

Of note, WC1 receptors are not the only scavenger receptors expressed in bovine tissues. As mentioned earlier, SR-A, SR-E, and SR-J were initially cloned from bovine lung and endothelial cells, respectively. Table IV refers only to scavenger receptors exclusively described in bovine tissues.

Consensus member nomenclature—The proposed definition for this group is transmembrane proteins with multiple SRCR domains.

SR-I1. The protein is currently known as CD163A or M130 [prototype]; the gene name is CD163.

SR-I1.1. The protein is the CD163 long tail variant 1.

SR-I1.2. The protein is the CD163 long tail variant 2.

SR-I2. The protein is currently known as CD163B or M160; the gene name is CD163L1.

SR-I3. The protein is currently known as CD163c- α or SCART1; the gene name is cd163l1.

SR-I4. The protein is currently known as murine SCART2.

SR-I5 to SR-I17. The proteins currently correspond to bovine WC1-1 to WC1-13.

Class J scavenger receptors: SR-J

The receptor for advanced glycation end products (RAGE) is currently the only member of the class J scavenger receptor group. It is a member of the Ig superfamily of cell surface molecules capable of interacting with a broad spectrum of ligands, including advanced glycation end products, high mobility group protein box-1 (HMGB1), and the S-100 protein. RAGE is composed of a single transmembrane domain that connects the amino-terminal ligand recognition and binding ectodomain with a short cytoplasmic domain. The extracellular domain also contains three Ig-like regions. RAGE is a pattern recognition receptor predominantly involved in the recognition of endogenous molecules released in the context of infection, physiological stress, or chronic inflammation (123). Signaling through RAGE mediates processes such as inflammation, oxidative stress, and apoptosis, which can lead to pathologies including atherosclerosis, diabetes, cancer, neurodegenerative disease, and stroke (124). RAGE signaling also mediates developmental processes in neurons such as cell migration and differentiation. However, studies assessing brain pathology in Alzheimer's disease revealed that diseased brains express higher levels of RAGE in the vasculature and on neurons compared with normal brains (23). In addition to the membrane-bound form, a circulating portion of the protein, known as "soluble" RAGE (termed sRAGE) also has biological properties. Elevated levels of soluble RAGE and HMGB1 expression were observed in patients with myocarditis compared with healthy controls (125). Moreover, RAGE-deficient mice were observed to be protected from developing troponin I-induced autoimmune myocarditis, with a concomitant decrease in several inflammatory mediators (126).

Consensus member nomenclature

SR-J1. The protein is currently known as the membrane-bound form of RAGE/AGER; the gene name is AGER.

SR-J1.1. This terminology will be used to designate the soluble form of RAGE.

Class K scavenger receptors: SR-K

Several lines of evidence suggest that CD44, a cell surface glycoprotein best known as a receptor for hyaluronan, should be included within the family of scavenger receptors.

Although CD44 is not structurally homologous to any of the known scavenger receptors, it nonetheless shares structural features in common with them. It is a multidomain, single-span transmembrane protein with only a modest cytosolic tail with limited direct signaling capabilities (127). The N-terminal extracellular domain consists of motifs that serve as docking sites for various ligands that include hyaluronan, extracellular matrix glycoproteins and proteoglycans, growth factors, cytokines, and matrix metalloproteinases. The C-terminal cytoplasmic tail binds to cytoskeletal elements and interacts with signaling molecules, which are members of the Src family kinases (128). Like other scavenger receptors, it is susceptible to shedding by exofacial proteases (129) and these cleavage products are thought to be biologically active (130).

CD44 also has functional properties in common with other scavenger receptors. It has a broad repertoire of ligands that include both host (131–133) and microbial (134) molecules.

It associates with a diverse range of other proteins, including TGF- β RI (135), CD3 (136), TLR 2 (137), and CD14 (138), and thereby engages with multiple intracellular signaling pathways, probably through formation of heteromultimeric signaling complexes and interactions with the actin cytoskeleton (139). Like other scavenger receptors, it contributes to the clearance of its extracellular matrix ligands via endocytosis (140). CD44 also contributes to pattern recognition (141) and innate immunity (142). As with other scavenger receptors, the pro- and anti-inflammatory contributions of CD44 are highly dependent on cell type and context (143–145). Along with its immunologic roles, CD44 also makes important, overlapping contributions to cellular migration (146), polarization (147), and homeostasis (148).

In light of these structural and functional similarities with other scavenger receptors, we have included CD44 as a distinct class of scavenger receptor, class M.

Consensus member nomenclature

SR-K1. The protein is currently known as CD44; the gene name is CD44.

Class L scavenger receptors: SR-L

SR-L1 is one of the largest members of the LDLR gene family (149). It is also known as LDLR-related protein 1 (LRP1), α_2 -macroglobulin receptor (A2MR), or CD91 (150). The gene family to which SR-L1 belongs includes as its best known member the LDLR, which serves as the principal clearance receptor for plasma cholesterol. The first members of this ancient family likely appeared with the first metazoans, as they are present in the most primitive multicellular eukaryotic organism that survives on the planet to this day. Remarkably, despite being separated from present day mammals by ~1 billion years of evolution, their exon/intron structure has been conserved virtually perfectly throughout time. Moreover, within the mammalian kingdom, SR-L1 in particular exhibits a remarkable degree of primary amino acid sequence conservation, which further emphasizes the unique importance of SR-L1, and several other members of the gene family, that is, LRP1b, LRP2, LRP4, LRP8, and very LDLR (termed VLDLR) for the survival of all species (151). LRP2 is involved in the reuptake of albumin, vitamin D-binding protein, and retinol-binding protein. LRP2 is defective in Donnai-Barrow syndrome (152).

Although LRP1/SR-L1 and other family members have been investigated intensely for a quarter century, we still do not have a complete understanding of all their biological functions. Rather than transmitting an extracellular signal across the membrane itself, SR-L1 mostly regulates such signals by serving as an extracellular sensor that detects the presence of a myriad of cargo ligands or bioactive molecules in the extracellular space. By modifying or integrating signals transmitted through classical signal transducing receptors SR-L1 serves a double purpose: it clears and removes—or scavenges—the ligands from the surface or extracellular space and, at the same time, the presence of these ligands modifies the signals sent to the interior of the cell by the associated classical signal transduction receptors. In this way, the cell can receive a variety of signals, which include, for instance, the activity of extracellular protease activity, cellular adhesion, nutrient availability, and growth signals, and which help the cell to determine its status within the tissue and the

organism as a whole. Hence, serving as an extracellular sensor and simultaneously as a scavenger receptor for the same ligands are merely two faces of the same coin.

Currently, over 100 functionally diverse ligands are known to interact with LRP1/SR-L1. Many of them have also been found to bind to LRP1b and to LRP2, both closely related family members of similar size, but different tissue distribution. For most of these ligands, the exact biological conditions that are regulated by their receptor interactions, and in particular the coreceptors or signal transducers whose functions they modulate, remain unknown. Only some of these functions are briefly listed here and they include limb and brain development, synaptic homeostasis and amyloid clearance and removal (153), liver lipid and energy metabolism, synaptic transmission, thrombosis, hemostasis, the protection of vascular integrity by regulating reverse cholesterol transport, fibrosis and smooth muscle cell proliferation (154), regulation of the inflammatory response (e.g., through TLRs and defensins), and others (155).

Megalin (SR-L2), also known as LRP2, is a multiligand endocytic receptor that belongs to the LRP family of proteins. It is a large membrane glycoprotein that belongs in the LDLR family. It is expressed on the apical membrane of several epithelial cells such as kidney proximal tubule cells, lung, thyroid, gallbladder, and neuroepithelium, as well as steroid-responsive tissues such as epididymis, prostate, ovaries, and uterus (156, 157). The cytoplasmic domain is small and has been suggested to regulate its endocytic activities (158). SR-L2 is expressed at the blood brain barrier where it binds and internalizes a variety of ligands such as leptin, insulin, and amyloid β peptide (157). Mice that lacked megalin/SR-L2 in brain endothelial cells exhibited neurodegeneration and neuroinflammation, as well as obesity (157). Evidence has also been demonstrated for the role of peroxisome proliferator-activated receptors (termed PPARs) in regulating the expression of SR-L2 (158).

Consensus member nomenclature

SR-L1. The protein is currently known as LDLR-related protein 1 (LRP1). It is also known as α_2 -macroglobulin receptor (A2MR), apolipoprotein E receptor (APOER) or CD91; the gene name is *Lrp1*.

SR-L2. The protein is currently known as megalin or gp330; the gene name is *Lrp2*.

Other potential scavenger receptors: SR-Others

The SR-Others group refers to molecules with scavenger activity that do not belong to the current scavenger receptor classes, but belong to other receptor families. These receptors include CD11b/CD18 α . Classifying these receptors as scavenger receptors will require the input of the wider scientific community.

Another receptor that may be considered for inclusion as a scavenger receptor is CD14. CD14 is a GPI-linked protein found on monocytes and polymorphonuclear leukocytes. CD14 has a binding site for amphipathic molecules such as LPS and lipidated microbial cell wall fragments, as well as phosphatidylcholine, phosphatidylinositol, and other endogenous phospholipids. Binding of amphiphiles to CD14 is readily reversible, allowing cell surface

CD14 to transfer the amphiphiles into the plasma membrane bilayer, to MD2 or to TLR4. By this means, CD14 can serve as a shuttle for the clearance of amphiphilic molecules. Shuttling of bacterial LPS is particularly favored because the plasma protein, LPS-binding protein, serves as a lipid transfer protein that specifically transfers LPS monomers into CD14. The binding of LPS to CD14 catalyzed by the LPS-binding protein can then be followed by transfer of the LPS to MD2 or TLR4 or into the plasma membrane bilayer. CD14 thereby fulfills the definition of scavenger receptor by binding multiple ligands and serving for their clearance. CD14 may be shed from the surface and appears in plasma and other biological fluids at relatively high levels. This soluble CD14 retains the ability to bind amphiphiles and retains the ability to serve in the shuttling of lipids.

Receptors that are not included as scavenger receptors

The decision to exclude specific receptors for viruses which facilitate entry of viral particles into cells is based on their inability to meet most of the criteria of scavenger receptors. Three examples of such receptors are angiotensin-converting enzyme-2 (ACE2), aminopeptidase N, which is also known as CD13, and lysosomal integral membrane protein 2 (also known as SCARB2 or CD36b like-2).

ACE2 is a type I membrane-bound zinc metallopeptidase. It functions as a monooxypeptidase that cleaves the C-terminal amino acid from angiotensins I and II as well as from several other peptides at its extracellular catalytic domain (159). It is also the receptor for the SARS coronavirus (160). Although its peptidase activity is not specific to a single peptide substrate or peptide sequence, its function is to metabolize self rather than altered-self peptides. In its role as a virus receptor, the only ligand that is reported to bind to ACE2 with high affinity is the S1 domain of the SARS virus (160), suggestive of a high specificity, as opposed to broad-ligand specificity. Additionally, ACE2 enables the SARS virus to enter cells simply by serving in a passive role as an anchor to the cell membrane, allowing it to fuse with and penetrate the cell membrane to enter cells. Moreover, this internalization allows the virus to proliferate rather than to be eliminated or inactivated as would occur with a scavenger receptor.

Aminopeptidase N (CD13, E.C. 3.4.11.2) is a type I membrane-bound zinc metallopeptidase that cleaves the amino terminal from angiotensin III, enkephalins, and several other peptides at its extracellular catalytic domain (161). It is also the receptor for several coronaviruses: 229E, transmissible gastroenteritis virus, porcine epidemic diarrhea virus, and feline infectious peritonitis virus, which belong to the group I coronaviruses (160, 162). Similar to ACE2, its peptidase activity is not specific to a single peptide substrate or peptide sequence, and its function is to metabolize self rather than altered-self peptides. In its role as a virus receptor, it is still relatively selective, inasmuch as it binds a limited number of class I coronaviruses, in contrast to a broad-ligand specificity. Additionally, aminopeptidase N enables the viruses to enter cells also serving a passive role as an anchor to the cell membrane that allows the viruses to fuse with and penetrate the cell membrane to enter cells. As noted above, this internalization allows the virus to proliferate rather than to be inactivated, as would occur with a scavenger receptor.

Lysosomal integral membrane protein 2 (LIMP2, also known as SCARB2 or CD36b like-2) is a member of the CD36 superfamily of proteins and is a type III glycoprotein located in the membranes of lysosomes and endosomes and contains two transmembrane domains (163). Some of its functions include transporting the lysosomal protein glucocerebrosidase from the endoplasmic reticulum through the Golgi system to lysosomes where cellular degradation occurs (164). Reports have demonstrated that LIMP2 is a receptor for a variety of human enteroviruses such as EV71, coxsackievirus (CV) A16, CVA14, and CVA7 (165), supporting viral propagation and cytopathic effects. It is not known if LIMP2 has additional scavenger receptor like functions. Classifying LIMP2 as a potential scavenger receptor will be reassessed if additional information becomes available.

Conclusions

In addition to their functions as phagocytic receptors and innate immune recognition receptors, scavenger receptors also play an important role as regulators of inflammatory signaling. A schematic representation of the current scavenger receptors classes is provided in Fig. 2. These receptors have also been implicated in a number of physiological and pathological processes. These include interactions with other innate immune receptors, delivery of ligands to different cellular compartments, and Ag presentation. Furthermore, the role of these receptors in neurodegenerative diseases, diabetes, and other metabolic disorders, and their potential as targets for therapeutic interventions to treat various disorders, warrant further study. Having a standardized nomenclature as described herein will help to overcome current ambiguities in communication about specific scavenger receptors and, perhaps, remove a major barrier for researchers in other fields to enter the scavenger receptor field.

Acknowledgments

We thank Ethan Tyler from the National Institutes of Health Medical Arts for assistance with Fig. 2.

Abbreviations in this article

ACE2	angiotensin-converting enzyme-2
ASGPR	asialoglycoprotein receptor
CLEC	C-type lectin-like
EMARDD	early-onset myopathy, areflexia, respiratory distress and dysphagia
FEEL	fasciclin, EGF-like and lamin type EGF-like
HDL	high-density lipoprotein
LDL	low-density lipoprotein
LDLR	LDL receptor
NIAID	National Institute of Allergy and Infectious Diseases

NIH	National Institutes of Health
OxLDL	oxidized LDL
RAGE	receptor for advanced glycation end products
SARS	severe acute respiratory syndrome
SRCR	scavenger receptor cysteine-rich
SREC	scavenger receptor expressed by endothelial cells

References

1. Goldstein JL, Ho YK, Basu SK, Brown MS. Binding site on macrophages that mediates uptake and degradation of acetylated low density lipoprotein, producing massive cholesterol deposition. *Proc Natl Acad Sci USA*. 1979; 76:333–337. [PubMed: 218198]
2. Brown MS, Goldstein JL. Lipoprotein metabolism in the macrophage: implications for cholesterol deposition in atherosclerosis. *Annu Rev Bio-chem*. 1983; 52:223–261.
3. Russell DW, Yamamoto T, Schneider WJ, Slaughter CJ, Brown MS, Goldstein JL. cDNA cloning of the bovine low density lipoprotein receptor: feedback regulation of a receptor mRNA. *Proc Natl Acad Sci USA*. 1983; 80:7501–7505. [PubMed: 6143315]
4. Sege RD, Kozarsky K, Nelson DL, Krieger M. Expression and regulation of human low-density lipoprotein receptors in Chinese hamster ovary cells. *Nature*. 1984; 307:742–745. [PubMed: 6321999]
5. Krieger M. The other side of scavenger receptors: pattern recognition for host defense. *Curr Opin Lipidol*. 1997; 8:275–280. [PubMed: 9335951]
6. Gu BJ, Saunders BM, Petrou S, Wiley JS. P2X(7) is a scavenger receptor for apoptotic cells in the absence of its ligand, extracellular ATP. *J Immunol*. 2011; 187:2365–2375. [PubMed: 21821797]
7. Shibata M, Ishii J, Koizumi H, Shibata N, Dohmae N, Takio K, Adachi H, Tsujimoto M, Arai H. Type F scavenger receptor SREC-I interacts with advillin, a member of the gelsolin/villin family, and induces neurite-like outgrowth. *J Biol Chem*. 2004; 279:40084–40090. [PubMed: 15247299]
8. Murphy PM, Baggiolini M, Charo IF, Hébert CA, Horuk R, Matsushima K, Miller LH, Oppenheim JJ, Power CA. International union of pharmacology. XXII Nomenclature for chemokine receptors. *Pharmacol Rev*. 2000; 52:145–176. [PubMed: 10699158]
9. The American Society for Bone and Mineral Research President's Committee on Nomenclature. Proposed standard nomenclature for new tumor necrosis factor members involved in the regulation of bone resorption. *Bone*. 2000; 27:761–764. [PubMed: 11113385]
10. Klos A, Wende E, Wareham KJ, Monk PN. International Union of Pharmacology. LXXXVII Complement peptide C5a, C4a, and C3a receptors. *Pharmacol Rev*. 2013; 65:500–543. [Published erratum appears in 2014 *Pharmacol. Rev*. 66: 466.]. [PubMed: 23383423]
11. Lodge D. The history of the pharmacology and cloning of ionotropic glutamate receptors and the development of idiosyncratic nomenclature. *Neuropharmacology*. 2009; 56:6–21. [PubMed: 18765242]
12. Brown MS, Basu SK, Falck JR, Ho YK, Goldstein JL. The scavenger cell pathway for lipoprotein degradation: specificity of the binding site that mediates the uptake of negatively-charged LDL by macrophages. *J Supramol Struct*. 1980; 13:67–81. [PubMed: 6255257]
13. Fogelman AM, Haberland ME, Seager J, Hokom M, Edwards PA. Factors regulating the activities of the low density lipoprotein receptor and the scavenger receptor on human monocyte-macrophages. *J Lipid Res*. 1981; 22:1131–1141. [PubMed: 6271886]
14. Kodama T, Reddy P, Kishimoto C, Krieger M. Purification and characterization of a bovine acetyl low density lipoprotein receptor. *Proc Natl Acad Sci USA*. 1988; 85:9238–9242. [PubMed: 3194423]

15. Kodama T, Freeman M, Rohrer L, Zabrecky J, Matsudaira P, Krieger M. Type I macrophage scavenger receptor contains α -helical and collagen-like coiled coils. *Nature*. 1990; 343:531–535. [PubMed: 2300204]
16. Rohrer L, Freeman M, Kodama T, Penman M, Krieger M. Coiled-coil fibrous domains mediate ligand binding by macrophage scavenger receptor type II. *Nature*. 1990; 343:570–572. [PubMed: 2300208]
17. Dunne DW, Resnick D, Greenberg J, Krieger M, Joiner KA. The type I macrophage scavenger receptor binds to gram-positive bacteria and recognizes lipoteichoic acid. *Proc Natl Acad Sci USA*. 1994; 91:1863–1867. [PubMed: 8127896]
18. Krieger M, Acton S, Ashkenas J, Pearson A, Penman M, Resnick D. Molecular flypaper, host defense, and atherosclerosis. Structure, binding properties, and functions of macrophage scavenger receptors. *J Biol Chem*. 1993; 268:4569–4572. [PubMed: 8383115]
19. El Khoury J, Hickman SE, Thomas CA, Cao L, Silverstein SC, Loike JD. Scavenger receptor-mediated adhesion of microglia to beta-amyloid fibrils. *Nature*. 1996; 382:716–719. [PubMed: 8751442]
20. Means TK, Mylonakis E, Tampakakis E, Colvin RA, Seung E, Puckett L, Tai MF, Stewart CR, Pukkila-Worley R, Hickman SE, et al. Evolutionarily conserved recognition and innate immunity to fungal pathogens by the scavenger receptors SCARF1 and CD36. *J Exp Med*. 2009; 206:637–653. [PubMed: 19237602]
21. Canton J, Neculai D, Grinstein S. Scavenger receptors in homeostasis and immunity. *Nat Rev Immunol*. 2013; 13:621–634. [PubMed: 23928573]
22. Gordon S. Pattern recognition receptors: doubling up for the innate immune response. *Cell*. 2002; 111:927–930. [PubMed: 12507420]
23. Wilkinson K, El Khoury J. Microglial scavenger receptors and their roles in the pathogenesis of Alzheimer's disease. *Int J Alzheimers Dis*. 2012; 2012:489456. [PubMed: 22666621]
24. Fraser I, Hughes D, Gordon S. Divalent cation-independent macrophage adhesion inhibited by monoclonal antibody to murine scavenger receptor. *Nature*. 1993; 364:343–346. [PubMed: 8332192]
25. Suzuki H, Kurihara Y, Takeya M, Kamada N, Kataoka M, Jishage K, Ueda O, Sakaguchi H, Higashi T, Suzuki T, et al. A role for macrophage scavenger receptors in atherosclerosis and susceptibility to infection. *Nature*. 1997; 386:292–296. [PubMed: 9069289]
26. Elomaa O, Kangas M, Sahlberg C, Tuukkanen J, Sormunen R, Liakka A, Thesleff I, Kraal G, Tryggvason K. Cloning of a novel bacteria-binding receptor structurally related to scavenger receptors and expressed in a subset of macrophages. *Cell*. 1995; 80:603–609. [PubMed: 7867067]
27. Jiang Y, Oliver P, Davies KE, Platt N. Identification and characterization of murine SCARA5, a novel class A scavenger receptor that is expressed by populations of epithelial cells. *J Biol Chem*. 2006; 281:11834–11845. [PubMed: 16407294]
28. Huang J, Zheng DL, Qin FS, Cheng N, Chen H, Wan BB, Wang YP, Xiao HS, Han ZG. Genetic and epigenetic silencing of SCARA5 may contribute to human hepatocellular carcinoma by activating FAK signaling. *J Clin Invest*. 2010; 120:223–241. [PubMed: 20038795]
29. Pugin J, Heumann ID, Tomasz A, Kravchenko VV, Akamatsu Y, Nishijima M, Glauser MP, Tobias PS, Ulevitch RJ. CD14 is a pattern recognition receptor. *Immunity*. 1994; 1:509–516. [PubMed: 7534618]
30. Stahl PD, Ezekowitz RA. The mannose receptor is a pattern recognition receptor involved in host defense. *Curr Opin Immunol*. 1998; 10:50–55. [PubMed: 9523111]
31. Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, Protter AA. CD36 is a receptor for oxidized low density lipoprotein. *J Biol Chem*. 1993; 268:11811–11816. [PubMed: 7685021]
32. Prabhudas M, Bowdish D, Drickamer K, Febbraio M, Herz J, Kobzik L, Krieger M, Loike J, Means TK, Moestrup SK, et al. Standardizing scavenger receptor nomenclature. *J Immunol*. 2014; 192:1997–2006. [PubMed: 24563502]
33. Hughes DA, Fraser IP, Gordon S. Murine macrophage scavenger receptor: in vivo expression and function as receptor for macrophage adhesion in lymphoid and non-lymphoid organs. *Eur J Immunol*. 1995; 25:466–473. [PubMed: 7875210]

34. Becker M, Cotena A, Gordon S, Platt N. Expression of the class A macrophage scavenger receptor on specific subpopulations of murine dendritic cells limits their endotoxin response. *Eur J Immunol.* 2006; 36:950–960. [PubMed: 16552714]
35. Sarrias MR, Grønlund J, Padilla O, Madsen J, Holmskov U, Lozano F. The scavenger receptor cysteine-rich (SRCR) domain: an ancient and highly conserved protein module of the innate immune system. *Crit Rev Immunol.* 2004; 24:1–37. [PubMed: 14995912]
36. Resnick D, Pearson A, Krieger M. The SRCR superfamily: a family reminiscent of the Ig superfamily. *Trends Biochem Sci.* 1994; 19:5–8. [PubMed: 8140623]
37. Gough PJ, Greaves DR, Suzuki H, Hakkinen T, Hiltunen MO, Turunen M, Herttuala SY, Kodama T, Gordon S. Analysis of macrophage scavenger receptor (SR-A) expression in human aortic atherosclerotic lesions. *Arterioscler Thromb Vasc Biol.* 1999; 19:461–471. [PubMed: 10073945]
38. Plüddemann A, Neyen C, Gordon S. Macrophage scavenger receptors and host-derived ligands. *Methods.* 2007; 43:207–217. [PubMed: 17920517]
39. Nakamura K, Funakoshi H, Miyamoto K, Tokunaga F, Nakamura T. Molecular cloning and functional characterization of a human scavenger receptor with C-type lectin (SRCL), a novel member of a scavenger receptor family. *Biochem Biophys Res Commun.* 2001; 280:1028–1035. [PubMed: 11162630]
40. Ohtani K, Suzuki Y, Eda S, Kawai T, Kase T, Keshi H, Sakai Y, Fukuoh A, Sakamoto T, Itabe H, et al. The membrane-type collectin CL-P1 is a scavenger receptor on vascular endothelial cells. *J Biol Chem.* 2001; 276:44222–44228. [PubMed: 11564734]
41. Ingersoll MA, Spanbroek R, Lottaz C, Gautier EL, Frankenberger M, Hoffmann R, Lang R, Haniffa M, Collin M, Tacke F, et al. Comparison of gene expression profiles between human and mouse monocyte subsets. *Blood.* 2010; 115:e10–e19. [PubMed: 19965649]
42. Frenkel D, Wilkinson K, Zhao L, Hickman SE, Means TK, Puckett L, Farfara D, Kingery ND, Weiner HL, El Khoury J. Scara1 deficiency impairs clearance of soluble amyloid- β by mononuclear phagocytes and accelerates Alzheimer's-like disease progression. *Nat Commun.* 2013; 4:2030. [PubMed: 23799536]
43. Berwin B, Hart JP, Rice S, Gass C, Pizzo SV, Post SR, Nicchitta CV. Scavenger receptor-A mediates gp96/GRP94 and calreticulin internalization by antigen-presenting cells. *EMBO J.* 2003; 22:6127–6136. [PubMed: 14609958]
44. Hampton RY, Golenbock DT, Penman M, Krieger M, Raetz CR. Recognition and plasma clearance of endotoxin by scavenger receptors. *Nature.* 1991; 352:342–344. [PubMed: 1852209]
45. Acton S, Resnick D, Freeman M, Ekkel Y, Ashkenas J, Krieger M. The collagenous domains of macrophage scavenger receptors and complement component C1q mediate their similar, but not identical, binding specificities for polyanionic ligands. *J Biol Chem.* 1993; 268:3530–3537. [PubMed: 8429029]
46. Whelan FJ, Meehan CJ, Golding GB, McConkey BJ, Bowdish DM. The evolution of the class A scavenger receptors. *BMC Evol Biol.* 2012; 12:227. [PubMed: 23181696]
47. Selman L, Skjodt K, Nielsen O, Floridon C, Holmskov U, Hansen S. Expression and tissue localization of collectin placenta 1 (CL-P1, SRCL) in human tissues. *Mol Immunol.* 2008; 45:3278–3288. [PubMed: 18423602]
48. Arredouani M, Yang Z, Ning Y, Qin G, Soininen R, Tryggvason K, Kobzik L. The scavenger receptor MARCO is required for lung defense against pneumococcal pneumonia and inhaled particles. *J Exp Med.* 2004; 200:267–272. [PubMed: 15263032]
49. van der Laan LJW, Döpp EA, Haworth R, Pikkarainen T, Kangas M, Elomaa O, Dijkstra CD, Gordon S, Tryggvason K, Kraal G. Regulation and functional involvement of macrophage scavenger receptor MARCO in clearance of bacteria *in vivo*. *J Immunol.* 1999; 162:939–947. [PubMed: 9916718]
50. Asch AS, Barnwell J, Silverstein RL, Nachman RL. Isolation of the thrombospondin membrane receptor. *J Clin Invest.* 1987; 79:1054–1061. [PubMed: 2435757]
51. Ockenhouse CF, Chulay JD. *Plasmodium falciparum* sequestration: OKM5 antigen (CD36) mediates cytoadherence of parasitized erythrocytes to a myelomonocytic cell line. *J Infect Dis.* 1988; 157:584–588. [PubMed: 2449507]

52. Coburn CT, Knapp FF Jr, Febbraio M, Beets AL, Silverstein RL, Abumrad NA. Defective uptake and utilization of long chain fatty acids in muscle and adipose tissues of CD36 knockout mice. *J Biol Chem.* 2000; 275:32523–32529. [PubMed: 10913136]
53. Gu X, Trigatti B, Xu S, Acton S, Babitt J, Krieger M. The efficient cellular uptake of high density lipoprotein lipids via scavenger receptor class B type I requires not only receptor-mediated surface binding but also receptor-specific lipid transfer mediated by its extracellular domain. *J Biol Chem.* 1998; 273:26338–26348. [PubMed: 9756864]
54. Stuart LM, Deng J, Silver JM, Takahashi K, Tseng AA, Hennessy EJ, Ezekowitz RAB, Moore KJ. Response to *Staphylococcus aureus* requires CD36-mediated phagocytosis triggered by the COOH-terminal cytoplasmic domain. *J Cell Biol.* 2005; 170:477–485. [PubMed: 16061696]
55. Stewart CR, Stuart LM, Wilkinson K, van Gils JM, Deng J, Halle A, Rayner KJ, Boyer L, Zhong R, Frazier WA, et al. CD36 ligands promote sterile inflammation through assembly of a Toll-like receptor 4 and 6 heterodimer. *Nat Immunol.* 2010; 11:155–161. [PubMed: 20037584]
56. Acton SL, Scherer PE, Lodish HF, Krieger M. Expression cloning of SR-BI, a CD36-related class B scavenger receptor. *J Biol Chem.* 1994; 269:21003–21009. [PubMed: 7520436]
57. Rigotti A, Acton SL, Krieger M. The class B scavenger receptors SR-BI and CD36 are receptors for anionic phospholipids. *J Biol Chem.* 1995; 270:16221–16224. [PubMed: 7541795]
58. Acton S, Rigotti A, Landschulz KT, Xu S, Hobbs HH, Krieger M. Identification of scavenger receptor SR-BI as a high density lipoprotein receptor. *Science.* 1996; 271:518–520. [PubMed: 8560269]
59. Krieger M. Charting the fate of the “good cholesterol”: identification and characterization of the high-density lipoprotein receptor SR-BI. *Annu Rev Bio-chem.* 1999; 68:523–558.
60. Ashraf MZ, Gupta N. Scavenger receptors: implications in atherothrombotic disorders. *Int J Biochem Cell Biol.* 2011; 43:697–700. [PubMed: 21292024]
61. Braun A, Trigatti BL, Post MJ, Sato K, Simons M, Edelberg JM, Rosenberg RD, Schrenzel M, Krieger M. Loss of SR-BI expression leads to the early onset of occlusive atherosclerotic coronary artery disease, spontaneous myocardial infarctions, severe cardiac dysfunction, and premature death in apolipoprotein E-deficient mice. *Circ Res.* 2002; 90:270–276. [PubMed: 11861414]
62. Covey SD, Krieger M, Wang W, Penman M, Trigatti BL. Scavenger receptor class B type I-mediated protection against atherosclerosis in LDL receptor-negative mice involves its expression in bone marrow-derived cells. *Arterioscler Thromb Vasc Biol.* 2003; 23:1589–1594. [PubMed: 12829524]
63. Webb NR, Connell PM, Graf GA, Smart EJ, de Villiers WJS, de Beer FC, van der Westhuyzen DR. SR-BII, an isoform of the scavenger receptor BI containing an alternate cytoplasmic tail, mediates lipid transfer between high density lipoprotein and cells. *J Biol Chem.* 1998; 273:15241–15248. [PubMed: 9614139]
64. Song L, Lee C, Schindler C. Deletion of the murine scavenger receptor CD68. *J Lipid Res.* 2011; 52:1542–1550. [PubMed: 21572087]
65. Ramprasad MP, Terpstra V, Kondratenko N, Quehenberger O, Steinberg D. Cell surface expression of mouse macrosialin and human CD68 and their role as macrophage receptors for oxidized low density lipoprotein. *Proc Natl Acad Sci USA.* 1996; 93:14833–14838. [PubMed: 8962141]
66. Hao L, Klein J, Nei M. Heterogeneous but conserved natural killer receptor gene complexes in four major orders of mammals. *Proc Natl Acad Sci USA.* 2006; 103:3192–3197. [PubMed: 16492762]
67. Yoshimoto R, Fujita Y, Kakino A, Iwamoto S, Takaya T, Sawamura T. The discovery of LOX-1, its ligands and clinical significance. *Cardiovasc Drugs Ther.* 2011; 25:379–391. [PubMed: 21805404]
68. Sawamura T, Kume N, Aoyama T, Moriwaki H, Hoshikawa H, Aiba Y, Tanaka T, Miwa S, Katsura Y, Kita T, Masaki T. An endothelial receptor for oxidized low-density lipoprotein. *Nature.* 1997; 386:73–77. [PubMed: 9052782]
69. Biocca S, Filesi I, Mango R, Maggiore L, Baldini F, Vecchione L, Viola A, Citro G, Federici G, Romeo F, Novelli G. The splice variant LOXIN inhibits LOX-1 receptor function through hetero-oligomerization. *J Mol Cell Cardiol.* 2008; 44:561–570. [PubMed: 18191942]
70. Herre J, Willment JA, Gordon S, Brown GD. The role of Dectin-1 in antifungal immunity. *Crit Rev Immunol.* 2004; 24:193–203. [PubMed: 15482254]


71. Drummond RA, Brown GD. The role of Dectin-1 in the host defence against fungal infections. *Curr Opin Microbiol.* 2011; 14:392–399. [PubMed: 21803640]
72. Reid DM, Montoya M, Taylor PR, Borrow P, Gordon S, Brown GD, Wong SY. Expression of the beta-glucan receptor, Dectin-1, on murine leukocytes in situ correlates with its function in pathogen recognition and reveals potential roles in leukocyte interactions. *J Leukoc Biol.* 2004; 76:86–94. [PubMed: 15107454]
73. Wollenberg A, Mommaas M, Oppel T, Schottdorf EM, Günther S, Moderer M. Expression and function of the mannose receptor CD206 on epidermal dendritic cells in inflammatory skin diseases. *J Invest Dermatol.* 2002; 118:327–334. [PubMed: 11841552]
74. Ashwell G, Morell AG. The role of surface carbohydrates in the hepatic recognition and transport of circulating glycoproteins. *Adv Enzymol Relat Areas Mol Biol.* 1974; 41:99–128. [PubMed: 4609051]
75. Ashwell G, Steer CJ. Hepatic recognition and catabolism of serum glycoproteins. *JAMA.* 1981; 246:2358–2364. [PubMed: 6271991]
76. Stockert RJ. The asialoglycoprotein receptor: relationships between structure, function, and expression. *Physiol Rev.* 1995; 75:591–609. [PubMed: 7624395]
77. Weigel PH. Endocytosis and function of the hepatic asialoglycoprotein receptor. *Subcell Biochem.* 1993; 19:125–161. [PubMed: 8385817]
78. Casey CA, Kragoskowsky SL, Sorrell MF, Tuma DJ. Chronic ethanol administration impairs the binding and endocytosis of asialo-orosomucoid in isolated hepatocytes. *J Biol Chem.* 1987; 262:2704–2710. [PubMed: 3818618]
79. Debanne MT, Evans WH, Flint N, Regoeczi E. Receptor-rich intracellular membrane vesicles transporting asialotransferrin and insulin in liver. *Nature.* 1982; 298:398–400. [PubMed: 6283390]
80. Tomana M, Phillips JO, Kulhavy R, Mestecky J. Carbohydrate-mediated clearance of secretory IgA from the circulation. *Mol Immunol.* 1985; 22:887–892. [PubMed: 2413348]
81. Dini L, Autuori F, Lentini A, Oliverio S, Piacentini M. The clearance of apoptotic cells in the liver is mediated by the asialoglycoprotein receptor. *FEBS Lett.* 1992; 296:174–178. [PubMed: 1370803]
82. Aziz-Seible RS, Lee SM, Kharbanda KK, McVicker BL, Casey CA. Ethanol feeding potentiates the pro-inflammatory response of Kupffer cells to cellular fibronectin. *Alcohol Clin Exp Res.* 2011; 35:717–725. [PubMed: 21223308]
83. Rotundo RF, Rebres RA, Mckeown-Longo PJ, Blumenstock FA, Saba TM. Circulating cellular fibronectin may be a natural ligand for the hepatic asialoglycoprotein receptor: possible pathway for fibronectin deposition and turnover in the rat liver. *Hepatology.* 1998; 28:475–485. [PubMed: 9696014]
84. Pike AF, Kramer NI, Blaauboer BJ, Seinen W, Brands R. A novel hypothesis for an alkaline phosphatase ‘rescue’ mechanism in the hepatic acute phase immune response. *Biochim Biophys Acta.* 2013; 1832:2044–2056. [PubMed: 23899605]
85. Meijer DK, Scholtens HB, Hardonk MJ. The role of the liver in clearance of glycoproteins from the general circulation, with special reference to intestinal alkaline phosphatase. *Pharm Weekbl Sci.* 1982; 4:57–70. [PubMed: 6180375]
86. Bertolino P, Bowen DG, McCaughan GW, Fazekas de St Groth B. Antigen-specific primary activation of CD8+ T cells within the liver. *J Immunol.* 2001; 166:5430–5438. [PubMed: 11313380]
87. Guy CS, Rankin SL, Michalak TI. Hepatocyte cytotoxicity is facilitated by asialoglycoprotein receptor. *Hepatology.* 2011; 54:1043–1050. [PubMed: 21656538]
88. McVicker BL, Tuma DJ, Kharbanda KK, Kubik JL, Casey CA. Effect of chronic ethanol administration on the in vitro production of proinflammatory cytokines by rat Kupffer cells in the presence of apoptotic cells. *Alcohol Clin Exp Res.* 2007; 31:122–129. [PubMed: 17207110]
89. McVicker BL, Tuma DJ, Kubik JA, Hindemith AM, Baldwin CR, Casey CA. The effect of ethanol on asialoglycoprotein receptor-mediated phagocytosis of apoptotic cells by rat hepatocytes. *Hepatology.* 2002; 36:1478–1487. [PubMed: 12447874]

90. McVicker BL, Thiele GM, Casey CA, Osna NA, Tuma DJ. Susceptibility to T cell-mediated liver injury is enhanced in asialoglycoprotein receptor-deficient mice. *Int Immunopharmacol.* 2013; 16:17–26. [PubMed: 23538026]
91. McVicker BL, Thiele GM, Tuma DJ, Casey CA. Hepatocyte-mediated cytotoxicity and host defense mechanisms in the alcohol-injured liver. *Hepatol Int.* 2014; 8(Suppl. 2):432–438. [PubMed: 26201322]
92. Bridge RGJ, Kubik JL, Casey CA, Tuma DJ, McVicker BL. The attenuation of inflammatory liver injury by alkaline phosphatase is impaired in livers lacking asialoglycoprotein receptors: role of the major hepatic trafficking system in detoxification processes. *Hepatology.* 2015; 62:200A.
93. Newton DP, Casey TD, Mc CA, Vicker BL. Inflammatory liver injury is associated with altered alkaline phosphatase activity and hepatic asialoglycoprotein receptor function. *Hepatology.* 2014; 60:229A.
94. Braun JR, Willnow TE, Ishibashi S, Ashwell G, Herz J. The major subunit of the asialoglycoprotein receptor is expressed on the hepatocellular surface in mice lacking the minor receptor subunit. *J Biol Chem.* 1996; 271:21160–21166. [PubMed: 8702886]
95. Adachi H, Tsujimoto M, Arai H, Inoue K. Expression cloning of a novel scavenger receptor from human endothelial cells. *J Biol Chem.* 1997; 272:31217–31220. [PubMed: 9395444]
96. Ramirez-Ortiz ZG, Pendergraft WF III, Prasad A, Byrne MH, Iram T, Blanchette CJ, Luster AD, Hacoheh N, El Khoury J, Means TK. The scavenger receptor SCARF1 mediates the clearance of apoptotic cells and prevents autoimmunity. *Nat Immunol.* 2013; 14:917–926. [PubMed: 23892722]
97. Ishii J, Adachi H, Aoki J, Koizumi H, Tomita S, Suzuki T, Tsujimoto M, Inoue K, Arai H. SREC-II, a new member of the scavenger receptor type F family, trans-interacts with SREC-I through its extracellular domain. *J Biol Chem.* 2002; 277:39696–39702. [PubMed: 12154095]
98. Iram T, Ramirez-Ortiz Z, Byrne MH, Coleman UA, Kingery ND, Means TK, Frenkel D, El Khoury J. Megf10 is a receptor for C1Q that mediates clearance of apoptotic cells by astrocytes. *J Neurosci.* 2016; 36:5185–5192. [PubMed: 27170117]
99. Shimaoka T, Kume N, Minami M, Hayashida K, Kataoka H, Kita T, Yonehara S. Molecular cloning of a novel scavenger receptor for oxidized low density lipoprotein, SR-PSOX, on macrophages. *J Biol Chem.* 2000; 275:40663–40666. [PubMed: 11060282]
100. Matloubian M, David A, Engel S, Ryan JE, Cyster JG. A transmembrane CXC chemokine is a ligand for HIV-coreceptor Bonzo. *Nat Immunol.* 2000; 1:298–304. [PubMed: 11017100]
101. Sheikine Y, Sirsjo A. CXCL16/SR-PSOX—a friend or a foe in atherosclerosis? *Atherosclerosis.* 2008; 197:487–495. [PubMed: 18191863]
102. Wilbanks A, Zondlo SC, Murphy K, Mak S, Soler D, Langdon P, Andrew DP, Wu L, Briskin M. Expression cloning of the STRL33/BONZO/TYMSTRligand reveals elements of CC, CXC, and CX3C chemokines. *J Immunol.* 2001; 166:5145–5154. [PubMed: 11290797]
103. Fukumoto N, Shimaoka T, Fujimura H, Sakoda S, Tanaka M, Kita T, Yonehara S. Critical roles of CXC chemokine ligand 16/scavenger receptor that binds phosphatidylserine and oxidized lipoprotein in the pathogenesis of both acute and adoptive transfer experimental autoimmune encephalomyelitis. *J Immunol.* 2004; 173:1620–1627. [PubMed: 15265890]
104. Shimaoka T, Nakayama T, Hieshima K, Kume N, Fukumoto N, Minami M, Hayashida K, Kita T, Yoshie O, Yonehara S. Chemokines generally exhibit scavenger receptor activity through their receptor-binding domain. *J Biol Chem.* 2004; 279:26807–26810. [PubMed: 15128749]
105. Park SY, Jung MY, Lee SJ, Kang KB, Gratchev A, Riabov V, Kzhyshkowska J, Kim IS. Stabilin-1 mediates phosphatidylserine-dependent clearance of cell corpses in alternatively activated macrophages. *J Cell Sci.* 2009; 122:3365–3373. [PubMed: 19726632]
106. Park SY, Jung MY, Kim HJ, Lee SJ, Kim SY, Lee BH, Kwon TH, Park RW, Kim IS. Rapid cell corpse clearance by stabilin-2, a membrane phosphatidylserine receptor. *Cell Death Differ.* 2008; 15:192–201. [PubMed: 17962816]
107. Schledzewski K, Géraud C, Arnold B, Wang S, Gröne HJ, Kempf T, Wollert KC, Straub BK, Schirmacher P, Demory A, et al. Deficiency of liver sinusoidal scavenger receptors stabilin-1 and -2 in mice causes glomerulofibrotic nephropathy via impaired hepatic clearance of noxious blood factors. *J Clin Invest.* 2011; 121:703–714. [PubMed: 21293057]

108. Rodamilans B, Muñoz IG, Bragado-Nilsson E, Sarrias MR, Padilla O, Blanco FJ, Lozano F, Montoya G. Crystal structure of the third extracellular domain of CD5 reveals the fold of a group B scavenger cysteine-rich receptor domain. *J Biol Chem.* 2007; 282:12669–12677. [PubMed: 17322294]
109. Kristiansen M, Graversen JH, Jacobsen C, Sonne O, Hoffman HJ, Law SK, Moestrup SK. Identification of the haemoglobin scavenger receptor. *Nature.* 2001; 409:198–201. [PubMed: 11196644]
110. Thomsen JH, Etzerodt A, Svendsen P, Moestrup SK. The haptoglobin-CD163-heme oxygenase-1 pathway for hemoglobin scavenging. *Oxid Med Cell Longev.* 2013; doi: 10.1155/2013/523652
111. Fabrick BO, van Bruggen R, Deng DM, Ligtenberg AJ, Nazmi K, Schornagel K, Vloet RP, Dijkstra CD, van den Berg TK. The macrophage scavenger receptor CD163 functions as an innate immune sensor for bacteria. *Blood.* 2009; 113:887–892. [PubMed: 18849484]
112. Nielsen MJ, Madsen M, Møller HJ, Moestrup SK. The macrophage scavenger receptor CD163: endocytic properties of cytoplasmic tail variants. *J Leukoc Biol.* 2006; 79:837–845. [PubMed: 16434690]
113. Telfer JC, Baldwin CL. Bovine gamma delta T cells and the function of gamma delta T cell specific WC1 co-receptors. *Cell Immunol.* 2015; 296:76–86. [PubMed: 26008759]
114. Moeller JB, Nielsen MJ, Reichhardt MP, Schlosser A, Sorensen GL, Nielsen O, Tornøe I, Grønlund J, Nielsen ME, Jørgensen JS, et al. CD163-L1 is an endocytic macrophage protein strongly regulated by mediators in the inflammatory response. *J Immunol.* 2012; 188:2399–2409. [PubMed: 22279103]
115. Holm D, Fink DR, Steffensen MA, Schlosser A, Nielsen O, Moeller JB, Holmskov U. Characterization of a novel human scavenger receptor cysteine-rich molecule SCART1 expressed by lymphocytes. *Immunobiology.* 2013; 218:408–417. [PubMed: 22795646]
116. Chen C, Herzig CT, Alexander LJ, Keele JW, McDanel TG, Telfer JC, Baldwin CL. Gene number determination and genetic polymorphism of the gamma delta T cell co-receptor WC1 genes. *BMC Genet.* 2012; 13:86. [PubMed: 23072335]
117. Herzig CT, Baldwin CL. Genomic organization and classification of the bovine WC1 genes and expression by peripheral blood gamma delta T cells. *BMC Genomics.* 2009; 10:191. [PubMed: 19393067]
118. Herzig CT, Waters RW, Baldwin CL, Telfer JC. Evolution of the CD163 family and its relationship to the bovine gamma delta T cell co-receptor WC1. *BMC Evol Biol.* 2010; 10:181. [PubMed: 20550670]
119. Blumerman SL, Herzig CT, Baldwin CL. WC1+ gammadelta T cell memory population is induced by killed bacterial vaccine. *Eur J Immunol.* 2007; 37:1204–1216. [PubMed: 17429840]
120. Rogers AN, Vanburen DG, Hedblom EE, Tilahun ME, Telfer JC, Baldwin CL. Gammadelta T cell function varies with the expressed WC1 coreceptor. *J Immunol.* 2005; 174:3386–3393. [PubMed: 15749871]
121. Wang F, Herzig CT, Chen C, Hsu H, Baldwin CL, Telfer JC. Scavenger receptor WC1 contributes to the $\gamma\delta$ T cell response to *Leptospira*. *Mol Immunol.* 2011; 48:801–809. [PubMed: 21227509]
122. Hsu H, Chen C, Nennering A, Holz L, Baldwin CL, Telfer JC. WC1 is a hybrid $\gamma\delta$ TCR coreceptor and pattern recognition receptor for pathogenic bacteria. *J Immunol.* 2015; 194:2280–2288. [PubMed: 25632007]
123. Ibrahim ZA, Armour CL, Phipps S, Sukkar MB. RAGE and TLRs: relatives, friends or neighbours? *Mol Immunol.* 2013; 56:739–744. [PubMed: 23954397]
124. Lee EJ, Park JH. Receptor for advanced glycation endproducts (RAGE), its ligands, and soluble RAGE: potential biomarkers for diagnosis and therapeutic targets for human renal diseases. *Genomics Inform.* 2013; 11:224–229. [PubMed: 24465234]
125. Bangert A, Andrassy M, Müller AM, Bockstahler M, Fischer A, Volz CH, Leib C, Göser S, Korkmaz-Icöz S, Zittrich S, et al. Critical role of RAGE and HMGB1 in inflammatory heart disease. *Proc Natl Acad Sci USA.* 2016; 113:E155–E164. [PubMed: 26715748]
126. Xie J, Méndez JD, Méndez-Valenzuela V, Aguilar-Hernández MM. Cellular signalling of the receptor for advanced glycation end products (RAGE). *Cell Signal.* 2013; 25:2185–2197. [PubMed: 23838007]

127. Naor D, Sionov RV, Ish-Shalom D. CD44: structure, function, and association with the malignant process. *Adv Cancer Res.* 1997; 71:241–319. [PubMed: 9111868]
128. Dzwonek J, Wilczynski GM. CD44: molecular interactions, signaling and functions in the nervous system. *Front Cell Neurosci.* 2015; 9:175. [PubMed: 25999819]
129. Anderegg U, Eichenberg T, Parthaune T, Haiduk C, Saalbach A, Milkova L, Ludwig A, Grosche J, Averbeck M, Gebhardt C, et al. ADAM10 is the constitutive functional sheddase of CD44 in human melanoma cells. *J Invest Dermatol.* 2009; 129:1471–1482. [PubMed: 18971959]
130. Päll T, Pink A, Kasak L, Turkina M, Anderson W, Valkna A, Kogerman P. Soluble CD44 interacts with intermediate filament protein vimentin on endothelial cell surface. *PLoS One.* 2011; 6:e29305. [PubMed: 22216242]
131. Culty M, Nguyen HA, Underhill CB. The hyaluronan receptor (CD44) participates in the uptake and degradation of hyaluronan. *J Cell Biol.* 1992; 116:1055–1062. [PubMed: 1370836]
132. Ehnis T, Dieterich W, Bauer M, Lampe B, Schuppan D. A chondroitin/dermatan sulfate form of CD44 is a receptor for collagen XIV (undulin). *Exp Cell Res.* 1996; 229:388–397. [PubMed: 8986622]
133. Turley E, Moore D. Hyaluronate binding proteins also bind to fibronectin, laminin and collagen. *Biochem Biophys Res Commun.* 1984; 121:808–814. [PubMed: 6204647]
134. Cywes C, Wessels MR. Group A *Streptococcus* tissue invasion by CD44-mediated cell signalling. *Nature.* 2001; 414:648–652. [PubMed: 11740562]
135. Wakahara K, Kobayashi H, Yagyu T, Matsuzaki H, Kondo T, Kurita N, Sekino H, Inagaki K, Suzuki M, Kanayama N, Terao T. Bikunin down-regulates heterodimerization between CD44 and growth factor receptors and subsequently suppresses agonist-mediated signaling. *J Cell Biochem.* 2005; 94:995–1009. [PubMed: 15597342]
136. Dianzani U, Bragardo M, Tosti A, Ruggeri L, Volpi I, Casucci M, Bottarel F, Feito MJ, Bonisconi S, Velardi A. CD44 signaling through p56lck involves lateral association with CD4 in human CD4+ T cells. *Int Immunol.* 1999; 11:1085–1092. [PubMed: 10383941]
137. Kawana H, Karaki H, Higashi M, Miyazaki M, Hilberg F, Kitagawa M, Harigaya K. CD44 suppresses TLR-mediated inflammation. *J Immunol.* 2008; 180:4235–4245. [PubMed: 18322236]
138. Taylor KR, Yamasaki K, Radek KA, Di Nardo A, Goodarzi H, Golenbock D, Beutler B, Gallo RL. Recognition of hyaluronan released in sterile injury involves a unique receptor complex dependent on Toll-like receptor 4, CD44, and MD-2. *J Biol Chem.* 2007; 282:18265–18275. [PubMed: 17400552]
139. Ponta H, Sherman L, Herrlich PA. CD44: from adhesion molecules to signalling regulators. *Nat Rev Mol Cell Biol.* 2003; 4:33–45. [PubMed: 12511867]
140. Thankamony SP, Knudson W. Acylation of CD44 and its association with lipid rafts are required for receptor and hyaluronan endocytosis. *J Biol Chem.* 2006; 281:34601–34609. [PubMed: 16945930]
141. Ruppert SM, Hawn TR, Arrigoni A, Wight TN, Bollyky PL. Tissue integrity signals communicated by high-molecular weight hyaluronan and the resolution of inflammation. *Immunol Res.* 2014; 58:186–192. [PubMed: 24614953]
142. Jiang D, Liang J, Noble PW. Hyaluronan as an immune regulator in human diseases. *Physiol Rev.* 2011; 91:221–264. [PubMed: 21248167]
143. Teder P, Vandivier RW, Jiang D, Liang J, Cohn L, Puré E, Henson PM, Noble PW. Resolution of lung inflammation by CD44. *Science.* 2002; 296:155–158. [PubMed: 11935029]
144. Draffin JE, McFarlane S, Hill A, Johnston PG, Waugh DJ. CD44 potentiates the adherence of metastatic prostate and breast cancer cells to bone marrow endothelial cells. *Cancer Res.* 2004; 64:5702–5711. [PubMed: 15313910]
145. Bollyky PL, Wu RP, Falk BA, Lord JD, Long SA, Preisinger A, Teng B, Holt GE, Standifer NE, Braun KR, et al. ECM components guide IL-10 producing regulatory T-cell (TR1) induction from effector memory T-cell precursors. *Proc Natl Acad Sci USA.* 2011; 108:7938–7943. [PubMed: 21518860]
146. Marhaba R, Zöller M. CD44 in cancer progression: adhesion, migration and growth regulation. *J Mol Histol.* 2004; 35:211–231. [PubMed: 15339042]

147. Guan H, Nagarkatti PS, Nagarkatti M. Role of CD44 in the differentiation of Th1 and Th2 cells: CD44-deficiency enhances the development of Th2 effectors in response to sheep RBC and chicken ovalbumin. *J Immunol.* 2009; 183:172–180. [PubMed: 19542428]
148. Bollyky PL, Falk BA, Long SA, Preisinger A, Braun KR, Wu RP, Evanko SP, Buckner JH, Wight TN, Nepom GT. CD44 co-stimulation promotes FoxP3+ regulatory T cell persistence and function via production of IL-2, IL-10, and TGF-beta. *J Immunol.* 2009; 183:2232–2241. [PubMed: 19635906]
149. Krieger M, Herz J. Structures and functions of multiligand lipoprotein receptors: macrophage scavenger receptors and LDL receptor-related protein (LRP). *Annu Rev Biochem.* 1994; 63:601–637. [PubMed: 7979249]
150. Herz J, Strickland DK. LRP: a multifunctional scavenger and signaling receptor. *J Clin Invest.* 2001; 108:779–784. [PubMed: 11560943]
151. Dieckmann M, Dietrich MF, Herz J. Lipoprotein receptors—an evolutionarily ancient multifunctional receptor family. *Biol Chem.* 2010; 391:1341–1363. [PubMed: 20868222]
152. Dachy A, Paquot F, Debray G, Bovy C, Christensen EI, Collard L, Jouret F. In-depth phenotyping of a Donnai-Barrow patient helps clarify proximal tubule dysfunction. *Pediatr Nephrol.* 2015; 30:1027–1031. [PubMed: 25822460]
153. Kanekiyo T, Bu G. The low-density lipoprotein receptor-related protein 1 and amyloid- β clearance in Alzheimer's disease. *Front Aging Neurosci.* 2014; 6:93. [PubMed: 24904407]
154. Strickland DK, Au DT, Cunfer P, Muratoglu SC. Low-density lipoprotein receptor-related protein-1: role in the regulation of vascular integrity. *Arterioscler Thromb Vasc Biol.* 2014; 34:487–498. [PubMed: 24504736]
155. Gonias SL, Campana WM. LDL receptor-related protein-1: a regulator of inflammation in atherosclerosis, cancer, and injury to the nervous system. *Am J Pathol.* 2014; 184:18–27. [PubMed: 24128688]
156. Oh SJ, Kim TH, Lim JM, Jeong JW. Progesterone induces expression of Lrp2 in the murine uterus. *Biochem Biophys Res Commun.* 2013; 441:175–179. [PubMed: 24140060]
157. Bartolome F, Antequera D, Tavares E, Pascual C, Maldonado R, Camins A, Carro E. Obesity and neuroinflammatory phenotype in mice lacking endothelial megalin. *J Neuroinflammation.* 2017; 14:26. [PubMed: 28143489]
158. Cabezas F, Lagos J, Céspedes C, Vio CP, Bronfman M, Marzolo MP. Megalin/LRP2 expression is induced by peroxisome proliferator-activated receptor -alpha and -gamma: implications for PPARs' roles in renal function. *PLoS One.* 2011; 6:e16794. [PubMed: 21311715]
159. Warner FJ, Smith AI, Hooper NM, Turner AJ. Angiotensin-converting enzyme-2: a molecular and cellular perspective. *Cell Mol Life Sci.* 2004; 61:2704–2713. [PubMed: 15549171]
160. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature.* 2003; 426:450–454. [PubMed: 14647384]
161. Danziger RS. Aminopeptidase N in arterial hypertension. *Heart Fail Rev.* 2008; 13:293–298. [PubMed: 18008160]
162. Cong Y, Li X, Bai Y, Lv X, Herrler G, Enjuanes L, Zhou X, Qu B, Meng F, Cong C, et al. Porcine aminopeptidase N mediated polarized infection by porcine epidemic diarrhea virus in target cells. *Virology.* 2015; 478:1–8. [PubMed: 25681796]
163. Yamayoshi S, Yamashita Y, Li J, Hanagata N, Minowa T, Takemura T, Koike S. Scavenger receptor B2 is a cellular receptor for enterovirus 71. *Nat Med.* 2009; 15:798–801. [PubMed: 19543282]
164. Neculai D, Schwake M, Ravichandran M, Zunke F, Collins RF, Peters J, Neculai M, Plumb J, Loppnau P, Pizarro JC, et al. Structure of LIMP-2 provides functional insights with implications for SR-BI and CD36. *Nature.* 2013; 504:172–176. [PubMed: 24162852]
165. Yamayoshi S, Iizuka S, Yamashita T, Minagawa H, Mizuta K, Okamoto M, Nishimura H, Sanjoh K, Katsushima N, Itagaki T, et al. Human SCARB2-dependent infection by coxsackievirus A7, A14, and A16 and enterovirus 71. *J Virol.* 2012; 86:5686–5696. [PubMed: 22438546]

**FIGURE 1.**

Proposed scavenger receptor nomenclature formula. SR stands for scavenger receptor. SR is followed by a hyphen, then a capital letter represents the class of scavenger receptor (A–J), followed by an Arabic numeral representing the type of molecule within the class (the numbering is based on the order in which the molecules were identified). Alternatively spliced forms of a molecule will be designated as 1.1, 1.2, etc. For existing spliced variants, the longest variant in terms of amino acid sequence will be given the first number.




FIGURE 2. Schematic representation of different classes of scavenger receptors. Mammalian scavenger receptors are divided into 12 classes. The various protein or carbohydrate domains are identified in the key within the figure. Class C is not represented here because it is present only in *Drosophila melanogaster*.

Table I

Workshop participants and their areas of expertise

Participant	Affiliation	Area of Expertise
Dawn Bowdish	McMaster University, Canada	Macrophage scavenger receptors in host defense
Kurt Drickamer	Imperial College, U.K.	C-type lectin and glycan recognition
Joseph El Khoury	Massachusetts General Hospital, Harvard Medical School, MA	Macrophage and microglial scavenger receptors in inflammation
Maria Febbraio	University of Alberta, Canada	CD36 signaling
Joachim Herz	University of Texas Southwestern Medical Center, TX	Functions of LDLR-related proteins
Lester Kobzik	Harvard School of Public Health, MA	MARCO and lung inflammation
Monty Krieger	Massachusetts Institute of Technology, MA	Scavenger receptors in host defense and lipoprotein and lipid metabolism
John Loike	Columbia University, NY	Scavenger receptors in neuropathology
Terry Means	Massachusetts General Hospital, Harvard Medical School, MA	Scavenger receptors and antifungal immune mechanisms
Soren Moestrup	Aarhus University, Denmark	Hemoglobin scavenger receptor
Philip Murphy	NIAID, NIH, MD	G-protein coupled receptors in leukocyte trafficking
Steven Post	University of Arkansas for Medical Sciences, AR	Scavenger receptor A and atherosclerosis
Tatsuya Sawamura	Shinshu University, Japan	Oxidized LDLR (LOX-1) and vascular inflammation
Samuel Silverstein	Columbia University, NY	Roles of scavenger receptors on mononuclear phagocytes in chronic inflammatory diseases
Xiang-Yang Wang	Virginia Commonwealth University, VA	Scavenger receptors in immune modulation and host response
Cynthia Baldwin	University of Massachusetts, Amherst, MA	$\gamma\delta$ T cells and WC1 coreceptor family in the bovine model
Paul Bollyky	Stanford University, CA	Immune dysregulation in wound healing, biofilms
Benita McVicker	University of Nebraska, NE	Hepatocellular-lymphocyte interactions
Robert Speth	Nova Southeastern University, FL	Regulation of the cardiovascular system
Janice Telfer	University of Massachusetts, Amherst, MA	Immune development; characterization of SRCR family of receptors in cattle
Geoffrey Thiele	University of Nebraska, NE	Peripheral vascular disease, cardiovascular disease, and immune tolerance
Samuel Wright	Cardiovascular Therapeutics, PA	Cardiovascular disease, atherosclerosis, and inflammation

Table II

Summary of current mouse scavenger receptor nomenclature and proposed changes

Current NCBI Gene Name	Alternative Names	Consensus Nomenclature	Accession Number	NCBI Gene ID	Chromosome
MSR1	SR-AI, SCARA1	SR-AI	NM_031195	20288	8
Alternatively spliced form of SR-AI	SR-AII	SR-AI.1	NM_001113326	20288	8
MARCO	SCARA2, Ly112	SR-A6	NM_010766	17167	1
COLLEC12	SCARA4, SRCL1, SRCLII, SRCLIII, CL-PI	SR-A4	NM_130449	140792	18
SCARA5	TESR	SR-A5	NM_028903	71145	14
CD36	SCARB3, PAS4	SR-B2	NM_001159558	12491	5
SCARB1	SR-BI, CD36L1	SR-B1	NM_016741	20778	5
Alternatively spliced form of SR-B1	SR-BII	SR-B1.1	NM_001205082	20778	5
CD68	gp110, SCARD1, Macrostialin	SR-D1	NM_009853	12514	11
OLR1	LOX-1, SCARE1	SR-E1	NM_138648	108078	6
CLEC7A	DECTIN-1	SR-E2	NM_020008	56644	6
MRC1	CD206, MR	SR-E3	NM_008625	17533	2
ASGPR	ASGR1, HL-1	SR-E4	NM_001291131	11889	11
SCARF1	SREC-1	SR-F1	NM_001004157	380713	11
MEGF10	GM331	SR-F2	NM_1001979	70417	18
CXCL16	SR-PSOX	SR-G1	NM_023158	66102	11
STAB1	FEEL-1	SR-H1	NM_138672	192187	14
STAB2	FEEL-2	SR-H2	NM_138673	192188	10
CD163	<i>MI30</i> , <i>CD163A</i>	SR-I1	NM_053094	93671	6
CD163L1	M160, CD163b	SR-I2	NM_172909	244233	7
mSCART1	mFLJ00268, CD163c- α	SR-I3	NM_001128146	244234	7
mSCART2	SCART2	SR-I4	NM_175533	244238	7
RAGE (membrane form)	AGER	SR-J1	NM_007425	11596	17
RAGE (soluble form)	AGER (soluble)	SR-J1.1	(AB207883)	11596	17
CD44	Ly24, Pgp-1, HERMES	SR-K1	NM_001039151	12505	2
LRP1	CD91, A2MIR	SR-L1	NM_008512	16971	10
LRP2	Megalyn, gp330	SR-L2	NM_001081088	14725	2
SRCRB4D		To be named	NM_001160366	109267	5

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Current NCBI Gene Name	Alternative Names	Consensus Nomenclature	Accession Number	NCBI Gene ID	Chromosome
SSC5D		To be named	NM_173008	269855	7
CD14		To be named	NM_009841	12475	18
Ly75	CD205, DEC-205	To be named	NM_013825	17076	2
CD207	Langerin	To be named	NM_144943	246278	6
CD209a	DC-SIGN, CLEC4L	To be named	NM_133238	170786	8

NCBI, National Center for Biotechnology Information.

Table III

Summary of current human scavenger receptor nomenclature and proposed changes

Current NCBI Gene Name	Alternative Names	Consensus Nomenclature	Accession Number	NCBI Gene ID	Chromosome
MSR1	SR-AI, CD204, SCARA1	SR-A1	NM_138715	4481	8
Alternatively spliced form of SR-AI	SR-AII	SR-A1.1	NM_002445	2960	8
MARCO	SCARA2	SR-A6	NM_006770	8685	2
SCARA3	MSRL1	SR-A3	NM_016240	51435	8
COLEC12	SCARA4, SRCL1, SRCLII, CL-P1	SR-A4	NM_130386	81035	18
SCARA5	TESR, NET33	SR-A5	NM_173833	286133	8
CD36	SCARB3, FAT, GPIV, PAS4	SR-B2	NM_001001548	948	7
SCARB1	SR-BI, CD36L1	SR-B1	NM_005505	949	12
CD68	gp110, SCARD1, LAMP4	SR-D1	NM_001251	968	17
OLR1	LOX-1, SCARE1, CLEC8A	SR-E1	NM_002543	4973	12
Alternatively spliced form of SRE-1	LOXIN	SR-E1.1	NM_001172632	4973	12
CLEC7A	Dectin-1, SCARE2, CD369	SR-E2	NM_197947	64581	12
CD206/MRC1	Mannose receptor 1	SR-E3	NM_002438	4360	10
ASGPR	ASGR1, CLEC4HI, HL-1	SR-E4	NM_001197216	432	17
SCARF1	SREC-1	SR-F1	NM_003693	8578	17
MEGF10	EMARDD	SR-F2	NM_032446	84466	5
CXCL16	SR-PSOX	SR-G1	NM_001100812	58191	17
STAB1	FEEL-1	SR-H1	NM_015136	608560	3
STAB2	FEEL-2	SR-H2	NM_017564	55576	12
CD163	M130, CD163A	SR-I1	NM_004244	9332	12
CD163L1	CD163B, M160	SR-I2	NM_001297650	283316	12
SCART1	CD163c- α	SR-I3	NR_002934.3	619207	10
RAGE (membrane form)	AGER	SR-J1	NM_001136	177	6
RAGE (soluble form)	AGER	SR-J1.1	(AB061668)	177	6
CD44	Pgp-1	SR-K1	NM_000610	960	11
LRP1	A2MR, APOER, CD91	SR-L1	NM_002332	4035	12
LRP2	Megalin, gp330	SR-L2	NM_004525	4036	2
SRCRB4D		To be named	NM_080744	136853	7

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Current NCBI Gene Name	Alternative Names	Consensus Nomenclature	Accession Number	NCBI Gene ID	Chromosome
SSC5D		To be named	NM_001144950	284297	19
CD14		To be named	NM_000591	929	5
Ly75/CD205		To be named	NM_002349	4065	2
CD207/Langerin		To be named	NM_015717	50489	2
CD209/DC-SIGN	CLEC4L	To be named	NM_021155	30835	19

NCBI, National Center for Biotechnology Information.

Table IV

Bovine scavenger receptors and proposed nomenclature changes

Current NCBI Gene Name	Alternative Names	Consensus Nomenclature	Accession Number	NCBI Gene ID	Chromosome
CD163	CD163A	SR-II	NM_001163413	GLEAN_00453, BGD:BT10376, 533844	5
CD163c- α	SCART1	SR-I3		GLEAN_14183	Unplaced
WC1-1		SR-I5	FJ031186	GLEAN_13183, BGD:BT11207, 783109	5
WC1-2		SR-I6	HQ687669	GLEAN_13182	5
WC1-3	BoWC1.1	SR-I7	FJ031191	GLEAN_13181	5
WC1-4	BoWC1.2	SR-I8	FJ031202, XM_001250405.1	GLEAN_13179, 751809	5
WC1-5		SR-I9	NM_176651	GLEAN_13176, 338056	5
WC1-6		SR-II0	JN234380.1	GLEAN_00457/GLEAN_00458	5
WC1-7		SR-II1	NM_001281912	GLEAN_00456	5
WC1-8		SR-II2	GU320289	GLEAN_12186	5
WC1-9		SR-II3	FJ031208.1	GLEAN_12191	5
WC1-10		SR-II4	XM_010805564	GLEAN_12192	5
WC1-11		SR-II5	FJ031209	GLEAN_12182	5
WC1-12		SR-II6	NM_001281917	GLEAN_09902	5
WC1-13		SR-II7	FJ031187	GLEAN_12187, BGD:BT10947 785733	5

NCBI, National Center for Biotechnology Information.