

Oral Magnesium Supplementation in Chronic Kidney Disease Stages 3 and 4: Efficacy, Safety, and Effect on Serum Calcification Propensity—A Prospective Randomized Double-Blinded Placebo-Controlled Clinical Trial



Iain Bressendorff¹, Ditte Hansen^{2,3}, Morten Schou⁴, Burton Silver⁵, Andreas Pasch⁶, Pierre Bouchelouche⁷, Lise Pedersen⁸, Lars Melholt Rasmussen⁹ and Lisbet Brandt¹

¹Department of Cardiology, Nephrology and Endocrinology, Nordsjællands Hospital, Hillerød, Denmark; ²Department of Nephrology, Herlev and Gentofte Hospital, Herlev, Denmark; ³Division of Nephrology, Roskilde Hospital, Roskilde, Denmark; ⁴Department of Cardiology, Herlev and Gentofte Hospital, Herlev, Denmark; ⁵IntraCellular Diagnostics Inc., Medford, Oregon, USA; ⁶Department of Clinical Research, University of Bern, Bern, Switzerland; ⁷Department of Clinical Biochemistry, Køge Hospital, Køge, Denmark; ⁸Department of Clinical Biochemistry, Holbæk Hospital, Holbæk, Denmark; and ⁹Department of Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense, Denmark

Introduction: Chronic kidney disease (CKD) is associated with high cardiovascular morbidity and mortality. Recent evidence suggests that increases in both serum and intracellular magnesium (Mg) can slow or even prevent the development of vascular calcification seen in CKD. Serum calcification propensity (T_{50}) is a novel functional test, which is associated with all-cause mortality in CKD and measures the ability of serum to delay the formation of crystalline nanoparticles. Theoretically, increasing serum Mg should improve T_{50} and thereby reduce the propensity towards ectopic calcification.

Methods: We conducted a randomized placebo-controlled double-blinded clinical trial to investigate the safety of 2 different doses of oral Mg supplementation in subjects with CKD stages 3 and 4 as well as their effects on intracellular Mg and T_{50} . Thirty-six subjects with CKD stages 3 and 4 were randomized to one of 3 groups (placebo, elemental Mg 15 mmol/d or elemental Mg 30 mmol/d) given as slow-release Mg hydroxide and followed for 8 weeks.

Results: Thirty-four subjects completed the trial. Intracellular Mg remained stable throughout the trial despite significant increases in both serum and urine Mg. T_{50} increased significantly by 40 min from 256 ± 60 (mean \pm SD) to 296 ± 64 minutes (95% confidence interval, 11–70, $P < 0.05$) in the Mg 30 mmol/d group after 8 weeks. No serious adverse events related to the study medication were reported during the study.

Discussion: Oral Mg supplementation was safe and well tolerated in CKD stages 3 and 4 and improved T_{50} , but did not increase intracellular Mg. Further studies are needed to investigate the long-term effects of Mg supplementation in CKD stage 3 and 4 and whether improvement in calcification propensity is related to clinical endpoints.

Kidney Int Rep (2017) 2, 380–389; <http://dx.doi.org/10.1016/j.ekir.2016.12.008>

KEYWORDS: calcification propensity; chronic kidney disease; magnesium

© 2016 International Society of Nephrology. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Chronic kidney disease (CKD) affects approximately 13% of the general population¹ and is associated with increased risk of cardiovascular disease (CVD) because of both traditional and nontraditional CVD risk

factors.² Among the nontraditional CVD risk factors, disturbances in mineral and bone disease are associated with vascular calcification,³ which is associated with increased cardiovascular mortality in end-stage renal disease.⁴ Key factors in this regard are phosphate (PO_4) and calcium (Ca), which can precipitate and induce an osteogenic transformation in the vascular smooth muscle cells of the arteries causing them to calcify and stiffen.³

Correspondence: Iain Bressendorff, Department of Cardiology, Nephrology and Endocrinology, Nordsjællands Hospital, Dyrehavevej 29, DK-3400, Hillerød, Denmark. E-mail: iain@bressendorff.com

Received 29 September 2016; revised 29 November 2016; accepted 23 December 2016; published online 31 December 2016

Epidemiological studies have found associations between higher levels of serum magnesium (Mg) and improved survival among patients suffering from CKD⁵ and end-stage renal disease,^{6–13} and higher levels of serum Mg (sMg) are associated with reduced progression of CKD.^{5,14} These associations are thought to be mediated by an antagonistic effect of Mg on the procalcifying milieu in CKD.¹⁵ *In vitro* calcifications induced by Ca and high concentrations of PO_4 can be prevented or reversed by adding or increasing Mg, which appears to be mediated by both upregulation of factors that inhibit calcification and downregulation of factors that promote calcification.^{16–20} Also, 2 small clinical trials of Mg supplementation in end-stage renal disease have shown reduced progression of vascular calcification.^{21,22} Thus, Mg supplementation could potentially be a therapeutic option to attenuate the progression of vascular calcification in CKD. Because studies have shown that influx of Mg into the vascular smooth muscle cell is important for prevention of vascular calcification,^{16,18,19} measurement of any effect of Mg therapy on intracellular Mg (iMg) would be of interest. However, because of the renal excretion of Mg, there is a risk that Mg therapy might result in toxic hypermagnesemia in patients with reduced kidney function. So far, no studies have examined the effect of oral Mg supplementation on iMg in subjects with CKD stages 3 and 4, or the safety of this intervention.

Serum calcification propensity (T_{50}) is a novel functional test, which determines the ability of serum to resist Ca/PO_4 precipitation by measuring the time-point of conversion from primary to secondary calciprotein particle.²³ Previous studies have shown that low T_{50} predicts all-cause mortality in CKD stages 3 and 4²⁴ and kidney transplant recipients,^{25,26} as well as graft failure in kidney transplant recipients.²⁵ T_{50} is believed to reflect the propensity towards ectopic calcification, although so far low T_{50} has not been directly associated with progression of vascular calcification. *In vitro* data indicate that Mg improves T_{50} ,²³ but so far this has not been examined in a clinical trial.

We conducted a randomized placebo-controlled double-blinded clinical trial to investigate the efficacy and safety of 2 different doses of oral Mg supplementation on iMg in subjects with CKD stages 3 and 4, as well as any effects on T_{50} . We hypothesized that greater doses of Mg supplementation would result in a greater increase in iMg compared with placebo as well as improving T_{50} in subjects with CKD stages 3 and 4 and low or low-normal sMg .

METHODS

Subjects

Subjects were recruited between October 2014 and February 2015 from the outpatient clinic at the Division of Nephrology, Roskilde County Hospital, Denmark. All patients were screened before planned visits to the clinic and those matching trial criteria were offered participation in the trial. Inclusion criteria were age > 18 years, estimated glomerular filtration rate (eGFR) < 60 ml/min per 1.73 m², total sMg < 0.82 mmol/l, safe contraceptive treatment in women of childbearing age, and written informed consent. Exclusion criteria were current treatment with hemodialysis or peritoneal dialysis, kidney transplant recipient, treatment with Mg-containing medication or supplements, parathyroid hormone (PTH) > 66 pmol/l, cancer, other medical condition that in the opinion of the investigators would prohibit participation in the trial, pregnancy or breastfeeding, allergies to any contents of the study medication, and participation in other interventional trials.

Design

The trial was an investigator-initiated double-blinded placebo-controlled clinical trial in which subjects were randomized in a ratio of 1:1:1 to 8 weeks of oral treatment with either placebo twice daily, slow-release Mg hydroxide 360 mg (equivalent to 15 mmol of elemental Mg) (Mablet, Gunnar Kjems ApS, Copenhagen, Denmark) once daily and placebo once daily, or slow-release Mg hydroxide 360 mg twice daily. Mg hydroxide and placebo tablets were identical in appearance, constituents, and containers, and did not contain Ca. Study medication was packed in consecutively numbered containers according to a computer generated block-randomized randomization list, and was administered to subjects consecutively as they entered the trial. Subjects and investigators were blinded to the study medication during the course of the trial.

At weeks 0, 4, and 8, sublingual epithelial cells were sampled for iMg and nonfasting blood samples, 24-hour urine samples were collected, and an electrocardiogram was performed. Subjects were instructed to maintain their usual diets and not to begin treatment with Mg-containing medication or supplements for the duration of the trial. Adherence was assessed by pill count at week 8.

The primary endpoint was a change in iMg as assessed by energy dispersive X-ray microanalysis of sublingual epithelial cells,²⁷ as this has been shown previously to correlate with iMg in human atrial cardiomyocytes.²⁸ Based on previous studies,^{29–33} an

increase in iMg of 2.0 mEq/l with a SD of 2.0 mEq/l was considered clinically relevant. With a probability of type 1 error (α) of 5% and power of 80% ($1 - \beta$) a sample size of 10 subjects per group would be necessary to detect a difference of 2.0 mEq/l. A dropout rate of 15% was anticipated, and therefore 36 subjects in total were included in the trial (12 subjects per group). Dropouts were not replaced.

Secondary endpoints were blood levels of total Mg, PO_4 , ionized Ca, PTH, 25-hydroxy vitamin D₃ (25-OH-D₃), T_{50} , fetuin-A, albumin, bicarbonate, intact fibroblast growth factor 23 (FGF23), eGFR, 24-hour urinary Mg (uMg) and PO_4 (uPO_4), and electrocardiogram-measured QT_c interval.

Ethics

The trial is in compliance with the Helsinki Declaration II of 1975, revised 1983, and was approved 5 May 2014 by the Danish National Committee on Biomedical Research Ethics (SJ-398) and the Danish Data Protection Agency. The trial was registered at [ClinicalTrials.gov](https://clinicaltrials.gov) (NCT02216877). All subjects gave written informed consent before initiating the trial.

Laboratory Analysis

iMg was measured in sublingual epithelial cells scraped from the mucosa adjacent to the frenulum and immediately fixed on a carbon slide with cytology fixative. The slides were examined with a scanning electron microscope (FEI; Thermo Fisher Scientific, Waltham, Massachusetts), and suitable cells were identified. iMg was measured with radiographic analysis of individual epithelial cells (EXA; Intracellular Diagnostics, Medford, Oregon) (normal range 34.0–42.0 mEq/l). Reported values are the mean of 5 to 10 cells per subject; a specimen was rejected if variance exceeded 2%. This method is used to assess total cellular magnesium and cannot differentiate free Mg from bound species.

Blood samples were drawn from subjects in non-fasting states. Samples for measurement of T_{50} , fetuin-A, and FGF23 were collected and immediately stored at $-80^\circ C$. The analyses were performed in bulk in the same analytical run so as to eliminate interassay variation. Blood levels of total Mg, ionized Ca, PO_4 , PTH, 25-OH-D₃, creatinine, hemoglobin, bicarbonate, and albumin were measured on a routine basis by the same standardized methods at the local department of clinical biochemistry. Urine samples were collected over 24 hours in 2.5-liter containers, which contained 100 ml of 2.5% hydrochloric acid. uMg and uPO_4 were analyzed on a routine basis by the same standardized analysis at the local laboratory.

T_{50} was measured using a standardized method previously described.²³ FGF23 was measured using the

human intact FGF23 ELISA kit 60-6500 (Immutopics, San Clemente, CA), and fetuin-A was measured using the human intact Quantikine human Fetuin A ELISA kit DFTA00 (R&D Systems, Minneapolis, MN). PTH was measured using the second-generation intact PTH immunoassay (reference values 1.5–7.6 $\rho mol/l$) (ADVIA Centaur, Siemens, Erlangen, Germany). eGFR was calculated using the CKD-EPI formula.

Statistical Analysis

The biostatistical evaluation was performed blinded using SPSS version 22.0.0.0 (IBM Corporation, Armonk, NY). Continuous data were described as mean \pm SD for data of a normal distribution, and as median and interquartile range (25th–75th percentile) for data of a non-normal distribution. For assessment of within-group changes, a one-way analysis of variance with repeated measures or a Friedman test (both with *post hoc* Bonferroni correction for multiple measures) was applied for data with Normal or non-Normal distributions, respectively. For assessment of between-group changes of iMg , sMg , uMg , and T_{50} , a mixed 2-way analysis of variance with repeated measures was applied. All tests were 2-sided tests and a $P < 0.05$ was considered statistically significant.

RESULTS

Ninety-two subjects met inclusion criteria for the trial, and of these, 36 agreed to participate (Figure 1). Demographic characteristics of trial subjects are displayed in Table 1. According to tablet count at final follow-up visit compliance $>95\%$ was achieved in all subjects.




Figure 1. CONSORT diagram. Mg, magnesium.

Table 1. Demographic information

Characteristics	Total (n = 34)	Placebo (n = 12)	Mg 15 mmol/d (n = 11)	Mg 30 mmol/d (n = 11)
Male sex, no. (%)	28 (82)	12 (100)	9 (82)	7 (64)
White race, no. (%)	33 (97)	11 (92)	11 (100)	11 (100)
Age (yr)	65.9 ± 8.7	70.4 ± 7.7	61.4 ± 10.2	65.4 ± 5.8
Body mass index (kg/m ²)	27.9 ± 4.5	27.5 ± 3.4	27.8 ± 3.0	28.4 ± 6.6
Smoker				
Active, no. (%)	9 (26)	2 (16)	4 (36)	3 (27)
Previous, no. (%)	15 (44)	8 (67)	4 (36)	3 (27)
Never, no. (%)	10 (29)	2 (17)	3 (27)	5 (46)
Comorbidities				
Diabetes mellitus				
Type 1, no. (%)	0 (0)	0 (0)	0 (0)	0 (0)
Type 2, no. (%)	11 (32)	3 (25)	4 (36)	4 (36)
Coronary artery disease, no. (%)	6 (18)	4 (33)	1 (9)	1 (9)
Heart failure, no. (%)	1 (3)	1 (8)	0 (0)	0 (0)
Hypertension, no. (%)	32 (94)	10 (83)	11 (100)	11 (100)
Dyslipidemia, no. (%)	23 (68)	10 (83)	6 (55)	7 (64)
Cerebrovascular disease, no. (%)	5 (15)	2 (17)	1 (9)	2 (18)
Peripheral artery disease, no. (%)	3 (9)	0 (0)	2 (18)	1 (9)
Cause of CKD				
Diabetes mellitus type 2, no. (%)	6 (18)	2 (17)	1 (9)	3 (27)
Hypertension, no. (%)	8 (24)	4 (33)	3 (27)	1 (9)
Chronic glomerulonephritis, no. (%)	9 (27)	5 (42)	4 (36)	0 (0)
Polycystic kidney disease, no. (%)	2 (6)	0 (0)	0 (0)	2 (18)
Interstitial nephropathy, no. (%)	1 (3)	0 (0)	0 (0)	1 (9)
Other, no. (%)	8 (24)	1 (8)	3 (27)	4 (36)
Medical therapy				
ACE inhibitor, no. (%)	12 (35)	5 (42)	3 (27)	4 (36)
ARB, no. (%)	13 (38)	3 (25)	6 (55)	4 (36)
Beta-blocker, no. (%)	16 (47)	5 (42)	5 (46)	6 (55)
Calcium-channel blocker, no. (%)	21 (62)	7 (58)	6 (55)	8 (73)
Loop diuretic, no. (%)	10 (29)	5 (42)	2 (18)	3 (27)
Thiazide-like diuretic, no. (%)	7 (21)	1 (8)	4 (36)	2 (18)
Statin, no. (%)	19 (56)	8 (67)	5 (46)	6 (55)
Phosphate binder, no. (%)	3 (9)	2 (17)	1 (9)	0 (0)
25-Hydroxy vitamin D, no. (%)	18 (53)	10 (83)	5 (46)	3 (27)
1,25-Dihydroxy vitamin D, no. (%)	2 (6)	1 (8)	1 (9)	0 (0)
Sodium bicarbonate, no. (%)	3 (9)	2 (17)	1 (9)	0 (0)
PPI, no. (%)	15 (44)	4 (33)	7 (64)	3 (27)

Table 1. (Continued)

Characteristics	Total (n = 34)	Placebo (n = 12)	Mg 15 mmol/d (n = 11)	Mg 30 mmol/d (n = 11)
Amiloride, no. (%)	2 (6)	0 (0)	1 (9)	1 (9)
Spironolactone, no. (%)	2 (6)	1 (8)	0 (0)	1 (9)
CNI, no. (%)	1 (3)	1 (8)	0 (0)	0 (0)
Systolic blood pressure (mm Hg)	138.4 ± 14.7	138.8 ± 14.7	139.1 ± 16.5	137.5 ± 14.1
Diastolic blood pressure (mm Hg)	82.3 ± 9.7	82.3 ± 9.3	83.3 ± 11.4	81.3 ± 9.2
Heart rate beats/min	75.6 ± 14.3	73.9 ± 10.5	78.5 ± 19.4	74.6 ± 12.94
eGFR _{CKD-EPI} (ml/min per 1.73 m ²)	32.6 ± 12.1	29.6 ± 14.4	36.2 ± 12.8	32.3 ± 8.4
Proteinuria (g/d)	1.63 ± 1.38	1.87 ± 1.33	1.76 ± 1.54	1.20 ± 1.30
Hemoglobin (mmol/l)	8.09 ± 0.86	8.15 ± 0.92	8.01 ± 0.96	8.09 ± 0.75
Albumin (g/l)	37.5 ± 3.1	36.2 ± 3.1	38.2 ± 3.2	38.4 ± 2.7

ACE, angiotensin converting enzyme; ARB, angiotensin 2 receptor blocker; CKD, chronic kidney disease; CNI, calcineurin inhibitor; eGFR, estimated glomerular filtration rate; Mg, magnesium; PPI, proton pump inhibitor.

The primary study endpoint, i Mg, did not change significantly compared with week 0 in any of the groups at either week 4 or week 8 (Table 2 and Figure 2a). s Mg and u Mg increased significantly from week 0 to week 4 in both Mg treatment groups, whereas there was no change in the placebo group (Table 2 and Figure 2b and c). At week 8, both s Mg and u Mg increased further compared with week 0 in the Mg 30 mmol/d group, whereas in the Mg 15 mmol/d group, only u Mg was increased compared with week 0 and the change in s Mg was no longer significant. There were no significant changes in s Mg or u Mg in the placebo group from week 0 to weeks 4 and 8. The effect of time and intervention on i Mg was not statistically significant (Table 3), but there was a significant effect of time and intervention on both s Mg and u Mg (Table 3).

T_{50} increased significantly in the Mg 30 mmol/d group compared with baseline at both weeks 4 and 8, while only increasing significantly within the Mg 15 mmol/d group compared with baseline at week 4, and with no significant changes within the placebo group (Table 4 and Figure 3). There was also a significant effect of time, and intervention on T_{50} and *post hoc* tests revealed that Mg 30 mmol/d produced significantly greater changes to T_{50} over time compared with placebo or Mg 15 mmol/d (Table 3). Factors other than Mg known to affect T_{50} (fetuin-A, PO_4 , ionized Ca, albumin, bicarbonate) did not change significantly in either group throughout the trial (Table 4).

eGFR did not change significantly throughout the trial, and there were no changes in blood analyses of mineral metabolism or u PO₄ in any of the treatment groups (Table 5). There was a significant effect of time on change

Table 2. Treatment effect on magnesium parameters, one-way ANOVA with repeated measures, and Bonferroni-corrected *post hoc* tests

Treatment group	Week 0	Week 4	Week 8	P value for time effect	Difference Week 0 versus week 4	Difference Week 0 versus week 8
Placebo						
_i Mg (mEq/l)	35.3 ± 1.9	34.4 ± 2.5	36.2 ± 2.7	0.11	-0.9 (-3.1 to 1.4)	0.9 (-0.9 to 2.7)
_s Mg (mmol/l)	0.830 ± 0.084	0.837 ± 0.100	0.808 ± 0.100	0.50	0.007 (-0.063 to 0.076)	-0.023 (-0.112 to 0.067)
_u Mg (mmol/d)	3.83 ± 1.62	3.92 ± 1.60	3.70 ± 2.82	0.78	0.09 (-0.70 to 0.87)	-0.13 (-1.41 to 1.14)
Mg 15 mmol/d						
_i Mg (mEq/l)	35.3 ± 2.7	36.0 ± 2.3	34.9 ± 2.9	0.40	0.7 (-1.3 to 2.7)	-0.4 (-2.8 to 2.0)
_s Mg (mmol/l)	0.790 ± 0.119	0.873 ± 0.086	0.863 ± 0.110	0.04*	0.083 (0.041 to 0.144)**	0.073 (-0.031 to 0.176)
_u Mg (mmol/d)	3.51 ± 1.33	5.00 ± 1.67	4.72 ± 1.75	0.003*	1.49 (0.68 to 2.31)**	1.21 (-0.15 to 2.58)
Mg 30 mmol/d						
_i Mg (mEq/l)	35.4 ± 2.7	36.2 ± 2.8	36.8 ± 2.5	0.30	0.9 (-2.2 to 3.9)	1.5 (-0.9 to 3.9)
_s Mg (mmol/l)	0.739 ± 0.101	0.844 ± 0.103	0.848 ± 0.099	0.001*	0.105 (0.024 to 0.185)**	0.109 (0.026 to 0.192)**
_u Mg (mmol/d)	3.16 ± 1.23	5.08 ± 1.68	5.43 ± 2.08	0.001*	1.91 (0.84 to 2.99)**	2.27 (0.67 to 3.87)**

Reported as mean ± SD for weeks 0, 4, and 8, and mean change with 95% confidence interval for differences between time points.

ANOVA, analysis of variance; CI, confidence interval; Mg, magnesium; _iMg, intracellular magnesium; _{ion}Mg, ionized magnesium; _sMg, serum magnesium; _uMg, urine magnesium.

*P < 0.05.

**P < 0.05 after Bonferroni adjustment for multiple comparisons.

in QT_c on electrocardiogram in the Mg 30 mmol/d group, but this change lost significance in *post hoc* tests (Table 5).

During the course of the trial, one subject in the Mg 15 mmol/d group suffered a transitory ischemic attack

and withdrew from the trial, and one subject in the Mg 30 mmol/d group withdrew because of difficulties swallowing the study medication. Further, one subject in the Mg 30 mmol/d group had side effects from the




Figure 2. (a) Mean change in intracellular magnesium compared with baseline. Error bars = 95% confidence interval. (b) Mean change in serum magnesium compared with baseline. Error bars = 95% confidence interval. *P < 0.05 after Bonferroni adjustment for multiple comparisons. (c) Mean change in 24-hour urine magnesium compared with baseline. Error bars = 95% confidence interval. *P < 0.05 after Bonferroni adjustment for multiple comparisons.

Table 3. Change compared with baseline for magnesium parameters and serum calcification propensity, 2-way mixed ANOVA with repeated measures

	Change at week 4 Mean (CI; P value)	Change at week 8 Mean (CI; P value)
Mg 15 mmol/d versus placebo		
iMg (mEq/l)	1.59 (−4.60 to 1.42; P = 0.41)	−1.32 (−3.97 to 1.34; P = 0.45)
sMg (mmol/l)	0.076 (−0.010 to 0.162; P = 0.09)	0.095 (−0.017 to 0.207; P = 0.11)
uMg (mmol/d)	1.55 (0.45 to 2.65; P = 0.005)*	1.59 (−0.09 to 3.27; P = 0.07)
T ₅₀ (min)	47.7 (1.3 to 94.1; P = 0.43)	29.1 (−17.6 to 75.9; P = 0.29)
Mg 30 mmol/d versus placebo		
iMg (mEq/l)	1.74 (−1.27 to 4.75; P = 0.34)	0.53 (−2.12 to 3.18; P = 0.88)
sMg (mmol/l)	0.098 (0.012 to 0.184; P = 0.02)*	0.132 (0.020 to 0.243; P = 0.02)*
uMg (mmol/d)	1.97 (0.90 to 3.04; P < 0.001)*	2.85 (1.21 to 4.49; P = 0.001)*
T ₅₀ (min)	59.6 (13.2 to 106.1; P = 0.01)*	56.2 (9.5 to 102.9; P = 0.02)*
Mg 30 mmol/d versus Mg 15 mmol/d		
iMg (mEq/l)	0.15 (−2.93 to 3.22; P = 0.99)	1.85 (−0.86 to 4.55; P = 0.23)
sMg (mmol/l)	0.022 (−0.067 to 0.110; P = 0.82)	0.036 (−0.078 to 0.151; P = 0.72)
uMg (mmol/d)	0.42 (−0.68 to 1.52; P = 0.62)	1.26 (−0.39 to 2.90; P = 0.16)
T ₅₀ (min)	11.9 (−35.5 to 59.3; P = 0.81)	27.1 (−20.7 to 74.8; P = 0.36)

P values for the effect of time and intervention: iMg, P = 0.133; sMg, P = 0.016; uMg, P = 0.009; T₅₀, P = 0.011.

ANOVA, analysis of variance; CI, confidence interval; Mg, magnesium; iMg, intracellular magnesium; sMg, serum magnesium; uMg, urine magnesium; T₅₀, serum calcification propensity. *P < 0.05.

medication similar to previously experience from other medications (abdominal cramps, flushing, and palpitations), and intolerance to one of the tablet constituents was suspected (most likely talcum). There were 2 cases of community-acquired pneumonia in the placebo group that required hospital admissions, and one case

of asthma exacerbation in the Mg 15 mmol/d group that required treatment with corticosteroids. None of the adverse events were considered to be due to the effects of Mg. Incidence of loosening of stool was identical (2 in each treatment group), but none of the study participants experienced frank diarrhea.

Table 4. Treatment effect on serum calcification propensity and related factors, one-way ANOVA with repeated measures or Friedman test, both with Bonferroni-adjusted *post hoc* tests

Treatment group	Week 0	Week 4	Week 8	P value for time effect	Difference Week 0 versus week 4	Difference Week 0 versus week 8
Placebo						
T ₅₀ (min)	298.4 ± 80.8	270.9 ± 75.4	282.6 ± 70.0	0.23	−27.5 (−76.7 to 21.7)	−15.8 (−57.2 to 25.6)
ionCa (mmol/l)	1.210 ± 0.054	1.206 ± 0.017	1.230 ± 0.041	0.84	−0.005 (−0.046 to 0.036)	−0.007 (−0.044 to 0.029)
sPO ₄ (mmol/l)	1.040 ± 0.202	1.118 ± 0.246	1.050 ± 0.258	0.33	0.078 (−0.054 to 0.210)	0.010 (−0.115 to 0.135)
Fetuin-A (g/l)	0.21 (0.16; 0.26)	0.24 (0.13; 0.49)	0.23 (0.13; 0.49)	0.92	0.03	0.02
Albumin (g/l)	36.2 ± 3.1	35.2 ± 4.2	36.1 ± 3.2	0.39	−1.0 (−3.5 to 1.5)	−0.1 (−2.0 to 1.8)
HCO ₃ (mmol/l)	24.4 ± 3.4	24.1 ± 2.5	25.2 ± 2.6	0.42	−0.3 (−3.2 to 2.5)	0.8 (−1.4 to 3.1)
Mg 15 mmol/d						
T ₅₀ (min)	263.5 ± 59.1	281.2 ± 39.0	276.7 ± 35.0	0.40	17.7 (−18.1 to 53.6)	13.3 (−29.2 to 55.8)
ionCa (mmol/l)	1.193 ± 0.048	1.191 ± 0.041	1.179 ± 0.050	0.45	−0.002 (−0.030 to 0.026)	−0.014 (−0.055 to 0.027)
sPO ₄ (mmol/l)	1.082 ± 0.239	1.085 ± 0.234	1.085 ± 0.188	0.99	0.003 (−0.197 to 0.202)	0.003 (−0.232 to 0.238)
Fetuin-A (g/l)	0.22 (0.11; 0.31)	0.28 (0.11; 0.42)	0.27 (0.15; 0.55)	0.53	0.06	0.05
Albumin (g/l)	38.2 ± 3.2	36.8 ± 3.0	37.5 ± 2.7	0.13	−1.4 (−3.3 to 0.6)	−0.7 (−2.6 to 1.2)
HCO ₃ (mmol/l)	25.0 ± 3.7	25.1 ± 3.5	25.7 ± 3.1	0.42	0.1 (−1.4 to 1.5)	0.7 (−0.8 to 2.3)
Mg 30 mmol/d						
T ₅₀ (min)	256.0 ± 60.4	285.6 ± 69.1	296.4 ± 63.9	0.004*	29.6 (6.8 to 52.5)**	40.4 (10.9 to 69.9)**
ionCa (mmol/l)	1.20 ± 0.04	1.21 ± 0.04	1.22 ± 0.06	0.18	−0.01 (−0.04 to 0.02)	−0.03 (−0.08 to 0.03)
sPO ₄ (mmol/l)	1.09 ± 0.18	1.09 ± 0.19	1.05 ± 0.20	0.62	−0.00 (−0.08 to 0.08)	−0.04 (−0.18 to 0.10)
Fetuin-A (g/l)	0.33 (0.11; 0.50)	0.20 (0.13; 0.38)	0.36 (0.22; 0.47)	0.72	−0.13	0.03
Albumin (g/l)	38.4 ± 2.7	38.0 ± 2.1	38.3 ± 1.9	0.77	0.4 (−1.5 to 2.2)	0.1 (−1.5 to 1.7)
HCO ₃ (mmol/l)	25.3 ± 1.8	26.2 ± 2.7	26.7 ± 2.0	0.08	0.9 (0.8 to 2.6)	1.4 (−0.3 to 3.2)

Reported as mean ± SD or median and interquartile range for weeks 0, 4, and 8 (as relevant), and change with 95% confidence interval or absolute change for differences between time points (as relevant).

ANOVA, analysis of variance; CI, confidence interval; ionCa, ionized calcium; HCO₃, bicarbonate; Mg, magnesium; sPO₄, serum phosphate; T₅₀, serum calcification propensity.

*P < 0.05.

**P < 0.05 after Bonferroni adjustment for multiple comparisons.




Figure 3. Mean change in serum calcification propensity compared with baseline. Error bars = 95% confidence interval. **P* < 0.05 after Bonferroni adjustment for multiple comparisons.

None of the adverse events were attributed to Mg supplementation.

DISCUSSION

The main finding of this clinical trial is that oral Mg supplementation using slow-release Mg hydroxide

30 mmol/d does not affect _iMg as assessed by energy dispersive X-ray microanalysis after 8 weeks of treatment, despite significant increases in _sMg and _uMg. Furthermore, this trial has shown for the first time that calcification propensity *T*₅₀ can be improved by oral Mg supplementation in CKD stages 3 and 4.

Study participants were included based on low to low-normal levels of _sMg, and it is plausible that subjects may have been Mg-depleted based on the low _iMg and _uMg in all groups at baseline. Thus, Mg supplementation would first have to replenish total body Mg stores before “spilling over” into the serum, which could explain the relatively modest increase in _sMg and _uMg. We speculate that _iMg stores increased in other compartments of the body (e.g., bone or muscle), and only later in epithelial cells from the sublingual mucosa, which would account for the lack of a treatment effect on _iMg.

Despite 8 weeks of Mg supplementation with 15 mmol/d or 30 mmol/d on top of dietary Mg intake, _uMg was only 4.72 mmol/d and 5.43 mmol/d in the 2 groups at week 8, respectively. Assuming that approximately one third of ingested Mg is available

Table 5. Treatment effect on mineral metabolism and kidney function, one-way ANOVA with repeated measures or Friedman test

Treatment group	Week 0	Week 4	Week 8	<i>P</i> value for time effect	Difference Week 0 versus week 4	Difference Week 0 versus week 8
Placebo						
eGFR (ml/min)	29.6 ± 14.4	28.1 ± 13.6	29.6 ± 12.6	0.38	-1.5 (-5.1 to 2.1)	0.0 (-3.7 to 3.7)
_{ion} Ca (mmol/l)	1.210 ± 0.054	1.206 ± 0.017	1.230 ± 0.041	0.84	-0.005 (-0.046 to 0.036)	-0.007 (-0.044 to 0.029)
_s PO ₄ (mmol/l)	1.040 ± 0.202	1.118 ± 0.246	1.050 ± 0.258	0.33	0.078 (-0.054 to 0.210)	0.010 (-0.115 to 0.135)
_u PO ₄ (mg/d)	2428 ± 751	2588 ± 860	2458 ± 365	0.82	159 (-854 to 1173)	30 (-651 to 711)
PTH (pmol/l)	10.4 (6.5; 16.8)	8.3 (7.0; 16.9)	11.6 (7.1; 23.3)	0.26	-2.1	1.2
25-D ₃ (nmol/l)	61.0 (48.5; 78.5)	66.5 (53.3; 78.0)	60.5 (56.3; 72.8)	0.34	5.5	-0.5
FGF-23 (IU)	98.0 (54.3; 173.8)	153.0 (68.3; 277.0)	112.0 (60.5; 357.3)	0.56	55	14
QT _c (ms)	408.3 ± 25.7	413.7 ± 19.7	414.8 ± 14.9	0.74	5.4 (-23.5 to 34.3)	6.5 (-17.7 to 30.7)
Mg 15 mmol/d						
eGFR (ml/min)	36.2 ± 12.8	38.6 ± 15.1	36.6 ± 14.0	0.17	2.5 (-1.8 to 6.7)	0.4 (-3.2 to 3.9)
_{ion} Ca (mmol/l)	1.193 ± 0.048	1.191 ± 0.041	1.179 ± 0.050	0.45	-0.002 (-0.030 to 0.026)	-0.014 (-0.055 to 0.027)
_s PO ₄ (mmol/l)	1.082 ± 0.239	1.085 ± 0.234	1.085 ± 0.188	0.99	0.003 (-0.197 to 0.202)	0.003 (-0.232 to 0.238)
_u PO ₄ (mg/d)	2281 ± 880	2381 ± 647	2450 ± 1045	0.64	100 (-414 to 614)	169 (-353 to 691)
PTH (pmol/l)	9.3 (4.5; 18.3)	10.1 (5.3; 13.9)	13.8 (5.7; 17.3)	0.34	0.8	4.5
25-D ₃ (nmol/l)	46.5 (35.5; 65.3)	49.0 (35.8; 54.8)	40.5 (33.8; 63.3)	0.67	3.5	-6
FGF-23 (IU)	62 (53; 112)	58 (42; 105)	59 (56; 111)	0.61	-4	-3
QT _c (ms)	421.2 ± 30.3	418.5 ± 23.1	415.9 ± 30.9	0.73	-2.7 (-17.2 to 11.8)	-5.3 (-27.8 to 17.2)
Mg 30 mmol/d						
eGFR (ml/min)	32.3 ± 8.4	32.6 ± 9.6	31.8 ± 9.6	0.86	-0.3 (-3.7 to 3.1)	0.5 (-3.6 to 4.5)
_{ion} Ca (mmol/l)	1.20 ± 0.04	1.21 ± 0.04	1.22 ± 0.06	0.18	-0.01 (-0.04 to 0.02)	-0.03 (-0.08 to 0.03)
_s PO ₄ (mmol/l)	1.09 ± 0.18	1.09 ± 0.19	1.05 ± 0.20	0.62	-0.00 (-0.08 to 0.08)	0.04 (-0.10 to 0.18)
_u PO ₄ (mg/d)	2126 ± 537	2027 ± 741	2071 ± 726	0.86	99 (-577 to 774)	55 (-366 to 476)
PTH (pmol/l)	13.2 (3.3; 17.9)	15.1 (3.4; 19.9)	13.0 (5.8; 15.5)	0.81	1.9	-0.2
25-D ₃ (nmol/l)	63.5 (44.3; 81.0)	64.0 (39.3; 79.3)	58.0 (40.8; 73.0)	0.38	0.5	-5.5
FGF-23 (IU)	75 (62; 110)	69 (57; 93)	72 (47; 96)	0.40	-6	-3
QT _c (ms)	427.6 ± 21.2	430.3 ± 22.3	418.3 ± 26.0	0.03*	-2.8 (-16.4 to 10.9)	-9.2 (-4.8 to 23.3)

Reported as mean ± SD or median and interquartile range for weeks 0, 4, and 8 (as relevant), and change with 95% confidence interval or absolute change for differences between time points (as relevant).

ANOVA, analysis of variance; 25-D₃, 25-hydroxy vitamin D₃; CI, confidence interval; _{ion}Ca, ionized calcium; _sCa, total serum calcium; eGFR, estimated glomerular filtration rate; FGF-23, intact fibroblast growth factor 23; Mg, magnesium; _sPO₄, serum phosphate; PTH, parathyroid hormone; _uPO₄, urine phosphate; QT_c, corrected QT interval.

**P* < 0.05.

for absorption, $_{\text{u}}\text{Mg}$ would be expected to be at least 5 mmol/d and 10 mmol/d for the 2 groups, if Mg intake was balanced by urinary excretion. The relatively small increases in $_{\text{u}}\text{Mg}$ in the active treatment groups suggest either reduced Mg absorption in the gut, Mg retention due to total body Mg depletion, or Mg retention due to compromised renal Mg excretion. Proton pump inhibitors reduce intestinal Mg absorption, and were used by 64% and 27% of subjects in the 15 mmol/d and 30 mmol/d groups, respectively. Thus, reduced absorption of Mg might have blunted the effect of Mg supplementation. Faecal Mg was not measured in this trial, so it was not possible to assess Mg absorption in the gut.

This is the first clinical trial to show that T_{50} is improved by Mg supplementation in CKD stages 3 and 4. Notably, there were no significant changes to any of the other factors known to affect T_{50} , suggesting that the change in T_{50} was caused by the increase in $_{\text{s}}\text{Mg}$ alone. Indeed, adding Mg to serum *ex vivo* is known to increase T_{50} .²³ The final measurement of T_{50} in the group treated with Mg 30 mmol/d was similar to that of healthy adults without CKD,³⁴ and the intervention shifted the mean T_{50} about half a SD for patients with predialysis CKD, which according to previous data might lead to a 20% risk reduction in all-cause mortality.²⁴

Mg delays the formation of secondary calciprotein particles in the T_{50} test.²³ A recent *in vitro* study convincingly demonstrated that secondary (but not primary) calciprotein particles induce calcification of vascular smooth muscle cells,³⁵ suggesting that delaying the time until formation of secondary calciprotein particles (i.e., increasing T_{50}) might improve calcification propensity via this mechanism. Thus, T_{50} appears to be a *modifiable* risk factor and might be useful not only for risk assessment, but also for monitoring and guiding therapy directed at inhibition of Ca/PO_4 crystal formation. Whether the improvement of T_{50} observed in this trial is sustained over a longer treatment period, and whether this translates to a reduction in clinically relevant outcomes (e.g., cardiovascular events or all-cause mortality) will have to be studied in a future randomized clinical trial, which may use individualized and combined interventions aimed at improving T_{50} .

Mg supplementation might be expected to lower $_{\text{s}}\text{PO}_4$ in the same manner as Mg-containing PO_4 binders. We found no significant differences in $_{\text{s}}\text{PO}_4$, $_{\text{u}}\text{PO}_4$ or intact FGF23 in any of the groups, suggesting that Mg supplementation with the formulation used in this trial does not affect PO_4 homeostasis in any clinically meaningful way. However, the sample size and increases in $_{\text{s}}\text{Mg}$ were probably not powered to

definitively rule out any interaction and larger study populations would be needed to answer this question.

Mg supplementation has previously been a concern in CKD and end-stage renal disease because of the risk of hypermagnesemia due to impaired renal excretion of Mg. There is general consensus that hypermagnesemia is likely not symptomatic at $_{\text{s}}\text{Mg} < 2.0 \text{ mmol/l}$ ³⁶ and that serious complications only occur at $_{\text{s}}\text{Mg} > 3.0 \text{ mmol/l}$. Based on this trial, 8 weeks of Mg supplementation with slow-release Mg hydroxide is safe at doses up to 30 mmol/d in subjects with low or low-normal $_{\text{s}}\text{Mg}$ and eGFR down to 15 ml/min per 1.73 m^2 (the lowest eGFR of any subject in this trial). However, whether there are risks associated with longer treatment duration or in subjects with higher levels of $_{\text{s}}\text{Mg}$ was not assessed in this trial.

There are limitations and strengths to the current trial. Limitations include that energy dispersive X-ray microanalysis of sublingual epithelial cells was used as a proxy for $_{\text{i}}\text{Mg}$, and not bone, vascular, or muscle biopsy, which are the gold standards for measurement of Mg levels in these tissues. Also, the method has not previously been applied to patients with CKD, which might have confounded the results. Furthermore, the sample size was small and the trial was of a short duration. Lastly, the formulation of slow-release Mg hydroxide used in this trial was chosen because it is widely used and readily available in Denmark. It is possible that other Mg formulations with greater bioavailability would have produced different effects on measures of Mg and on T_{50} . Strengths of the trial include its single-center, randomized controlled and blinded study design.

In conclusion, 8 weeks of Mg supplementation using slow-release Mg hydroxide 30 mmol/d did not affect $_{\text{i}}\text{Mg}$ in sublingual epithelial cells. Mg supplementation improved T_{50} , which might aid in treating systemic calcification propensity. Mg supplementation was safe and well tolerated with no adverse events related to Mg treatment and no incidences of symptomatic hypermagnesemia. Slow-release Mg hydroxide 30 mmol/d might therefore be useful for increasing $_{\text{s}}\text{Mg}$ in future clinical trials addressing vascular calcification or calcification propensity in CKD stages 3 and 4.

DISCLOSURE

BS is the Research Director and President of IntraCellular Diagnostics, Inc., Medford, OR, USA. AP is one of the co-inventors of the T_{50} test and is the CEO and co-founder of Calciscon AG, Bern, Switzerland, which specializes in performing the T_{50} test. Gunnar Kjems ApS provided the study medication free of charge, but had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. All the other authors declared no competing interests.

REFERENCES

- Coresh J, Selvin E, Stevens LA, et al. Prevalence of chronic kidney disease in the United States. *JAMA*. 2007;298:2038–2047.
- Go AS, Chertow GM, Fan D, et al. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med*. 2004;351:1296–1305.
- Moe SM, Chen NX. Mechanisms of vascular calcification in chronic kidney disease. *J Am Soc Nephrol*. 2008;19:213–216.
- London GM, Guerin AP, Marchais SJ, et al. Arterial media calcification in end-stage renal disease: impact on all-cause and cardiovascular mortality. *Nephrol Dial Transplant*. 2003;18:1731–1740.
- Van Laecke S, Nagler EV, Verbeke F, et al. Hypomagnesemia and the risk of death and GFR decline in chronic kidney disease. *Am J Med*. 2013;126:825–831.
- Sakaguchi Y, Fujii N, Shoji T, et al. Hypomagnesemia is a significant predictor of cardiovascular and non-cardiovascular mortality in patients undergoing hemodialysis. *Kidney Int*. 2014;85:174–181.
- Sakaguchi Y, Fujii N, Shoji T, et al. Magnesium modifies the cardiovascular mortality risk associated with hyperphosphatemia in patients undergoing hemodialysis: a cohort study. *PLoS One*. 2014;9:e116273.
- Kanbay M, Yilmaz MI, Apetrii M, et al. Relationship between serum magnesium levels and cardiovascular events in chronic kidney disease patients. *Am J Nephrol*. 2012;36:228–237.
- Ishimura E, Okuno S, Yamakawa T, et al. Serum magnesium concentration is a significant predictor of mortality in maintenance hemodialysis patients. *Magnes Res*. 2007;20:237–244.
- Joao Matias P, Azevedo A, Laranjinha I, et al. Lower serum magnesium is associated with cardiovascular risk factors and mortality in haemodialysis patients. *Blood Purif*. 2014;38:244–252.
- Fein P, Weiss S, Ramos F, et al. Serum magnesium concentration is a significant predictor of mortality in peritoneal dialysis patients. *Adv Perit Dial*. 2014;30:90–93.
- Kurita N, Akizawa T, Fukagawa M, et al. Contribution of dysregulated serum magnesium to mortality in hemodialysis patients with secondary hyperparathyroidism: a 3-year cohort study. *Clin Kidney J*. 2015;8:744–752.
- de Roij van Zuijdewijn CL, Grooteman MP, Bots ML, et al. Serum magnesium and sudden death in European hemodialysis patients. *PLoS One*. 2015;10:e0143104.
- Sakaguchi Y, Iwatani H, Hamano T, et al. Magnesium modifies the association between serum phosphate and the risk of progression to end-stage kidney disease in patients with non-diabetic chronic kidney disease. *Kidney Int*. 2015;88:833–842.
- Massy ZA, Druke TB. Magnesium and cardiovascular complications of chronic kidney disease. *Nat Rev Nephrol*. 2015;11:432–442.
- Montezano AC, Zimmerman D, Yusuf H, et al. Vascular smooth muscle cell differentiation to an osteogenic phenotype involves TRPM7 modulation by magnesium. *Hypertension*. 2010;56:453–462.
- Kircelli F, Peter ME, Sevinc Ok E, et al. Magnesium reduces calcification in bovine vascular smooth muscle cells in a dose-dependent manner. *Nephrol Dial Transplant*. 2012;27:514–521.
- Louvet L, Buchel J, Steppan S, et al. Magnesium prevents phosphate-induced calcification in human aortic vascular smooth muscle cells. *Nephrol Dial Transplant*. 2013;28:869–878.
- Montes de Oca A, Guerrero F, Martinez-Moreno JM, et al. Magnesium inhibits Wnt/beta-catenin activity and reverses the osteogenic transformation of vascular smooth muscle cells. *PLoS One*. 2014;9:e89525.
- Zelt JG, McCabe KM, Svajger B, et al. Magnesium modifies the impact of calcitriol treatment on vascular calcification in experimental chronic kidney disease. *J Pharmacol Exp Ther*. 2015;355:451–462.
- Turgut F, Kanbay M, Metin MR, et al. Magnesium supplementation helps to improve carotid intima media thickness in patients on hemodialysis. *Int Urol Nephrol*. 2008;40:1075–1082.
- Mortazavi M, Moeinzadeh F, Saadatnia M, et al. Effect of magnesium supplementation on carotid intima-media thickness and flow-mediated dilatation among hemodialysis patients: a double-blind, randomized, placebo-controlled trial. *Eur Neurol*. 2013;69:309–316.
- Pasch A, Farese S, Graber S, et al. Nanoparticle-based test measures overall propensity for calcification in serum. *J Am Soc Nephrol*. 2012;23:1744–1752.
- Smith ER, Ford ML, Tomlinson LA, et al. Serum calcification propensity predicts all-cause mortality in predialysis CKD. *J Am Soc Nephrol*. 2014;25:339–348.
- Keyzer CA, de Borst MH, van den Berg E, et al. Calcification propensity and survival among renal transplant recipients. *J Am Soc Nephrol*. 2016;27:239–248.
- Dahle DO, Asberg A, Hartmann A, et al. Serum calcification propensity is a strong and independent determinant of cardiac and all-cause mortality in kidney transplant recipients. *Am J Transplant*. 2016;16:204–212.
- Silver BB. Development of cellular magnesium nano-analysis in treatment of clinical magnesium deficiency. *J Am Coll Nutr*. 2004;23:732S–737S.
- Haigney MC, Silver B, Tanglao E, et al. Noninvasive measurement of tissue magnesium and correlation with cardiac levels. *Circulation*. 1995;92:2190–2197.
- Shechter M, Sharir M, Labrador MJ, et al. Oral magnesium therapy improves endothelial function in patients with coronary artery disease. *Circulation*. 2000;102:2353–2358.
- Shechter M, Bairey Merz CN, Stuehlinger HG, et al. Effects of oral magnesium therapy on exercise tolerance, exercise-induced chest pain, and quality of life in patients with coronary artery disease. *Am J Cardiol*. 2003;91:517–521.
- McBride BF, Min B, Kluger J, et al. An evaluation of the impact of oral magnesium lactate on the corrected QT interval of patients receiving sotalol or dofetilide to prevent atrial or ventricular tachyarrhythmia recurrence. *Ann Noninvasive Electrocardiol*. 2006;11:163–169.
- Pokan R, Hofmann P, von Duvillard SP, et al. Oral magnesium therapy, exercise heart rate, exercise tolerance, and

- myocardial function in coronary artery disease patients. *Br J Sports Med.* 2006;40:773–778.
33. Shechter M, Saad T, Shechter A, et al. Comparison of magnesium status using X-ray dispersion analysis following magnesium oxide and magnesium citrate treatment of healthy subjects. *Magnes Res.* 2012;25:28–39.
 34. de Seigneux S, Ponte B, Berchtold L, et al. Living kidney donation does not adversely affect serum calcification propensity and markers of vascular stiffness. *Transpl Int.* 2015;28:1074–1080.
 35. Aghagolzadeh P, Bachtler M, Bijarnia R, et al. Calcification of vascular smooth muscle cells is induced by secondary calcium-protein particles and enhanced by tumor necrosis factor- α . *Atherosclerosis.* 2016;251:404–414.
 36. Jahn-Dechent W, Ketteler M. Magnesium basics. *Clin Kidney J.* 2012;5(Suppl 1):i3–i14.