



Original Article

## Change in hand dexterity and habitual gait speed reflects cognitive decline over time in healthy older adults: a longitudinal study

TAKUMI ABE, MS<sup>1, 2)\*</sup>, YUKI SOMA, PhD<sup>3)</sup>, NARUKI KITANO, PhD<sup>4)</sup>, TAKASHI JINDO, PhD<sup>4)</sup>, AYANE SATO, MS<sup>1)</sup>, KENJI TSUNODA, PhD<sup>5)</sup>, TAISHI TSUJI, PhD<sup>6)</sup>, TOMOHIRO OKURA, PhD<sup>7)</sup>

<sup>1)</sup> Doctoral Program in Physical Education, Health and Sport Sciences, University of Tsukuba: 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8577 Japan

<sup>2)</sup> Japan Society for the Promotion of Science, Japan

<sup>3)</sup> Department of Social Medicine, Hirosaki University Graduate School of Medicine, Japan

<sup>4)</sup> Physical Fitness Research Institute, Meiji Yasuda Life Foundation of Health and Welfare, Japan

<sup>5)</sup> Faculty of Social Welfare, Yamaguchi Prefectural University, Japan

<sup>6)</sup> Center for Preventive Medical Sciences, Chiba University, Japan

<sup>7)</sup> Faculty of Health and Sport Sciences, University of Tsukuba, Japan

**Abstract.** [Purpose] There is a relationship between physical and cognitive functions; therefore, impairment of physical function would mean cognitive decline. This study aimed to investigate the association between change in physical and cognitive functions. [Subjects and Methods] Participants were 169 healthy community-dwelling older adults who attend the survey after three years from baseline (mean age,  $72.4 \pm 4.8$  years). Grip strength, one-leg standing balance, five-times-sit-to-stand test, timed up and go, 5-m habitual walk, and a peg-moving task were used to evaluate physical performance. Five cognitive function tests were used to assess attention, memory, visuospatial function, verbal fluency, and reasoning. Cognitive function was defined as the cumulative score of these tests. [Results] At baseline, five-times-sit-to-stand test, timed up and go, and hand dexterity were independently associated with cognitive function. In longitudinal analyses, changes in habitual walking speed and hand dexterity were significantly associated with change in cognitive function. [Conclusion] Deterioration of specific physical function, such as hand dexterity and walking ability, may be associated with progression of cognitive decline. Decreasing extent of daily functions, such as hand dexterity and walking ability, can be useful indices to grasp changes in cognitive function.

**Key words:** Cognition, Physical function, Performance test

(This article was submitted Jun. 5, 2017, and was accepted Jun. 26, 2017)

### INTRODUCTION

Physical and cognitive functions are important for maintaining independent living in old age<sup>1)</sup>. Previous epidemiological studies with cross-sectional and longitudinal designs have reported a relationship between physical and cognitive functions in older adults<sup>2–8)</sup>. In general, older adults with poor physical performance exhibit low cognitive function.

Advancing age leads to a decline in physical performance. A Japanese prospective study, which included grip strength, usual gait speed, and one-leg standing balance as physical performance tests, showed that the change in physical performance with increasing age depends on those tests<sup>9)</sup>. For example, although change in usual gait speed remains almost unchanged from the age of 65 to 75 years, one-leg standing balance decreases by approximately 10 seconds in that period. Based on the

\*Corresponding author. Takumi Abe (E-mail: s1530485@u.tsukuba.ac.jp)

©2017 The Society of Physical Therapy Science. Published by IPEC Inc.



This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <http://creativecommons.org/licenses/by-nc-nd/4.0/>)

implications, the longitudinal association between change in physical and cognitive functions would also depend on physical performance tests. A previous systematic review pointed out that the findings on the association between change in physical and cognitive functions are relatively small because few studies have reported the association<sup>10</sup>. Therefore, it is necessary to examine whether a change in physical performance is associated with a change in cognitive function.

Previous studies have predominantly used a small number of performance tests to examine the relationship between physical and cognitive functions<sup>3, 6, 11, 12</sup>. As stated above, each physical performance has a different decreasing rate. A comprehensive evaluation of physical function (i.e., using various physical performance tests), as opposed to the use of a single performance test such as a walking test, is required to determine the association between changes in physical and cognitive functions in detail.

This study aimed to longitudinally investigate the association between changes in physical and cognitive functions with various physical performance tests. This would reveal the changes in physical function that correspond to/indicate changes in cognitive function. We hypothesized that physical function can be divided into two types: performance predominately requiring muscle strength, such as upper and lower extremity muscle strength, and performance that is less reliant on muscle strength, such as hand dexterity. The latter type of physical function has previously been shown to be strongly associated with cognitive function.

## **SUBJECTS AND METHODS**

This study included participants of the cohort study known as the “Kasama study.” The detail of the Kasama study was described previously<sup>13</sup>. Briefly, the Kasama study includes follow-up and new participants aged  $\geq 65$  without long-term care insurance. Physical and cognitive functions are primary measurements. We defined the first participation in the survey from 2009 to 2012 as baseline, with the follow-up period set at three years. Although 604 participants who have complete data were identified from the sampling of the baseline period of this study, we excluded data based on following criteria: loss to follow-up ( $n=431$ ); missing data at follow up period ( $n=4$ ). The final sample included 169 participants. Follow-up data could be obtained from only 28.0% of the participants. We assume that most participants included for the analyses are healthy and highly health-conscious because they have maintained a good health status for three years and because they responded to the call for participation in our study. This study was approved by the Ethical Committee of University of Tsukuba (Ref No., Tai 21–25 and Tai 23–36, 26–132). Details of this study were explained to participants prior to enrollment. Informed written consent was obtained from all participants.

To measure grip strength, participants were asked to demonstrate grip strength by squeezing a hand dynamometer (TKK5401, Takei Scientific Instruments Co. Ltd., Niigata, Japan) one after the other. The highest score of two trials for each hand was used for analysis. In measuring one-leg balance with eyes open, we recorded the time for which participants were able to stand on one leg up to a maximum of 60 sec to evaluate static balance. Participants were allowed to choose the lifted leg. The longest time of two trials was used for analysis. In measuring five-times-sit-to-stand movement test, participants were asked to sit on a chair with arms folded over their chests and then to stand up and sit down from a chair five times as fast as possible. The shorter time of two trials was used for analysis. The timed up and go test (TUG) is a commonly used performance test to evaluate functional mobility or dynamic balance. The distance between the chair and marker was 3 m. According to the modified method<sup>14</sup>, participants were instructed to rise from the chair, turn around a marker, and sit down on the chair as fast as possible. We recorded the time taken to complete the task, with the faster time of two trials used for analysis. In measuring 5-m habitual walk, we created a 11-m walking path with a 5-m central part for measurements. Participants were asked to walk at their usual pace along the walking path two times, with the shorter time being recorded. Peg-moving task was used to evaluate hand dexterity. We used a peg board (TKK1306, Takei Scientific Instruments Co. Ltd., Niigata, Japan). Participants were asked to move two pegs simultaneously from the distal board to proximal board using both hands as fast as possible. We recorded the time required for participants to finish moving 48 pegs.

We used five cognitive function tests, known as the “5 Cog,” to evaluate cognitive function<sup>15, 16</sup>. This included five tests. The position Stroop, category cued recall, clock drawing, category fluency, and similarity tests that are part of the Wechsler Adult Intelligence Scale-Revised were included for evaluating attention, memory, visuospatial function, verbal fluency, and reasoning respectively. Participants were asked to perform the 5 Cog. We calculated total 5-Cog scores to define cognitive function in this study.

Using a self-questionnaire, we obtained the demographic information: age; gender; education; medical history of stroke, diabetes, and heart disease; current medications. Depression symptoms were evaluated using the Geriatric Depression Scale<sup>17</sup>. Height and weight were used to calculate body mass index according to the following formula:  $[\text{kg}/\text{m}^2]$ .

The mean and standard deviation of continuous variables and the ratio of categorical variables at baseline were described as descriptive data. Change in physical and cognitive functions were defined as the amount of change from baseline to the end of the follow-up period (minus indicates function decline). We analyzed the following associations: 1) the cross-sectional association between physical and cognitive functions; 2) the longitudinal association between change in physical and cognitive functions. Multiple regression analysis was used to determine associations. In the crude model, cognitive function was used as the dependent variable and physical function as the independent variable. The adjusted model included age, gender, body mass index, education ( $<12$  or  $\geq 12$ ), depressive mood ( $<6$  or  $\geq 6$ ), smoking status (none, past, current), current medications,

**Table 1.** Characteristics of the participants

		All (n=169)
		Mean ± SD
Characteristics		
Age	(year)	72.4 ± 4.8
Women	n, (%)	80 (47.3)
Height	(cm)	156.7 ± 8.3
Weight	(kg)	57.2 ± 9.0
Body mass index	(kg / m <sup>2</sup> )	23.2 ± 2.6
Education	(year)	
<12	n, (%)	113 (66.9)
12≤	n, (%)	56 (33.1)
Depression mood		
<6	n, (%)	136 (80.5)
6≤	n, (%)	33 (19.5)
Smoking status		
None	n, (%)	125 (74.0)
Past	n, (%)	40 (23.7)
Current	n, (%)	4 (2.4)
Current medications	n, (%)	39 (29.8)
Medical history of stroke	n, (%)	6 (3.6)
Medical history of diabetes	n, (%)	20 (11.8)
Medical history of heart disease	n, (%)	29 (17.2)
Physical function		
Grip strength	(kg)	30.1 ± 7.7
One-leg standing balance	(sec)	37.4 ± 21.3
Five-times-sit-to-stand test	(sec)	7.5 ± 2.1
Timed up and go	(sec)	5.7 ± 1.0
5-m habitual walk	(sec)	3.6 ± 0.5
Peg-moving task	(sec)	37.2 ± 5.5
Cognitive function		
Total score	(point)	67.5 ± 15.7

**Table 2.** Change in physical and cognitive functions

		All (n=169)
		Mean ± SD
Physical function		
ΔGrip strength	(kg)	-1.11 ± 2.79
ΔOne-leg standing balance	(sec)	-4.06 ± 19.76
ΔFive-times-sit-to-stand test	(sec)	-0.27 ± 2.07
ΔTimed up and go	(sec)	-0.21 ± 0.88
Δ5-m habitual walk	(sec)	-0.13 ± 0.65
ΔPeg-moving task	(sec)	0.40 ± 3.97
Cognitive function		
ΔTotal score	(point)	5.72 ± 9.21

Δ: amount of change from baseline to the end of the follow-up period (minus indicates function decline).

medical history of stroke, diabetes, and heart disease as covariates. P-values <0.05 were considered to be statistically significant. All analyses were conducted using IBM SPSS Statistics for Windows, Version 21.0 (IBM Corp., Armonk, NY, USA).

## RESULTS

Participant characteristics at baseline are shown in Table 1. Changes in physical and cognitive functions are presented in Table 2. Table 3 shows the cross-sectional association between physical and cognitive functions. All measurements were significantly associated with cognitive function in the crude model, except the grip strength. However, the five-times-sit-to-stand test ( $\beta=-0.222$ ,  $p=0.001$ ), TUG ( $\beta=-0.169$ ,  $p=0.030$ ), and peg-moving task ( $\beta=-0.213$ ,  $p=0.005$ ) remained significantly associated with cognitive function after the inclusion of covariates. In longitudinal analyses, changes in the five-times-sit-to-stand test, 5-m habitual walk, and peg-moving task performance were found to be significantly associated with change in cognitive function in the crude model (Table 4). In the adjusted model, these associations were also observed in the 5-m habitual walk ( $\Delta$  5-m habitual walk,  $\beta=0.174$ ,  $p=0.031$ ) and peg-moving task ( $\Delta$  peg-moving task,  $\beta=0.172$ ,  $p=0.029$ ).

## DISCUSSION

This study investigated the association between physical and cognitive functions from two perspectives. First, the evaluation of cross-sectional associations revealed that the five-times-sit-to-stand test, TUG, and peg-moving task were independently associated with cognitive function. Next, we confirmed a significant longitudinal association between change in 5-m habitual walk and the peg-moving task and change in cognitive function.

**Table 3.** The association between physical and cognitive functions at baseline

	Crude		Adjusted	
	$\beta$	p value	$\beta$	p value
Grip strength	0.068	0.383	0.181	0.073
One-leg standing balance	0.294	0.001	0.123	0.075
Five-times-sit-to-stand test	-0.402	<0.001	-0.222	0.001
Timed up and go	-0.379	<0.001	-0.169	0.030
5-m habitual walk	-0.231	0.003	-0.115	0.099
Peg moving task	-0.381	<0.001	-0.213	0.005

Adjusted for age, gender, body mass index, education (<12 or  $\geq$ 12), depressive mood (<6 or  $\geq$ 6), smoking status (none, past, current), current medications, medical history of stroke, diabetes, and heart disease.

**Table 4.** The association between change in physical and cognitive functions

	Crude		Adjusted	
	$\beta$	p value	$\beta$	p value
$\Delta$ Grip strength	0.022	0.774	0.010	0.904
$\Delta$ One-leg standing balance	0.103	0.182	0.094	0.251
$\Delta$ Five-times-sit-to-stand test	0.165	0.033	0.148	0.060
$\Delta$ Timed up and go	-0.149	0.053	0.056	0.492
$\Delta$ 5-m habitual walk	0.194	0.011	0.174	0.031
$\Delta$ Peg-moving task	0.180	0.019	0.172	0.029

Adjusted for age, gender, body mass index, education (<12 or  $\geq$ 12), depressive mood (<6 or  $\geq$ 6), smoking status (none, past, current), current medications, medical history of stroke, diabetes, and heart disease.

A cross-sectional association between physical and cognitive functions has been reported by previous epidemiological studies<sup>2-5</sup>). In the five-times-sit-to-stand test, our results supported previous studies<sup>2, 4</sup>). As for TUG and hand dexterity, there is relatively little evidence. It has been confirmed that TUG is associated with executive function in older adults with mild cognitive impairment (MCI)<sup>18</sup>). Most participants in this study were without MCI or dementia because they did not use long-term care insurance. We obtained results that were similar to those obtained from older adults with MCI. Evaluations of hand dexterity were not included in previous epidemiological studies previous epidemiology studies<sup>2-5</sup>). A review focused on hand motor function and demonstrated that hand dexterity is associated with cognitive function<sup>19</sup>). Schröter et al.<sup>20</sup>) reported that participants with mild cognitive impairment or probable Alzheimer's disease have decreased hand motor function compared with participants without cognitive decline. In corroboration with these studies, hand dexterity was found to be significantly associated with cognitive function in this study. Except for hand dexterity, some physical functions were not found to be associated with cognitive function even adjustment for covariates, indicating that variables included as covariates influenced these associations. Therefore, as a novel finding, it is likely that TUG and hand dexterity as well as the five-times-sit-to-stand test are more closely associated with cognitive function than physical function, such as grip strength or lower extremity function, in cross-sectional analyses.

The relation between changes in physical and cognitive functions has been demonstrated by previous longitudinal studies<sup>8, 21</sup>). The performance tests used in this study, for example, the TUG or peg-moving task, differed from those used in previous studies. Accordingly, our results demonstrate that change in physical function, such as hand dexterity and walking ability, are associated with change in cognitive function in the adjusted model. In particular, cross-sectional and longitudinal associations were observed between hand dexterity and cognitive function in this study. These findings indicate that a decline in hand dexterity may accompany cognitive decline and suggest that hand motor function is connected to brain function<sup>22</sup>). Of measures of physical function, hand motor function appears to be most relevant to cognitive function. Watson et al.<sup>12</sup>) reported declining cognitive function to be associated with change in walking ability. In corroboration with their study, a significant relation between changes in walking ability and cognitive function was observed in this study. Furthermore, gait speed at baseline is known to be a predictor of incident dementia<sup>7</sup>). Our findings indicate that rapid decreases in gait speed may be a sign of cognitive decline even if older adults do not have a slow gait speed at baseline.

This study has some limitations. First, it is possible that the follow-up period was insufficient for healthy older adults as changes in cognitive function from baseline to the end of the follow-up period were relatively small. Accordingly, we were unable to study the development of mild cognitive impairment or dementia. Future long-term observation studies with large

sample sizes are required to overcome this limitation. Next, our study cannot deny a sampling bias because our sample consisted of some individuals who attended the follow-up survey. Older adults with a lower level of cognitive function or bad health status tend to be non-responders<sup>23</sup>; that is, participants who were included in the analysis had good health conditions. Therefore, our results can only be generalized to healthy older adults. Finally, we were unable to investigate potentially confounding factors, such as nutrition. Few studies have indicated a longitudinal association between physical function and cognitive functions. The results of this study are of importance to this field as we obtained new findings by comprehensively measuring physical function in a longitudinal study.

In conclusion, performance tests that were associated with cognitive function had differences between the cross-sectional and longitudinal analyses. Declines in specific physical functions are associated with cognitive function, particularly hand dexterity and walking ability, and may reflect progressing cognitive decline. The peg-moving task, which showed significant association with cognitive function in the cross-sectional and longitudinal analyses, would enable measuring not only hand dexterity but also cognitive function in healthy older adults.

## REFERENCES

- 1) Rajan KB, Hebert LE, Scherr P, et al.: Cognitive and physical functions as determinants of delayed age at onset and progression of disability. *J Gerontol A Biol Sci Med Sci*, 2012, 67: 1419–1426. [[Medline](#)] [[CrossRef](#)]
- 2) Bullain SS, Corrada MM, Shah BA, et al.: Poor physical performance and dementia in the oldest old: the 90+ study. *JAMA Neurol*, 2013, 70: 107–113. [[Medline](#)] [[CrossRef](#)]
- 3) Fitzpatrick AL, Buchanan CK, Nahin RL, et al. Ginkgo Evaluation of Memory (GEM) Study Investigators: Associations of gait speed and other measures of physical function with cognition in a healthy cohort of elderly persons. *J Gerontol A Biol Sci Med Sci*, 2007, 62: 1244–1251. [[Medline](#)] [[CrossRef](#)]
- 4) Narazaki K, Matsuo E, Honda T, et al.: Physical fitness measures as potential markers of low cognitive function in Japanese community-dwelling older adults without apparent cognitive problems. *J Sports Sci Med*, 2014, 13: 590–596. [[Medline](#)]
- 5) Won H, Singh DK, Din NC, et al.: Relationship between physical performance and cognitive performance measures among community-dwelling older adults. *Clin Epidemiol*, 2014, 6: 343–350. [[Medline](#)]
- 6) Atkinson HH, Rosano C, Simonsick EM, et al. Health ABC study: Cognitive function, gait speed decline, and comorbidities: the health, aging and body composition study. *J Gerontol A Biol Sci Med Sci*, 2007, 62: 844–850. [[Medline](#)] [[CrossRef](#)]
- 7) Wang L, Larson EB, Bowen JD, et al.: Performance-based physical function and future dementia in older people. *Arch Intern Med*, 2006, 166: 1115–1120. [[Medline](#)] [[CrossRef](#)]
- 8) Atkinson HH, Rapp SR, Williamson JD, et al.: The relationship between cognitive function and physical performance in older women: results from the women's health initiative memory study. *J Gerontol A Biol Sci Med Sci*, 2010, 65: 300–306. [[Medline](#)] [[CrossRef](#)]
- 9) Taniguchi Y, Fujiwara Y, Murayama H, et al.: Prospective study of trajectories of physical performance and mortality among community-dwelling older Japanese. *J Gerontol A Biol Sci Med Sci*, 2016, 71: 1492–1499. [[Medline](#)] [[CrossRef](#)]
- 10) Clouston SA, Brewster P, Kuh D, et al.: The dynamic relationship between physical function and cognition in longitudinal aging cohorts. *Epidemiol Rev*, 2013, 35: 33–50. [[Medline](#)] [[CrossRef](#)]
- 11) Soumaré A, Tavernier B, Aléprouvitch A, et al.: A cross-sectional and longitudinal study of the relationship between walking speed and cognitive function in community-dwelling elderly people. *J Gerontol A Biol Sci Med Sci*, 2009, 64: 1058–1065. [[Medline](#)] [[CrossRef](#)]
- 12) Watson NL, Rosano C, Boudreau RM, et al. Health ABC Study: Executive function, memory, and gait speed decline in well-functioning older adults. *J Gerontol A Biol Sci Med Sci*, 2010, 65: 1093–1100. [[Medline](#)] [[CrossRef](#)]
- 13) Okura T, Tsuji T, Tsunoda K, et al.: Study protocol and overview of the Kasama study: creating a comprehensive, community-based system for preventive nursing care and supporting successful aging. *J Phys Fit Sport Med*, 2017, 6: 49–57.
- 14) Shumway-Cook A, Brauer S, Woollacott M: Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Phys Ther*, 2000, 80: 896–903. [[Medline](#)]
- 15) Miyamoto M, Kodama C, Kinoshita T, et al.: Dementia and mild cognitive impairment among non-responders to a community survey. *J Clin Neurosci*, 2009, 16: 270–276. [[Medline](#)] [[CrossRef](#)]
- 16) Sato D, Seko C, Hashitomi T, et al.: Differential effects of water-based exercise on the cognitive function in independent elderly adults. *Aging Clin Exp Res*, 2015, 27: 149–159. [[Medline](#)] [[CrossRef](#)]
- 17) Sheikh JI, Yesavage JA: Geriatric Depression Scale (GDS). *Clin Gerontol*, 1986, 5: 165–173.
- 18) McGough EL, Kelly VE, Logsdon RG, et al.: Associations between physical performance and executive function in older adults with mild cognitive impairment: gait speed and the timed “up & go” test. *Phys Ther*, 2011, 91: 1198–1207. [[Medline](#)] [[CrossRef](#)]
- 19) Scherder E, Dekker W, Eggermont L: Higher-level hand motor function in aging and (preclinical) dementia: its relationship with (instrumental) activities of daily life—a mini-review. *Gerontology*, 2008, 54: 333–341. [[Medline](#)] [[CrossRef](#)]
- 20) Schröter A, Mergl R, Bürger K, et al.: Kinematic analysis of handwriting movements in patients with Alzheimer's disease, mild cognitive impairment, depression and healthy subjects. *Dement Geriatr Cogn Disord*, 2003, 15: 132–142. [[Medline](#)] [[CrossRef](#)]
- 21) Tabbarah M, Crimmins EM, Seeman TE: The relationship between cognitive and physical performance: MacArthur studies of successful aging. *J Gerontol A Biol Sci Med Sci*, 2002, 57: M228–M235. [[Medline](#)] [[CrossRef](#)]
- 22) Koppelmans V, Hirsiger S, Méritlat S, et al.: Cerebellar gray and white matter volume and their relation with age and manual motor performance in healthy older adults. *Hum Brain Mapp*, 2015, 36: 2352–2363. [[Medline](#)] [[CrossRef](#)]
- 23) Launer LJ, Wind AW, Deeg DJ: Nonresponse pattern and bias in a community-based cross-sectional study of cognitive functioning among the elderly. *Am J Epidemiol*, 1994, 139: 803–812. [[Medline](#)] [[CrossRef](#)]