



Published in final edited form as:

Cancer Metastasis Rev. 2017 September ; 36(3): 463–473. doi:10.1007/s10555-017-9687-8.

EGFR-Targeted Therapies in the Post-Genomic Era

Mary Jue Xu, Daniel E. Johnson, and Jennifer R. Grandis

Department of Otolaryngology, University of California San Francisco, 2380 Sutter Street, San Francisco, CA 94113, USA

Abstract

Over ninety percent of head and neck cancers overexpress the epidermal growth factor receptor (EGFR). In diverse tumor types, EGFR overexpression has been associated with poorer prognosis and outcomes. Therapies targeting EGFR include monoclonal antibodies, tyrosine kinase inhibitors, phosphatidylinositol 3-kinase (PI3K) inhibitors, and antisense gene therapy. Few EGFR-targeted therapeutics are approved for clinical use. The monoclonal antibody cetuximab is Food and Drug Administration (FDA) approved EGFR-targeted therapy, yet has exhibited modest benefit in clinical trials. The humanized monoclonal antibody nimotuzumab is also approved for head and neck cancers in Cuba, Argentina, Colombia, Peru, India, Ukraine, Ivory Coast and Gabon in addition to nasopharyngeal cancers in China. Few other EGFR-targeted therapeutics for head and neck cancers have led to as significant responses as seen in lung carcinomas, for instance. Recent genome sequencing of head and neck tumors has helped identify patient subgroups with improved response to EGFR inhibitors, for example cetuximab in patients with the KRAS-variant and the tyrosine kinase inhibitor erlotinib for tumors harboring MAPK1^{E322K} mutations. Genome sequencing has furthermore broadened our understanding of dysregulated pathways, holding the potential to enhance the benefit derived from therapies targeting EGFR.

Keywords

EGFR; head and neck SCC; genomics; cetuximab

1. Introduction

Stanley Cohen's discovery of epidermal growth factor (EGF) was awarded the 1986 Nobel Prize, heralding the development of EGFR-targeted therapeutics [1]. In diverse tumor types including head and neck, bladder, ovarian and cervical cancers, EGFR overexpression has been associated with poorer prognosis and outcomes [2–4]. In 2004, the FDA initially approved the monoclonal antibody cetuximab for metastatic colorectal cancer. Its use was expanded to head and neck squamous cell carcinomas (HNSCC) in 2006. Cetuximab remains the only EGFR-directed treatment FDA-approved for head and neck cancers. Here,

Corresponding Author, Dr. Jennifer R. Grandis, American Cancer Society Professor, Associate Vice Chancellor, Clinical and Translational Research Director, Clinical and Translational Science Institute Professor, Otolaryngology – Head and Neck Surgery, 550 16th. Street, San Francisco, CA 94158, Phone: 415-514-8060, Jennifer.Grandis@ucsf.edu.

Conflict of Interest

The authors declare that they have no conflict of interest.

we review EGFR-targeted therapies and highlight insights from recent genomic research relevant to head and neck cancers.

2. Receptor Pathway and Function

2.1 EGFR Structure

EGFR, also called HER1 or ErbB1, was the first member of the ErbB family of tyrosine kinase receptors discovered [5]. This family also includes HER2/neu (ErbB2), HER3 (ErbB3), and HER4 (ErbB4). The 170 kDa EGFR receptor spans the membrane once and contains extracellular, transmembrane, and intracellular regions. The extracellular component is comprised of 4 domains. Domains I and III are leucine rich and structurally similar to domains found in the insulin receptor [6], a cell surface receptor known to share downstream signaling pathways with EGFR [7,8]. Domains II and IV are cysteine rich and similar to laminin [9]. The intracellular region harbors the intrinsic tyrosine kinase activity of EGFR. Existing in both closed monomer and open dimer conformations [10], EGFR is composed of twenty percent carbohydrates, with N-linked glycosylation affecting receptor structure and stability; increased glycosylation stabilizes and drives the equilibrium towards the extended conformation [4,11,12].

2.2 EGFR Pathway

Epidermal growth factor (EGF), transforming growth factor- α (TGF- α), amphiregulin, heparin-binding EGFR, and betacellulin are among the ligands which bind to domains I and III of EGFR. Subsequent exposure of domain II results in receptor dimerization via disulfide bonds. After dimerization at the cell surface, autophosphorylation of tyrosine residues in the cytoplasmic region provides docking sites for signal transducers, including proteins such as Ras, to bind and initiate intracellular signaling cascades and gene transcription [4,13]. Downstream signaling cascades of EGFR can be broadly divided into the following pathways: RAS/RAF/MEK/MAPK/ERK, phosphatidylinositol 3-kinase (PI3K) and Akt, protein kinase C (PKC), Src, and the JAK/STAT pathways (Figure 1) [14]. These extensively studied signaling cascades influence gene expression, proliferation, angiogenesis, apoptosis inhibition, cell motility, metastasis, adhesion, and angiogenesis [4,15].

2.3 EGFR Function in Normal Physiology and Cancer

Indisputably, EGFR possesses a critical role in development and differentiation, particularly in epithelial and glial cells. Highly expressed in the basal layer of the epidermis and the outer root sheath of hair cells, EGFR influences migration and differentiation of keratinocytes and hair follicle development. Mouse models expressing mutant EGFR develop papillomas and squamous cell carcinomas (SCC) [16]. In neurons, EGFR regulates migration and neurodegeneration, with mutations leading to glioma-like tumors in murine models [16]. Furthermore, in lung tissue, EGFR influences maturation of type II pneumocytes; following lung damage these cells proliferate into type I pneumocytes, and replace damaged tissue.

In head and neck cancers, EGFR is overexpressed in over 90% of tumors and correlates with poorer outcomes [17,18]. In tissue from 91 HNSCC patients, tumor EGFR level was a

statistically significant predictor of disease-free survival (DFS) ($p=0.0001$) along with tumor site and TGF- α level [17]. In the large phase III RTOG 9003 trial evaluating radiation regimens, retrospective subset analysis of 155 patients reinforced the correlation between EGFR expression and decreased overall survival (OS) along with increased local-regional relapse [19]. In addition to overall increased expression, EGFR copy number was associated with a 91% (20/22 patients) 5-year mortality compared to 29% (30/102 patients) in patients with a normal copy number [20]. Similar associations exist for breast, lung, and other tumor types [4,21,22].

3. EGFR Targeted Therapies

Until the development of targeted therapeutics, chemotherapy for head and neck cancers was predominated by non-specific inhibitors of cellular division and proliferation. FDA-approved therapies included cisplatin, methotrexate, 5-fluorouracil (5-FU), bleomycin, and docetaxel, all of which produced clinical response rates ranging from 20–40% [22]. Common side effects included dysphagia, odynophagia, nausea, vomiting, and hematologic suppression [22,23]. EGFR-targeted therapies approved and under-development include monoclonal antibodies (Table I), tyrosine kinase inhibitors (Table II), PI3K inhibitors, and antisense gene therapy.

3.1 Monoclonal Antibodies

In 2006, cetuximab was the first targeted treatment for head and neck cancers approved by the FDA (Table I). A chimeric murine antibody linked to human IgG, cetuximab was approved in combination with radiation (XRT) in locally advanced (LA) disease, as a single agent for recurrent or metastatic HNSCC after failure of platinum therapies, and in combination with 5-FU and platinum based therapies for first-line recurrent or metastatic HNSCC [14]. In addition to inhibiting ligand binding, alternative mechanisms of action involve initiating receptor endocytosis, activating antibody-dependent cell-mediated cytotoxicity (ADCC), and inhibiting repair of radiation-induced damage [23,24].

In clinical care, cetuximab improved patient outcomes when combined with radiotherapy (Table III). Randomized, phase III, multicenter trials assessing the addition of cetuximab to radiotherapy noted increased local-regional control and increased median OS from 29.3 months (95% CI 20.6–41.4) to 49.0 months (95% CI 32.8–69.5) [25,26]. Importantly, patients experienced unchanged rates of treatment-related toxicities. However, higher grade of acneiform rash, a common side effect, was associated with improved OS and thought indicative of an inflammatory response [26].

Cetuximab also conferred additional benefit in combination with chemotherapy (Table III). In a phase II multicenter study, patients with recurrent or metastatic HNSCC were started on cetuximab therapy; cisplatin was subsequently added following disease progression. Of the 103 patients, 46% benefited from cetuximab with either disease control or stabilization with a mean time to progression of 70 days [27]. Similarly, in a phase III trial, addition of cetuximab to platinum-based and 5-FU therapies increased median OS from 7.4 months to 10.1 months and progression-free survival (PFS) from 3.3 months to 5.6 months [28]. Though the improvements observed were modest, these trials prompted FDA approval for

rates of 55% to 75% for patients harboring an EGFR tyrosine kinase domain mutation [38,39].

A new generation of TKIs with multiple targets and irreversible binding have shown clinical potential in HNSCC. Afatinib, an irreversible inhibitor of EGFR, HER2, and HER4 kinases exhibited comparable outcomes to cetuximab. In a randomized, phase II study assessing afatinib versus cetuximab for treatment of recurrent or metastatic HNSCC in 124 patients, median OS was 35.9 weeks with afatinib and 47.1 weeks for cetuximab ($p = 0.78$) [40]. Following treatment failure in each arm, patients were transferred to the other treatment arm, during which disease control was 38.9% with afatinib and 18.8% with cetuximab. In light of these promising results, a phase III trial involving 483 patients following treatment failure on platinum-based therapy noted improved PFS with use of afatinib (median 2.6 months) compared to methotrexate (median 1.7 months) for second-line treatment (hazard ratio (HR) 0.80, 95% CI 0.65–0.98, $p=0.03$) [41].

Dacomitinib, another irreversible multi-targeted TKI, and lapatinib, an oral reversible inhibitor of EGFR and HER2, have exhibited limited effects in early studies [42–46].

3.3 Phosphatidylinositol 3-kinase (PI3K) Inhibitors

PI3K mutations are prevalent in head and neck cancers, noted in 34% of HPV negative HNSCC and 56% of HPV positive samples [46,47]. Buparlisib is an oral, pan-PI3K inhibitor noted to modestly improve PFS in recurrent and metastatic head and neck cancer patients (Table III). In a phase II trial of 158 patients assessing buparlisib as a second-line therapy following progression on platinum-based chemotherapy, buparlisib improved median PFS to 4.6 months with buparlisib and paclitaxel compared to 3.5 months with placebo and paclitaxel (HR 0.65, 95% CI 0.45–0.95) [49]. Of note, 46% of patients were previously treated with EGFR-targeted therapy. Future studies in varying patient populations may elicit more marked improvements in survival.

3.4 Antisense Gene Therapy

Antisense therapy centers on inhibiting messenger RNA (mRNA) by binding complementary, engineered nucleic acids. This is thought to lead to inhibition of transcription, splicing, and mRNA modification. An additional mechanism described is RNase H-mediated cleavage [50].

EGFR-targeted antisense therapy has completed early phase clinical testing. In a phase I trial of 17 HNSCC patients, antisense DNA targeting EGFR was directly injected into patients' tumors. Seven patients demonstrated either stable or clinically responsive disease noted by decreased tumor volume [51]. A phase I/II trial combining EGFR antisense with radiation and cetuximab was recently completed (NCT01592721). Future research will also need to address systemic activity of EGFR-targeted antisense activity.

4. Insights from Genomic Research

Despite the widespread overexpression of EGFR in cancers, cetuximab treatment leads to only a modest response in HNSCC [52]. As a novel tool, genome sequencing has

restructured our understanding of dysregulated pathways and provided deeper insight into EGFR-targeted therapies.

Given the broad landscape of mutations in HNSCC, mutations in four major classes of proteins/pathways have been identified: 1) mitogenic pathways (PI3K/mTOR), 2) differentiation and NOTCH pathways, 3) regulators of cell cycle proliferation through p16 and cyclin D1, and 4) regulators of apoptosis, including p53, whose loss of function is found almost universally in smoking-related HNSCC (Table IV) [47]. Whole-exome sequencing of 151 head and neck tumor samples revealed that, aside from p53, the PI3K pathway, which promotes mitogenic signaling, was the most commonly mutated pathway, with mutations occurring in 30.5% of samples [47,48]. Additional sequencing efforts discovered novel mutations in NOTCH1, functioning as a tumor suppressor gene [53,54].

Sequencing of HNSCC tumors has not identified recurrent EGFR driving mutations. In contrast to NSCLC in which EGFR mutations are clustered in exons 18–21, the region encoding the tyrosine kinase domain, EGFR mutations in head and neck cancers appear more dispersed across the gene (Figure 2) [54]. Chang *et al.* (2016) assessed 11,119 human tumor samples and 41 types of cancers to create an algorithm identifying frequently mutated residues; hot spots were noted in HRAS and PIK3CA in head and neck cancers but not in EGFR [55]. Perhaps lack of recurrent EGFR mutations contributes to the limited effects of TKIs and cetuximab in HNSCC. In contrast, TKIs for the treatment of NSCLC which harbor tyrosine kinase domain mutations exhibit RECIST response rates of 55% to 75% [39].

With the lack of driving mutations and the global upregulation of EGFR, the vast landscape of mutations implicates co-activation of additional pathways. Notably the *KRAS*-variant germline and MAPK1^{E322K} mutation were highlighted in recent literature. Patients harboring a germline mutation in the micro-RNA binding site of *KRAS* have poorer overall survival [57]. Surprisingly, in a phase III trial in which cetuximab did not confer benefit when added to chemoradiation in unselected HNSCC patients [29], patients with the *KRAS*-variant (70 of 413 patients tested) had increased OS in the first two years following treatment with cetuximab (HR 0.19; 95% CI, 0.04–0.86; *P* = 0.03) [57]. This improvement in survival from cetuximab was not seen for wild-type *KRAS* patients. In *KRAS*-variant patients, TFG-β1 was found to be upregulated; this cytokine has been implicated in suppressing antitumor immunity through regulatory T-cell induction [58]. Authors of this study proposed that through ADCC and improved dendritic cell priming of cytotoxic T lymphocytes, cetuximab bolstered the antitumor immunity otherwise inhibited in *KRAS*-variant patients [57].

In addition to the *KRAS*-variant, genomic sequencing revealed that tumor samples from a patient with a MAPK1^{E322K} mutation were exquisitely sensitive to EGFR TKIs. In a window-of-opportunity clinical trial, a patient with a stage IVA tongue carcinoma who received a 13-day course of erlotinib experienced remarkable disease reduction from initial clinical T1N2c disease with bulky lymphadenopathy to pathological T1N0 disease. Following surgery, the patient has remained disease-free for more than 4 years without additional treatment [59]. No EGFR mutation was identified in this patient. However, the patient's MAPK1^{E322K} mutation was studied in in vitro and in vivo models and found to be

associated with upregulation of amphiregulin and stimulation of an autocrine feedback loop involving EGFR, ERK, and amphiregulin. Remarkably, upregulated amphiregulin increased tumor sensitivity to erlotinib, an effect emphasized by the loss of erlotinib sensitivity following amphiregulin knockdown in MAPK1^{E322K} models [60].

Improved response to EGFR inhibitors (cetuximab in HNSCC tumors with the *KRAS*-variant and erlotinib in HNSCCs harboring MAPK1^{E322K} mutations) emphasizes the importance of patient selection for EGFR-targeted therapies. These studies suggest that genomic sequencing will further elicit predictive biomarkers of EGFR therapeutic response and deepen our understanding of EGFR-related cellular dysfunction that can be exploited in the clinic.

In summary, the clinical benefit of EGFR-targeted therapies in head and neck tumors has been more modest than expected given the near universal upregulation of EGFR. No dominant EGFR driver mutation has been discovered in HNSCC as in NSCLC, and *KRAS* mutations do not clearly indicate endogenous cetuximab resistance as they have in colon cancer. Most HNSCC cohorts sequenced to date have been performed on primary tumors without accompanying information on cetuximab treatment and clinical outcome. The coexistence of multiple deregulated pathways, in the absence of driver EGFR mutations, strongly supports the co-activation of alternative signaling pathways as a mechanism of *de novo* or acquired cetuximab resistance. As with *KRAS*-variant tumors and MAPK1^{E322K} mutations, opportunities to exploit these pathways may lead to improved patient selection and therapeutic strategies.

5. Conclusion

Cetuximab remains the only FDA approved EGFR-targeted therapy for HNSCC and provides improved survival in a subset of patients when used in combination with chemotherapy or radiation. However, long-term survival rates for head and neck cancers have remained unchanged despite increased use of EGFR-targeted therapies. Continued genomic research understanding the dysregulated and co-activated pathways will improve patient selection and future EGFR-targeted strategies.

Acknowledgments

This work was supported by National Institutes of Health grants R01 DE24728 (DEJ), P50CA097190 (DEJ and JRG), and R01 DE023685 (JRG), and American Cancer Society grant CRP-13-308-06-COUN (JRG).

References

1. Cohen S. Purification of a nerve-growth promoting protein from the mouse salivary gland and its neuro-cytotoxic antiserum. *Proceedings of the National Academy of Sciences of the United States of America*. 1960; 46(3):302–11. [PubMed: 16578483]
2. Wong AJ, Bigner SH, Bigner DD, Kinzler KW, Hamilton SR, Vogelstein B. Increased expression of the epidermal growth factor receptor gene in malignant gliomas is invariably associated with gene amplification. *Proceedings of the National Academy of Sciences of the United States of America*. 1987; 84(19):6899–903. [PubMed: 3477813]
3. Nicholson RI, Gee JM, Harper ME. EGFR and cancer prognosis. *European Journal of Cancer*. 2001; 37(Suppl 4):S9–15. [PubMed: 11597399]

4. Herbst RS. Review of epidermal growth factor receptor biology. *International Journal of Radiation Oncology, Biology, Physics*. 2004; 59(2 Suppl):21–6.
5. Cohen S. Isolation of a mouse submaxillary gland protein accelerating incisor eruption and eyelid opening in the new-born animal. *The Journal of Biological Chemistry*. 1962; 237:1555–62. [PubMed: 13880319]
6. Ward CW, Garrett TP. The relationship between the L1 and L2 domains of the insulin and epidermal growth factor receptors and leucine-rich repeat modules. *BMC Bioinformatics*. 2001; 2:4. [PubMed: 11504559]
7. van der Veeken J, Oliveira S, Schiffelers RM, Storm G, van Bergen En Henegouwen PM, Roovers RC. Crosstalk between epidermal growth factor receptor- and insulin-like growth factor-1 receptor signaling: implications for cancer therapy. *Current Cancer Drug Targets*. 2009; 9(6):748–60. [PubMed: 19754359]
8. Roudabush FL, Pierce KL, Maudsley S, Khan KD, Luttrell LM. Transactivation of the EGF receptor mediates IGF-1-stimulated shc phosphorylation and ERK1/2 activation in COS-7 cells. *Journal of Biological Chemistry*. 2000; 275(29):22583–9. [PubMed: 10807918]
9. Ward CW, Hoyne PA, Flegg RH. Insulin and epidermal growth factor receptors contain the cysteine repeat motif found in the tumor necrosis factor receptor. *Proteins*. 1995; 22(2):141–53. [PubMed: 7567962]
10. Ferguson KM, Berger MB, Mendrola JM, Cho HS, Leahy DJ, Lemmon MA. EGF activates its receptor by removing interactions that autoinhibit ectodomain dimerization. *Molecular Cell*. 2003; 11(2):507–17. [PubMed: 12620237]
11. Cummings RD, Soderquist AM, Carpenter G. The oligosaccharide moieties of the epidermal growth factor receptor in A-431 cells. Presence of complex- type N-linked chains that contain terminal N-acetylgalactosamine residues. *Journal Biological Chemistry*. 1985; 260(22):11944–52.
12. Whitson KB, Whitson SR, Red-Brewer ML, McCoy AJ, Vitali AA, Walker F, et al. Functional effects of glycosylation at Asn-579 of the epidermal growth factor receptor. *Biochemistry*. 2005; 44(45):14920–31. [PubMed: 16274239]
13. Yewale C, Baradia D, Vhora I, Patil S, Misra A. Epidermal growth factor receptor targeting in cancer: a review of trends and strategies. *Biomaterials*. 2013; 34(34):8690–707. [PubMed: 23953842]
14. Chong CR, Jänne PA. The quest to overcome resistance to EGFR-targeted therapies in cancer. *Nature Medicine*. 2013; 19(11):1389–400.
15. Goldman CK, Kim J, Wong WL, King V, Brock T, Gillespie GY. Epidermal growth factor stimulates vascular endothelial growth factor production by human malignant glioma cells: a model of glioblastoma multiforme pathophysiology. *Molecular Biology of the Cell*. 1993; 4(1):121–33. [PubMed: 7680247]
16. Sibilio M, Kroismayr R, Lichtenberger BM, Natarajan A, Hecking M, Holcman M. The epidermal growth factor receptor: from development to tumorigenesis. *Differentiation*. 2007; 75(9):770–87. [PubMed: 17999740]
17. Rubin Grandis J, Melhem MF, Gooding WE, Day R, Holst VA, Wagener MM, et al. Levels of TGF- α and EGFR protein in head and neck squamous cell carcinoma and patient survival. *Journal of the National Cancer Institute*. 1998; 90(11):824–32. [PubMed: 9625170]
18. Grandis JR, Tweardy DJ. Elevated levels of transforming growth factor alpha and epidermal growth factor receptor messenger RNA are early markers of carcinogenesis in head and neck cancer. *Cancer Research*. 1993; 53(15):3579–84. [PubMed: 8339264]
19. Ang KK, Berkey BA, Tu X, Zhang HZ, Katz R, Hammond EH, et al. Impact of epidermal growth factor receptor expression on survival and pattern of relapse in patients with advanced head and neck carcinoma. *Cancer Research*. 2002; 62(24):7350–7356. [PubMed: 12499279]
20. Temam S, Kawaguchi H, El-Naggar AK, Jelinek J, Tang H, Liu DD, et al. Epidermal growth factor receptor copy number alterations correlate with poor clinical outcome in patients with head and neck squamous cancer. *Journal of Clinical Oncology*. 2007; 25:2164–2170. [PubMed: 17538160]
21. Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science*. 1987; 235(4785):177–82. [PubMed: 3798106]

22. Franklin WA, Veve R, Hirsch FR, Helfrich BA, Bunn PA Jr. Epidermal growth factor receptor family in lung cancer and premalignancy. *Seminars in Oncology*. 2002; 29(1 Suppl 4):3–14.
23. Wen Y, Grandis JR. Emerging drugs for head and neck cancer. *Expert Opinion on Emerging Drugs*. 2015; 20(2):313–29. [PubMed: 25826749]
24. Chen DJ, Nirodi CS. The epidermal growth factor receptor: a role in repair of radiation-induced DNA damage. *Clinical Cancer Research*. 2007; 13(22 Pt 1):6555–60. [PubMed: 18006754]
25. Bonner JA, Harari PM, Giralt J, Azarnia N, Shin DM, Cohen RB, et al. Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *The New England Journal of Medicine*. 2006; 354(6):567–78. [PubMed: 16467544]
26. Bonner JA, Harari PM, Giralt J, Cohen RB, Jones CU, Sur RK, et al. Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncology*. 2010; 11(1): 21–8. [PubMed: 19897418]
27. Vermorken JB, Trigo J, Hitt R, Koralewski P, Diaz-Rubio E, Rolland F, et al. Open-label, uncontrolled, multicenter phase II study to evaluate the efficacy and toxicity of cetuximab as a single agent in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck who failed to respond to platinum-based therapy. *Journal of Clinical Oncology*. 2007; 25(16): 2171–7. [PubMed: 17538161]
28. Vermorken JB, Mesia R, Rivera F, Remenar E, Kawecki A, Rottey S, et al. Platinum-based chemotherapy plus cetuximab in head and neck cancer. *The New England Journal of Medicine*. 2008; 359(11):1116–27. [PubMed: 18784101]
29. Ang KK, Zhang Q, Rosenthal DI, Nguyen-Tan PF, Sherman EJ, Weber RS, et al. Randomized phase III trial of concurrent accelerated radiation plus cisplatin with or without cetuximab for stage III to IV head and neck carcinoma: RTOG 0522. *Journal of Clinical Oncology*. 2014; 32(27): 2940–50. [PubMed: 25154822]
30. Vermorken JB, Stöhlmacher-Williams J, Davidenko I, Licitra L, Winkvist E, Villanueva C, et al. Cisplatin and fluorouracil with or without panitumumab in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck (SPECTRUM): an open-label phase 3 randomised trial. *Lancet Oncology*. 2013; 14(8):697–710. [PubMed: 23746666]
31. Mesía R, Henke M, Fortin A, Minn H, Yunes Ancona AC, Cmelak A, et al. Chemoradiotherapy with or without panitumumab in patients with unresected, locally advanced squamous-cell carcinoma of the head and neck (CONCERT-1): a randomised, controlled, open-label phase 2 trial. *Lancet Oncology*. 2015; 16(2):208–20. [PubMed: 25596660]
32. Machiels JP, Subramanian S, Ruzsa A, Repassy G, Lifirenko I, Flygare A, et al. Zolatumumab plus best supportive care versus best supportive care alone in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck after failure of platinum-based chemotherapy: an open-label, randomised phase 3 trial. *Lancet Oncology*. 2011; 12(4):333–43. [PubMed: 21377930]
33. Rodríguez MO, Rivero TC, del Castillo Bahi R, Muchuli CR, Bilbao MA, Vinageras EN, et al. Nimotuzumab plus radiotherapy for unresectable squamous-cell carcinoma of the head and neck. *Cancer Biology & Therapy*. 2010; 9(5):343–9. [PubMed: 20448462]
34. Basavaraj C, Sierra P, Shivu J, Melarkode R, Montero E, Nair P. Nimotuzumab with chemoradiation confers a survival advantage in treatment-naïve head and neck tumors over expressing EGFR. *Cancer Biology & Therapy*. 2010; 10(7):673–81. [PubMed: 20647773]
35. Fayette J, Wirth L, Oprean C, Udrea A, Jimeno A, Rischin D, et al. Randomized Phase II Study of Duligotuzumab (MEHD7945A) vs. Cetuximab in Squamous Cell Carcinoma of the Head and Neck (MEHGAN Study). *Frontiers in Oncology*. 2016; 6:232. [PubMed: 27843803]
36. Martins RG, Parvathaneni U, Bauman JE, Sharma AK, Raez LE, Papagikos MA, et al. Cisplatin and radiotherapy with or without erlotinib in locally advanced squamous cell carcinoma of the head and neck: A randomized phase II trial. *Journal of Clinical Oncology*. 2013; 31(11):1415–21. [PubMed: 23460709]
37. Argiris A, Ghebremichael M, Gilbert J, Lee JW, Sachidanandam K, Kolesar JM, et al. Phase III randomized, placebo-controlled trial of docetaxel with or without gefitinib in recurrent or metastatic head and neck cancer: An Eastern Cooperative Oncology Group trial. *Journal of Clinical Oncology*. 2013; 31(11):1405–14. [PubMed: 23460714]

38. Paez JG, Jänne PA, Lee JC, Tracy S, Greulich H, Gabriel S, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science*. 2004; 304(5676):1497–500. [PubMed: 15118125]
39. Afghahi A, Sledge GW Jr. Targeted Therapy for Cancer in the Genomic Era. *Cancer Journal*. 2015; 21(4):294–8.
40. Seiwert TY, Fayette J, Cupissol D, Del Campo JM, Clement PM, Hitt R, et al. A randomized, phase II study of afatinib versus cetuximab in metastatic or recurrent squamous cell carcinoma of the head and neck. *Annals of Oncology*. 2014; 25(9):1813–20. [PubMed: 24928832]
41. Machiels JP, Haddad RI, Fayette J, Licitra LF, Tahara M, Vermorken JB, et al. Afatinib versus methotrexate as second-line treatment in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck progressing on or after platinum-based therapy (LUX-Head & Neck 1): an open-label, randomised phase 3 trial. *Lancet Oncology*. 2015; 16(5):583–94. [PubMed: 25892145]
42. Harrington K, Berrier A, Robinson M, Remenar E, Housset M, de Mendoza FH, et al. Randomised Phase II study of oral lapatinib combined with chemoradiotherapy in patients with advanced squamous cell carcinoma of the head and neck: rationale for future randomised trials in human papilloma virus-negative disease. *European Journal of Cancer*. 2013; 49(7):1609–18. [PubMed: 23265705]
43. Harrington K, Temam S, Mehanna H, D'Cruz A, Jain M, D'Onofrio I, et al. Postoperative Adjuvant Lapatinib and Concurrent Chemoradiotherapy Followed by Maintenance Lapatinib Monotherapy in High-Risk Patients With Resected Squamous Cell Carcinoma of the Head and Neck: A Phase III, Randomized, Double-Blind, Placebo-Controlled Study. *Journal of Clinical Oncology*. 2015; 33(35):4202–9. 10. [PubMed: 26527790]
44. Del Campo JM, Hitt R, Sebastian P, Carracedo C, Lokanatha D, Bourhis J, et al. Effects of lapatinib monotherapy: results of a randomised phase II study in therapy-naive patients with locally advanced squamous cell carcinoma of the head and neck. *British Journal of Cancer*. 2011; 105(5):618–27. [PubMed: 21829197]
45. de Souza JA, Davis DW, Zhang Y, Khattri A, Seiwert TY, Aktolga S, et al. A phase II study of lapatinib in recurrent/metastatic squamous cell carcinoma of the head and neck. *Clinical Cancer Research*. 2012; 18(8):2336–43. [PubMed: 22371453]
46. Abdul Razak AR, Soulières D, Laurie SA, Hotte SJ, Singh S, Winquist E, et al. A phase II trial of dacomitinib, an oral pan-human EGF receptor (HER) inhibitor, as first-line treatment in recurrent and/or metastatic squamous-cell carcinoma of the head and neck. *Annals of Oncology*. 2013; 24(3):761–9. [PubMed: 23108949]
47. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015; 517(7536):576–82. [PubMed: 25631445]
48. Lui VW, Hedberg ML, Li H, Vangara BS, Pendleton K, Zeng Y, et al. Frequent mutation of the PI3K pathway in head and neck cancer defines predictive biomarkers. *Cancer Discovery*. 2013; 3(7):761–9. [PubMed: 23619167]
49. Soulières D, Faivre S, Mesía R, Remenár É, Li SH, Karpenko A, et al. Buparlisib and paclitaxel in patients with platinum-pretreatment recurrent or metastatic squamous cell carcinoma of the head and neck (BERIL-1): a randomised, double-blind, placebo-controlled phase 2 trial. *Lancet Oncology*. 2017; 18(3):323–335. [PubMed: 28131786]
50. Gleave ME, Monia BP. Antisense therapy for cancer. *Nature Reviews Cancer*. 2005; 5(6):468–79. [PubMed: 15905854]
51. Lai SY, Koppikar P, Thomas SM, Childs EE, Egloff AM, Seethala RR, et al. Intratumoral epidermal growth factor receptor antisense DNA therapy in head and neck cancer: first human application and potential antitumor mechanisms. *Journal of Clinical Oncology*. 2009; 27(8):1235–42. [PubMed: 19204206]
52. Hammerman PS, Hayes DN, Grandis JR. Therapeutic insights from genomic studies of head and neck squamous cell carcinomas. *Cancer Discovery*. 2015; 5(3):239–44. [PubMed: 25643909]
53. Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, et al. The mutational landscape of head and neck squamous cell carcinoma. *Science*. 2011; 333(6046):1157–60. [PubMed: 21798893]

54. Agrawal N, Frederick MJ, Pickering CR, Bettegowda C, Chang K, Li RJ, et al. Exome sequencing of head and neck squamous cell carcinoma reveals inactivating mutations in NOTCH1. *Science*. 2011; 333(6046):1154–7. [PubMed: 21798897]
55. Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR, et al. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature*. 2014; 505(7484):495–501. [PubMed: 24390350]
56. Chang MT, Asthana S, Gao SP, Lee BH, Chapman JS, Kandoth C, et al. Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. *Nature Biotechnology*. 2016; 34(2):155–63.
57. Weidhaas JB, Harris J, Schae D, Chen AM, Chin R, Axelrod R, et al. The KRAS-Variant and Cetuximab Response in Head and Neck Squamous Cell Cancer: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncology*. 2016; 3(4):483–491.
58. Zdanov S, Mandapathil M, Abu Eid R, Adamson-Fadeyi S, Wilson W, Qian J, et al. Mutant KRAS Conversion of Conventional T Cells into Regulatory T Cells. *Cancer Immunology Research*. 2016; 4(4):354–65. [PubMed: 26880715]
59. Van Allen EM, Lui VW, Egloff AM, Goetz EM, Li H, Johnson JT, et al. Genomic Correlate of Exceptional Erlotinib Response in Head and Neck Squamous Cell Carcinoma. *JAMA Oncology*. 2015; 1(2):238–44. [PubMed: 26181029]
60. Wen Y, Li H, Zeng Y, Wen W, Pendleton KP, Lui VW, et al. MAPK1E322K mutation increases head and neck squamous cell carcinoma sensitivity to erlotinib through enhanced secretion of amphiregulin. *Oncotarget*. 2016; 7(17):23300–11. [PubMed: 27004400]




Figure 1. Epidermal growth factor receptor downstream signaling pathways include RAS/RAF/MEK/ MAPK/ERK, phosphatidylinositol 3-kinase(PI3K) and Akt, protein kinase C (PKC), Src, and the JAK/STAT pathways. Subsequent signaling cascades influence gene expression, proliferation, angiogenesis, apoptosis inhibition, cell motility, metastasis, adhesion, and angiogenesis.




Figure 2. EGFR Mutation Patterns

Mutation patterns in EGFR across tumor types. EGFR mutations appear recurrent and localized in lung cancer and glioblastoma multiforme, in contrast to the pattern seen in head and neck carcinomas. Missense mutations, represented by circles, are colored by degree of conservation of base pair; dark green is conserved and white is not conserved [51].

Table I

EGFR-targeted Monoclonal Antibodies

Compound	Company	Description	Approval and Clinical Indications
Cetuximab Erbitux (IMC-C225)	ImClone Systems Incorporated Bristol-Myers Squibb Eli Lilly Merck KGaA	Chimeric, murine antibody and human IgG1	2004: FDA approval for metastatic colorectal 2006: FDA approval for use in combination with XRT for locally or regionally advanced HNSCC or as monotherapy for platinum refractory, recurrent, or metastatic HNSCC 2009: FDA approval for KRAS wild type colorectal cancer 2011: FDA approval for use as first-line treatment in combination with platinum based chemotherapeutics and 5-FU for recurrent local-regional or metastatic HNSCC 2006: FDA approval for metastatic CRC
Panitumumab Vectibix (ABX-EGF)	Amgen Takeda	Humanized mAb	2007: European Medicines Agency approval for use in combination with FOLFIRI chemotherapy for metastatic colon cancer 2008: Health Canada approval for refractory EGFR-expressive metastatic CRC with wild type KRAS 2014: FDA approval in combination with FOLFOX for first line treatment of wild type KRAS CRC 2006: Approval for HNSCC in India
Nimotuzumab	YM Biosciences	Humanized mAb	2008: Approval in combination with XRT for NPC in China Phase II and III studies for cancers including HNSCC, esophageal, gastric, CRC, and gliomas
Zalutumumab Genmab	Genmab MATOS Pharma	Human IgG1	Phase I, II, and III for HNSCC, NSCLC, and CRC
Duligotuzumab	Roche	Humanized dual EGFR/HER3 mAb	Phase I and II studies in HNSCC

CRC, colorectal cancer. FDA, Food and Drug Administration. FOLFOX, a chemotherapy combination of leucovorin, fluorouracil, and oxaliplatin. HNSCC, head and neck squamous cell carcinoma. NPC, nasopharyngeal carcinoma. XRT, radiotherapy.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table II

EGFR-targeted Tyrosine Kinase Inhibitors

Compound	Company	Description	Approval and Clinical Indications
Gefitinib Iressa (ZD1839)	AstraZeneca Pharmaceuticals	Reversible binding EGFR specific Oral medicine	2003: advanced or metastatic NSCLC
Erlotinib Tarceva (OSI-774)	Genentech Astellas	Reversible binding EGFR specific	2004: locally advanced or metastatic NSCLC; approved in combination with gemcitabine for locally advanced or metastatic pancreatic cancer 2007- in combination for breast cancer patient on capecitabine
Lapatinib Tykerb	GlaxoSmithKline	Reversible binding Inhibition of HER2/neu and EGFR	2010- in combination with an aromatase inhibitor for HER2 and hormone receptor positive metastatic breast cancer
Afatinib	Boehringer Ingelheim Pharmaceuticals	Irreversible Pan-ErbB binding	2013: first-line treatment of metastatic NSCLC with EGFR exon 19 deletions or exon 21 (L858R) substitutions
Dasatinib (Sprycel)	Bristol-Myers Squibb	c-Scr kinases; thought to interfere with nuclear localization and of EGFR (Raju 2012)	2006: adult chromosome-positive chronic myelogenous leukemia (CP-CML) for which imatinib was ineffective 2010: newly diagnosed CP-CML
Dacomitinib	Pfizer	Irreversible Pan-ErbB binding	Phase I, II and III trials for cancers including HNSCC, NSCLC, and glioblastoma multiforme
ASP8273	Astellas Pharma	Irreversible binding Affinity higher for EGFR activating and T790M mutations compared to wild type	Phase I, II, and III trials in NSCLC and solid malignancies

Table III

Select Clinical Trials of EGFR Therapeutics

Lead Author (Year Published)	Phase	Study	Patient and Disease Demographics				EGFR expression (% of total patients)	Outcomes
			Number	Stage	Tumor Characteristics	Number		
Bonner et al. (2006)	III	High-dose XRT with and without cetuximab	424	III or IV	Nonmetastatic, measurable SCC Oropharynx (56%), hypopharynx (17%), larynx (27%)	79	<ul style="list-style-type: none"> - Addition of cetuximab improved median duration of local-regional control (24.4 months cetuximab with XRT vs. 14.9 months XRT alone) - Five-year OS 5-year overall survival 45.6% on cetuximab and XRT versus 36.4% with radiotherapy alone - Median OS 49.0 on cetuximab and XRT 49.0 months (95% CI 32.8–69.5) versus 29.3 months (20.6–41.4) on XRT alone - Presence of cetuximab-induced rash was associated with improved survival 	
Bonner et al. (2010)	III	High-dose XRT with and without cetuximab	424	III or IV	Nonmetastatic, measurable SCC Oropharynx (56%), hypopharynx (17%), larynx (27%)	79	<ul style="list-style-type: none"> - 103 patients started on single agent cetuximab; 53 progressed and began combination therapy - Median OS 178 days in the single-agent phase - During the single-agent phase, disease was controlled (any response or stabilized disease) in 46% of patients and time to progression 70 days - During the dual-agent phase, disease was controlled in 26% of patients and time to progression 50 days 	
Vermorken et al. (2007)	II	Update on Bonner et al. (2006) study Treatment with cetuximab and subsequent combination of cetuximab and platinum therapy in the setting of disease progression Open-label, no control arm	103	III or IV	Recurrent or metastatic SCC Pharynx (38%), larynx (20%), paranasal sinuses (3%); other (39%) Progression after a 2-6 cycles of platinum-based therapy	97	<ul style="list-style-type: none"> - Addition of cetuximab lead to an increased median OS from 7.4 months to 10.1 months - Addition of cetuximab increased the progression-free survival time from 3.3 months to 5.6 months 	
Vermorken et al. (2008)	III	Platinum-based therapy and fluorouracil with and without cetuximab	442	NR	Metastatic or local-regionally recurrent SCC Oral cavity (20%), oropharynx (34%), hypopharynx (14%), larynx (25%), other (7%)	92		

Lead Author (Year Published)	Patient and Disease Demographics					EGFR expression (% of total patients)	Outcomes
	Phase	Study	Number	Stage	Tumor Characteristics		
Ang et al. (2014)	III	Cisplatin-based C-XRT with and without cetuximab	891	III or IV	Oropharynx (70%), hypopharynx (7%), larynx (23%)	NR	<ul style="list-style-type: none"> - Cetuximab did not improve 3-year OS (72.9% without cetuximab vs. 75.8% with cetuximab; P = .32) - Addition of cetuximab led to increased radiation treatment interruptions and increased severe grade mucositis, rash, fatigue, anorexia, and hypokalemia - Patients with HPV+ tumor had improved survival with 3-year OS 85.6% vs. with HPV- tumors 60.1% (p < 0.001)
Weidhass et al. (2016)	III	Cisplatin-based C-XRT with and without cetuximab Patients subcategorized by KRAS mutation	413	III or IV	Oropharynx (72%), hypopharynx/larynx (28%) 17% KRAS-variant (70/413)	NR	<ul style="list-style-type: none"> - Cetuximab improved one- and two-year OS with in patients with KRAS-variant (HR, 0.19; 95% CI, 0.04-0.86; p = 0.03)
Mesia et al. (2015)	II	Cisplatin-based C-XRT compared to a dose-reduced cisplatin-based C-XRT with panitumumab	150	III or IV	Oral cavity (9%), oropharynx (53%), hypopharynx (19%), larynx (18%) Locally advanced SCC	NR	<ul style="list-style-type: none"> - Panitumumab did not improve two-year local-regional control (68% without vs. 61% with panitumumab) - Addition of panitumumab was associated with increased rates of grade 3-4 mucosal inflammation, dysphagia, and radiation-related skin toxicity
Fayette et al. (2016)	II	Duligotuzumab compared to cetuximab following progressing on/after cisplatin-based chemotherapy	121	III or IV	Recurrent or metastatic SCC Oral cavity (29%), oropharynx (30%), hypopharynx (10%), larynx (16%), unspecified (10%), unknown (6%)	NR	<ul style="list-style-type: none"> - OS was statistically similar between duligotuzumab (7.2 months) compared to cetuximab (8.7 months; HR 1.15, 90% CI 0.81-1.63) - Expression level of neuregulin 1 (NRG1, ligand to HER3) nor ERBB3 expression (encodes HER3) did not influence response lai
Martins et al. (2013)	II	Cisplatin and XRT with and without erlotinib Randomized	204	III or IV	Locally advanced SCC Oral cavity (7%), oropharynx (67%), hypopharynx (6%), larynx (18%), nasopharynx (1%), other (1%)	4/90 samples assessed had EGFR amplification	<ul style="list-style-type: none"> - Addition of erlotinib did not increase toxicity - The TKI erlotinib did not confer additional tumor response or survival

Lead Author (Year Published)	Patient and Disease Demographics					Outcomes
	Phase	Study	Number	Stage	Tumor Characteristics	
Argiris et al. (2013)	III	Docetaxel with or without gefitinib	270	NR	Recurrent or metastatic SCC	<ul style="list-style-type: none"> - The TKI gefitinib did not lead to improved survival or outcomes
		Randomized			Oral cavity (22%), oropharynx (33%), larynx (26%), multiple (5%), other (14%)	
Kim et al. (2015)	II	Dacomitinib monotherapy	48	NR	Local-regionally recurrent or metastatic SCC	<ul style="list-style-type: none"> - 20.8% (10) of patients with partial response and 65% (31) of patients with stable disease - OS 6.6 months and PFS 3.9 months - in the cohort, the patients with PI3K pathway mutations
Machiels et al. (2015)	III	Afatinib or methotrexate as a second-line therapy following prior platinum-based therapy and disease progression	483	NR	Oral cavity (37%), oropharynx (23%), hypopharynx (17%), larynx (19%), maxillary sinus (4%) Recurrent or metastatic SCC Progression after or on platinum-based therapy	<ul style="list-style-type: none"> - PFS improved with afatinib (median 2.6 months) compared to methotrexate (median 1.7 months), hazard ratio 0.80 (95% CI 0.65-0.98, p=0.03) - Of note, 59% of patients were previously treated with EGFR-targeted therapy
Harrington et al. (2015)	III	Adjuvant C-XRT with lapatinib or placebo followed by 1 year of lapatinib or placebo	688	II, III, IVA	Surgical margin <5mm or ECE Oral cavity (41%), oropharynx (19%), hypopharynx (13%), larynx (23%), multiple sites (4%)	<ul style="list-style-type: none"> - Addition of lapatinib did not improve overall survival (HR 0.96, 95% CI 0.73 to 1.25) nor disease free survival (HR 1.10, 0.85 to 1.43) - Lapatinib was associated with increased grade 3-4 adverse events (75%) compared to placebo (67%, p=0.019)
Soulières et al. (2017)	II	Buparlisib, oral pan-PI3K inhibitor, or placebo with paclitaxel as second-line therapy after progression with platinum-based treatment	158	NR	Recurrent or metastatic SCC Progression after or on platinum-based therapy Oral cavity (29%), oropharynx (28%), hypopharynx (18%), larynx (16%), nasopharynx (3%), other/unknown (6%)	<ul style="list-style-type: none"> - Median PFS was improved with second-line buparlisib and paclitaxel (4.6 months) compared to placebo and paclitaxel (3.5 months; HR 0.65 [95% CI 0.45-0.95]) - Of note, 46% of patients were previously treated with EGFR-targeted therapy
BERIL-1						

C-XRT, chemoradiotherapy. ECE, extracapsular extension. HR, hazard ratio. NR, not recorded. OS, overall survival. PFS, progression-free survival. SCC, squamous cell carcinoma. XRT, radiotherapy.

Table IV

Frequency of mutations in commonly deregulated head and neck cancer pathways

Cell Cycle		RTK/RAS/PI(3)K		Cell Death		Differentiation	
Gene	HPV +	Gene	HPV -	Gene	HPV -	Gene	HPV +
Pathway	Red	Pathway	Red	Pathway	Red	Pathway	Red
	Red		Red		Red		Red
	Blue		Blue		Blue		Blue
	Red		Red		Red		Red
	Blue		Blue		Blue		Blue
	Red		Red		Red		Red
	Blue		Blue		Blue		Blue
	Red		Red		Red		Red
	Blue		Blue		Blue		Blue