

SPIRITUALITY AND SPIRITUAL WELLBEING

The World Health Organization in 1948 defined health as a complete sense of physical, mental, and social wellbeing. In 1998 the definition was revised to include the importance of spiritual wellbeing and the connection between mind and body in the healing process. The word spirit comes from the Latin *'spiritus'* meaning breath, and spirituality has been described as a search for existential meaning regarding a power other than the self that is not necessarily called 'God'. This search for meaning includes a need for identity and relatedness, and is sometimes expressed as a need to feel loved or accepted and of having a sense of self-worth. Spirituality is about acceptance, integration, and wholeness, and is the essence of what it means to be human.

SPIRITUAL CARE

Spiritual care can be defined as care that recognises and responds to the needs of the human spirit when faced with trauma, ill health, or sadness. It includes the need for meaning and self-worth, to express oneself, for faith support or simply for a sensitive listener. It is usually given in a one-to-one relationship, is person centred, and makes no assumptions about personal conviction or life orientation.

SPIRITUAL NEEDS ASSESSMENT AND ASSESSMENT TOOLS

Rumbold in a review of spiritual needs assessment suggested that it should include: respect for the patients' perspectives and privacy, involvement of all members of the interdisciplinary team, clear documentation of needs and strategic responses to these needs, integration of strategies into an overall care plan, provision of a shared framework for continuity of care between community agencies and inpatient services, and not conflating spiritual issues with religious practice.¹

Many assessment models or tools have been proposed. These are mainly for use by healthcare professionals and chaplains, and one of the most commonly used is the F (Faith), I (Importance), C (Community), and A (Address in care or action) or FICA/HOPE tool, which was developed by a group of primary care physicians in the US and validated in the City of Hope Medical Centre

"Spiritual care ... recognises and responds to the needs of the human spirit when faced with trauma, ill health, or sadness."

by the George Washington Institute for Spirituality and Health. The FICA tool allows doctors to assess the severity of spiritual distress, identify a patient's spiritual resources of strength, and integrate that information into the clinical treatment plan.² It is a model that could be readily adapted for use in UK general practice.

ARGUMENTS AGAINST USING ASSESSMENT TOOLS

Arguments against using tools to assess spiritual needs include the view that the vocabulary of the spirit belongs to a language of depth and meaning. It is argued that this unfolds in the context of relationship rather than by dispassionate analysis, and that the best way of obtaining information about spiritual needs is by listening to patients' stories and encouraging them to articulate their concerns. Spiritual care is not a set of prescribed and proscribed roles but a series of highly fluid interpersonal processes in the context of mutually recognised human values and experiences. Sight, speech, touch, and presence are the ways in which healthcare professionals impact on patients' spiritual wellbeing regardless of their awareness or intent.

RESULTS OF SPIRITUAL INTERVENTIONS

Several studies have demonstrated positive outcomes from spiritual interventions. Patients with type 2 diabetes and/or cardiovascular disease have been shown to incorporate spirituality into their self-management routines with a positive effect on their health and wellbeing,³ and patients receiving talking therapy from a chaplain

have been shown to experience similar improvements in wellbeing as patients who were prescribed antidepressants.⁴ Good spiritual care of patients who are terminally ill has also been shown to be associated with less aggressive end-of-life care and greater quality of life near death.⁵

Results of systematic literature reviews have been less convincing. A review of randomised controlled trials of faith-based psychological therapies for anxiety and depression found that the trials were typically small and susceptible to significant bias.⁶ Another review investigating spirituality as a complementary treatment in mental health care concluded that, due to the diversity of protocols and outcomes, and lack of standardisation of interventions, it was not possible to give a definite answer one way or the other.⁷ A review of five randomised controlled trials investigated if the presence of a spiritual care counsellor in palliative care teams helped patients feel emotionally supported. However, the results were inconclusive.⁸

WHY ARE GPs AND OTHER HEALTHCARE PROFESSIONALS RELUCTANT TO PROVIDE SPIRITUAL CARE?

In a qualitative evidence synthesis Vermandere *et al* concluded that most GPs saw it as their role to identify and assess patients' spiritual needs. However, they struggled with spiritual language and experienced feelings of discomfort and fear that patients would refuse to engage in the discussion.⁹ Although nurses have traditionally provided spiritual care, reservations have been expressed

"GPs ... struggled with spiritual language and experienced feelings of discomfort and fear that patients would refuse to engage in the discussion."

“... healthcare professionals should ... include spiritual beliefs when discussing care with dying patients and those close to them.”

regarding their involvement in spiritual assessment.

A recent hospital-based study suggested there was strong agreement among clinical and non-clinical staff on the importance of delivering spiritual care but uncertainty in the ability to recognise and meet the spiritual needs of patients.¹⁰ In a qualitative analysis healthcare professionals had difficulty in formulating descriptions of spiritual care, which was in marked contrast to the importance they attached to this aspect of holistic care. The reasons given for being reluctant to provide spiritual care included: lack of available time, lack of training and expertise, and a sense that others could do a better job.¹¹

DO CHAPLAINS HAVE A ROLE IN GENERAL PRACTICE?

Chaplains act as spiritual care providers, emotion facilitators, grief counsellors, and cultural and religious experts. They are increasingly becoming involved in spiritual care in the community and in general practice. In a pilot study community clergy who received training with healthcare professionals and chaplains enhanced the quality of care they provided to patients dying at home who wished to receive spiritual support¹² and, in a previously mentioned study, it was suggested that primary care chaplaincy could be considered as an alternative to cognitive behavioural therapy.⁴ Perhaps in future the GP will be regarded as the generalist in spiritual care who refers appropriate patients to the chaplain, who is the specialist in spiritual issues.

RECENT NICE GUIDANCE ON SPIRITUAL CARE

In 2014, an audit of end-of-life care by the Royal College of Physicians reported that nearly half of all deaths in England occurred in hospital and that spiritual issues were documented in only one in seven patients.¹³ The National Institute for Health and Care Excellence¹⁴ has issued new guidance on the spiritual care of dying patients, which states that only by attending to a patient's spiritual, cultural, religious, and social

ADDRESS FOR CORRESPONDENCE

Ian J Hamilton

Institute of Health and Wellbeing, University of Glasgow, 1 Lilybank Gardens, Glasgow, G12 8RZ, UK.

E-mail: ijdhamilton@doctors.org.uk

preferences can truly individualised care be delivered, and all healthcare professionals should now include spiritual beliefs when discussing care with dying patients and those close to them.

CONCLUSION

Spiritual care is not religious (although religious care that is given in the context of the shared religious beliefs, values, liturgies, and lifestyle of a faith community may be spiritual) and it is an important aspect of holistic patient care. There are two challenges: first, developing instruments that will more accurately assess the effectiveness of spiritual interventions⁹ and, second, persuading GPs and other healthcare professionals of the importance of providing good spiritual care.

Ian J Hamilton,

Researcher, Institute of Health and Wellbeing, University of Glasgow, Glasgow.

Jillian Morrison,

Professor, Institute of Health and Wellbeing, University of Glasgow, Glasgow.

Sara Macdonald,

Lecturer, Institute of Health and Wellbeing, University of Glasgow, Glasgow.

Ethical approval

It was not necessary to obtain approval for the study and no ethics committee approval was sought.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

DOI: <https://doi.org/10.3399/bjgp17X693845>

REFERENCES

1. Rumbold BD. A review of spiritual assessment in health care practice. *Med J Aust* 2007; **186**(10 Suppl): S60–S62.
2. Puchalski CM. Formal and informal spiritual assessment. *Asian Pac J Cancer Prev* 2010; **11**(Suppl 1): 51–57.
3. Unantenne N, Warren N, Canaway R, Manderson L. The strength to cope: spirituality and faith in chronic disease. *J Relig Health* 2013; **52**(4): 1147–1161.
4. Macdonald G. Primary care chaplaincy: a valid talking therapy? *Br J Gen Pract* 2017; DOI: 10.3399/bjgp17X689221.
5. Balboni TA, Balboni MJ, Enzinger AC, et al. Provision of spiritual support to patients with advanced cancer by religious communities and associations with medical care at the end of life. *JAMA Intern Med* 2013; **173**(12): 1109–1117.
6. Anderson N, Heywood-Everett S, Siddiqi N, et al. Faith-adapted psychological therapies for depression and anxiety: systematic review and meta-analysis. *J Affect Disord* 2015; **176**: 183–196.
7. Anderson N, Heywood-Everett S, Siddiqi N, et al. Religious and spiritual interventions in mental health care: a systematic review and meta-analysis of randomised controlled clinical trials. *Psychol Med* 2015; **45**(14): 2937–2949.
8. Candy B, Jones L, Varagunam M, et al. Spiritual and religious interventions for well-being of adults in the terminal phase of disease. *Cochrane Database Syst Rev* 2012; **5**: CD007544. DOI: 10.1002/14651858.CD007544.pub 2.
9. Vermandere M, De Lepeleire J, Smeets L, et al. Spirituality in general practice: a qualitative evidence synthesis. *Br J Gen Pract* 2011; DOI: 10.3999/bjgp11X606663.
10. Austin PD, Macleod R, Siddall PJ, et al. The ability of hospital staff to recognise and meet patients' spiritual needs: a pilot study. *Journal for the Study of Spirituality* 2016; **6**(1): 20–37. <http://dx.doi.org/10.1080/20440243.2016.1158453> [accessed 20 Sep 2017].
11. Selby D, Seccaraccia D, Huth J, et al. A qualitative analysis of a healthcare professional's understanding and approach to management of spiritual distress in an acute care setting. *J Palliat Med* 2016; **19**(11): 1197–1204.
12. Goodhead A, Speck P, Selman L. 'I think you just learnt as you went along' — community clergy's experience of and attitudes towards caring for dying people: a pilot study. *Palliat Med* 2016; **30**(7): 674–683.
13. Royal College of Physicians. *National Care of the Dying Audit of Hospitals*. <https://www.rcplondon.ac.uk/projects/outputs/national-care-dying-audit-hospitals> [accessed 25 Oct 2017].
14. National Institute of Health and Care Excellence. *Care of dying adults in the last days of life. NICE guideline [NG31]*. 2015. <https://www.nice.org.uk/guidance/ng31> [accessed 25 Oct 2017].