



Published in final edited form as:

*Science*. 2017 September 01; 357(6354): 912–916. doi:10.1126/science.aan0677.

## The intestinal microbiota regulates body composition through NFIL3 and the circadian clock

Yuhao Wang<sup>1</sup>, Zheng Kuang<sup>1</sup>, Xiaofei Yu<sup>1, #</sup>, Kelly A. Ruhn<sup>1</sup>, Masato Kubo<sup>2, 3</sup>, and Lora V. Hooper<sup>1, 4, \*</sup>

<sup>1</sup>Department of Immunology, The University of Texas Southwestern Medical Center, Dallas, TX 75390

<sup>2</sup>Laboratory for Cytokine Regulation, RIKEN Center for Integrative Medical Sciences (IMS), RIKEN Yokohama Institute, Suehiro-cho 1-7-22, Tsurumi, Yokohama, Kanagawa 230-0045, Japan

<sup>3</sup>Division of Molecular Pathology, Research Institute for Biomedical Science, Tokyo University of Science, 2669 Yamazaki, Noda-shi, Chiba, 278-0022, Japan

<sup>4</sup>The Howard Hughes Medical Institute, The University of Texas Southwestern Medical Center, Dallas, TX 75390

### Abstract

The intestinal microbiota has been identified as an environmental factor that markedly impacts energy storage and body fat accumulation, yet the underlying mechanisms remain unclear. Here we show that the microbiota regulates body composition through the circadian transcription factor NFIL3. *Nfil3* transcription oscillates diurnally in intestinal epithelial cells and the amplitude of the circadian oscillation is controlled by the microbiota through group 3 innate lymphoid cells (ILC3), STAT3, and the epithelial cell circadian clock. NFIL3 controls expression of a circadian lipid metabolic program and regulates lipid absorption and export in intestinal epithelial cells. These findings provide mechanistic insight into how the intestinal microbiota regulates body composition and establish NFIL3 as an essential molecular link among the microbiota, the circadian clock, and host metabolism.

---

The worldwide obesity epidemic presents a pressing public health crisis. More than 2.1 billion individuals are overweight or obese throughout the world, and ~3.4 million deaths are caused each year by obesity-related disease (1). Consequently, there is an urgent need to identify host and environmental factors that regulate human metabolism and energy homeostasis.

---

\*To whom correspondence should be addressed: lora.hooper@utsouthwestern.edu.

#Current address: Laboratory of Molecular Genetics, The Rockefeller University, New York, NY 10065

Supporting Online Material

Materials and Methods

Table S1

Figures S1 to S12

References 32-37

The intestinal microbiota is an environmental factor that markedly impacts mammalian body composition. The microbiota promotes energy storage in adipose tissue, and thus microbiologically sterile (germ-free) mice have less body fat relative to conventionally-raised mice (2). This is because in part the microbiota enhance energy harvest from the host diet (3) and promote storage of that energy in adipose tissue (2). Less is known about how microbial regulation of host metabolic pathways might also impact energy storage and body composition.

Many host metabolic pathways are synchronized with day-night light cycles through the circadian clock. The mammalian circadian clock is a network of transcription factors, present in all cells of the body, that drives rhythmic ~24-hour oscillations in gene expression. Synchronization of metabolism with the clock couples energetically expensive metabolic pathways to the availability of dietary substrates, optimizing energy utilization. Emerging evidence indicates that the microbiota interacts with the circadian clock in ways that profoundly impact host metabolism, and that disrupting these interactions can lead to obesity and other metabolic diseases (4, 5). However, little is known about the mechanisms that govern microbiota interactions with the circadian clock and how these interactions alter host metabolism.

NFIL3, also known as E4BP4, is a basic leucine zipper transcription factor that is expressed in a variety of immune cells and controls immune functions that vary by cell type (6, 7). We found that small intestinal epithelial cells also expressed NFIL3, and that expression was markedly reduced in germ-free mice (Fig. 1A). This accorded with prior findings in antibiotic-treated mice (8), and suggested that epithelial NFIL3 might regulate a physiological activity that is responsive to the intestinal microbiota.

To identify the physiological functions of NFIL3 in intestinal epithelial cells, we generated an epithelial cell-specific *Nfil3* knockout mouse (*Nfil3*<sup>IEC</sup>). *Nfil3*<sup>IEC</sup> mice raised on a chow diet weighed less than their *Nfil3*<sup>fl/fl</sup> littermates (Fig. 1B, fig. S1A) and had reduced body fat and increased lean body mass relative to *Nfil3*<sup>fl/fl</sup> littermates (fig. S1B,C). The body composition differences were not due to off-target effects of CRE expression (fig. S1D,E), differences in food intake (fig. S2A), physical activity (fig. S2B), or energy utilization (fig. S3A–C), which were similar between the two groups. We also did not detect differences in the response to intestinal injury (fig. S4A–D) or in the expression of key inflammatory cytokines and antimicrobial proteins (fig. S5A–E).

When placed on a high fat, Western-style diet (HFD) for 10 weeks, both *Nfil3*<sup>fl/fl</sup> and *Nfil3*<sup>IEC</sup> mice gained weight (Fig. 1B). However, the *Nfil3*<sup>IEC</sup> mice maintained lower body weights (Fig. 1B) and had lower body fat percentages (Fig. 1C) and higher lean body mass percentages (fig. S6A). The *Nfil3*<sup>IEC</sup> mice also had lower epididymal fat pad weights (Fig. 1D, fig. S6B), and were protected from elevated blood triglycerides (Fig. 1E), liver fat accumulation (Fig. 1F), lowered glucose tolerance (Fig. 1G), and increased insulin resistance (Fig. 1G). 16S ribosomal RNA sequencing of fecal microbiotas indicated that the metabolic differences between *Nfil3*<sup>fl/fl</sup> and *Nfil3*<sup>IEC</sup> mice were not due to differences in microbiota taxonomic composition (fig. S7). Thus, epithelial NFIL3 regulates lipid storage and body composition in mice.

Because epithelial NFIL3 expression is microbiota-dependent we sought to determine if NFIL3-dependent body fat accumulation also depends on the microbiota. Depletion of the microbiota through antibiotic treatment produced lower fat and higher lean body mass in the *Nfil3<sup>fl/fl</sup>* mice, resulting in body compositions that were not significantly different from those of *Nfil3<sup>IEC</sup>* mice (Fig. 1H, fig. S8). Thus, high fat diet-induced body fat accumulation requires both NFIL3 and a microbiota.

We next studied the mechanism by which the microbiota regulates NFIL3 expression. *Nfil3* transcription is controlled by the circadian clock (9) and accordingly, *Nfil3* transcript abundance oscillated diurnally in an enriched population of small intestinal epithelial cells acquired by laser capture microdissection (Fig. 2A). Comparison of *Nfil3* expression levels across a circadian cycle in wild-type germ-free and conventional mice revealed that the microbiota is required for maximal *Nfil3* expression and thus governs the amplitude of *Nfil3* transcriptional rhythms (Fig. 2A). This was reflected in protein expression levels (Fig. 2B) and accords with prior findings in antibiotic-treated mice (8).

*Nfil3* expression is directly regulated by the core circadian clock transcriptional repressor REV-ERB $\alpha$ . In T cells and liver cells, REV-ERB $\alpha$  binds to a consensus sequence in the *Nfil3* gene locus and represses transcription, resulting in a rhythmic diurnal *Nfil3* expression pattern (6, 10). *Rev-erba* transcript and protein abundance also oscillated diurnally in intestinal epithelial cells but was higher in germ-free than in conventional mice (Fig. 2C,D). This suggested that REV-ERB $\alpha$  governs circadian rhythmicity in *Nfil3* expression, and that the microbiota might induce *Nfil3* expression by repressing *Rev-erba* expression. Indeed, REV-ERB $\alpha$  bound directly to the *Nfil3* promoter in intestinal epithelial cells as assessed by chromatin immunoprecipitation (ChIP) assay (Fig. 2E). Further, *Nfil3* expression in antibiotic-treated *Rev-erba<sup>-/-</sup>* mice was similar to that in conventional wild-type mice (Fig. 2F). Thus, microbiota regulation of *Nfil3* expression is REV-ERB $\alpha$ -dependent, and the microbiota elevates *Nfil3* expression by repressing *Rev-erba* expression.

We next sought to determine how bacterial signals are relayed to the epithelial circadian clock to regulate *Rev-erba* and *Nfil3* expression. Intestinal epithelial cells sense the microbiota through Toll-like receptors (TLR) and their common signaling adaptor MyD88 to regulate expression of key genes (11). We therefore tested if MyD88 is required for microbiota regulation of *Rev-erba* and *Nfil3* expression in epithelial cells. We quantified *Rev-erba* and *Nfil3* transcripts in epithelial cells from *Myd88<sup>-/-</sup>* mice and wild-type littermates at Zeitgeber time (ZT4), which is when *Nfil3* expression is near-peak in conventional mice (Fig. 2A). Epithelial *Rev-erba* expression was increased and *Nfil3* expression was consequently reduced to germ-free levels in *Myd88<sup>-/-</sup>* mice (Fig. 2G), indicating that microbiota regulation of the *Rev-erba*-*Nfil3* cascade requires MyD88. While epithelial MyD88 was dispensable for microbiota-induced alterations in *Rev-erba* and *Nfil3* expression, *Myd88* expression in a CD11c<sup>+</sup> cell population (which includes some dendritic cells) was required for maximal repression of *Rev-erba* and induction of *Nfil3* (Fig. 2G), suggesting a requirement for dendritic cells (DCs).

To further test for DC involvement, we used a mouse model of DC depletion in which *Diphtheria* toxin receptor (DTR) is expressed under the control of the *Cd11c* promoter (12).

On administration of *Diphtheria* toxin (DT), CD11c<sup>+</sup> cells are selectively killed (12). On CD11c<sup>+</sup> cell depletion, *Rev-erba* expression was increased and *Nfil3* expression was decreased (Fig. 2H), supporting a requirement for DCs in microbiota induction of *Nfil3*.

Previous studies have identified a subepithelial cellular signaling relay in the small intestine that captures microbiota signals and passes them to epithelial cells to alter expression of key epithelial cell genes (13, 14). In this circuit, bacteria activate TLR-MyD88 signaling in DCs, and the bacterial signals are relayed from DC to group 3 innate lymphoid cells (ILC3) through the cytokine interleukin-23 (IL-23). ILC3 then signal to the epithelium through the production of IL-22 (15).

To determine if ILCs were required for microbiota-induced *Nfil3* expression we studied ID2-deficient mice (*Id2<sup>gfp/gfp</sup>*), which lack all known ILC subsets (16, 17). *Id2<sup>gfp/gfp</sup>* mice showed increased *Rev-erba* expression and consequent decreased *Nfil3* expression in the small intestinal epithelium (Fig. 2H), consistent with a requirement for ILCs in regulating *Rev-erba* expression. *Rag1<sup>-/-</sup>* mice, which lack T and B cells, showed decreased epithelial *Rev-erba* expression and increased *Nfil3* expression (Fig. 2H). The fact that *Nfil3* expression was higher in *Rag1<sup>-/-</sup>* mice than in conventional wild-type mice is likely due to the elevated bacterial loads as well as aberrant expansion of ILC3 in the small intestines of these mice (18), and indicates that T and B cells are not required for induction of *Nfil3* expression. ROR $\gamma$ t-deficient (*Rorc<sup>gfp/gfp</sup>*) mice (19), which lack both TH17 cells and ILC3, showed circadian *Rev-erba* and *Nfil3* expression patterns (Fig. 2I,J) that were similar to those of germ-free mice (Fig. 2A,B). This establishes that ILC3 are required for microbiota induction of *Nfil3* expression through *Rev-erba*. Treatment of *Myd88<sup>-/-</sup>* mice with recombinant IL-23 or IL-22 restored epithelial *Rev-erba* and *Nfil3* expression to wild-type conventional levels (Fig. 2K), further supporting the idea that the subepithelial DC-ILC3 circuit relays microbiota signals to epithelial cells to regulate *Nfil3* expression.

The intestinal DC-ILC3 circuit can be triggered by flagellin or lipopolysaccharide (LPS), which is present in the outer membranes of Gram-negative bacteria (20,21). Accordingly, treatment of germ-free mice with flagellin and LPS decreased *Rev-erba* expression and increased *Nfil3* expression (fig. S9A,B). Further, *Rev-erba* expression decreased and *Nfil3* expression increased when we monoassociated germ-free mice with Gram-negative, flagellated bacterial species, including *Salmonella typhimurium* and *Escherichia coli* (fig. S9A,B). *Rev-erba* and *Nfil3* expression were not markedly altered by monoassociation with the Gram-positive species *Enterococcus faecalis* or with the Gram-negative non-flagellated species *Bacteroides thetaiotaomicron*, suggesting that *Nfil3* expression is selectively activated by Gram-negative, motile bacteria. This is likely because such bacteria produce both flagellin and LPS, and because they can readily penetrate the intestinal epithelial barrier and contact lamina propria DCs.

We next sought to identify epithelial cell-intrinsic pathways downstream of IL-22 that regulate *Rev-erba* and *Nfil3* expression. The transcription factor STAT3 is a key response element downstream of the IL-22 receptor (IL-22R)(14). Activation of the IL-22R leads to phosphorylation of STAT3, which then binds to the promoters of its target genes and either activates or inhibits their transcription. Indeed, ChIP analysis of intestinal epithelial cells

showed STAT3 binding to the *Rev-erba* promoter (Fig. 3A,B). STAT3 binding was markedly reduced in epithelial cells from germ-free mice (Fig. 3A,B). Co-transfection of STAT3-encoding vectors with luciferase reporters fused to the *Rev-erba* promoter resulted in decreased luciferase activity in HEK-293T cells (Fig. 3C), and expression of a dominant active form of STAT3 further inhibited the luciferase activity (Fig. 3C). Thus, STAT3 binds to the *Rev-erba* promoter and inhibits its transcription.

We further tested the role of STAT3 in repressing *Rev-erba* transcription using cultured intestinal organoids. Addition of recombinant IL-22 to organoid cultures resulted in STAT3 phosphorylation (Fig. 3D), supporting prior findings that IL-22 activates STAT3 in intestinal epithelial cells (14). At the same time, expression of *Rev-erba* was decreased and expression of *Nfil3* was increased, compared to controls (Fig. 3E,F). In contrast, when we added a STAT3 phosphorylation inhibitor (Stattic) (22) to organoids together with recombinant IL-22, STAT3 phosphorylation was inhibited (Fig. 3D), and *Rev-erba* and *Nfil3* showed expression levels similar to those of the controls (Fig. 3E,F). These findings further support the idea that STAT3 is a transcriptional repressor of *Rev-erba*.

We next investigated whether STAT3 is required for microbiota repression of epithelial *Rev-erba* expression *in vivo*. We generated a mouse with an epithelial cell-specific deletion of *Stat3* (*Stat3*<sup>IEC</sup>) and assayed for *Rev-erba* and *Nfil3* expression in intestinal epithelial cells. Consistent with our *in vitro* findings, the *Stat3*<sup>IEC</sup> mice showed increased expression of epithelial *Rev-erba* and decreased expression of *Nfil3* as compared to their *Stat3*<sup>fl/fl</sup> littermates (Fig. 3G,H). The expression of *Stat3* and the activation of STAT3 in intestinal epithelial cells did not exhibit diurnal rhythms (fig. S10A,B), indicating that STAT3 does not generate the rhythmicity in *Rev-erba* and *Nfil3* expression. Instead, our findings suggest that diurnal rhythms in NFIL3 expression are generated by the circadian clock through REV-ERBA, while the amplitude of these rhythms is fine-tuned by the microbiota through STAT3.

To understand the mechanism by which epithelial NFIL3 regulates fat storage and body composition, we compared the transcriptomes of epithelial cells from *Nfil3*<sup>fl/fl</sup> and *Nfil3*<sup>IEC</sup> mice. We performed an RNAseq analysis on intestinal epithelial cells at multiple time points across the 24-hour day-night light cycle. We identified 33 transcripts having differential abundances in *Nfil3*<sup>fl/fl</sup> and *Nfil3*<sup>IEC</sup> mice, noting that expression of a number of these genes was diurnally rhythmic in *Nfil3*<sup>fl/fl</sup> mice (Fig. 4A). Rhythmic expression of the clock genes *Bmal1* (*Arnt1*), *Per2*, and *Nr1d1* (*Rev-erba*) was maintained in *Nfil3*<sup>IEC</sup> mice (Fig. 4A), indicating that the core clock mechanism remains intact. Seventeen of the transcripts encoded proteins that are known to function in lipid uptake and metabolism (Fig. 4A). These included *Cd36*, encoding a transporter that imports dietary fatty acids into cells (23); *Scd1*, encoding a stearyl-CoA-desaturase 1 (24); *Cyp2e1*, encoding a fatty acid hydroxylase (25); and *Fabp4*, encoding a fatty acid binding protein (26). Deletion of each of these genes protects against HFD-induced obesity and/or insulin-resistance (24–27), suggesting that their lowered expression in *Nfil3*<sup>IEC</sup> mice could account in part for the metabolic phenotypes of these mice.

We further analyzed *Cd36* and *Scd1* transcripts by qRT-PCR, confirming that expression of both genes required epithelial NFIL3 and the microbiota (Fig. 4B). Western blot analysis

confirmed that CD36 protein levels were reduced in germ-free and *Nfil3*<sup>IEC</sup> mice (Fig. 4C). Additionally, expression of *Cd36* and *Scd1* was reduced in *Id2*<sup>gfp/gfp</sup> and *Stat3*<sup>IEC</sup> mice (Fig. 4D,E), consistent with microbiota regulation of *Nfil3* expression through ILC and epithelial STAT3. Thus, the microbiota regulates an NFIL3-dependent lipid metabolic program that is intrinsic to intestinal epithelial cells.

These findings accord with the key functions of intestinal epithelial cells in fatty acid uptake and metabolism. Following absorption by enterocytes, fatty acids are processed and packaged into chylomicrons for export into the circulation. These processes show circadian rhythmicity, resulting in diurnal variations in circulating lipids (28), and disruption of rhythms in chylomicron export, as seen in mice lacking Nocturnin, is also associated with a lean phenotype (29). Thus, we hypothesized that the metabolic phenotypes of *Nfil3*<sup>IEC</sup> mice might arise from reduced epithelial cell uptake and processing of dietary lipids with consequent lowered export of lipids to the circulation for storage in adipose tissue.

We tested this idea by visualizing lipid stores in small intestinal tissues from HFD-fed *Nfil3*<sup>IEC</sup> and *Nfil3*<sup>fl/fl</sup> mice. Lipids were detected with oil red O staining of small intestinal sections, revealing that the intestinal epithelial cells of *Nfil3*<sup>fl/fl</sup> mice harbor abundant lipids (Fig. 4F). By contrast, there was lowered detection of lipids in intestinal epithelial cells from *Nfil3*<sup>IEC</sup> mice (Fig. 4F). *Nfil3*<sup>IEC</sup> mice also showed reduced oil red O staining in subepithelial intestinal tissues, suggesting reduced export of lipids into the lymphatic capillaries that transport packaged lipids to the bloodstream (fig. S11). Consistent with these findings, lipid concentrations were also lower in intestinal epithelial cells (Fig. 4G) but higher in feces (Fig. 4H) from *Nfil3*<sup>IEC</sup> mice as compared to *Nfil3*<sup>fl/fl</sup> mice. Thus, epithelial NFIL3 regulates lipid absorption and export in intestinal epithelial cells, potentially explaining why *Nfil3*<sup>IEC</sup> mice have a limited body fat gain on a high fat diet.

Here we have shown that NFIL3 is an essential molecular link among the microbiota, the circadian clock, and host metabolism. Our findings show that the microbiota acts through NFIL3 to regulate lipid uptake and storage, thus providing new insight into how the intestinal microbiota regulates host metabolism and body composition. Further, we have shown that the ILC3-STAT3 signaling relay forms an essential conduit between the microbiota and the epithelial circadian clock, thus identifying key molecular circuitry through which the microbiota interacts with the clock (fig. S12). These results potentially provide a deeper understanding of why perturbing microbiota-clock interactions can lead to metabolic disease (4, 5). Our studies could also help to explain why circadian clock disruptions in humans, arising from shift work or international travel, are associated with an increased occurrence of metabolic diseases including obesity, diabetes, and cardiovascular disease (30, 31). Ultimately, our findings could lead to new strategies for treating metabolic disease by targeting NFIL3, STAT3, the microbiota, or the circadian clock.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank C.L. Behrendt-Boyd, Tess Leal, and Brian Hassell for assistance with mouse experiments and other members of the Hooper laboratory for fruitful discussions and critical reading of the manuscript. This work was supported by NIH R01 DK070855 (L.V.H.), a Burroughs Wellcome Foundation Investigators in the Pathogenesis of Infectious Diseases Award (L.V.H.), the Welch Foundation (L.V.H.) and the Howard Hughes Medical Institute (L.V.H.). All data and code to understand and assess the conclusions of this research are available in the main text, supplementary materials and via the GEO repository with accession number GSE100339.

## References

1. Ng M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014; 384:766–781. [PubMed: 24880830]
2. Bäckhed F, et al. The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA*. 2004; 101:15718–15723. [PubMed: 15505215]
3. Turnbaugh PJ, et al. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*. 2006; 444:1027–131. [PubMed: 17183312]
4. Thaïss CA, et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell*. 2014; 159:514–529. [PubMed: 25417104]
5. Leone V, et al. Effects of diurnal variation of gut microbes and high-fat feeding on host circadian clock function and metabolism. *Cell Host Microbe*. 2015; 17:681–689. [PubMed: 25891358]
6. Yu X, et al. T<sub>H</sub>17 cell differentiation is regulated by the circadian clock. *Science*. 2013; 342:727–730. [PubMed: 24202171]
7. Yu X, et al. The basic leucine zipper transcription factor NFIL3 directs the development of a common innate lymphoid cell precursor. *eLife*. 2014; 3:e04406.
8. Mukherji A, Kobiita A, Ye T, Chambon P. Homeostasis in intestinal epithelium is orchestrated by the circadian clock and microbiota cues transduced by TLRs. *Cell*. 2013; 153:812–827. [PubMed: 23663780]
9. Mitsui S, Yamaguchi S, Matsuo T, Ishida Y, Okamura H. Antagonistic role of E4BP4 and PAR proteins in the circadian oscillatory mechanism. *Genes Dev*. 2001; 15:995–1006. [PubMed: 11316793]
10. Duez H, et al. Regulation of bile acid synthesis by the nuclear receptor Rev-erba. *Gastroenterology*. 2008; 135:689–698. [PubMed: 18565334]
11. Vaishnava S, et al. The antibacterial lectin RegIII $\gamma$  promotes the spatial segregation of microbiota and host in the intestine. *Science*. 2011; 334:255–258. [PubMed: 21998396]
12. Jung S, et al. In vivo depletion of CD11c<sup>+</sup> dendritic cells abrogates priming of CD8<sup>+</sup> T cells by exogenous cell-associated antigens. *Immunity*. 2002; 17:211–220. [PubMed: 12196292]
13. Sanos SL, et al. ROR $\gamma$ t and commensal microflora are required for the differentiation of mucosal interleukin 22–producing NKp46<sup>+</sup> cells. *Nat Immunol*. 2009; 10:83–91. [PubMed: 19029903]
14. Sano T, et al. An IL-23R/IL-22 circuit regulates epithelial serum amyloid A to promote local effector Th17 responses. *Cell*. 2015; 163:381–393. [PubMed: 26411290]
15. Sonnenberg GF, Fouser LA, Artis D. Functional biology of the IL-22-IL-22R pathway in regulating immunity and inflammation at barrier surfaces. *Adv Immunol*. 2010; 107:1–29. [PubMed: 21034969]
16. Boos MD, Yokota Y, Eberl G, Kee BL. Mature natural killer cell and lymphoid tissue-inducing cell development requires Id2-mediated suppression of E protein activity. *J Exp Med*. 2007; 204:1119–1130. [PubMed: 17452521]
17. Savage AK, et al. The transcription factor PLZF directs the effector program of the NKT cell lineage. *Immunity*. 2008; 29:391–403. [PubMed: 18703361]
18. Sawa S, et al. ROR $\gamma$ t<sup>+</sup> innate lymphoid cells regulate intestinal homeostasis by integrating negative signals from the symbiotic microbiota. *Nat Immunol*. 2011; 12:320–326. [PubMed: 21336274]

19. Eberl G, et al. An essential function for the nuclear receptor ROR $\gamma$ t in the generation of fetal lymphoid tissue inducer cells. *Nat Immunol.* 2004; 5:64–73. [PubMed: 14691482]
20. Kinnebrew MA, et al. Interleukin 23 Production by Intestinal CD103+CD11b+ Dendritic Cells in Response to Bacterial Flagellin Enhances Mucosal Innate Immune Defense. *Immunity.* 2012; 36:276–287. [PubMed: 22306017]
21. Pickard JM, et al. Rapid fucosylation of intestinal epithelium sustains host-commensal symbiosis in sickness. *Nature.* 2014; 514:638–641. [PubMed: 25274297]
22. Lindemans CA, et al. Interleukin-22 promotes intestinal-stem-cell-mediated epithelial regeneration. *Nature.* 2015; 528:560–564. [PubMed: 26649819]
23. Coburn CT, et al. Defective uptake and utilization of long chain fatty acids in muscle and adipose tissues of CD36 knockout mice. *J Biol Chem.* 2000; 275:32523–32529. [PubMed: 10913136]
24. Cohen P, et al. Role for stearoyl-CoA desaturase-1 in leptin-mediated weight loss. *Science.* 2002; 297:240–243. [PubMed: 12114623]
25. Zong H, Armoni M, Harel C, Karnieli E, Pessin JE. Cytochrome P-450 CYP2E1 knockout mice are protected against high-fat diet-induced obesity and insulin resistance. *AJP: Endocrinol Metabol.* 2012; 302:E532–9.
26. Hotamisligil GS, et al. Uncoupling of obesity from insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid binding protein. *Science.* 1996; 274:1377–1379. [PubMed: 8910278]
27. Cai L, Wang Z, Ji A, Meyer JM, van der Westhuyzen DR. Scavenger receptor CD36 expression contributes to adipose tissue inflammation and cell death in diet-induced obesity. *PLoS ONE.* 2012; 7:e36785. [PubMed: 22615812]
28. Pan X, Hussain MM. Clock is important for food and circadian regulation of macronutrient absorption in mice. *J Lipid Res.* 2009; 50:1800–1813. [PubMed: 19387090]
29. Douris N, et al. Nocturnin regulates circadian trafficking of dietary lipid in intestinal enterocytes. *Curr Biol.* 2011; 21:1347–1355. [PubMed: 21820310]
30. Buxton OM, et al. Adverse metabolic consequences in humans of prolonged sleep restriction combined with circadian disruption. *Sci Transl Med.* 2012; 4:129ra43–129ra43.
31. Scheer FAJL, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci USA.* 2009; 106:4453–4458. [PubMed: 19255424]
32. Motomura Y, et al. The transcription factor E4BP4 regulates the production of IL-10 and IL-13 in CD4<sup>+</sup> T cells. *Nat Immunol.* 2011; 12:450–459. [PubMed: 21460847]
33. Madison BB, et al. Cis elements of the villin gene control expression in restricted domains of the vertical (crypt) and horizontal (duodenum, cecum) axes of the intestine. *J Biol Chem.* 2002; 277:33275–33283. [PubMed: 12065599]
34. Moh A, et al. Role of STAT3 in liver regeneration: survival, DNA synthesis, inflammatory reaction and liver mass recovery. *Lab Invest.* 2007; 87:1018–1028. [PubMed: 17660847]
35. Cash HL, Whitham CV, Behrendt CL, Hooper LV. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science.* 2006; 313:1126–1130. [PubMed: 16931762]
36. Stappenbeck TS, Hooper LV, Manchester JK, Wong MH, Gordon JI. Laser capture microdissection of mouse intestine: characterizing mRNA and protein expression, and profiling intermediary metabolism in specified cell populations. *Meth Enzymol.* 2002; 356:167–196. [PubMed: 12418197]
37. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc.* 2008; 3:1101–1108. [PubMed: 18546601]



**One sentence summary**


The intestinal microbiota regulates body composition through the transcription factor NFIL3, which controls a circadian lipid metabolic pathway in the intestine.

Author Manuscript


Author Manuscript

Author Manuscript

Author Manuscript




**Figure 1. *Nfil3*<sup>IEC</sup> mice are resistant to high fat diet (HFD)-induced obesity**  
**(A)** qRT-PCR analysis of *Nfil3* transcript abundance in small intestinal epithelial cells recovered by laser capture microdissection from conventional (conv) and germ-free mice.  
**(B)** Age-matched *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice were co-housed and placed on a high fat diet (HFD) for 10 weeks. Body weight was measured before and after diet switching. **(C)** Body fat percentages of mice in **(B)**. **(D)** Epididymal fat pad weight, **(E)** serum triglyceride concentration, **(F)** hematoxylin & eosin (H&E) staining of liver (scale bar=100 μm), and **(G)** glucose tolerance and insulin tolerance tests. **(H)** Body fat percentage of mice treated with or without antibiotics after switching to HFD. All data represent two independent experiments with 4–8 mice per group. Male mice were used in all experiments. Means±SEM are plotted; statistics were performed with Student's t-test or one-way ANOVA. \*p<0.05; \*\*p<0.01; \*\*\*p<0.001; ns, not significant.



**Figure 2. The microbiota induces epithelial NFIL3 expression through the circadian clock factor REV-ERBa and a DC-ILC3 signaling relay**


(A–D) qRT-PCR analysis of *Nfil3* (A) and *Rev-erba* (C) transcript abundance in small intestinal epithelial cells from germ-free (dotted line) and conventional mice (solid line) across a 24-hour day-night light cycle. Western blot analysis of NFIL3 (B) and REV-ERBa (D) was performed on small intestinal epithelial cells isolated from conventional or germ-free mice, Lamin B is the loading control. (E) Chromatin immunoprecipitation assay on intestinal epithelial cells using immunoglobulin G (IgG) or anti-REV-ERBa antibody.

Precipitated fragments of the *Nfil3* promoter were detected by qRT-PCR. **(F)** qRT-PCR analysis of epithelial *Nfil3* expression in conventional wild-type, antibiotic (Abx)-treated wild-type or Abx-treated *Rev-erba*<sup>-/-</sup> mice. **(G)** qRT-PCR analysis of epithelial *Rev-erba* and *Nfil3* expression in germ-free and conventional wild-type mice and conventional *Myd88*<sup>fl/fl</sup>, *Myd88*<sup>-/-</sup>, *Myd88*<sup>IEC</sup> (epithelial cell-specific knockout) and *Myd88*<sup>DC</sup> (DC-specific knockout) mice. **(H)** qRT-PCR analysis of epithelial *Rev-erba* and *Nfil3* expression in germ-free and conventional wild-type mice, conventional *Cd11c*-DTR mice that were untreated or treated with *Diphtheria* toxin (DT), *Id2*<sup>gfp/gfp</sup> and *Rag1*<sup>-/-</sup> mice. **(I,J)** qRT-PCR analysis of epithelial *Nfil3* **(I)** and *Rev-erba* **(J)** expression in *Rorc*<sup>+/+</sup> (solid line) and *Rorc*<sup>gfp/gfp</sup> (dotted line) mice. **(K)** qRT-PCR analysis of epithelial *Rev-erba* and *Nfil3* expression in *Myd88*<sup>-/-</sup> mice treated with recombinant IL-23, IL-22 or vehicle. Data in E,F,G,H, and K were collected at ZT4. N=3–8 mice per group. Means±SEM are plotted; statistics were performed with Student's t-test or one-way ANOVA. \*p<0.05; \*\*p<0.01; \*\*\*p<0.001; ns, not significant; conv, conventional; ZT, Zeitgeber time.



**Figure 3. STAT3 represses *Rev-erba* transcription by binding directly to its promoter**

(A) Schematic of the *Rev-erba* gene promoter. (B) ChIP analysis of intestinal epithelial cells from conventional (conv) or germ-free mice using immunoglobulin G (IgG) or anti-STAT3 antibody. Precipitated fragments of the *Rev-erba* promoter or control exon were detected by qRT-PCR. (C) Luciferase reporter assay. A 504 bp fragment of *Rev-erba* promoter was fused to a firefly luciferase reporter. HEK-293T cells were transfected with reporters and either empty vector, a wild-type STAT3-encoding vector (*Stat3-wt*), or a dominant active STAT3-encoding vector (*Stat3-c*). (D) Western-blot of total STAT3, phosphorylated STAT3 (p-STAT3) in small intestinal organoids treated with IL-22 and/or the STAT3 inhibitor Static. Mito 70 is the loading control. (E,F) qRT-PCR analysis of *Rev-erba* (E) and *Nfil3* (F) expression in small intestinal organoids treated with IL-22 and/or Static. (G,H) qRT-PCR analysis of epithelial *Rev-erba* (G) and *Nfil3* (H) expression in *Stat3<sup>fl/fl</sup>* and *Stat3<sup>IEC</sup>* mice at ZT4. N=3–8 samples per group. Means±SEM are plotted; statistics were performed with Student's t-test or one-way ANOVA. \*p<0.05; \*\*p<0.01; \*\*\*p<0.001; \*\*\*\*p<0.0001; ns, not significant.



**Figure 4. Epithelial NFIL3 controls expression of a circadian lipid metabolic program and regulates lipid absorption in intestinal epithelial cells**

(A) RNAseq analysis of epithelial cell transcripts in *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>ΔIEC</sup>* mice across a circadian cycle. The heatmap visualizes expression levels of the 33 genes that have altered expression in *Nfil3<sup>ΔIEC</sup>* mice as compared to *Nfil3<sup>fl/fl</sup>* mice. Genes encoding proteins that function in lipid metabolism are highlighted in blue. Top panels show sustained circadian expression of the core clock genes *Bmal1* (*Arntl*), *Per2*, and *Nr1d1* (*Rev-erba*) in *Nfil3<sup>ΔIEC</sup>* mice. (B) qRT-PCR analysis of epithelial *Cd36* and *Scd1* expression in germ-free wild-type

(wt) and conventional *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice at ZT4. **(C)** Western blot of epithelial CD36 in germ-free (gf) and conventional wild-type (conv) mice, and in conventional *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice. All mice were fed a HFD. Mice were sacrificed at ZT4.  $\alpha$ -tubulin is the loading control. **(D,E)** qRT-PCR analysis of epithelial *Cd36* and *Scd1* expression in conventional wild-type (wt) and ID2-deficient (*Id2<sup>gfp/gfp</sup>*) mice **(D)** and *Stat3<sup>fl/fl</sup>* and *Stat3<sup>IEC</sup>* mice **(E)** at ZT4. **(F)** Oil red O detection of lipids in the small intestines of *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice fed on HFD. Nuclei were stained with Methyl Green. Scale bar=40  $\mu$ m. **(G)** Total lipid concentrations in isolated small intestinal epithelial cells from *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice fed on HFD. **(H)** Total neutral lipid concentrations in feces of *Nfil3<sup>fl/fl</sup>* and *Nfil3<sup>IEC</sup>* mice fed on HFD. Data in B,D,E,G,H have N=5-12 mice per group. Means $\pm$ SEM are plotted; statistics were performed with Student's t-test or one-way ANOVA. \*p<0.05; \*\*p<0.01; ns, not significant; ZT, Zeitgeber time