

## ORIGINAL ARTICLE

# Mitochondrial UQCRB as a new molecular prognostic biomarker of human colorectal cancer

Hyun-Chul Kim<sup>1,2</sup>, Junghwa Chang<sup>1</sup>, Hannah S Lee<sup>1</sup> and Ho Jeong Kwon<sup>1,3</sup>

Ubiquinol cytochrome *c* reductase binding protein (UQCRB) is important for mitochondrial complex III stability, electron transport, cellular oxygen sensing and angiogenesis. However, its potential as a prognostic marker in colorectal cancer (CRC) remains unclear. The aim of this study was to determine whether UQCRB can be used as a diagnostic molecular marker for CRC. The correlation between the expression of three genes (*UQCRB*, *UQCRFS1* and *MT-CYB*) in the mitochondrial respiratory chain complex III and clinico-pathological features was determined. Compared to non-tumor tissues, *UQCRB* gene expression was upregulated in CRC tissues. Gene and protein expression of the genes were positively correlated. Copy number variation (CNV) differences in *UQCRB* were observed in CRC tissues (1.32-fold) compared to non-tumor tissues. The CNV of *UQCRB* in CRC tissues increased proportionally with gene expression and clinical stage. Single-nucleotide polymorphisms in the 3'-untranslated region of *UQCRB* (rs7836698 and rs10504961) were investigated, and the rs7836698 polymorphism was associated with CRC clinical stage. DNA methylation of the *UQCRB* promoter revealed that most CRC patients had high methylation levels (12/15 patients) in CRC tissues compared to non-tumor tissues. UQCRB overexpression and CNV gain were correlated with specific CRC clinico-pathological features, indicating clinical significance as a prognostic predictor in CRC. Gene structural factors may be more important than gene transcription repression factors with respect to DNA methylation in UQCRB overexpression. Our results provide novel insights into the critical role of UQCRB in regulating CRC, supporting UQCRB as a new candidate for the development of diagnostics for CRC patients.

*Experimental & Molecular Medicine* (2017) 49, e391; doi:10.1038/emm.2017.152; published online 17 November 2017

## INTRODUCTION

Colorectal cancer (CRC) is a common malignancy worldwide and the leading cause of cancer-related death.<sup>1</sup> Moreover, CRC is the fourth leading cause of cancer-related deaths in South Korea. Although the prevalence of CRC is lower in South Korea than in Western industrialized nations, the number of patients with CRC has increased rapidly over the past several decades.<sup>2–4</sup> Despite recent advances in the early diagnosis and treatment of CRC, the overall prognosis for CRC patients remains poor. Thus, novel and effective targets must be identified to improve therapeutic efficacy and clinical outcomes.

The mitochondrion is the main energy-producing organelle of the cell, bearing five oxidative phosphorylation complexes that regulate ATP synthesis.<sup>5–6</sup> Of the five oxidative phosphorylation complexes, complex III carries out electron transport, ubiquinol radical stabilization and cellular oxygen sensing.<sup>7</sup> Mitochondrial complex III (cytochrome *bc1* complex)

transfers electrons from ubiquinol to cytochrome *c* and contributes to the generation of an electrochemical proton gradient.<sup>8–10</sup> Notably, mitochondrial complex III deficiency is associated with lactic acidosis, hypoglycemia, encephalopathy, among other disorders in humans.<sup>11</sup>

The *UQCRB*, *UQCRFS1* (encoding Rieske iron–sulfur protein) and *MT-CYB* (encoding cytochrome *b*) genes are components of complex III. Ubiquinol cytochrome *c* reductase binding (UQCRB) is the homolog of the yeast Qcr7 subunit, which is located in the matrix-inner mitochondrial membrane interphase and is thought to interact with cytochrome *b* in the early stages of the assembly pathway.<sup>12</sup> Qcr7 incorporation is essential for the stabilization of heme *b*.<sup>13</sup> UQCRB plays an important role in *MT-CYB* stability, explaining the low amounts of *b*-type cytochrome found in the mitochondria of CRC patients.

UQCRB is a functional gene with nuclear encoding of a 13.3-kDa subunit of mitochondrial complex III; UQCRB is

<sup>1</sup>Chemical Genomics Global Research Laboratory, Department of Biotechnology, College of Life Science and Biotechnology, Yonsei University, Seoul, Korea; <sup>2</sup>Department of Molecular Biology, Seoul Medical Science Institute/Seoul Clinical Laboratories, Yongin, Korea and <sup>3</sup>Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Korea

Correspondence: Professor HJ Kwon, Chemical Genomics Global Research Laboratory, Department of Biotechnology, College of Life Science and Biotechnology, Yonsei University, 50 Yonsei-ro, Seodaemun-gu, Seoul 120-749, Korea.

E-mail: kwonhj@yonsei.ac.kr

Received 12 January 2017; revised 21 April 2017; accepted 26 April 2017

located at the human chromosome location 8q22 and participates in stabilization of the ubisemiquinone radical through hydrophobic interactions with ubiquinone.<sup>14,15</sup> In addition, UQCRB is found in the respiratory chains of all aerobic organisms, as well as in the electron transfer systems of chloroplasts and photosynthetic bacteria.<sup>16,17</sup> Deletion of the gene encoding UQCRB causes a defect in complex III function, resulting in hypoglycemia and lactic acidosis. Disorders of complex III are comparatively rare, but they present as a clinically heterogeneous group of diseases.<sup>18</sup>

We previously demonstrated that loss of function of UQCRB by either genetic and pharmacological means inhibited angiogenesis, indicating that UQCRB plays a key role in this process and is a prognostic marker of angiogenesis and mitochondria-related diseases.<sup>19,20</sup> However, few studies have examined its clinical role in CRC. Thus, we investigated the clinical roles of UQCRB in CRC using genetic/proteomic analysis of human CRC tissues.

## MATERIALS AND METHODS

### Study participant data and tissue samples

This study was conducted on a total of 15 human CRC tissues and matched adjacent non-tumor tissues, which were histopathologically and clinically diagnosed at the Severance Hospital, Seoul, Korea, from November 2012 to December 2012. Disease-free tissue samples served as control groups. Tissues were flash-frozen in liquid nitrogen immediately after resection. Colorectal tissues were classified according to the World Health Organization tumor classification system. We obtained Institutional Review Board approval for the use of the CRC tissues from the Yonsei University Institutional Review Board (IRB No. 1040917-201408-BR-211-02E). The demographic and pathologic characteristics of the CRC patients are summarized in Table 1.

### Analysis of gene expression (quantitative real-time RT-PCR)

Total RNA from 15 pairs of CRC tissues and non-tumor tissues was extracted using aRNeasy Mini Kit from Qiagen (Hilden, Germany) according to the manufacturer's instructions. RNA integrity was spectrophotometrically confirmed by electrophoresis on a 2% agarose gel. First-strand cDNA was synthesized from 0.5 µg of DNase-treated total RNA using the SuperScript III First-Strand Synthesis System (Invitrogen, Carlsbad, CA, USA) with random hexamers.

Real-time PCR was carried out using an ABI 7500 Fast Real-Time PCR System (Applied Biosystems, Foster City, CA, USA). The RNA expression levels of *UQCRB*, *UQCRFS1* and *MT-CYB* in CRC tissues with respect to their normal adjacent tissues were investigated by TaqMan Gene Expression Assay with a commercial primer/probe product (Applied Biosystems). The TaqMan Gene Expression Assay primers used for *UQCRB*, *UQCRFS1* and *MTCYB* were Hs01890823\_s1, Hs00705563\_s1 and Hs02596867\_s1, respectively. Human *GAPDH* was used as an internal control. PCR was carried out using 10 µl of 2× TaqMan Universal PCR Master Mix, 1 µl of TaqMan Gene Expression Assay, 8 µl of RNase-free water and 1 µl of cDNA template. All reactions were conducted in duplicate along with no-template controls. The thermal cycling conditions comprised an initial step at 95 °C for 10 min, followed by 40 cycles at 95 °C for 10 s and 60 °C for 40 s. Relative expression was calculated using the  $2^{-\Delta\Delta Ct}$  method.<sup>21</sup>

### Protein expression

Immunohistochemistry was performed to investigate alterations in protein expression in CRC tissues. The tissue sections, which were 4-µm thick from paraffin-embedded blocks, were mounted on coated slides and placed in drying racks. Subsequently, these samples were deparaffinized and rehydrated using xylene and graded alcohol washes. The sections were heated in 10 mM citrate buffer (pH 6.0) in a pressure boiler for 10 min to achieve antigen recovery. Endogenous peroxidase activity was blocked with 3% hydrogen peroxide. The primary antibody was rabbit polyclonal UQCRB antibody(1:200, HPA043060; Sigma-Aldrich, St Louis, MO, USA), which was applied overnight at 4 °C. After washing, tissue sections were treated with anti-rabbit secondary antibody (Sigma-Aldrich) for 30 min, followed by further incubation using the horseradish peroxidase complex method following the manufacturer's instructions. The sections were developed with 3,3'-diaminobenzidine as the chromogen (DAKO, Carpinteria, CA, USA), and the microscopic images were captured digitally. As negative controls, we replaced the primary antibody with 5% fetal bovine serum.

The immunohistochemistry results were semiquantitatively examined by two independent observers who were blinded to the patient data. Staining intensity and the percentage of the stained cells were used as criteria for evaluation. A measure of scoring according to staining intensity (0, negative; I, weak; II, moderate; III, strong) and the percentage of stained cells (0, none; 1, <10%; 2, 10–50%; 3, >50%) was carried out to divide the samples into two groups: a high-expression group (moderate/strong, stain rated 2 and 3) and B low-expression group (negative/weak, stain rated 0 and 1).<sup>22</sup>

### Copy number variation

Total genomic DNA from 15 pairs of CRC tissues and non-tumor tissues was extracted using aQIAamp DNA Mini Kit from Qiagen according to the manufacturer's instructions. The relative changes in *UQCRB* and *BRUNOLA* (*CELF4*)<sup>23</sup> copy numbers between CRC tissues and non-tumor tissues were determined by real-time PCR and the TaqMan Copy Number Assay (Applied Biosystems). The TaqMan assay primers used for *UQCRB* and *BRUNOLA* were Hs02791612\_cn and Hs06439304\_cn, respectively. PCR was performed in a total volume of 25 µl in each well, which contained 12.5 µl of TaqMan Universal MasterMix (Applied Biosystems), 1 µl of each primer (12.5 pM) and 25 ng of genomic DNA. The thermal cycling conditions included an initial step at 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s and 60 °C for 60 s.

For all PCR assays, cycle threshold (Ct) numbers were established using 7500 Software v2.0.6 (Applied Biosystems), and the copy numbers, which were normalized against a reference gene (RNase P, two copies in a diploid genome) and the calibrator (normal sample), were determined using the  $2^{-\Delta\Delta Ct}$  method. A two-fold increase or decrease in the copy number of *UQCRB* and *BRUNOLA* in CRC tissues compared to the corresponding normal sample within the pair was considered an amplification or a deletion, respectively. Copy number variations (CNVs) were calculated using Applied Biosystems CopyCaller software (Applied Biosystems).

### Single-nucleotide polymorphism

Single-nucleotide polymorphisms (SNPs) were verified by high-throughput bi-directional Sanger DNA sequencing using the 3130×1 genetic analyzer (Applied Biosystems) for high-quality resolution of polymorphisms, insertions and deletions in the coding sequence (333 base pairs, NG\_008237), marker rs7836698 and marker rs10504961.<sup>24</sup>

**Table 1 Clinical characteristics of 15 CRC patients at the time of diagnosis**

Parameter	Number of patients	% of patients	Range
<i>Gender</i>			
Male	10	66.7	
Female	5	33.3	
<i>Age at diagnosis (year)</i>			
<65	9	60.0	39–64
≥65	6	40.0	65–79
<i>Localization</i>			
Rectum	5	33.3	
Colon	9	60.0	
Cecum	1	6.7	
<i>Tumor size (cm)</i>			
<3	1	6.7	2.7–3
3–5	9	60.0	3–5
>5	5	33.3	5–8.45
<i>Clinical (TNM) stage</i>			
I	0	0	
II	6	40.0	
III	8	53.3	
IV	1	6.7	

Abbreviations: CRC, colorectal cancer; TNM, tumor-node metastasis.

Major/minor alleles, the location and minor allele frequency of two SNPs (rs7836698 and rs10504961) were C/T, the 5'-untranslated region (UTR), and T-0.42 and T-0.50, respectively.

### DNA methylation

Genomic DNA isolated from CRC tissues and non-tumor tissues was analyzed for DNA methylation in the promoter region of the human *UQCRB* gene using the EpiTect Methyl II PCR Array System according to the supplier's instructions (Qiagen). Genomic DNA (250 ng) from tissues was digested using aEpiTect methyl II DNA restriction Kit (Qiagen) according to the manufacturer's instructions. The *UQCRB* gene EpiTect methyl II Custom PCR array (Qiagen) was applied. Real-time PCR was performed using RT<sup>2</sup> SYBR Green ROX qPCR Mastermix (Qiagen) according to the manufacturer's recommendations in the ABI 7500 Fast Real-Time PCR System (Applied Biosystems). For this analysis, the relative concentrations of differentially methylated DNA (specifically, hypermethylated, unmethylated and intermediary methylated DNA) were determined by comparing the amount of each digest with that of a mock digest. For each sample, data were expressed as the sum of the percent of hypermediary and intermediary methylated DNA. Ct values were used to calculate the percentages of methylated and unmethylated DNA according to the recommendations of the manufacturer (Qiagen).

### Statistical analysis

*UQCRB* expression and CNV profiles in the different groups are shown as the mean and standard error of the mean. Statistical calculations were conducted using the GraphPad Prism 5 (GraphPad, La Jolla, CA, USA) and MedCalc software packages ([https://www.](https://www.medcalc.org/)

[medcalc.org/](https://www.medcalc.org/)).<sup>25</sup> To compute *P*-values, Tukey's multiple comparison test and one-way analysis of variance tests were used. A *P*-value of <0.05 was considered statistically significant. Each experiment was repeated at least twice.

## RESULTS


### *UQCRB* gene expression

The *UQCRB*, *UQCRFS1* and *MT-CYB* genes are important components in mitochondrial complex III.<sup>12,13</sup> Real-time PCR analysis showed that *UQCRB*, *UQCRFS1* and *MT-CYB* mRNA expression levels were relatively higher in CRC tissues (T) than in the matched adjacent non-tumor tissues (N) (Figure 1a–c). Expression levels were normalized to *GAPDH*. *UQCRB* and *UQCRFS1* mRNA expression levels were significantly upregulated in all 15 pairs (100%, 15/15 patients) of CRC tissues compared to non-tumor tissues (Figure 1a and b). *UQCRB* showed the highest mRNA expression level among the three target genes. The CRC tissues/non-tumor tissues (T/N) ratio of *UQCRB* mRNA expression was >1.3-fold in all samples, and the highest ratio was 20.61-fold (Figure 1a). By contrast, *MT-CYB* mRNA expression was upregulated in 13 pairs (86.6%, 13/15 patients) of CRC tissues compared to non-tumor tissues and was downregulated only in two pairs (13.3%, 2/15 patients) of CRC patients (Figure 1c). However, no significant difference was observed in the expression of other genes (*UQCRB* and *UQCRFS1*). *UQCRB* gene expression showed the highest expression (average log T/N, 0.78) of complex III-related three genes. On average, the three complex III-related genes (*UQCRB*, *UQCRFS1* and *MT-CYB*) were confirmed to show high gene expression. The average log T/N expression of *UQCRFS1* and *MT-CYB* were 0.51 and 0.31, respectively (Figure 1d).


We used the area under the receiver operative characteristic curve (AUC) to evaluate the predictive efficacy of RNA expression. To evaluate the prediction power of the three genes in complex III, we compared the AUC of models developed with *UQCRB* and other genes. The AUC of *UQCRB* expression was 0.991 (95% CI, 0.868–1.000; *P*<0.0001). The Youden index, sensitivity and specificity of *UQCRB* expression were 0.933%, 93.3% and 100.0%, respectively. The AUC values of *UQCRFS1* and *MT-CYB* were 1.000 (95% CI, 0.884–1.000; *P*<0.0001) and 0.836 (95% CI, 0.655–0.945; *P*<0.0001), respectively. The Youden index, sensitivity and specificity of *UQCRFS1* expression were 1.000%, 100.0% and 100.0%, respectively. However, the Youden index, sensitivity and specificity for *MT-CYB* expression were 0.600%, 86.7% and 73.3%, respectively (Figure 2). From here forward, we decided to focus on *UQCRB* in the following study because *UQCRB* was significantly overexpressed among the complex III-related three genes (*UQCRB*, *UQCRFS1* and *MT-CYB*).

### *UQCRB* protein expression

*UQCRB* protein was overexpressed within the tumor tissues examined in this study, with *UQCRB* staining mainly found in the cytoplasm. According to our definition, the rate of high *UQCRB* expression in CRC tissues was 13.3% (2/15 patients),




**Figure 1** Logarithm of the ratio of colorectal cancer (CRC) tissues (T)/matched adjacent non-tumor tissues (N) for RNA levels of *UQCRB* (a), *UQCRFS1* (b) and *MT-CYB* (c). (d) Expression level of complex III genes in CRC. Gene expression levels were normalized to that of *GAPDH*. Relative expression was calculated using the  $2^{-\Delta\Delta Ct}$  method. Data are presented as the mean  $\pm$  s.e.m. *P*-values are represented as \**P*<0.05, \*\**P*<0.01 and \*\*\**P*<0.001.



**Figure 2** Analysis of ROC curve of *UQCRB* (a), *UQCRFS1* (b) and *MT-CYB* (c) expression. The AUC of individual candidate genes was calculated to diagnose colorectal cancer (CRC). AUC was calculated using the MedCalc analysis program.

whereas the rate of high UQCRB expression in non-tumor tissues was 0% (0/15 patients). As shown in Figure 3, the UQCRB protein of two patients (patient no. 5 and no. 13) was overexpressed in CRC tissues compared to its expression in non-tumor tissues. CRC tissues (Figure 3c and d) showed moderate to strong cytoplasmic staining of UQCRB, whereas non-tumor tissues (Figure 3a and b) showed either weak staining or no staining. Compared to non-tumor tissues,

UQCRB protein of CRC tissues was highly expressed. Furthermore, two high-expression tissues in immunohistochemistry analysis were consistent with high gene expression (Figures 1 and 3). A positive correlation was observed between gene expression and protein expression. These results indicate that genetic and proteomic expression of UQCRB is upregulated in CRC tissues compared to that in non-tumor tissues.



**Figure 3** Detection of UQCRB protein expression in representative (patients 5 and 13) colorectal cancer (CRC) tissues (T) and non-tumor tissues (N) as assessed by immunohistochemical staining with UQCRB-specific antibodies. CRC tissues (c, d) showed moderate to strong cytoplasmic staining of UQCRB, whereas non-tumor tissues (a, b) demonstrated either weak staining or no staining. Compared to non-tumor tissues, UQCRB protein of CRC tissues was highly expressed (original magnification;  $\times 400$ ).

Interestingly, there was no association between protein expression and other factors, such as gender, age, tumor localization, tumor size and clinical stage.

#### UQCRB gene CNV

CNV varies individually in human genomes at the genetic level. To determine the clinical stage-related changes in the genetic variations of *UQCRB*, we investigated the level of CNV in CRC tissues. *UQCRB* and *BRUNOLA* copy numbers were calculated and normalized to the endogenous reference gene RNase P, which is known to be present in two copies in a diploid genome. TaqMan assay Copy Number Assays designed to specifically target *UQCRB* exon 4 and *BRUNOLA* intron 2 sequences were tested on genomic DNA from 15 CRC tissues paired with non-tumor tissues.

Figure 4 shows the results of the calculated *UQCRB* copy numbers in the tested CRC tissues/non-tumor tissues. The *UQCRB* copy number was significantly upregulated in 14 (14/15 patients, 93.3%) pairs of CRC tissues compared to its copy number in non-tumor tissues (Figure 4a). In addition, the *BRUNOLA* copy number, representing a negative control gene in CRC, was upregulated in only five (5/15 patients, 33.3%) pairs of CRC tissues compared to that in non-tumor tissues (Figure 4a). Comparative analysis indicated that the average log T/N of the *UQCRB* and *BRUNOLA* copy number levels were 0.11 and  $-0.07$ , respectively. The AUC of *UQCRB* CNV was 0.891 (95% CI, 0.723–0.975;  $P < 0.0001$ ). The Youden index, sensitivity and specificity of *UQCRB* CNV were 0.867%, 86.7%


and 100.0%, respectively, whereas the AUC of *BRUNOLA* was 0.664 (95% CI, 0.457–0.872;  $P < 0.0001$ ; Figure 4b).

#### Relationship between gene expression and CNV


To explore whether this change in *UQCRB* gene expression was associated with a corresponding change in *UQCRB* copy number, the amount of mRNA of *UQCRB* and the corresponding *UQCRB* copy number were compared in CRC tissues/non-tumor tissues. As shown in Figure 5, the relationships between *UQCRB* gene expression, CNV and clinical stages were analyzed in 15 CRC patients. Gene expression was proportionately correlated with CNV and clinical stages. The samples included six cases of clinical stage II (40%, 6/15 patients), eight cases of clinical stage III (53.3%, 8/15 patients) and one case of clinical stage IV (6.6%, 1/15 patients). The numerical correlation of all samples was 0.105, but it was  $-0.033$  in clinical stage II, 0.279 in clinical stage III and 0.161 in clinical stage III–IV. Although a limited number of samples were available for this analysis, we predict a high correlation for *UQCRB* gene expression with its CNV in a low clinical stage (stage II) for early diagnosis. However, the highest correlation was observed for a high clinical stage (stage III). These data confirm a high numerical correlation with a high clinical stage rather than a low clinical stage.

#### SNP analysis of *UQCRB* gene

SNP analysis results of the *UQCRB* CDS (NG\_008237.1; 105..123, 2408..2479, 3695..3861, 4503..4580) were not noticed. We analyzed the two polymorphisms in the 3'-UTR region of



**Figure 4** Gene copy number variation. (a) *UQCRB* and *BRUNOL4* gene copy number alterations in patients with colorectal cancer (CRC). *UQCRB* copy number; logarithm T/N of patient No. 1 shows deletion, whereas the others show amplification. (b) Analysis of ROC curve of *UQCRB*, and *BRUNOL4* CNV. The AUC of genes was calculated to diagnose CRC. AUC was calculated using the MedCalc analysis program.



**Figure 5** Correlation coefficient was utilized to describe the association between gene expression and copy number in colorectal cancer (CRC) tissues per non-tumor tissues for *UQCRB*. Gene expression is shown on the left y axis; copy number is shown on the right y axis.

*UQCRB* (rs7836698 and rs10504961). Two SNPs (rs7836698 and rs10504961) were already known as prognostic markers for CRC.<sup>24</sup> As shown in Table 2, the marker rs7836698 in *UQCRB* showed an increased TT genotype in CRC tissues compared to

non-tumor tissues. The TT genotype-to-CC genotype ratios of rs7836698 in CRC tissues and non-tumor tissues were 83.3% and 7.7%, respectively. In particular, the marker rs7836698 was significantly associated with CRC tumor cases compared to non-tumor tissues. The rs7836698 polymorphism was associated with clinical stage in CRC. However, the marker rs10504961 showed no difference between CRC tissues and non-tumor tissues. Interestingly, carriers of the two *UQCRB* polymorphisms, the TT carriers, were ~2-fold more likely to develop tumors in the colon than in the rectum.

#### DNA methylation analysis of *UQCRB*

Cancer initiation and progression is regulated by both genetic and epigenetic modifications. Accordingly, we assessed epigenetic modifications of *UQCRB* and analyzed the classical dependence of DNA methylation in the gene promoter regions with that of gene expression. In a total of 30 samples, we determined *UQCRB* differential methylation in CRC tissues and non-tumor tissues from 15 patients with CRC. Through quantitative PCR-based DNA methylation analysis, we found that *UQCRB* was significantly more methylated in CRC tissues than in non-tumor tissues. Notably, CRC tissues in 12 of 15 patients showed a higher degree of methylation than non-tumor tissues (Figure 6a). The average degree of methylation in CRC tissues was 2.59%, whereas this value was 1.49% in non-tumor tissues. The methylation rate in the *UQCRB* promoter region was 1.7-fold higher in CRC tissues than in non-tumor tissues. Accordingly, there was a significant correlation between

gene expression and methylation level of the *UQCRB* promoter (Figure 6b). To verify the result, however, alternative methods, such as bisulfite-pyrossequencing analysis, will be conducted in a following study.

**Table 2 Association of two polymorphisms in *UQCRB* gene**


SNP	Tissue case	Genotype	Clinical (TNM) stage			Total no. (%)
			II	III	IV	
rs7836698	CRC tissues	CC	3	2	1	6 (40)
		CT	1	3		4 (26.6)
		TT	2	3		5 (33.3)
	Non-tumor tissues	CC	5	7	1	13 (86.6)
		CT		1		1 (6.6)
		TT	1			1 (6.6)
rs10504961	CRC tissues	CC	4	4	1	9 (60)
		CT	1	3		4 (26.6)
		TT	1	1		2 (13.3)
	Non-tumor tissues	CC	4	4	1	9 (60)
		CT	1	4		5 (33.3)
		TT	1			1 (6.6)

Abbreviations: CRC, colorectal cancer; TNM, tumor-node metastasis. Major/minor allele and location of two SNPs (rs7836698 and rs10504961) were C/T and 5'-UTR, respectively.

## DISCUSSION

Mitochondrial complex III deficiency is one of the least common oxidative phosphorylation defects associated with mitochondrial disease.<sup>26</sup> However, complex III deficiency is a genetic condition that can affect several parts of the body, including the brain, kidneys, liver, heart and skeletal muscles. Signs and symptoms of complex III deficiency typically begin in infancy, but they can appear later.<sup>27,28</sup> Many studies have suggested that hereditary defects in *UQCRB* of complex III cause several mitochondrial diseases, such as hypoglycemia, lactic acidosis, myopathy and cardiomyopathy.<sup>18,29</sup>

Few studies have examined *UQCRB* in CRC. The primary purpose of this study was to explore the relationship between the expression of *UQCRB* and the clinico-pathologic features of CRC. In the present study, the expression of three representative genes (*UQCRB*, *UQCRFS1* and *MT-CYB*) associated with complex III in CRC was examined and assessed for its potential clinical value in CRC development and progression. The expression of the three genes was increased significantly in CRC tissues. Furthermore, we demonstrated that *UQCRB* expression was upregulated at both the gene and protein levels in CRC tissues compared to matched adjacent non-tumor tissues. High levels of *UQCRB* expression were more frequently observed in CRC patients with poor pathologic differentiation or in those with advanced stages. We demonstrated that high *UQCRB* expression was correlated with



**Figure 6** DNA methylation of *UQCRB* promoter. (a) DNA methylation level of *UQCRB* promoter region from patients. (b) The correlation was utilized to describe the association between the gene expression and methylation level of the *UQCRB* promoter region.

CRC tissues, whereas low UQCRB expression was correlated with non-tumor tissues.

Owens *et al.*<sup>30</sup> quantified the mitochondrial oxidative phosphorylation activities of all five complexes and confirmed the activity and gene expression of complex III subunits in human breast cancer cell lines and primary tumors. The activity of complex III of breast cancer cell lines showed large differences compared to normal activity. In addition, *UQCRFS1* overexpression was detected in breast tumors compared to matched normal subjects. These studies directly demonstrated the contribution of impaired mitochondrial oxidative phosphorylation complex III to breast tumorigenesis.<sup>30</sup> In agreement with the results of previous studies, our results directly demonstrate the contribution of mitochondrial oxidative phosphorylation complex III to colon tumorigenesis. In particular, we found that unlike other genes, *MY-CYB* showed negative logarithm figures of the T/N ratio in clinical stage II, indicating that among the three genes, *MT-CYB* showed differences in clinical stages II and III. Moreover, patients with high gene expression generally showed an increased risk of distant metastasis and advanced clinical stage. These data demonstrate that complex III plays an important role in CRC progression and metastasis.

A previous study showed that *UQCRFS1* was overexpressed (14/40 patients, 35%) in breast cancer.<sup>30</sup> The expression of mitochondrial-encoded genes, including cytochrome B (complex III), generally decreased in myelodysplastic syndromes (MDS) and acute myeloid leukemia-MDS, whereas the mitochondrial DNA copy number increased rather than being reduced. Overall, MDS patients showing a decrease in mitochondrial gene expression displayed a stoichiometric imbalance in mitochondrial-encoded genes. The authors of that study concluded that dysregulated mitochondrial gene expression extends beyond a simple age-related effect and is compatible with the putative role of mitochondrial dysfunction in MDS pathophysiology.<sup>31</sup>

Notably, there were significant correlations between UQCRB expression and tumor types, although we detected no relationship between UQCRB expression and tumor size. The expression of the three genes may also be associated with the localization (rectum, colon and cecum) of CRC. We found that gene expression was high in most of the rectum samples of CRC (5/5 patients, 100%). Accordingly, UQCRB could potentially be used as a biomarker of rectal cancer.

The AUC is a standard analytical tool for evaluating diagnostic tests.<sup>32</sup> Although we were unable to accurately determine the AUC because of our limited sample number, we analyzed AUC with our tested CRC tissues ( $n = 15$ ). Genetic and proteomic expression of UQCRB was upregulated. The concordance rate between high expression of the *UQCRB* gene (100%, 15/15 patients) and UQCRB protein was 13.3% (2/15 patients).

*UQCRFS1* expression in additional matched normal and breast tumors revealed overexpression (35%, 14/40 patients) in breast tumors. *UQCRFS1* knockdown in breast tumor led to decreased mitochondrial membrane potential, and a reduction

in Matrigel invasion mediated by reduced reactive oxygen species levels, coinciding with decreased expression of NADPH oxidase involved in reactive oxygen species production.<sup>30</sup>

Cancer biomarkers in mitochondrial genetic/proteomic studies have not been extensively studied. Deregulation of the mitochondrial tumor suppressor genes (*SIRT3*, *SIRT4* and *MTUS1*) was detected in head and neck squamous cell carcinoma.<sup>33</sup> Sirtuins, a family of orthologs of yeast silent information regulator 3 (*Sirt3*) and 4 (*Sirt4*), are important tumor suppressor genes located in the mitochondria.<sup>34–37</sup> However, UQCRB was reported to be frequently downregulated in endometrial stromal cells.<sup>38</sup> In addition, the expression level of UQCRB was significantly decreased in Pallister–Killian syndrome samples, and an inverse linear correlation was observed between the level of miR-1224 and the UQCRB expression level.<sup>39</sup>

We observed that the *UQCRB* copy number was upregulated in CRC. Gene expression was proportionately correlated with CNV and clinical stage. In particular, a high numerical correlation was detected in clinical stage III. Similarly, *UQCRB* CNV appears to be more suitable as a candidate marker in CRC. A high correlation between expression and CNV has been reported in ovarian cancer.<sup>40</sup> The *BRUNOLA* copy number was downregulated, as expected. Poulgiannis *et al.*<sup>23</sup> demonstrated that *BRUNOLA* (encoding Bruno-like 4 splicing factor) copy number loss is an independent prognostic indicator in CRC. Lu *et al.*<sup>41</sup> evaluated CNV and gene expression to identify genes reproducibly associated with tumorigenesis. CNV regions identified 475 genes differentially expressed between tumor and normal tissues and showed concordance between CNV and gene expression changes. Based on combined gene expression profiles and CNV, the identified genes/pathways may serve as prognostic biomarkers for lung tumorigenesis. Wrzeszczynski *et al.*<sup>42</sup> evaluated gene expression data and predicted 156 genes with CNV and then correlated changes in expression. Among these genes, *UQCRB* was identified within The Cancer Genome Atlas (<http://tcga.cancer.gov>) data set and Memorial Sloan-Kettering Cancer Center data set in ovarian cancer. *UQCRB* is overexpressed in liver cancer cells, and CNV was found in cervical cancer and may be used as a cellular marker for cervical cancer diagnosis.<sup>42</sup> Moreover, altered copy number and changes in *UQCRB* gene expression were identified in specimens of hepatocellular carcinoma, and pancreatic ductal adenocarcinoma from microarray studies indicated that UQCRB plays a significant role in tumorigenesis.<sup>43,44</sup>

In the present study, we also demonstrated that for the *UQCRB* rs7836698 polymorphism, the TT genotype was associated with tumorigenesis in CRC tissues, but not in non-tumor tissues. There was no significant correlation between rs7836698 polymorphisms and gender, age, tumor localization, or tumor size. However, the TT genotype and CT genotype were more likely to be increased in stage III than in stage II. The genotype change at rs7836698 between CRC tissues and non-tumor tissues was proportional to clinical stage. Thus, rs7836698 polymorphisms are closely associated



with biomarkers of CRC. Two SNPs (rs7836698 and rs10504961) in the 3'-UTR of *UQCRB* were associated with overall survival of CRC patients, suggesting that SNPs identified by multivariate analysis can be used as independent prognostic factors for CRC.<sup>24</sup> As small numbers of analyzed SNPs strongly influence statistical analysis, a larger number of patient samples will be collected in the following study to obtain statistically significant results.

To determine whether the clinical stage-related increase in DNA methylation of the *UQCRB* promoter is a universal aging phenomenon, we measured the level of DNA methylation in the promoter of *UQCRB*. Clinical stage was not found to affect DNA methylation in *UQCRB*. In addition, there was no difference in the level of DNA methylation of the *UQCRB* promoter between elderly ( $\geq 65$  years) and young subjects ( $< 65$  years). However, the methylation rate in CRC tissues was higher than in non-tumor tissues. DNA methylation is a potential mechanism for regulating the expression of a gene. Cytosine residues in *CG* dinucleotides are targets for DNA methylation, and gene expression is typically reduced when DNA methylation occurs at a promoter. Accordingly, the level of DNA methylation was related to gene expression. Our results demonstrated that methylation level is highly correlated with mRNA expression, which is inversely proportional in CRC ( $R^2 = 0.1926$ ). However, the statistical correlation between the expression level and the methylation level was lower than the correlation between the expression level and the CNV level. Moarii *et al.*<sup>45</sup> suggested that the high correlation between promoter DNA methylation and gene expression differentiates tumor tissues from non-tumor tissues. The gene expression of a few specific genes, particularly transcription factors, is associated with DNA methylation in a tissue-dependent manner. In addition, both methylation and CNV are important requisite predictors of gene expression variation.<sup>45</sup> Ling *et al.*<sup>46</sup> suggested that the relative mRNA level of *UQCRB* was also reduced in muscle in the elderly compared to in young twins. The level of DNA methylation was also found to be related to gene expression. *UQCRB* was among the genes showing reduced expression in diabetic muscle. Age-related DNA methylation of *UQCRB* increases, whereas expression of *UQCRB* decreases with aging. DNA methylation is associated with an age-related decrease in gene expression in human muscle in some, but not all, genes. Changes in the DNA methylation and expression of all genes associated with SNPs (amplified and deleted genes) were observed in ovarian cancer tumors. In addition, tumor suppressor and oncogenic features of these modalities and correlation analysis with expression were determined. Genes showing a strong correlation for methylation-dependent expression changes exhibited varying copy number aberrations.<sup>46</sup>

A previous study demonstrated the extent of genomic and epigenetic alterations for known tumor suppressors and oncogenes and used these defined features to identify potential ovarian cancer gene candidates.<sup>42</sup> DNA methylation alterations are a significant feature in the cancer genome.<sup>47,48</sup> SNP amplification and promoter hypomethylation may play a role

in the overexpression of oncogenes. By contrast, a particularly well-known oncogene may be deleted in a specific cancer, decreasing its pathogenic role within that cancer or an individual sample. Interestingly, for highly amplified genes, a high level of methylation accompanied by low expression indicates altered tumor suppressor function in the cancer cell. In highly amplified genes, low-level methylation and high expression indicate oncogenic features in a cancer cell. As one of the largest epigenetic studies of CRC, as well as the first investigation of *UQCRB* methylation in CRC, our findings highlight the potential for *UQCRB* methylation as a biomarker of CRC.

Recent reports have revealed genetic variations in *UQCRB* in several cancers, including hepatocellular carcinoma, ovarian cancer, pancreatic ductal adenocarcinoma and CRC. By contrast, our *UQCRB* study, which used multiple tools, was the first study of CRC. In addition, numerous studies detected *UQCRB* expression at the gene level, whereas our study focused on its expression at the gene and protein levels. In addition, we detected *UQCRB* SNPs and conducted epigenetic analysis at the gene level. Our results may be useful for predictive diagnosis in which *UQCRB* expression correlates with CRC patients' clinical features. However, the precise mechanism by which *UQCRB* predicts CRC development remains unclear. Thus, further studies, including overexpression and knock-down of *UQCRB* expression in CRC cells, are needed to explore the mechanism by which *UQCRB* is involved in the development and progression of CRC and its exact regulating pathway *in vitro* and *in vivo*.

We previously reported that *UQCRB* plays a role in angiogenesis and that the developed cell-permeable PTD-*UQCRB* can be utilized as a pro-angiogenic agent.<sup>49</sup> In addition, we provided a molecular basis for *UQCRB*-related biological processes and revealed the potential key roles of *UQCRB* in angiogenesis and mitochondria-mediated metabolic disorders.<sup>50</sup> In this study, we found that *UQCRB* expression and CNV were significantly upregulated in CRC tissues compared to non-tumor tissues and were associated with clinical stage and pathologic differentiation. Representative CRC biomarkers discovered during the recent years continue to be closely examined: MSI, chromosome 18q loss of heterozygosity (18qLOH), *p53*, *KRAS*, *BRAF*, *NRAS*, *PIK3CA* mutations, *PTEN* expression, *UGT1A1* gene polymorphism and ezrin protein. Mutations in the tumor suppressor gene *p53* (chromosome region 17p13) occur in 50–70% of all CRC.<sup>51</sup> Together with these known CRC biomarkers, multivariate analyses of this study demonstrated that *UQCRB* can be used as a novel prognostic biomarker of human CRC.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### ACKNOWLEDGEMENTS

This work was partly supported by grants from the National Research Foundation of Korea, which is funded by the Korean government (MSIP; 2012M3A9D1054520, 2015K1A1A2028365,

2015M3A9B6027818, 2015M3A9C4076321); the Brain Korea 21 Plus Project, Republic of Korea; and the Seoul Medical Science Institute in 2015. The transcriptome analysis data of UQCRB overexpressing cells are available at Korean BioInformation Center (KBR20171018\_0000001~KBR20171018\_0000336).

- 1 Smith RA, Andrews K, Brooks D, DeSantis CE, Fedewa SA, Lortet-Tieulent J *et al*. Cancer screening in the United States, 2016: a review of current American Cancer Society Guidelines and current issues in cancer screening. *CA Cancer J Clin* 2016; **66**: 96–114.
- 2 Kim DH. Risk factors of colorectal cancer. *J Korean Soc Coloproctol* 2009; **25**: 356–362.
- 3 Yang SK, Yun S, Kim JH, Park JY, Kim HY, Kim YH *et al*. Epidemiology of inflammatory bowel disease in the Songpa-Kangdong district, Seoul, Korea, 1986–2005: a KASID study. *Inflamm Bowel Dis* 2008; **14**: 542–549.
- 4 Park HC, Shin A, Kim BW, Jung KW, Won YJ, Oh JH *et al*. Data on the characteristics and the survival of Korean patients with colorectal cancer from the Korea central cancer registry. *Ann Coloproctol* 2013; **29**: 144–149.
- 5 DiMauro S, Schon EA. Mitochondrial respiratory-chain diseases. *N Engl J Med* 2003; **348**: 2656–2668.
- 6 Heo HJ, Kim HK, Youm JB, Cho SW, Song IS, Lee SY *et al*. Mitochondrial pyruvate dehydrogenase phosphatase 1 regulates the early differentiation of cardiomyocytes from mouse embryonic stem cells. *Exp Mol Med* 2016; **48**: e254.
- 7 Crivellone MD, Wu M, Tzagoloff A. Assembly of the mitochondrial membrane system. Analysis of structural mutants of the yeast coenzyme QH<sub>2</sub>-cytochrome C reductase complex. *J Biol Chem* 1988; **263**: 14323–14333.
- 8 Berry EA, Guergova-Kuras M, Huang LS, Crofts AR. Structure and function of cytochrome *bc* complexes. *Annu Rev Biochem* 2000; **69**: 1005–1075.
- 9 Darrouzet E, Moser CC, Dutton PL, Daldal F. Large scale domain movement in cytochrome *bc*<sub>1</sub>: a new device for electron transfer in proteins. *Trends Biochem Sci* 2001; **26**: 445–451.
- 10 Trumpower BL. The protonmotive Q cycle. Energy transduction by coupling of proton translation by the cytochrome *bc*<sub>1</sub> complex. *J Biol Chem* 1990; **265**: 11409–11412.
- 11 Lonlay PD, Valnot I, Barrientos A, Gorbatyuk M, Tzagoloff A, Taanman JW *et al*. A mutant mitochondrial respiratory chain assembly protein causes complex III deficiency in patients with tubulopathy, encephalopathy and liver failure. *Nat Genet* 2001; **29**: 57–60.
- 12 Zara V, Conte L, Trumpower BL. Biogenesis of the yeast cytochrome *bc*<sub>1</sub> complex. *Biochim Biophys Acta* 2009; **1793**: 89–96.
- 13 Hildenbeutel M, Hegg EL, Stephan K, Gruschke S, Meunier B, Ott M. Assembly factors monitor sequential hemylation of cytochrome *b* to regulate mitochondrial translation. *J Cell Biol* 2014; **205**: 511–524.
- 14 Suzuki H, Hosokawa Y, Toda H, Nishikimi M, Ozawa T. Cloning and sequencing of a cDNA for human mitochondrial ubiquinone-binding protein of complex III. *Biochem Biophys Res Commun* 1988; **156**: 987–994.
- 15 Malaney S, Heng HH, Tsui LC, Shi XM, Robinson BH. Localization of the human gene encoding the 13.3-kDa subunit of mitochondrial complex III (UQCRB) to 8q22 by in situ hybridization. *J Cytogenet Genome Res* 1996; **73**: 297–299.
- 16 Mulder W, Scholten IH, van Roon H, Grivell LA. Isolation and characterisation of the linked genes APA2 and QCR7, coding for Ap4A phosphorylase II and the 14 kDa subunit VII of the mitochondrial *bc*<sub>1</sub>-complex in the yeast *Kluyveromyces lactis*. *Biochim Biophys Acta* 1994; **1219**: 719–723.
- 17 Hemrika W, Jong MD, Berden JA, Grivell LA. The C-terminus of the 14-kDa subunit of ubiquinol-cytochrome-c oxidoreductase of the yeast *Saccharomyces cerevisiae* is involved in the assembly of a functional enzyme. *Eur J Biochem* 1994; **220**: 569–576.
- 18 Haut S, Brivet M, Touati G, Rustin P, Lebon S, Garcia-Cazorla A *et al*. A deletion in the human QP-C gene causes a complex III deficiency resulting in hypoglycemia and lactic acidosis. *Hum Genet* 2003; **113**: 118–122.
- 19 Cho YS, Jung HJ, Seok SH, Payumo AY, Chen JK, Kwon HJ. Functional inhibition of UQCRB suppresses angiogenesis in zebrafish. *Biochem Biophys Res Commun* 2013; **433**: 396–400.
- 20 Jung HJ, Shim JS, Lee J, Song YM, Park KC, Choi SH *et al*. Terpestacin inhibits tumor angiogenesis by targeting UQCRB of mitochondrial complex III and suppressing hypoxia-induced reactive oxygen species production and cellular oxygen sensing. *J Biol Chem* 2010; **285**: 11584–11595.
- 21 Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> method. *Methods* 2001; **25**: 402–408.
- 22 Huang Y, Yu H, Lei H, Xie C, Zhong Y. Matrix metalloproteinase 7 is a useful marker for 5-fluorouracil-based adjuvant chemotherapy in stage II and stage III colorectal cancer patients. *Med Oncol* 2014; **31**: 824.
- 23 Pouligiannis G, Ichimura K, Hamoudi RA, Luo F, Leung SY, Yuen ST *et al*. Prognostic relevance of DNA copy number changes in colorectal cancer. *J Pathol* 2010; **220**: 338–347.
- 24 Lascorz J, Bevier M, Schonfels WV, Kalthoff H, Aselmann H, Beckmann J *et al*. Polymorphisms in the mitochondrial oxidative phosphorylation chain genes as prognostic markers for colorectal cancer. *BMC Med Genet* 2012; **13**: 31.
- 25 Kim W, Kim E, Lee S, Kim D, Chun J, Park KH *et al*. TFAP2C-mediated upregulation of TGFBR1 promotes lung tumorigenesis and epithelial-mesenchymal transition. *Exp Mol Med* 2016; **48**: e273.
- 26 Fernández-Vizarrá E, Zeviani E. Nuclear gene mutations as the cause of mitochondrial complex III deficiency. *Front Genet* 2015; **6**: 134.
- 27 Blazquez A, Gil-Borlado MC, Moran M, Verdu A, Cazorla-Calleja MR, Martin MA *et al*. Infantile mitochondrial encephalomyopathy with unusual phenotype caused by a novel *BCS1L* mutation in an isolated complex III-deficient patient. *Neuromuscul Disord* 2009; **19**: 143–146.
- 28 De Meirleir L, Seneca S, Damis E, Sepulchre B, Hoorens A, Gerlo E *et al*. Clinical and diagnostic characteristics of complex III deficiency due to mutations in the *BCS1L* gene. *Am J Med Genet* 2003; **121A**: 126–131.
- 29 Borisov VB. Mutations in respiratory chain complexes and human diseases. *Ital J Biochem* 2004; **53**: 34–40.
- 30 Owens KM, Kulawiec M, Desouki MM, Vanniarajan A, Singh KK. Impaired OXPHOS complex III in breast cancer. *PLoS ONE* 2011; **11**: e23846.
- 31 Schildgen V, Wulfert M, Gattermann N. Impaired mitochondrial gene transcription in myelodysplastic syndromes and acute myeloid leukemia with myelodysplasia-related changes. *Exp Hematol* 2011; **39**: 666–675.
- 32 Zou KH, O'Malley AJ, Mauri L. Receiver-operating characteristic analysis for evaluating diagnostic tests and predictive models. *Circulation* 2007; **115**: 654–657.
- 33 Mahjabeen I, Kayani MA. Loss of mitochondrial tumor suppressor genes expression is associated with unfavorable clinical outcome in head and neck squamous cell carcinoma: Data from retrospective study. *PLoS ONE* 2016; **11**: e0146948.
- 34 Kim HS, Patel K, Muldoon-Jacobs K, Bisht KS, Aykin-Burns N, Pennington JD *et al*. SIRT3 is a mitochondria-localized tumor suppressor required for maintenance of mitochondrial integrity and metabolism during stress. *Cancer Cell* 2010; **17**: 41–52.
- 35 Hirschev MD, Shimazu T, Goetzman E, Jing E, Scher B, Lombard DB *et al*. SIRT3 regulates mitochondrial fatty-acid oxidation by reversible enzyme deacetylation. *Nature* 2010; **464**: 121–125.
- 36 Jeong SM, Xiao C, Finley LW, Lahusen T, Souza AL, Pierce K *et al*. SIRT4 has tumor-suppressive activity and regulates the cellular metabolic response to DNA damage by inhibiting mitochondrial glutamine metabolism. *Cancer Cell* 2013; **23**: 450–463.
- 37 Seibold S, Rudroff C, Weber M, Galle J, Wanner C, Marx M. Identification of a new tumor suppressor gene located at chromosome 8p21.3e-22. *FASEB J* 2003; **17**: 1180–1182.
- 38 Chen J, Gu L, Ni J, Hu P, Hu K, Shi YL. MiR-183 regulates ITGB1P expression and promotes invasion of endometrial stromal cells. *Biomed Res Int* 2015; **2015**: 340218.
- 39 Izumi K, Zhang Z, Kaur M, Krantz ID. 12p microRNA expression in fibroblast cell lines from probands with Pallister-Killian syndrome. *Chromosome Res* 2014; **22**: 453–461.
- 40 Bowtell DD. The genesis and evolution of high-grade serous ovarian cancer. *Nat Rev Cancer* 2010; **10**: 803–808.
- 41 Lu TP, Lai LC, Tsai MH, Chen PC, Hsu CP, Lee JM. Integrated analyses of copy number variations and gene expression in lung adenocarcinoma. *PLoS ONE* 2011; **6**: e24829.
- 42 Wrzeszczynski KO, Varadan V, Byrnes J, Lum E, Kamalakaran S, Levine DA *et al*. Identification of tumor suppressors and oncogenes from genomic and epigenetic features in ovarian cancer. *PLoS ONE* 2011; **6**: e28503.

- 43 Jia HL, Ye QH, Qin LX, Budhu A, Forgues M, Chen Y *et al*. Gene expression profiling reveals potential biomarkers of human hepatocellular carcinoma. *Clin Cancer Res* 2007; **13**: 1133–1139.
- 44 Harada T, Chelala C, Crnogorac-Jurcevic T, Lemoine NR. Genome-wide analysis of pancreatic cancer using microarray-based techniques. *Pancreatology* 2009; **9**: 13–24.
- 45 Moarii1 M, Boeva1 V, Vert JP, Reyat F. Changes in correlation between promoter methylation and gene expression in cancer. *BMC Genomics* 2015; **16**: 873.
- 46 Ling C, Poulsen P, Simonsson S, Rönn T, Holmkvist J, Almgren P. Genetic and epigenetic factors are associated with expression of respiratory chain component NDUFB6 in human skeletal muscle. *J Clin Invest* 2007; **117**: 3427–3435.
- 47 Ting AH, McGarvey KM, Baylin SB. The cancer epigenome—components and functional correlates. *Genes Dev* 2006; **20**: 3215–3231.
- 48 Sadikovic B, Al-Romaih K, Squire JA, Zielenska M. Cause and consequences of genetic and epigenetic alterations in human cancer. *Curr Genomics* 2008; **9**: 394–408.
- 49 Chang JH, Jung HJ, Park HJ, Cho SW, Lee SK, Kwon HJ. Cell-permeable mitochondrial ubiquinol–cytochrome c reductase binding protein induces angiogenesis *in vitro* and *in vivo*. *Cancer Lett* 2015; **366**: 52–60.
- 50 Chang JH, Jung HJ, Jeong SH, Kim HK, Han J, Kwon HJ. A mutation in the mitochondrial protein UQCRB promotes angiogenesis through the generation of mitochondrial reactive oxygen species. *Biochem Biophys Res Commun* 2014; **433**: 290–297.
- 51 Lech G, Słotwiński R, Słodkowski M, Krasnodębski IW. Colorectal cancer tumour markers and biomarkers: recent therapeutic advances. *World J Gastroenterol* 2016; **22**: 1745–1755.



**This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>**

© The Author(s) 2017