

High prevalence of relapse in children with Philadelphia-like acute lymphoblastic leukemia despite risk-adapted treatment

Acute lymphoblastic leukemia (ALL) remains a leading cause of cancer-related death in children and young adults. Since the 1960s, improvements in the treatment of children with ALL have led to 10-year survival rates now exceeding 85%.¹ Philadelphia-like (Ph-like) ALL is characterized by a gene expression profile similar to that of *BCR-ABL1* positive (Ph+) ALL but lacking the *BCR-ABL1* oncogene and similarly, patients experience poor outcome.²⁻⁴ Ph-like ALL is associated with a range of genetic alterations, particularly rearrangements, which activate cytokine receptor and kinase signalling.²⁻⁴ In 2011, the Australian and New Zealand Children's Haematology/Oncology Group (ANZ-CHOG) completed enrolment of patients to a minimal residual disease (MRD) intervention clinical trial, known as ALL8 (*clinicaltrials.gov Identifier: ACTRN12607000302459*). Children were stratified to high-risk regimens based on several criteria, including treatment failure or high MRD at day 79.⁵ Overall, 46 children with precursor B-ALL relapsed, and surprisingly 72% (33/46) of these patients were classified as medium-risk.⁵ In this retrospective study of ALL8, the frequency of patients with Ph-like ALL and their attendant genomic lesions were studied, and clinical outcomes were compared to those of non Ph-like B-ALL

patients. The incidence of Ph-like ALL was 11.7%, where the majority of these children were reported to be of Caucasian ethnicity and stratified as being either standard- or medium-risk. Significantly, 57.8% (11/19) of Ph-like ALL patients subsequently relapsed compared to 16% (26/143) who were not Ph-like, with significantly inferior event-free and overall survival ($P < 0.0001$ and $P = 0.003$, respectively).

Six hundred and fifty-six patients aged between one and eighteen years were evaluated for eligibility on the ANZ-CHOG ALL8 trial from 2002-2011. Ethical approval was obtained from each institutional Human Research Ethics Committee and parents or legal guardians gave written, informed consent. Two hundred and forty-five patients, selected on the basis of sample accessibility, were available for Ph-like ALL screening (*Online Supplementary Figure S1*). The mean age at diagnosis (6.4 yrs vs. 5.6 yrs, $P = 0.02$) and higher white cell counts (WCC) ($P < 0.001$), were significantly different between those available for analysis and those patients excluded, with the studied group also having a higher number of patients over ten years of age (23% vs. 16%) (*Online Supplementary Table S1*). Risk groups were similar in each cohort and, overall, event-free and relapse-free survival were not significantly different between both groups (*Online Supplementary Figure S2*).

The criteria for stratification to the high-risk group, based upon Berlin-Frankfurt-Münster (BFM) protocols, were the presence of *BCR-ABL1* or *MLL t(4;11)* translocation; poor prednisolone response at day eight; failure to achieve

Table 1A. Age, MRD, risk stratification, relapse status (A), rearrangements and variants of patients (B) identified with Ph-like ALL and *P2RY8-CRLF2*.

Patient code	Age at diagnosis	Final risk stratification	MRD at day 79	Relapsed	Relapse free survival (years)	SCT
A2489	5.2	High	2x10 ²	Y	1.77	Y
A5258	14.9	High	6x10 ³	N	4.56	Y
A1781	3.5	Medium	Negative	Y	4.34	N
A5243	16.3	Medium	Pos<10 ⁻⁴	Y	2.56	N
A1516	8.2	Medium	Pos<10 ⁻⁴	N*	1.91	N
A4513	13.2	Medium	Negative	N	5.46	N
A1725	12.7	Medium	Pos<10 ⁻⁴	N	5.07	N
A2497	16.7	Medium	Pos<10 ⁻⁴	Y	1.92	Y
A3019	14.9	Medium	Negative	Y	0.96	N
A5164	6.2	Medium	Negative	Y	2.26	Y
A1702	3	Medium	Negative	N	10.86	N
A1747	3.1	Medium	Pos<10 ⁻⁴	Y	2.18	Y
A2173	8.2	Medium	Negative	Y	2.15	N
A3100	15	Medium	Negative	Y	1.06	N
A5416	5.5	Medium	Negative	Y	2.12	Y
A3086	11	Medium	Negative	N	8.07	N
A5428	1.5	Medium	Negative	Y	2.54	Y
A2481	5.9	Standard	Negative	N	2.94	N
A2005	13.4	Medium	Pos<10 ⁻⁴	N	5.78	N
A2273	5	Medium	Negative	Y	3.54	Y
A2426	10.7	Standard	Negative	Y	4.51	N
A2517	4.9	Standard	Negative	Y	2.35	Y
A3239	2.1	Medium	Negative	N	5.15	N
A3700	2.7	Medium	Negative	N	6.90	N
A4964	12.9	High	5x10 ⁻⁴	N**	1.76	Y

remission by day 33 or high MRD ($>5 \times 10^{-4}$) at day 79 (Table 1A). Standard- and medium-risk patients received the same standard BFM four-drug induction chemotherapy regimen including a prednisolone pre-phase and intrathecal methotrexate. In addition to the four-drug protocol, high-risk patients received a further three novel intensive blocks of chemotherapy followed by stem cell transplant (SCT) in most cases.⁵ All *BCR-ABL1* positive patients also received imatinib.

Determination of Ph-like ALL has differed between cohorts. European studies have favored the term *BCR-ABL1*-like ALL and have used hierarchical clustering (HC) of an Affymetrix gene expression array based on a probe set of 110 genes designed to detect major pediatric ALL subtypes.³ In contrast, US studies have used a TaqMan Low Density Arrays (TLDA) based approach consisting of either eight or fifteen genes selected by Prediction Analysis for Microarrays (PAM) analysis.^{7,8} While there is overlap, the HC model identifies a greater proportion of patients as having a *BCR-ABL1*-like signature, but this approach does not directly identify causative fusions.^{9,10} Based on the US approach, we have designed a custom TLDA using nine genes to identify patients with Ph-like ALL.^{7,11}

The TLDA was used according to manufacturer's instructions (Thermo Fisher Scientific, MA, USA) to deter-

mine Ph-like status. Genes were selected based upon prior reports³ with *CRLF2*, *PDGFRB*, *ABL1*, *ABL2* and *EPOR* also included to aid identification of potential fusions (*Online Supplementary Methods*). Reverse transcription polymerase chain reaction (RT-PCR) followed by Sanger sequencing was performed using a panel of 30 known fusions on all TLDA positive cases and those with high *CRLF2* gene expression.⁴ Cases with high *CRLF2* expression were also subjected to fluorescent *in situ* hybridization (FISH) to confirm *IGH-CRLF2* fusions. Illumina TruSeq stranded library preparation for messenger ribonucleic acid sequencing (mRNA seq) on the Illumina NextSeq or HiSeq platforms was performed on all TLDA positive and high *CRLF2* cases, with the sole exception being that of a case with low RNA quality (*Online Supplementary Methods*). Patients were classified as having Ph-like ALL if a sample was TLDA positive.

Cytokine or kinase activating lesions have been identified in the majority of childhood and adolescent/young adults (AYA) with Ph-like ALL.¹² Of the 245 childhood B-ALL patients evaluated, eight patients were identified as being *BCR-ABL1* positive and 75 had an *ETV6-RUNX1* fusion, leaving 162 available for Ph-like screening. Nineteen patients (11.7%) were identified as having Ph-like ALL, as determined by TLDA. Rearrangements were identified in 17/19 patients (Table 1B).

Table 1B. Age, MRD, risk stratification, relapse status (A), rearrangements and variants of patients (B) identified with Ph-like ALL and *P2RY8-CRLF2*.


Patient code	Rearrangement	TLDA	mRNA sequencing	Variants detected by mRNA and/or Sanger sequencing	<i>IKZF1</i> deletion
A2489	<i>EBF1-PDGFRB</i>	Pos	Y	no variants detected	del 4-7
A5258	<i>EBF1-PDGFRB</i>	Pos	Y	no variants detected	del 4-7
A1781	<i>SSBP2-JAK2</i>	Pos	Y	no variants detected	del 4-8
A5243	<i>PAX5-JAK2</i>	Pos	Y	no variants detected	del 2-7
A1516	<i>IGH-EPOR</i>	Pos	Y	<i>SREBF1</i> pV580M	del 2-7
A4513	<i>IGH-EPOR</i>	Pos	Y	<i>ABL2</i> pP608S, <i>SREBF1</i> pV580M, <i>TYK2</i> pG363S, <i>RUNX1</i> pL29S	del 4-7
A1725	<i>IGH-CRLF2</i>	Pos	Y	<i>CRLF2</i> pF232C, <i>CDKN2A</i> pA148T	del 2-8
A2497	<i>IGH-CRLF2</i>	Pos	Y	<i>CREBBP</i> pN1940S	del 2-8
A3019	<i>IGH-CRLF2</i>	Pos	Y	<i>JAK2</i> pR683S	del 4-7
A5164	<i>IGH-CRLF2</i>	Pos	Y	<i>CRLF2</i> pF232C	del 4-7
A1702	<i>P2RY8-CRLF2</i>	Pos	Y	<i>JAK2</i> pI682F	del 4-7
A1747	<i>P2RY8-CRLF2</i>	Pos	Y	<i>JAK2</i> pT875N, <i>FLT3</i> pD324N	del 2-7
A2173	<i>P2RY8-CRLF2</i>	Pos	Y	<i>JAK2</i> pR683S	del 4-7
A3100	<i>P2RY8-CRLF2</i>	Pos	Y	<i>NRAS</i> pG13D, <i>JAK3</i> pP132T	del 4-7
A5416	<i>P2RY8-CRLF2</i>	Pos	Y	<i>JAK2</i> pR683S, <i>SET2D</i> pM1080I	del 2-8
A3086	<i>PSMG1-ERG</i>	Pos	Y	<i>PAX5</i> pG266E, <i>BRAF</i> pA31V	None
A5428	<i>PAX5-ZNF521</i>	Pos	Y	<i>ABL1</i> pS972L, <i>ABL2</i> pK909R, <i>CREBBP</i> pV1243I, <i>SREBF1</i> pV580M, <i>TYK2</i> pI684S, <i>RUNX1</i> pE395A	del 4-7
A2481	Unknown	Pos	Y	no variants detected	None
A2005	Unknown	Pos	Y	<i>IKZF1</i> pN159Y, <i>PTK2B</i> pT65R, <i>BRAF</i> pD594G	None
A2273	<i>P2RY8-CRLF2</i>	Neg	N	Poor RNA	del 4-7
A2426	<i>P2RY8-CRLF2</i>	Neg	Y	<i>KRAS</i> pA146P	None
A2517	<i>P2RY8-CRLF2</i>	Neg	Y	<i>TYK2</i> pR568W, <i>RUNX1</i> pL29S	del 4-7
A3239	<i>P2RY8-CRLF2</i>	Neg	Y	<i>CREBBP</i> pN1940S	None
A3700	<i>P2RY8-CRLF2</i>	Neg	Y	no variants detected	None
A4964	<i>P2RY8-CRLF2</i>	Neg	Y	<i>NRAS</i> pG12D, <i>SET2D</i> pM1080I	del 2-8

*Secondary malignancy (AML) at 1.91 years; **died in remission. MRD: minimal residual disease; TLDA: Taqman low density array; mRNA: messenger ribonucleic acid; Pos: positive result; Neg: negative result; Y: yes; N: no; del: deletion of exons; SCT: stem cell transplant.


Previous reports have suggested 27-60% of Ph-like ALL patients harbor rearrangements of *CRLF2*, with 50% of these demonstrating concomitant mutations in *JAK1* or *JAK2*.^{4,13} Similarly, the majority of ALL8 Ph-like ALL cases (9/19, 47%) harbored *CRLF2* rearrangements (*CRLF2r*), with 77.7% (7/9) demonstrating concomitant *JAK2* or *CRLF2* mutations. While there is some conjecture, the majority of studies have demonstrated *CRLF2* overexpression is significantly associated with poor outcome.^{14,15} Importantly, 77.7% (7/9) of ALL8 Ph-like ALL patients with a *CRLF2r*, subsequently relapsed. Interestingly, a further six TLDA negative patients harbored *CRLF2r*. Of these

patients three relapsed, and two subsequently underwent SCT. A fourth patient received a SCT but died in remission. The remaining identified fusions included *EBF1-PDGFRB* and *IGH-EPOR* (n=2), *PAX5-JAK2*, *SSBP2-JAK2*, *PAX5-ZNF521* and *PSMG1-ERG* in one patient each (Table 1B).

IKZF1 deletions are shown to be significantly associated with relapse risk in both Ph+ and Ph-like ALL, with the frequency reported to be between 27% and 69% in Ph-like ALL cases.^{3,12,13} Of note, the ALL8 cohort included B-ALL patients in all risk stratifications, whereas other studies only included high-risk patients, potentially limiting comparisons between groups.^{3,12,13} Herein, *IKZF1* deletions



Number at risk	0	1	2	3	4	5
<i>BCR-ABL1</i>	8	8	8	8	8	4
<i>ETV6-RUNX1</i>	75	75	75	70	60	51
Ph-like ALL	19	19	16	9	8	5
B-ALL	143	141	132	116	105	84



Number at risk	0	1	2	3	4	5
<i>BCR-ABL1</i>	8	8	8	8	8	5
<i>ETV6-RUNX1</i>	75	75	75	75	69	59
Ph-like ALL	19	19	19	16	14	11
B-ALL	143	141	140	133	123	95

Figure 1. Children with Ph-like ALL have inferior survival outcomes compared to other B-ALL patients. Kaplan-Meier analysis with log-rank statistic of A) event free survival and B) overall survival at five years post diagnosis. Children with Ph-like ALL are shown in red, *BCR-ABL1*+ patients in orange, *ETV6-RUNX1* in green and the remaining B-ALL children in blue. The number of patients at risk for the different B-ALL subtypes is shown below the graph at each year.

were significantly associated with Ph-like ALL (84% vs. 14%, $P < 0.0001$). One Ph-like ALL patient, for whom a fusion was not identified, harbored an *IKZF1* p.N159Y mutation detected by mRNA seq and validated in genomic DNA by PCR and Sanger sequencing.

Most studies of patients with Ph-like ALL have demonstrated significantly inferior outcomes, which may be improved with treatment intensification.^{3,4,12} In contrast to patients enrolled on Total Therapy XV, a study of risk-directed therapy based on MRD wherein no significant differences in outcome were reported,¹³ the ALL8 Ph-like ALL cohort demonstrated significantly inferior event-free ($P < 0.0001$) and overall survival ($P = 0.003$; Figure 1).

On the ALL8 protocol, only two patients with Ph-like ALL had a final high-risk classification (both *EBF1-PDGFRB*) as a result of high MRD at day 79. One non Ph-like *P2RY8-CRLF2* patient was also reclassified to high-risk as a result of day 79 MRD. All three cases had *IKZF1* deletions; one relapsed and a second died in remission. On ALL8, 72% (8/11) of patients classified as Ph-like ALL relapsed within six months of completing their two years of maintenance therapy (average time to relapse 2.1 years). At five years, the overall survival of Ph-like cases was 78% (15/19), indicating that many patients were salvaged by further therapy or SCT,⁵ but their survival rate was still significantly inferior to other B-ALL sub-groups. Similar to that observed in Total Therapy XV, ALL8 patients with Ph-like disease were twice as likely to undergo SCT.¹³

Herein, we demonstrate that despite a risk adjusted treatment approach, there remained a high rate of relapse among children in the ANZCHOG ALL8 study who were retrospectively identified as Ph-like. Of note, the MRD risk stratification used in this protocol did not identify all Ph-like ALL cases as high-risk. Finally, rapid identification of Ph-like disease may guide therapeutic intervention with rationally targeted therapies based on patient specific driving genomic lesions. Tyrosine kinase inhibitors are increasingly utilized in patients with ABL-class fusions, with current and future trials likely to inform drug efficacy in the case of other targets.

Susan L. Heatley,^{1,2,3} Teresa Sadras,^{1,2} Chung H. Kok,^{1,2} Eva Nievergall,^{1,2} Kelly Quek,¹ Phuong Dang,¹ Barbara McClure,¹ Nicola Venn,⁴ Sarah Moore,⁵ Jeffrey Suttle,⁵ Tamara Law,⁴ Anthea Ng,^{3,6} Walter Muskovic,⁴ Murray D. Norris,⁴ Tamas Revesz,^{2,3,7} Michael Osborn,⁴ Andrew S. Moore,^{3,9,10} Ram Suppiah,¹¹ Chris Fraser,^{3,9} Frank Alvaro,^{3,12} Timothy P. Hughes,^{1,2,13} Charles G. Mullighan,¹⁴ Glenn M. Marshall,^{3,4,6,8,15} Luciano Dalla Pozza,^{3,6} David T. Yeung,^{1,2,13} Rosemary Sutton^{3,4,8,15} and Deborah L. White^{1,2,3,8,16}

¹Cancer Theme, South Australian Health & Medical Research Institute (SAHMRI), Adelaide, SA, Australia; ²Discipline of Medicine, University of Adelaide, SA, Australia; ³Australian and New Zealand Children's Haematology/Oncology Group (ANZCHOG); ⁴Molecular Diagnostics, Children's Cancer Institute, Sydney, NSW, Australia; ⁵Genetics and Molecular Pathology, SA Pathology, Adelaide, SA, Australia; ⁶Sydney Children's Hospital Network, Sydney, NSW, Australia; ⁷Women's & Children's Hospital, Adelaide, SA, Australia; ⁸Australian Genomic Health Alliance (AGHA); ⁹Oncology Services Group, Children's Health Queensland Hospital and Health Service, Brisbane, QLD, Australia; ¹⁰The University of Queensland Diamantina Institute and UQ Child Health Research Centre, The University of Queensland, Brisbane, QLD, Australia; ¹¹Mater Children's Hospital Brisbane, QLD, Australia; ¹²John Hunter Children's Hospital, Newcastle, NSW, Australia; ¹³Division of

Haematology, SA Pathology, Adelaide, SA, Australia; ¹⁴Department of Pathology, St Jude Children's Research Hospital, Memphis, TN, USA; ¹⁵School of Women and Children's Health, University of New South Wales, Sydney, NSW, Australia and ¹⁶Discipline of Paediatrics, University of Adelaide, SA, Australia

Acknowledgments: we thank the staff at the Tissue Bank of the Children's Research Institute for their assistance.

Funding: National Health and Medical Research Council, Australia (APP1057746, APP1044884); Channel 7 Children's Research Fund, Adelaide, SA, Australia; Leukaemia Foundation, Australia; Cancer Council of South Australia, Adelaide, SA, Australia; Beat Cancer, Adelaide, SA, Australia.

Correspondence: deborah.white@sahmri.com
doi:10.3324/haematol.2016.162925

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.

References

- Pui CH, Pei D, Campana D, et al. A revised definition for cure of childhood acute lymphoblastic leukemia. *Leukemia*. 2014; 28(12):2336-2343.
- Mullighan CG, Su X, Zhang J, et al. Deletion of *IKZF1* and prognosis in acute lymphoblastic leukemia. *N Engl J Med*. 2009;360(5):470-480.
- Den Boer ML, van Slegtenhorst M, De Menezes RX, et al. A subtype of childhood acute lymphoblastic leukaemia with poor treatment outcome: a genome-wide classification study. *Lancet Oncol*. 2009; 10(2):125-134.
- Roberts KG, Morin RD, Zhang J, et al. Genetic alterations activating kinase and cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*. 2012;22(2):153-166.
- Marshall GM, Dalla Pozza L, Sutton R, et al. High-risk childhood acute lymphoblastic leukemia in first remission treated with novel intensive chemotherapy and allogeneic transplantation. *Leukemia*. 2013;27(7):1497-1503.
- Karsa M, Dalla Pozza L, Venn NC, et al. Improving the identification of high risk precursor B acute lymphoblastic leukemia patients with earlier quantification of minimal residual disease. *PLoS One*. 2013; 8(10):e76455.
- Reshmi SC, Harvey RC, Roberts KG, et al. Targetable kinase gene fusions in high-risk B-ALL: a study from the Children's Oncology Group. *Blood*. 2017;129(25):3352-3361.
- Harvey RC, Kang H, Roberts KG, et al. Development and validation of a highly sensitive and specific gene expression classifier to prospectively screen and identify B-Precursor Acute Lymphoblastic Leukemia (ALL) patients with a Philadelphia chromosome-like ("Ph-like" or "BCR-ABL1-Like") signature for therapeutic targeting and clinical intervention. *Blood*. 2013;122(21):826.
- Boer JM, Marchante JR, Evans WE, et al. BCR-ABL1-like cases in pediatric acute lymphoblastic leukemia: a comparison between DCOG/Erasmus MC and COG/St. Jude signatures. *Haematologica*. 2015;100(9):e354-357.
- Boer JM, Steeghs EM, Marchante JR, et al. Tyrosine kinase fusion genes in pediatric BCR-ABL1-like acute lymphoblastic leukemia. *Oncotarget*. 2017;8(3):4618-4628.
- Roberts KG, Gu Z, Payne-Turner D, et al. High frequency and poor outcome of Philadelphia chromosome-like acute lymphoblastic leukemia in adults. *J Clin Oncol*. 2017;35(4):394-401.
- Roberts KG, Li Y, Payne-Turner D, et al. Targetable kinase-activating lesions in Ph-like acute lymphoblastic leukemia. *N Engl J Med*. 2014; 371(11):1005-1015.
- Roberts KG, Pei D, Campana D, et al. Outcomes of children with BCR-ABL1-like acute lymphoblastic leukemia treated with risk-directed therapy based on the levels of minimal residual disease. *J Clin Oncol*. 2014;32(27):3012-3020.
- Harvey RC, Mullighan CG, Chen IM, et al. Rearrangement of *CRLF2* is associated with mutation of *JAK* kinases, alteration of *IKZF1*, Hispanic/Latino ethnicity, and a poor outcome in pediatric B-progenitor acute lymphoblastic leukemia. *Blood*. 2010;115(26):5312-5321.
- van der Veer A, Waanders E, Pieters R, et al. Independent prognostic value of BCR-ABL1-like signature and *IKZF1* deletion, but not high *CRLF2* expression, in children with B-cell precursor ALL. *Blood*. 2013;122(15):2622-2629.