



Published in final edited form as:

Eur Eat Disord Rev. 2017 November ; 25(6): 432–450. doi:10.1002/erv.2553.

The Science Behind the Academy for Eating Disorders' Nine Truths About Eating Disorders

Katherine Schaumberg, Ph.D.¹, Elisabeth Welch, Ph.D.², Lauren Breithaupt, M.A.^{2,3}, Christopher Hübel, M.D., M.Sc.^{2,4}, Jessica H. Baker, Ph.D.¹, Melissa A. Munn-Chernoff, Ph.D.¹, Zeynep Yilmaz, Ph.D.¹, Stefan Ehrlich, M.D., Ph.D.^{5,6}, Linda Mustelin, M.D., Ph.D., MPH^{1,7}, Ata Ghaderi, Ph.D.⁸, J. Andrew Hardaway, Ph.D.¹, Emily C. Bulik-Sullivan, B.A.⁹, Anna M. Hedman, Ph.D.², Andreas Jangmo, M.Sc.², Ida A.K. Nilsson, Ph.D.¹⁰, Camilla Wiklund, M.Sc.², Shuyang Yao, M.Sc.², Maria Seidel, M.Sc.^{5,6}, and Cynthia M. Bulik, Ph.D.^{1,2,11}

¹Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

²Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

³Department of Psychology, George Mason University, Fairfax, VA, USA ⁴MRC Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's

College London, United Kingdom ⁵Division of Psychological & Social Medicine and

Developmental Neurosciences, Faculty of Medicine, Technische Universität Dresden, Germany

⁶Eating Disorder Treatment and Research Center, Department of Child and Adolescent

Psychiatry, Faculty of Medicine, Technische Universität Dresden, Germany ⁷Department of Public

Health and Institute for Molecular Medicine Finland FIMM, University of Helsinki, Helsinki, Finland

⁸Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden ⁹Department of

Medicine, University of North Carolina at Chapel Hill, NC, USA ¹⁰Department of Molecular

Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden and Center for Molecular

Medicine, Karolinska Hospital ¹¹Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Abstract

Objective—In 2015, the Academy for Eating Disorders (AED) collaborated with international patient, advocacy, and parent organizations to craft the “*Nine Truths About Eating Disorders.*” This document has been translated into over 30 languages and has been distributed globally to replace outdated and erroneous stereotypes about eating disorders with factual information. In this paper, we review the state of the science supporting the *Nine Truths.*

Methods—The literature supporting each of the *Nine Truths* was reviewed, summarized, and richly annotated.

Results—Most of the *Nine Truths* arise from well-established foundations in the scientific literature. Additional evidence is required to further substantiate some of the assertions in the

document. Future investigations are needed in all areas to deepen our understanding of eating disorders, their causes, and their treatments.

Conclusions—The “*Nine Truths About Eating Disorders*” is a guiding document to accelerate global dissemination of accurate and evidence-informed information about eating disorders.

Keywords

anorexia nervosa; bulimia nervosa; binge eating disorder; medical; mortality; genetics; environment

Eating disorders are serious mental illnesses that affect millions of individuals worldwide regardless of race, age, nationality, or sex and incur considerable personal, familial, and societal costs. The cumulative lifetime risk by age 80 of anorexia nervosa (AN), bulimia nervosa (BN) and binge-eating disorder (BED) approximates 4.6% (Hudson, Hiripi, Pope, & Kessler, 2007). Inclusion of subthreshold eating disorder behaviors raises this estimate to nearly 10%. Despite the prevalence and toll that eating disorders exact on society, we lack comprehensive understanding of the etiology of eating disorders. We face significant limitations in our ability to prevent, detect, and treat this class of disorders. Stigma surrounding eating disorders has overshadowed the field for decades and has perpetuated misconceptions about their causes, hampered efforts at advancing knowledge, and misdirected lay understanding of these conditions. Perhaps most importantly, stigma surrounding eating disorders has prevented those in need from seeking help (Ali et al., 2017).

In May 2015, the Academy for Eating Disorders (AED) and several international advocacy organizations issued a document entitled “*Nine Truths About Eating Disorders*” (<http://www.aedweb.org/index.php/25-press-releases/163-press-release-aed-releases-nine-truths-about-eating-disorders?quot>). The AED focused on presenting truths rather than dispelling myths to introduce empirical evidence into the general knowledge base about eating disorders. The document has been translated into over 30 languages and is being disseminated worldwide to transform perceptions and understanding of eating disorders. In this paper, we present an overview of the empirical foundation upon which the *Nine Truths* rest to foster a more accurate understanding of the current state of scientific knowledge about eating disorders for patients, families, professionals, and the public.

The truths span a broad literature. In addition to the review of empirical studies, we also attend to modern theoretical and conceptual models and authoritative reviews to evaluate the current state of the science behind the *Nine Truths*. For each truth, we present supporting statements and a strength of evidence rating (Low, Moderate, or High; see Supplementary Table S1 & S2). A detailed summary of the evidence is presented in Supplementary Table S2. In addition to these tables, online supplementary materials provide a rich source of background information and all references for the main text presented there as an annotated bibliography.

Truth #1: Many people with eating disorders look healthy, yet may be extremely ill

1.1 Eating disorders are associated with significant somatic, psychosocial, and psychological risk

Eating disorders are associated with somatic complications in multiple organ systems including the cardiovascular, gastrointestinal, musculoskeletal, dermatologic, endocrine, hematological, and neurological systems (Mehler & Brown, 2015; Mehler & Rylander, 2015; Thornton et al., 2017) as well as psychiatric comorbidities (see Supplementary Table S3). The more chronic and severe the eating disorder, the greater the likelihood of serious somatic complications (Westmoreland, Krantz, & Mehler, 2016). However, severe complications can emerge at any time during the course of illness (Westmoreland et al., 2016). Furthermore, eating disorders are associated with a number of measurable psychological and neurocognitive traits (see Supplementary Table S4 and Statement 4.2).

1.2 Most individuals with eating disorders do not appear emaciated

Weight loss is a defining characteristic of AN, but not BN or BED. In fact, eating disorders are present in all BMI categories (Duncan, Ziobrowski, & Nicol, 2017; Flament et al., 2015), and AN is less common than the combined prevalence of other eating disorder diagnoses (Kessler et al., 2013; Lindvall Dahlgren & Wisting, 2016; Qian et al., 2013). On average, the BMI of individuals with AN is lower than the BMI of those with BN, which is lower than the BMI of those with BED. Yet, restrictive eating disorders also occur among normal- and overweight individuals and individuals with BN and BED can be normal weight, overweight, or obese (see 5.4).

1.3 Somatic, psychosocial, and psychological manifestations and comorbidities of eating disorders may be difficult to detect

Many serious somatic complications of eating disorders are not readily visible to lay observers or recognizable to the affected individual (see Supplementary Table S3). Even experienced healthcare professionals have difficulty accurately identifying complications or may misattribute their causes (Currin et al., 2007b; Currin, Schmidt, & Waller, 2007a; Currin, Waller, & Schmidt, 2009; Gaudiani & Mehler, 2016). From a broad perspective, eating disorders have also been neglected in research and funding in proportion to the public health burden that they incur (Geil, Schmidt, Fernandez-Aranda & Zipfel, 2017; Schmidt et al., 2016).

Individuals with eating disorders may fail to report the psychological components of eating disorders or have poor insight into their level of impairment (Dalle Grave, Calugi, & Marchesini, 2008; Griffiths, Mond, Murray, & Touyz, 2015; Nordbø et al., 2012; Santonastaso et al., 2009; Vandereycken, 2006a; Vandereycken, 2006b). However, psychological features are often present, even if at milder levels (Carter & Bewell-Weiss, 2011) with some variation across cultures (Lee, Lee, Ngai, Lee, & Wing, 2001; Pike & Dunne, 2015) and in younger patients (Carter & Bewell-Weiss, 2011; Norris et al., 2014) (see Supplementary Tables S3 & S4). Signs and symptoms of an eating disorder should

always be taken seriously and not dismissed or minimized. Immediate attention is warranted, and a comprehensive evaluation should be the first step in treatment planning (American Psychiatric Association, 2006; Hay et al., 2014; National Collaborating Centre for Mental Health, 2004).

1.4 Most individuals with eating disorders do not enter treatment; those who do often do so many years into the course of illness

Epidemiological studies across the world indicate that only a minority of individuals who meet diagnostic criteria for eating disorders seek treatment (Hoek & van Hoeken, 2003; Hudson et al., 2007; Keski-Rahkonen et al., 2009; Kessler et al., 2013; Preti et al., 2009; Twomey, Baldwin, Hopfe, & Cieza, 2015). Eating disorders thus remain undetected, and, even when detected, may not be viewed as serious issues warranting medical intervention (Keel & Brown, 2010).

Truth #1: Summary and future research directions

Confidence ratings: Moderate (1.3) to High (1.1; 1.2; 1.4) (see Supplementary Table S2)

- A healthy appearance and failure to acknowledge the severity of these illnesses can delay help-seeking and detection by friends, family, providers, and even patients themselves.
- Longitudinal research is needed to identify early signs of somatic complications and psychiatric comorbidities in eating disorders. A better understanding of prodromal signs and the illness trajectory will enable early detection.
- *Understanding educational needs for physicians and other front-line providers is necessary for broad dissemination of screening and educational tools. For more information on addressing eating disorders in clinical practice, see the AED Guide to Recognition and Management of Eating Disorders (<http://www.aedweb.org/index.php/education/eating-disorder-information/eating-disorder-information-13>).*

Truth #2: Families are not to blame, and can be the patients' and providers' best allies in treatment

2.1 Biological risk factors contribute to the development of eating disorders

Modern etiological models of psychiatric illnesses consider the bidirectional risk between biology and environment (see Truth #4 for summary of biological factors). The assertion that parental characteristics or family dynamics are necessary and sufficient for the development of eating disorders (i.e., “families are to blame”) represents an historical and dated model of psychopathology and disregards modern etiological conceptualizations of psychiatric risk. Accordingly, the first part of this truth, “families are not to blame,” is empirically and logically justified. This does not imply that evaluation of family functioning in eating disorders is without merit, as such studies may provide actionable information for providers, caregivers, and patients.

2.2 Prototypical family interaction patterns that exist premorbidly among families with eating disorders have not been identified

A critical methodological issue continues to plague studies of family functioning in eating disorders. Most studies are correlational/differential in nature, precluding causal interpretation. Moreover, the direction of causality has not been examined. Prospective longitudinal designs are necessary to determine whether interactions among family members exist premorbidly or are a consequence of the illness. Some prospective studies have investigated effects of parent and family functioning in predicting later eating disorder onset with mixed results. For example, some evidence suggests that parental factors predict later eating pathology (Johnson, Cohen, Kasen, & Brook, 2002; Nicholls & Viner, 2009; Shoebridge & Gowers, 2000); however, reviews have not identified consistent patterns of risk associated with parenting styles or family interactions (Campbell & Peebles, 2014; Eisler, 2005; Larsen, Strandberg-Larsen, Micali, & Andersen, 2015; le Grange, Lock, Loeb, & Nicholls, 2010; Strober & Humphrey, 1987; Yager, 1982). Indeed, greater family conflict, reduced parental alliance, and increased feelings of depression in families with a child suffering from AN might reflect an accommodation process in response to a severe and life-threatening condition (Sim et al., 2009). Investigations of parental factors have also been limited by lack of controls with other psychiatric disorders, measurement inconsistencies, and lack of statistical power. For example, certain adverse familial experiences such as sexual abuse may contribute to the risk of pathology in general, and are not eating disorder specific (Kendler et al., 2000).

2.3 Eating disorders place stress on families

Studies on the experience of caring for a patient with an eating disorder suggest a significant burden and negative impact on the health and well-being of caregivers—especially among mothers and partners (Anastasiadou, Medina-Pradas, Sepulveda, & Treasure, 2014; Kyriacou, Treasure, & Schmidt, 2008). Those caring for patients with AN have reported higher levels of distress than individuals caring for patients with psychoses (Treasure et al., 2001). Parents can initially perceive starvation to be deliberate, which evokes a strong emotional response, significant distress, and can lead to desperate responses in parents in the absence of clear guidance (Whitney et al., 2005). Attributions for these responses should consider the parent's desire to cease the starvation and save their child. Thus, assisting families in developing tools to deal effectively with an eating disorder is imperative. Distress associated with an eating disorder often extends beyond the identified patient. Stresses associated with having a psychiatrically ill child or partner, coupled with the responsibility for collaborating with providers in the treatment of individuals with eating disorders, underscore the importance of self-care for caregivers (Patel, Wheatcroft, Park, & Stein, 2002; Treasure & Nazar, 2016).

2.4 Family-based treatments have demonstrated effectiveness for the treatment of adolescent AN

Families and support systems are needed as patient allies during treatment (le Grange et al., 2010). The entire family is affected when dealing with chronic and severe illnesses such as AN. Familial organizational changes that emerge may serve to maintain AN and limit access

to adaptive resources the family possesses that are necessary to help overcome the eating disorder (Cook-Darzens, 2016; Eisler, 2005). Family-based treatment (FBT), whereby parents reassert control over the child's eating, is a promising approach to the treatment of adolescent AN and has some empirical support for the treatment of adolescent BN (Couturier, Kimber, & Szatmari, 2013; le Grange, Lock, Agras, Bryson, & Jo, 2015). FBT helps families recognize resources and knowledge they possessed prior to the onset of the disorder and re-implement them in the family system (Lock & le Grange, 2015). FBT is recommended by many national guidelines for the treatment of eating disorders in youth (Watson & Bulik, 2013) (see Supplementary Table S5).

The role of the family is also important for adults with eating disorders. Partners can be an asset in treatment of adults since they typically express a strong desire to help, yet fear that anything they do or say will inadvertently exacerbate the situation (Treasure & Nazar, 2016). Couple-based interventions for eating disorders leverage the power of relationships and engage the partner in the recovery process (Bulik, Baucom, Kirby, & Pisetsky, 2011; Kirby, Runfola, Fischer, Baucom, & Bulik, 2015; Schmidt et al., 2013). Initial results of couple-based interventions are promising and suggest that close support from a family member enhances treatment regardless of patient age. However, much of family and couple-based intervention research has focused on patients with AN; additional studies are required to confirm the benefit of engaging family members in the treatment of BN and BED (see Supplementary Table S5).

Truth #2: Summary and future research directions

Confidence ratings: Moderate (2.2; 2.3) to High (2.1; 2.4) (see Supplementary Table S2)

- Typical patterns of family functioning or structure that give rise to eating disorders have not been identified. Families are not to blame and in most cases can be the patients' and providers' best allies in treatment.
- *Reviews on family functioning in eating disorders point to the need for rigorous prospective designs to help understand how environmental variables, including family systems, may interact with biological risk (as discussed in Truth #7 & #8) to either heighten risk or buffer against the development of eating disorders.* (Larsen et al., 2015; Saltzman & Liechty, 2016).
- Eating disorders place stress on a family system, and future investigations that aim to reduce the burden on caregivers are necessary. Consideration of in-home care may be a useful direction for services.
- Families represent an important base of support for those in recovery, and the effectiveness of family-based treatments for adolescents highlights how parents and caregivers can be important allies in treatment. Future studies that build on this success by examining how families can be best integrated into care of older adolescents, adults, and those who binge eat are of great interest.

Truth #3: An eating disorder diagnosis is a health crisis that disrupts personal and family functioning

3.1 Eating disorders interfere with personal and family functioning. 3.2 Eating disorders produce financial burden. 3.3 In adolescence, eating disorders may lead to functional impairment and delays in healthy development. 3.4 In adulthood, eating disorders may interfere with intimate relationships, reproductive health, parenting, and health-related quality of life

Truth #3 is covered by statements in several other Truths. As discussed in Truth #1, an eating disorder represents a health crisis that affects every aspect of an individual's life. In addition to myriad psychiatric and somatic complications and comorbidities enumerated in Truth #1, eating disorders also lead to considerable psychological distress, as well as isolation, stigmatization, and difficulties with family and other interpersonal relationships (Ali et al., 2017; Caslini et al., 2016; Dimitropoulos, McCallum, Colasanto, Freeman, & Gadalla, 2016; van Langenberg, Sawyer, Le Grange, & Hughes, 2016). Further, eating disorders are associated with financial burden, delays in healthy development, functional impairment, and may interfere with social role functioning including intimate relationships, reproductive health, and parenting (see summaries in Supplementary Tables S2–4).

Truth #3: Summary and future research directions

Confidence ratings: Moderate (3.3; 3.4) to High (3.1; 3.2) (see Supplementary Table S2)

- Eating disorders clearly represent a health crisis (see Truth #1); the effects of which disrupt functioning beyond immediate complications of the eating disorder.
- Financial burden of eating disorders are significant, and they affect all areas of social and economic well-being, along with delaying or preventing healthy childhood and adolescent development.
- Future investigations that examine the true cost of eating disorders over the long-term are warranted. Longitudinal studies of eating disorders, including intervention studies, are encouraged to include secondary outcomes related to healthy development in youth, education, finances, employment, reproductive health, and overall quality of life.
- An empirical review of the literature on relationship, role functioning, and quality of life in eating disorders would advance understanding of how eating disorders influence these vital, but understudied, outcomes.

Truth #4: Eating disorders are not choices, but serious biologically influenced illnesses

4.1 Disordered eating behaviors can be guided by biological processes associated with automatic (unconscious) events

In vulnerable individuals, biological drives towards automaticity can provoke rigid habits to the point where individuals struggle to regain control over their dysregulated eating and physical activity (Steinglass & Walsh, 2016). For example, altered inhibitory control, the ability to refrain from engaging in prepotent automatic responses, has been shown across eating disorders subtypes (Collantoni et al., 2016; Galimberti, Martoni, Cavallini, Erzegovesi, & Bellodi, 2012) with the greatest support for bulimic subtypes (Lavagnino, Arnone, Cao, Soares, & Selvaraj, 2016; Wu, Hartmann, Skunde, Herzog, & Friederich, 2013) (see Supplementary Table S4 for a review of traits). Such findings are supported by a position paper that reviewed literature identifying alterations in neurobiological pathways related to reward and self-control associated with eating disorders (Wierenga et al., 2014). Further, a recent theoretical model identifies eating behaviors in AN as habitual behaviors, similar to compulsions in obsessive compulsive disorder, supported by case-control studies on neuropsychological and neuroimaging tasks (Godier et al., 2016; Steinglass & Walsh, 2016). Evidence from animal studies and human neuroimaging also supports some shared neurobiology in eating disorders and other habit-related disorders, including addiction (Kaye et al., 2013b; O'Hara, Campbell, & Schmidt, 2015).

4.2 Biologically-influenced, fundamental personality traits and cognitive styles are associated with eating disorders

Eating disorders are consistently associated with fundamental personality traits and cognitive styles. These traits are influenced by genetic factors, exist premorbidly, become exacerbated during acute stages of illness, persist after recovery, and/or may affect the prognosis of eating disorders. Some implicated traits are shared across disorders (e.g., weak central coherence, altered reward sensitivity, anxiety, difficulty with set shifting, altered interoceptive awareness), whereas others are more differentially associated with specific eating disorder phenotypes (e.g., harm avoidance in AN, negative urgency in BN) (see Supplementary Table S4 for overview of associated traits). Overall, identification of genetically influenced personality traits and cognitive styles may reveal core biological risk factors for the development of eating disorders.

4.3 Individuals with eating disorders may experience non-typical responses to eating and activity

Individuals with eating disorders may have distinct responses to energy restriction and food consumption. For example, individuals with AN may have a paradoxical response to negative energy balance (i.e., taking in less energy than one expends, (Bulik, 2016), such that caloric intake is associated with dysphoric mood (Frank, 2012), whereas caloric restriction evokes a calming, anxiolytic, or euphorogenic effect (Bulik, 2016; Kaye, 2008; Kaye, Wierenga, Bailer, Simmons, & Bischoff-Grethe, 2013a). Non-typical responses to other behaviors such as physical activity and purging (as both positively and negatively

reinforcing) are also reported in individuals with eating disorders (Berg et al., 2013; Giel et al., 2013; Kaye, 2008; Klein et al., 2010). Such processes highlight alterations from typical experiences of reinforcement as relevant to development and maintenance of eating disorders, and such patterns may be driven by variations in neurobiology.

4.4 Eating disorders are associated with dysregulation in neurotransmitter availability and function

Although the precise underlying neurobiology is not fully understood, findings of positron emission tomography (PET) and single-photon emission computed tomography (SPECT) implicate dysregulation in both dopaminergic (DA) and serotonergic (5-HT) systems in eating disorders (Culbert, Racine, & Klump, 2015; Kaye et al., 2013a; Kaye et al., 2013b; Kaye, 2008; Kessler, Hutson, Herman, & Potenza, 2016; Spies, Knudsen, Lanzenberger, & Kasper, 2015). These systems are central in rewarding aspects of food, motivation, executive functions, and the regulation of mood, satiety, and impulse control.

4.5 Brain structure and function differ between those with active eating disorders and unaffected individuals

Both human and animal studies have addressed the role of brain anatomy and function in eating disorder psychopathology through use of brain imaging techniques. Studies revealing deviations in structure, function, and activation in the brains of individuals with eating disorders are reviewed comprehensively in several publications (Frank, 2013; Frank, 2015a; Kaye, 2008; O'Hara et al., 2015; Seitz et al., 2014; Seitz, Herpertz-Dahlmann, & Konrad, 2016; Titova, Hjorth, Schiöth, & Brooks, 2013; Van den Eynde et al., 2012).

Structural neuroimaging studies in eating disorders have predominantly shown grey matter reductions in various brain regions that are most pronounced in patients with AN (Seitz et al., 2016). Associations with nutritional abnormalities have been repeatedly demonstrated and in AN volume reductions tend to quickly normalize with weight gain (Bernardoni et al., 2016; Seitz et al., 2016). Functional and structural neuroimaging studies in eating disorders provide evidence that aberrant frontostriatal neural circuitry may represent altered reward pathways, manifesting in impaired regulation of appetite, emotion, and self-control (Frank, 2015b; Friederich, Wu, Simon, & Herzog, 2013; Kaye, Wagner, Fudge, & Paulus, 2011; Kessler et al., 2016; Marsh et al., 2009; Marsh, Maia, & Peterson, 2009). Specifically, altered functioning of limbic regions together with either reduced or exaggerated 'top-down' cognitive control (via the prefrontal cortex) are seen as contributing to impulsive (e.g., BN, BED) or exaggerated self-control (e.g., AN) related symptoms/behaviors (Ehrlich et al., 2015; Friederich et al., 2013; Hege et al., 2015; Kaye & Strober, 2009; Kessler et al., 2016; King et al., 2016; Marsh et al., 2009; Sanders et al., 2015). Neuroimaging and behavioral findings suggestive of alterations in reward pathways have been shown across eating disorders (see Frank, 2015a for review). Findings are mixed regarding the direction of change and the subregions of the brain reward system, likely due to research design issues such as failure to control for nutritional and medication status, exercise, comorbidity, and inadequate sample sizes (Frank, 2015a).

The persistence of core eating disorder psychopathology may reflect not only preexisting neurobiological vulnerabilities, but also neuroadaptation (Treasure et al., 2015), whereby changes may occur in the brain as a consequence of prolonged eating disorder behaviors (e.g., binge eating or restriction). Adolescence, in particular, is associated with a host of neuronal changes, such as increased synaptogenesis, pruning, and myelination of frontal and limbic areas, which are involved in emotional processing and cognition (Benes, 1998; Blakemore & Choudhury, 2006; Tau & Peterson, 2010). A maturing brain may be particularly vulnerable to the insults caused by extreme food restriction or excessive exercise resulting in negative energy balance or highly variable energy consumption (binge-fast cycles).

Evidence from brain structure and function, though preliminary, advances support for the assertion that eating disorders are biologically influenced. Brain structure and function appears to be altered in the active disease state, though the exact nature and stability of differences requires further investigation. Even if brain structure and function differences only occur after an initial shift in eating behavior, these changes may highlight biologically-driven maintenance patterns that impede recovery.

4.6 Feeding and activity behavior is biologically regulated in animals

Animal models shed light on highly specific brain pathways implicated in eating disorder features, including restriction and binge eating. Controlled experiments have led to the development of animal models of hunger (Atasoy, Betley, Su, & Sternson, 2012) and binge eating (Murray, Tulloch, Chen, & Avena, 2015), providing evidence of neurobiological origins of eating disorders. In addition, an activity-based anorexia (ABA) rodent model highlights increased physical activity and reduced body weight in response to restricted food access in animals (Chowdhury, Chen, & Aoki, 2015). Using neural circuit-level approaches that enable activation or inhibition of anatomically and genetically defined brain pathways, like optogenetics and chemogenetics, multiple pathways have been identified that regulate different patterns of feeding behavior (Hardaway, Crowley, Bulik, & Kash, 2015; Sternson & Roth, 2014) (see Supplementary Table S7 for specific regions and nuclei). This approach elevates understanding of how discrete neural circuits control feeding and metabolism, and provides additional evidence of how feeding behavior may be biologically influenced. Further study is needed to determine whether these are therapeutic entry points into pathological models of eating disorders.

4.7 Endocrine changes are associated with eating disorder risk

The risk for eating disorders increases during reproductive milestones (e.g., puberty, pregnancy) and sex hormones play a role in this risk (Baker, Girdler, & Bulik, 2012; Klump, Keel, Sisk, & Burt, 2010). For example, AN in females typically develops around puberty and is rare before the pubertal transition. Earlier pubertal timing is also associated with increased eating disorder symptoms. Increases in estrogen at puberty are hypothesized to activate genes that influence eating disorder development (Culbert et al., 2015; Culbert, Racine, & Klump, 2016; Klump et al., 2010). The increased risk for eating disorder symptoms at puberty is not surprising given that puberty in females involves considerable

changes not only in sex hormones, but also in body composition and in neuropeptides that modulate metabolism (Loomba-Albrecht & Styne, 2009; Siervogel et al., 2003).

Pregnancy has also been suggested as both a risk and protective period for eating disorder symptoms. Women with acute AN and BN often report symptom improvement or remission during pregnancy, whereas pregnancy increases risk for relapse for those in remission from AN (Kimmel, Ferguson, Zerwas, Bulik, & Meltzer-Brody, 2016). Pregnancy may also mark a vulnerable time for BED onset (Bulik et al., 2007). Eating disorder symptoms fluctuate across the menstrual cycle in a manner that mirrors changes in sex hormones (Baker et al., 2012; Edler, Lipson, & Keel, 2007; Klump, Keel, Culbert, & Edler, 2008; Racine et al., 2012). Paralleling these findings, a direct association between diminishing estrogen and increasing progesterone levels and eating disorder symptoms has been observed (Edler et al., 2007; Klump et al., 2008). The menopause transition, which involves prolonged and erratic changes in sex hormones, may represent an additional vulnerability period for the development or re-emergence of an eating disorder (Baker & Runfola, 2016; Mangweth-Matzek et al., 2013).

Much less is known about the role of reproductive milestones and sex hormones in the risk for eating disorders in males. Some studies suggest that boys who experience either early or late puberty are at increased risk for eating disorder symptoms (Ricciardelli & McCabe, 2004). Testosterone may be a protective factor against eating disorder development, but findings are inconclusive (Baker et al., 2012).

In addition, aberrant blood and cerebrospinal fluid levels of various appetite-regulating peptides have been observed in individuals suffering from AN or BN (Monteleone & Maj, 2013). Most of these studies, however, are limited both by small sample sizes and their sampling process because plasma levels of appetite-regulating peptides may not reflect the concentrations in the central nervous system. Serum leptin levels have also been tied with eating disturbances. Serum leptin levels correspond with fat mass in healthy, energy-balanced humans (Hebebrand, Muller, Holtkamp, & Herpertz-Dahlmann, 2007). As would be expected due to their low BMI and fat mass, in acute stages of the illness, individuals with AN generally have low serum leptin levels (Föcker et al., 2011). The observed levels in AN are typically lower than those in BMI-matched healthy lean individuals, most likely due to differences in fat mass (Hebebrand et al., 2007). Intriguingly, hypoleptinemia in AN has also been associated with characteristic hyperactivity (Ehrlich et al., 2009; Holtkamp et al., 2006). Hypoleptinemia is considered to be a state biomarker for AN and together with BMI may represent a useful diagnostic test to distinguish constitutional thinness from AN (Föcker et al., 2011). Additional endocrine changes observed in eating disorders are presented in Supplementary Table S3.

Truth #4: Summary and future research directions

Confidence ratings: Moderate (4.3; 4.4; 4.7); Moderate to High (4.1); High (4.2, 4.5; 4.6) (see Supplementary Table S2)

- The precise nature of underlying biological signatures is an active area of investigation and evidence in support of Truth #4 is accumulating rapidly. In-

depth work concentrating on personality traits, cognition, neurobiology, brain anatomy and function, endocrinology, genomics and other -omics (see Truths #7 and 8) contributes to improved understanding of the biological underpinnings of eating disorders.

- Future research directions for this truth include:
 - examining neuropsychologically-based treatment approaches and outcomes;
 - treatment matching based on phenotypic psychobiological profiles;
 - evaluation of childhood behavioral and neurobiological traits;
 - systematic reviews on altered response to food and exercise in eating disorders and brain function;
 - additional investigation of neurotransmitter availability and function in eating disorders using methods including postmortem brain analyses, measures of cerebrospinal fluid, PET imaging, and magnetic imaging spectroscopy;
 - basic science and animal research to further probe neural circuitry associated with eating disorder risk;
 - further examination of the role of longitudinal endocrine changes in eating disorders, including the menopause transition along with the role of hormonal changes in men's eating disorder risk.

Truth #5: Eating disorders affect people of all genders, ages, races, ethnicities, body shapes and weights, sexual orientations, and socioeconomic statuses

5.1 Eating disorders affect both males and females

Since research on eating disorders has historically focused on women, the nosology of eating disorders has evolved based on female symptom profiles (Anderson & Bulik, 2004) and normative data on males are lacking (see Supplementary Figure S1 for lifetime prevalence of eating disorders by sex). Evidence indicates that, when diagnosed with eating disorders, men and women often differ in clinical characteristics (Núñez-Navarro, et al., 2012; Welch, Gharedi, & Swenne, 2015), though psychiatric comorbidities appear common across gender (Ulfvebrand et al., 2015). Available evidence also suggests that males may also be less likely to seek treatment (Striegel, Bedrosian, Wang, & Schwartz, 2012), less likely to be diagnosed with an eating disorder even when presenting with identical symptoms as females (Currin et al., 2007a), and less likely to access treatment even with similar clinical severity (Austin et al., 2008). When they do access psychological treatment, men may be at higher risk for attrition than women (Agüera et al., 2017).

5.2 Eating disorders occur across the lifespan

The typical age of onset of both AN and BN is in adolescence or early adulthood (Currin, Schmidt, Treasure, & Jick, 2005; Keski-Rahkonen et al., 2007; Keski-Rahkonen et al., 2009; Smink, van Hoeken, & Hoek, 2012; Zerwas et al., 2015). Childhood-onset AN is seen clinically from about age 7 years upwards, whereas BN before puberty is quite rare (Nicholls & Bryant-Waugh, 2009). Likewise, BED often begins in late adolescence or early adulthood (Hudson et al., 2007; Kessler et al., 2013; Mustelin, Raevuori, Hoek, Kaprio, & Keski-Rahkonen, 2015; Preti et al., 2009), though some people report that they began binge eating early childhood—even before going on their first diet (Grilo & Masheb, 2000). Overall, however, BED commonly begins later than AN and BN, with new cases steadily arising up to age 40–60 years in the population (Hudson et al., 2007; Preti et al., 2009).

Eating disorders in midlife are either recurring or persisting early-onset disorders or new late-onset disorders (Baker & Runfola, 2016; Gagne et al., 2012; Peat, Peyrerl, & Muehlenkamp, 2008). Research on eating disorders diagnosed outside of the typical age range is limited. Current evidence suggests that late-onset eating disorders (defined as after age 25) are associated with less severe eating disorder symptomatology and associated psychopathology, but an increased likelihood of premorbid obesity (Bueno et al., 2014). Bulimic symptoms in particular are relatively common in midlife women (Baker et al., 2017; Gagne et al., 2012), with one study finding that, among 2,000 women above age 50, 13% endorsed an eating disorder symptom (Gagne et al., 2012). Although the etiology of midlife eating disorders remains poorly understood, life events such as divorce, loss of family members, or somatic illness could serve as triggers (Kally & Cumella, 2008; Peat et al., 2008), and pregnancy or menopause with accompanying biological changes may increase vulnerability for onset or recurrence of eating disorders (Baker & Runfola, 2016; Baker et al., 2017; Bulik et al., 2007; Peat et al., 2008). Very little is known about eating disorders in men in midlife and beyond.

5.3 Eating disorders occur in all races and ethnicities

A review of community studies from 30 countries found no systematic association between ethnicity/race and eating disorder occurrence (see Supplementary Figure S2) (Mitchison & Hay, 2014). Although eating disorders were initially considered to be limited to Western culture, accumulating evidence ties eating disorders more generally to economic development, urbanization, and industrialization across the globe (Pike, Dunne, & Addai, 2013; Pike, Hoek, & Dunne, 2014). Rising incidences of eating disorders have been reported in numerous countries, particularly in Asia and the Middle East (Pike & Dunne, 2015; Pike et al., 2014). In the United States, the prevalence of eating disorders in ethnic and racial minority groups is similar to non-Latino whites, while ethnic minority groups more frequently report binge-eating behavior compared with non-Latino whites (Marques et al., 2011). AN has been found to be somewhat less common among Black than White Americans (Pike et al., 2013; Striegel-Moore & Franko, 2003). Importantly, racial and ethnic minorities are underrepresented in specialist eating disorder services, possibly due to underdetection in primary care (Striegel-Moore et al., 2003).

5.4 Eating disorders occur in individuals of all shapes and sizes

Weight and BMI can vary substantially across the different types of eating disorders. In a sample of over 3,000 adolescents, eating disorders were present in all BMI categories (Flament et al., 2015). Restrictive eating disorders in normal- and overweight individuals are increasingly being acknowledged. The DSM-5 facilitates the diagnosis of atypical AN in individuals who meet all criteria for AN with the exception of low weight (American Psychiatric Association, 2013). This diagnosis is appropriate, for example, in individuals who begin at high weights and lose weight precipitously. A substantial portion of treatment-seeking adolescents with restrictive eating disorders have a history of overweight or obesity (Lebow, Sim, & Kransdorf, 2015), and there is a well-established relationship among dietary restriction, obesity, and eating disorders (Field et al., 2003; Neumark-Sztainer et al., 2006). In a review of clinical trials of BN, baseline BMI was most commonly in the normal range (Berkman et al., 2006), whereas community studies indicate that BN is prevalent in overweight and obese adolescents (Flament et al., 2015) and predicts weight gain over time (Fairburn, Cooper, Doll, Norman, & O'Connor, 2000; Micali et al., 2015). Individuals with BED are commonly overweight or obese (Hudson et al., 2007; Kessler et al., 2013), yet a substantial minority of individuals with BED are normal-weight, particularly early in the course of illness (Fairburn et al., 2000; Mustelin et al., 2015) (see Statement 1.2 for additional information on BMI and eating disorders).

5.5 Eating disorders are present across different sexual orientations and gender identities

Homosexual orientation is regarded as a risk factor for eating disorders in men: gay and bisexual men report more body dissatisfaction and disordered eating, and are more likely to be diagnosed with an eating disorder than heterosexual men (Brown & Keel, 2012; French, Story, Remafedi, Resnick, & Blum, 1996; Russell & Keel, 2002). In women, the evidence on sexual orientation and disordered eating is mixed. Lower body dissatisfaction among homosexual women have been observed in some, but not all studies (Alvy, 2013; French et al., 1996; Moore & Keel, 2003; Morrison, Morrison, & Sager, 2004). In a population-based cohort of adolescents, unhealthy weight control behaviors (e.g., laxative use, fasting, and vomiting) were significantly more prevalent among sexual minority males and females than in their heterosexual peers (Hadland, Austin, Goodenow, & Calzo, 2014).

Most research on eating-related pathology has focused on cisgender individuals (i.e., those whose gender identity matches the sex they were assigned at birth). A study of over 280,000 American college students indicated that transgender individuals may have particularly high eating disorder risk: 16% of transgender youth reported being diagnosed with an eating disorder in the past year, compared with 2% and 4% of cisgender sexual minority men and women, respectively (Diemer, Grant, Munn-Chernoff, Patterson, & Duncan, 2015). Similarly, a study of Canadian transgender youth found high rates of endorsement of disordered eating behaviors (Watson, Veale, & Saewyc, 2016). Body dissatisfaction, in particular, appears to contribute to eating disorder risk in trans individuals as body dissatisfaction is elevated in transgender people compared to cisgender peers (Jones, Haycraft, Murjan, & Arcelus, 2016), with one study finding that trans males had levels of body dissatisfaction comparable to cisgender men with eating disorders (Witcomb et al., 2015).

5.6 There is no consistent association between socioeconomic status and risk for eating disorders

Although higher parental education has been associated with increased risk of being diagnosed with an eating disorder in registry studies (Ahrén et al., 2013; Goodman, Heshmati, & Koupil, 2014), evidence suggests that this association may be genetically rather than socially mediated (Duncan et al., 2017). No consistent association has been observed between socioeconomic status and risk of eating disorders (Mitchison & Hay, 2014). In Australian population surveys, both binge eating and purging increased more in low-income than high-income individuals during a 10-year time period, suggesting an ongoing shift in the demographics of disordered eating (Mitchison, Hay, Slewa-Younan, & Mond, 2014).

Truth #5: Summary and future research directions

Confidence ratings: Moderate (5.5; 5.6); Moderate to High (5.2); High (5.1; 5.3; 5.4) (see Supplementary Table S2)

- No dominant pattern of age, body size, sexual orientation or gender identity, race, ethnicity, or socioeconomic status is associated with eating disorder risk.
- Providers should remain vigilant to eating disorders in all individuals regardless of demographic characteristics.
- Further research is needed in the following areas:
 - Studies on socioeconomic status and eating disorders that clarify inconsistent patterns observed and proposed genetic associations.
 - Longitudinal studies that consider weight trajectories as they relate to eating disorder symptom development, as it is clear that individuals may develop eating disorders from any pre-morbid weight.
 - Research on eating disorders among sexual minorities that aid in the development of targeted prevention and intervention efforts, specifically longitudinal studies that examine how sexual and gender identity development in youth may impact eating disorder risk

Truth #6: Eating disorders carry an increased risk for both suicide and medical complications

6.1 Eating disorders are associated with premature death

The most significant medical complication of an eating disorder is premature death. The standardized mortality ratio (SMR) associated with AN ranges between 5.9 and 6.2, meaning the risk of death for individuals with AN is up to 6.2 times greater than the risk in the general population, and the weighted annual mortality rate of AN is reported as 5.1 per 1000 person years (Chesney, Goodwin, & Fazel, 2014; Papadopoulos, Ekblom, Brandt, & Ekselius, 2009). Additionally, for females with AN between the ages of 15–24 years old, the mortality rate is 12 times higher than the death rate of all other causes of death (Klump, Bulik, Kaye, Treasure, & Tyson, 2009). Notably, AN also has one of the highest mortality

rates of any psychiatric illness (Chesney et al., 2014), and one in five deaths in AN is attributable to suicide (Arcelus, Mitchell, Wales, & Nielsen, 2011).

The mortality rate for BN is also significantly elevated relative to the general population, with meta-analyses estimating the SMR for BN to be 1.9 (Chesney et al., 2014). For those with BN, mortality risk may increase with severity (Huas et al., 2013). One clinical follow-up study in Finland found the all-cause mortality hazard ratio for BED to be 1.77 (0.60, 5.27) (Suokas et al., 2013). Though similar in effect size to reported SMRs for BN, this hazard ratio for BED was not significant. With the inclusion of BED in the DSM-5, more studies on epidemiology, course, and outcome of BED are likely.

6.2 Risk of suicide is elevated in eating disorders

The risk of suicide attempts is also elevated in eating disorders. In the Swedish population born between 1979 and 2001, the odds ratio (OR) of suicide attempts was estimated to be 5.3 (95% CI: 5.0, 5.5) for any eating disorder, meaning that the risk of suicide attempts in people with eating disorders is 5.3 times the risk in individuals without an eating disorder. The ORs for suicide were 4.4 (95% CI: 4.1, 4.7) for AN and 6.3 (95% CI: 5.7, 6.9) for BN (Yao et al., 2016). Similar relative risks have been reported in the Danish population for the period between 1989 and 2006 (Zerwas et al., 2015). A large clinical study found that 35.6% of eating disorder patients had attempted suicide at least once, and patients with binge eating and/or purging behaviors were associated with an elevated risk for suicide attempts compared with patients without such behaviors (Fedorowicz et al., 2007; Foulon et al., 2007). In Sweden, 13.6% of women with a lifetime history of BED had at least one lifetime suicide attempt (Pisetsky, Thornton, Lichtenstein, Pedersen, & Bulik, 2013; Runfola, Thornton, Pisetsky, Bulik, & Birgegård, 2014).

Based on a meta-analysis, the *suicide-specific* SMR is 18.1 (95% CI: 11.5, 28.7) for AN (Keshaviah et al., 2014). Among female AN patients in specialized care, this ratio could be as high as 31.0 (95% CI: 21.0, 44.0) (Preti, Rocchi, Sisti, Camboni, & Miotto, 2011). The suicide-specific SMR is reported as 7.5 (95% CI: 1.6, 11.6) for BN (Preti et al., 2011) and no deaths by suicide in individuals with BED were reported; however, more data for BED are expected to emerge as recognition and reporting of BED increases. Familial co-aggregation of eating disorders and suicide attempt has been observed in nationwide population data (Yao et al., 2016). Two studies from Australia (Wade, Fairweather-Schmidt, Zhu, & Martin, 2015) and Sweden (Thornton, Welch, Munn-Chernoff, Lichtenstein, & Bulik, 2016) have reported that the co-occurrence of eating disorders and suicide may be in part due to shared genetic factors.

Whereas women with disordered eating in the community may be more likely to attempt suicide than males (Davison, Marshall-Fabien, & Gondara, 2014), no sex differences have been found for the risk of suicide attempts or death by suicide in eating disorders (Yao et al., 2016).

Truth #6: Summary and future research directions

Confidence ratings: High (6.1;6.2) (see Supplementary Table S2)

- Increased risk of premature death, including suicide, among eating disorders is well established; however, little is known about the mechanism underlying this association.
- *Future investigations should consider why eating disorders specifically display increased risk for suicide and examine how psychobiological models of suicide (Anestis et al., 2016) may pertain to those with eating disorders, including how unique complications associated with eating disorders, such as nutritional status, may influence risk as proposed by these models.*

Truth #7: Genes and environment play important roles in the development of eating disorders

7.1 Eating disorders run in families

Family, twin, and genetic research has established that eating disorders run in families and genes play a role in this familial pattern (Yilmaz, Hardaway, & Bulik, 2015). Familial history of AN increases the risk of AN development fourfold compared with the general population (Steinhausen, Jakobsen, Helenius, Munk-Jørgensen, & Strober, 2015). Furthermore, AN, BN, and eating disorder not otherwise specified (EDNOS) track together in families, suggesting a lack of specificity (Lilenfeld et al., 1998; Strober, Freeman, Lampert, Diamond, & Kaye, 2000). BED also aggregates in families independent of obesity (Fowler & Bulik, 1997; Hudson et al., 2006). Twin studies cannot identify which genes influence risk, but they have identified a strong genetic contribution in AN, BN, and BED. Specifically, 48–74% of the total variance in liability to AN, 55–62% to BN, and 39–45% to BED is attributable to genetic factors (Yilmaz et al., 2015).

7.2 Genes play a role in eating disorder risk

Genome-wide association studies (GWAS), which scan the entire genome in a hypothesis-free manner, and related approaches such as exome sequencing and whole genome sequencing have rapidly accelerated the field. The Eating Disorders Working Group of the Psychiatric Genomics Consortium (PGC-ED) recently identified the first genome-wide significant locus for AN (Duncan et al., 2017) in an area that harbors genes previously implicated in type 1 diabetes and other autoimmune disorders. We expect this will mark an inflection point in genomic discovery if AN follows the same progression of findings as other psychiatric disorders such as schizophrenia, where increased sample size has led to fruitful genomic discovery (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). GWAS represent a starting point for genomic discovery, as post-GWAS science reveals causative biological pathways and the functional significance of implicated genes and epigenetic enhancer regions. No GWAS of BN or BED have been conducted to date. In addition to GWAS approaches, familial linkage analysis with whole-genome and exome sequencing has identified two potential missense mutations (Cui et al., 2013), which evidence a connection with eating-disordered behaviors in a recent mouse model (Lutter et al., 2017).

7.3 Environmental factors play a role in eating disorder risk

Genes do not act alone: environment plays an important role. Cross-sectional and longitudinal twin studies also indicate that nonshared environmental factors account for variance in eating disorder symptoms. Cultural pressure for thinness has been identified as a specific risk factor for eating disorders, and clinical trials of interventions that reduce thin-ideal internalization have led to reductions in eating disorder symptoms (Culbert et al., 2015). While thin-ideal internalization may have some genetic influence, one longitudinal twin study indicates that nonshared environmental influences were most important in the etiology of thin-ideal internalization (Suisman et al., 2014).

7.4 Only a small portion of individuals exposed to environmental risk develop eating disorders

Dieting, drive for thinness, and portion size escalation are widespread in industrialized countries and may represent risk scenarios for the development of eating disorders (Jacobi, Hayward, de Zwaan, Kraemer, & Agras, 2004; Steenhuis & Vermeer, 2009; Striegel-Moore & Bulik, 2007); however, despite nearly ubiquitous exposure, threshold illnesses are disproportionately rare. A current hypothesis is that individuals *genetically predisposed* to eating disorders are most vulnerable to societal pressures and environmental insults. Eating disorders are “complex traits,” meaning that multiple genetic and environmental factors—each of small to moderate effect—act together to increase risk. Genetic and environmental factors may not only act in an additive manner, but may co-act in other ways (see Truth #8).

Truth #7: Summary and future research directions

Confidence rating: Moderate (7.4); Moderate to High (7.1); High (7.2; 7.3) (see Supplementary Table S2)

- Genomic discovery in AN is accelerating rapidly, but work on BN and BED is woefully behind. Very large sample sizes (in the tens of thousands) are key to discovering genetic variants associated with risk, and global cooperation is underway to achieve such sample sizes.
- Advances in genetic methodology, coupled with increasing knowledge about environmental risk factors, will provide a more complete and accurate picture of eating disorder etiology.

Truth #8: Genes alone do not predict who will develop eating disorders

8.1 Eating disorders do not follow Mendelian transmission patterns

Inheritance patterns for eating disorders do not follow the traditional Mendelian patterns where variation in one gene results in one disorder (e.g., Huntington’s chorea). Rather, hundreds (or perhaps thousands) of genes act in concert and are influenced by environmental factors. An individual’s risk is a composite of the cumulative number of genetic and environmental risk and protective factors to which they are exposed. This pattern is supported by several case-control studies examining candidate genes that show inconsistent effects (see Yilmaz et al., 2015 for a review).

8.2 Many cases of eating disorders are sporadic, meaning there is no known family member who suffers from an eating disorder

Family studies indicate that the relative risk for eating disorders is higher in family members of affected individuals; however, the majority of affected individuals have no known affected family members (Bould et al., 2015; Steinhausen et al., 2015; Strober et al., 2000). This literature is limited in that eating disorder history among relatives may not be fully known or accurately captured.

8.3 Genes and environment may co-act to influence risk for eating disorders

Genes represent probabilities in all complex traits, such as eating disorders. Individuals with a high genetic susceptibility for disordered eating may be protected by other factors, whereas individuals at relatively low genetic risk may be burdened with cumulative or extreme environmental insults leading to possible eating disorder development despite their favorable genetic profile. Understanding the role that genes and environment play in eating disorders requires a deep acceptance of probability and of uncertainty.

Genes and environment may co-act to influence risk for eating disorders (Trace, Baker, Peñas-Lledó, & Bulik, 2013). First, in most families, parents and extended family provide both genes and shared environment, meaning that these two factors are confounded. Second, individuals with a stronger genetic susceptibility for eating disorders might be more sensitive to environmental factors (dieting, bullying, teasing, or overeating). Whereas many adolescents may try dieting, only for a few does it serve as an environmental trigger for an underlying genetic predisposition. Third, an individual who is genetically predisposed to traits associated with eating disorders (e.g., perfectionism, persistence, high physical activity) can seek out environments that may serve as triggers (e.g., sports that have a lean body type ideal, certain social media content) (Carrotte, Vella, & Lim, 2015; Giel et al., 2016; Rousselet et al., 2017). This phenomenon is known as an active gene-environment correlation (Plomin, DeFries, & Loehlin, 1977). Genetic research combined with ambulatory assessment may help understand how environmental influences affect risk for eating disorders by pinpointing specificity of risk factors.

Rigorous studies of gene-environment interaction in eating disorders are sparse. Some developmental twin studies have examined gene-environment interaction (Culbert et al., 2015). For example, contribution of genetic risk to the emergence of dysfunctional eating attitudes and disordered eating varies with developmental stage, with higher genetic effects observed in mid-to-late adolescence and mid-to-late puberty (Culbert et al., 2015; Culbert, Burt, McGue, Iacono, & Klump, 2009; Klump, Burt, McGue, & Iacono, 2007). More sophisticated analytic techniques that examine the *interplay* between genetic risk and family environment indicate that the fit between an individual's genotype and his or her family environment may be relevant for eating disorder risk (Culbert et al., 2015). For example, following a report of a rare missense mutation being associated with the development of eating disorders, Lutter et al. (2017) found that group (vs. individually) housed transgenic female mice displayed irregular feeding and anxiety behaviors, preliminarily revealing both sex-specific and gene by environment effects. In human studies, large samples using genome-wide and phenome-wide data are required for credible conclusions.

Additional ways in which genes and environment interact are via mechanisms collectively called epigenetics—the modification of DNA, RNA, or proteins by biological or environmental factors. These mechanisms alter gene expression without changing the DNA sequence. Importantly, epigenetic changes such as DNA methylation are tissue specific and can rarely be directly studied in the brain. Therefore, it is important to determine whether epigenetic changes seen in blood are good proxies for epigenetic changes in brain (Walton et al., 2016). While additional research is needed, prevention efforts represent a promising area for the application of epigenetic findings. For example, prevention efforts may be most effective for certain non-genetic risk factors or during particularly vulnerable time periods.

Preliminary epigenetic studies have reported changes in dopaminergic genes and genes for proopiomelanocortin (*POMC*), cannabinoid receptor 1 (*CNR1*, also referred to as *CBI*), atrial natriuretic peptide (*NPPA*, also referred to as *ANP*), alpha synuclein (*SNCA*), and oxytocin receptor (*OXTR*) (Ehrlich et al., 2010; Ehrlich et al., 2012; Frieling et al., 2007; Frieling et al., 2008; Frieling et al., 2010; Kim, Kim, Kim, & Treasure, 2014; Schroeder et al., 2012). If replicated, epigenetic findings could make important contributions to understanding the role of non-DNA elements in eating disorder susceptibility.

Truth #8: Conclusions and future research directions

Confidence ratings: Low (8.2;8.3); Moderate (8.1) (see Supplementary Table S2)

- A complex interplay between genetic and environmental factors underlies the development of eating disorders.
- Future research on genetic pathways and their interplay with environmental factors has the potential to provide key understanding of the multiple and nuanced facets by which individuals may develop eating pathology.
- In the short-term, large population-based studies with both genotypic and phenotypic information to probe gene-environment interactions, along with case-control studies to examine potential epigenetic effects represent key areas for advancing knowledge regarding complex risk patterns.

9. Truth #9: Full recovery from an eating disorder is possible. Early detection and intervention are important

9.1 A substantial portion of individuals with eating disorders achieve recovery

Full recovery from an eating disorder is not only possible, but indeed probable. A substantial portion of individuals with eating disorders achieve recovery, some without seeking treatment (Eddy et al., 2016; Keel & Brown, 2010; Steinhausen & Weber, 2009; Steinhausen, 2009). Five-year clinical recovery rates have been estimated at 67% for AN (Keski-Rahkonen et al., 2007) and 55% for BN (Keski-Rahkonen et al., 2009) in community samples, and by 10 years after eating disorder onset 70% of individuals are recovered (Berkman, Lohr, & Bulik, 2007). Although recovery is attainable, there is a lack of consensus on the exact definition of recovery, making it difficult to compare recovery rates across studies (Bardone-Cone et al., 2010; Emanuelli, Waller, Jones-Chester, & Ostuzzi,

2012). Traditionally, these definitions focus on physical and behavioral recovery. Physical recovery refers to the resumption and maintenance of a healthy body weight and a normalization of all physical parameters affected by the eating disorder, whereas behavioral recovery means the absence of eating-disorder related behaviors such as food restriction, binge eating, and purging. Psychological recovery, including the attainment of normal attitudes toward food and the body, is important yet often overlooked. It has been proposed that full recovery is achieved only when patients are indistinguishable from healthy controls on all eating disorder related measures, including psychological aspects (Bardone-Cone et al., 2010). Although this definition may seem stringent, it is attainable. Full recovery from an eating disorder is possible, and given that lingering eating disorder attitudes predict relapse (Helverskov et al., 2010), the psychological component of recovery is clinically relevant.

9.2 Early detection and intervention may improve prognosis

For some, recovery from an eating disorder is possible without treatment; however, early detection and intervention are preferred for all eating disorders (Treasure et al., 2015). For AN, a longer duration of illness before presentation for treatment is associated with poor outcome (Keel & Brown, 2010; Pike, 1998; Richard, Bauer, & Kordy, 2005), and the probability of recovering decreases as a function of duration of illness, irrespective of treatment (Pike, 1998). For BN, some studies find that a longer duration of illness is associated with poor outcome, whereas others observe that severity of illness and additional psychiatric comorbidities are more significant predictors of outcome (Steinhausen & Weber, 2009). However, in general, the sooner an eating disorder is identified and treatment can begin, the better prognosis there is for full recovery.

9.3 Effective psychological interventions for eating disorders exist. Many, but not all, patients benefit. & 9.4 Medication can be an effective treatment component for eating disorders

Treatment for an eating disorder typically includes psychological treatment and may include medication (Zipfel, Giel, Bulik, Hay, & Schmidt, 2015). For AN, weight restoration is an essential first step in treatment. Inpatient renourishment for AN is typically directed by clinical guidelines that advocate for a “low and slow” approach, due to concerns about refeeding syndrome (Solomon & Kirby, 1990). However, this approach is being challenged in favor of more aggressive renourishment techniques, leading to shorter hospital stays and a favorable safety profile (Garber et al., 2013; Madden et al., 2015; Redgrave et al., 2015). Once medical stabilization of an eating disorder is established, patients may step down to other levels of care.

The evidence base has been thoroughly reviewed for psychotherapeutic and medication interventions for eating disorders. Supplementary Tables S5 & S6 provide an overview of psychotherapeutic and medication treatments. Well-established psychological treatments include family-based treatment for adolescents with AN along with cognitive behavioral therapy and interpersonal psychotherapy for adults with BED or BN; several more recently developed psychotherapeutic approaches have some support. Emerging, sophisticated methodologies that are responsive to treatment progress [e.g. Sequential Multi-Phase

Randomized Trials (SMART) designs], and more effective at probing specific treatment components [e.g. Multiphase Optimization Strategies (MOST) designs] may assist in moving interventions forward (Collins, Murphy, Nair & Strecher, 2005; Lei et al., 2012). Harnessing large volumes of available data from behavioral monitoring devices is also a major task that, as it is undertaken, has great potential to improve treatment outcomes. Advances in eating disorder treatment will likely include not simply increasing the *number* of available treatment approaches with empirical support, but will instead focus on identifying and targeting specific processes with large amounts of biological, cognitive, and behavioral data to enhance individual-level outcomes. In addition to treatment of active eating disorders, several prevention programs have been well-established and widely disseminated in recent years (Watson et al., 2016), and approaches such as train-the-trainer and web-based dissemination methods may offer ways to further increase the reach of effective eating disorder prevention.

Truth #9: Summary and future research directions

Confidence Ratings: Low (9.4 for AN); Moderate (9.2); High (9.1; 9.3; 9.4 for BN/BED) (see Supplementary Table S2)

- Some evidence-based treatments have proven efficacy.
- Increasing understanding of the mechanisms underlying eating disorders will facilitate the development of more effective and personalized prevention and treatment options, eventually leading to increased recovery rates and shorter recovery times.
- *Recovery from eating disorders can and does occur at any age and for those who do not achieve complete remission, quality of life and somatic status may be improved, monitored, and stabilized* (Treasure, Stein, & Maguire, 2015).
- Future research goals include:
 - development of strategies for early detection and intervention
 - development of a provider’s toolbox that includes psychological and pharmacological interventions that are effective for a range of eating disorders in diverse populations
 - drug development or repurposing investigations to target core biological pathology of AN
 - studies of long-term efficacy of medication interventions for all eating disorders
 - studies of the effectiveness of medications for eating disorders in community settings.

General conclusion

We summarize the available literature that led to the development of the “*Nine Truths About Eating Disorders*.” Eating disorders are not choices and do affect individuals from all walks of life. They result from a combination of biological (including genetic) and environmental

factors. Eating disorders increase the risk for suicide and medical complications, and interrupt personal and family functioning. Families are not to blame and can be critical sources of support in recovery.

Clearly, additional work is needed to better understand risk factors, course of illness, and treatment of eating disorders. Important for advancing science in this area is the ability to remain flexible in thinking about causal factors and acknowledge accumulating evidence underlying these truths to eliminate misconceptions that have plagued the field for decades. In addition, providers should be mindful of the multitude of ways eating disorders can arise and be especially vigilant to signs of somatic and psychiatric complications resulting from AN, BN, and BED. As scientists, providers, patients, family, and friends, we need to continue educating others in the community about these truths in order to detect and treat eating disorders as soon as possible.

Yet, the science of this field cannot be advanced in the absence of appropriate investment and financial support from organizations worldwide that fund research. A 2015 blog post by the former director of the US National Institute of Mental Health, Thomas Insel, MD, revealed how woefully underfunded research on eating disorders was relative to the disability-adjusted life years associated with the illnesses (http://www.nimh.nih.gov/funding/funding-strategy-for-research-grants/white-paper_149362.pdf). Despite the dire morbidity and mortality statistics, eating disorders continue to be low-priority illnesses, we contend, in part due to long-standing misconceptions about their causes and consequences. Funding is required for larger more definitive collaborative studies to avoid the confusion that arises from conflicting results from small, underfunded, underpowered, and nonreplicated investigations. Far too often, such small-budget studies are all that investigators can afford to conduct. Further, as discussed in a recent commentary focused on European initiatives and needs, few formal training structures are currently in place that allow new investigators to develop expertise in emerging technologies and transdisciplinary approaches that hold the greatest promise for advancing scientific understanding of eating disorders (Schmidt et al., 2016).

Science is constantly evolving, and novel methods will enhance our ability to clarify the etiology of eating disorders and to develop scientifically informed and effective treatments for these debilitating illnesses. With adequate support for science, emerging information will facilitate the refinement of the *Nine Truths* and may in fact uncover new truths. Ultimately, it is our hope that dissemination of the *Nine Truths* will serve to reduce stigma and misunderstanding, and, via their impact on science and practice, reduce illness burden, improve quality of life, and eliminate mortality from eating disorders.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding acknowledgements: National Institute of Mental Health [5T32MH076694 (Bulik); K01MH109782 (Yilmaz); K01MH106675 (Baker)]; Swedish Research Council/Vetenskapsrådet; DNR: 538-2013-8864 (Bulik); National Science Foundation/Vetenskapsrådet GROW Fellowship (Breithaupt).

References

- Ahrén JC, Chiesa F, Koupil I, Magnusson C, Dalman C, Goodman A. We are family--parents, siblings, and eating disorders in a prospective total-population study of 250,000 Swedish males and females. *International Journal of Eating Disorders*. 2013; 46:693–700. <https://doi.org/10.1002/eat.22146>. [PubMed: 23740699]
- Ali K, Farrer L, Fassnacht DB, Gulliver A, Bauer S, Griffiths KM. Perceived barriers and facilitators towards help-seeking for eating disorders: A systematic review. *International Journal of Eating Disorders*. 2017; 50:9–21. <https://doi.org/10.1002/eat.22598>. [PubMed: 27526643]
- Alvy LM. Do lesbian women have a better body image? Comparisons with heterosexual women and model of lesbian-specific factors. *Body Image*. 2013; 10:524–534. <https://doi.org/10.1016/j.bodyim.2013.06.002>. [PubMed: 23927850]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5. Arlington, VA: American Psychiatric Publishing; 2013.
- American Psychiatric Association. Practice guideline for the treatment of patients with eating disorders. 2006. Retrieved from http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/eatingdisorders.pdf
- Anastasiadou D, Medina-Pradas C, Sepulveda AR, Treasure J. A systematic review of family caregiving in eating disorders. *Eating Behaviors*. 2014; 15:464–477. <https://doi.org/10.1016/j.eatbeh.2014.06.001>. [PubMed: 25064301]
- Anderson CB, Bulik CM. Gender differences in compensatory behaviors, weight and shape salience, and drive for thinness. *Eating Behaviors*. 2004; 5:1–11. <https://doi.org/10.1016/j.eatbeh.2003.07.001>. [PubMed: 15000949]
- Anestis JC, Anestis MD, Rufino KA, Cramer RJ, Miller H, Khazem LR, Joiner TE. Understanding the relationship between suicidality and psychopathy: An examination of the Interpersonal-Psychological Theory of Suicidal Behavior. *Archives of Suicide Research*. 2016; 20:349–368. <https://doi.org/10.1080/13811118.2015.1048399>. [PubMed: 26954611]
- Arcelus J, Mitchell AJ, Wales J, Nielsen S. Mortality rates in patients with anorexia nervosa and other eating disorders. A meta-analysis of 36 studies. *Archives of General Psychiatry*. 2011; 68:724–731. <https://doi.org/10.1001/archgenpsychiatry.2011.74>. Authors meta-analyzed 36 quantitative studies covering the mortality of eating disorders. The weighted mortality rates (i.e., deaths per 1000 person-years) were 5.1 for AN, 1.7 for BN, and 3.3 for EDNOS. The standardized mortality ratios were 5.86 for AN, 1.93 for BN, and 1.92 for EDNOS. One in 5 individuals with AN who died had committed suicide. [PubMed: 21727255]
- Atasoy D, Betley JN, Su HH, Sternson SM. Deconstruction of a neural circuit for hunger. *Nature*. 2012; 488:172–177. <https://doi.org/10.1038/nature11270>. Demonstration that cell and pathway specific optogenetic activation of a pathway from Agouti related peptide-expressing neurons in the arcuate nucleus to the paraventricular nucleus of the hypothalamus induces hunger. [PubMed: 22801496]
- Austin SB, Ziyadeh NJ, Forman S, Prokop LA, Keliher A, Jacobs D. Screening high school students for eating disorders: Results of a national initiative. *Prevention of Chronic Disease*. 2008; 5:A114.
- Baker JH, Girdler SS, Bulik CM. The role of reproductive hormones in the development and maintenance of eating disorders. *Expert Review of Obstetrics and Gynecology*. 2012; 7:573–583. <https://doi.org/10.1586/eog.12.54>. [PubMed: 23585773]
- Baker JH, Peterson CM, Thornton L, Brownley KA, Bulik CM, Girdler SS, ... Bromberger JT. Reproductive and appetite hormones and bulimic symptoms during midlife. *European Eating Disorders Review*. 2017; 25:188–194. <https://doi.org/10.1002/erv.2510>. Authors compared bulimic symptoms in premenopausal and perimenopausal midlife women and examined the association between these symptoms and reproductive and appetite hormones. No mean differences in bulimic

symptoms were observed between premenopause and perimenopause. A significant positive association between leptin and binge eating was observed. [PubMed: 28276114]

- Baker JH, Runfola CD. Eating disorders in midlife women: A perimenopausal eating disorder. *Maturitas*. 2016; 85:112–116. <https://doi.org/10.1016/j.maturitas.2015.12.017>. [PubMed: 26857889]
- Bardone-Cone AM, Harney MB, Maldonado CR, Lawson MA, Robinson DP, Smith R, Tosh A. Defining recovery from an eating disorder: Conceptualization, validation, and examination of psychosocial functioning and psychiatric comorbidity. *Behavior Research and Therapy*. 2010; 48:194–202. <https://doi.org/10.1016/j.brat.2009.11.001>.
- Benes FM. Brain development, VII. Human brain growth spans decades. *American Journal of Psychiatry*. 1998; 155:1489. <https://doi.org/10.1176/ajp.155.11.1489>. [PubMed: 9812107]
- Berg KC, Crosby RD, Cao L, Peterson CB, Engel SG, Mitchell JE, Wonderlich SA. Facets of negative affect prior to and following binge-only, purge-only, and binge/purge events in women with bulimia nervosa. *Journal of Abnormal Psychology*. 2013; 122:111–118. <https://doi.org/10.1037/a0029703>. [PubMed: 22985015]
- Berkman ND, Bulik CM, Brownley KA, Lohr KN, Sedway JA, Rooks A, Gartlehner G. Management of eating disorders. Evidence Report/Technology Assessment. 2006; 135:1–166.
- Berkman ND, Lohr KN, Bulik CM. Outcomes of eating disorders: A systematic review of the literature. *International Journal of Eating Disorders*. 2007; 40:293–309. <https://doi.org/10.1002/eat.20369>. [PubMed: 17370291]
- Bernardoni F, King JA, Geisler D, Stein E, Jaite C, Nätsch D, ... Ehrlich S. Weight restoration therapy rapidly reverses cortical thinning in anorexia nervosa: A longitudinal study. *Neuroimage*. 2016; 130:214–222. <https://doi.org/10.1016/j.neuroimage.2016.02.003>. Grey matter reductions which are typically found in acutely underweight anorexia nervosa patients were found to be reversed at a rate much faster than previously thought upon successful weight gain. [PubMed: 26876474]
- Blakemore SJ, Choudhury S. Development of the adolescent brain: Implications for executive function and social cognition. *Journal of Child Psychology and Psychiatry*. 2006; 47:296–312. <https://doi.org/10.1111/j.1469-7610.2006.01611.x>. [PubMed: 16492261]
- Bould H, Sovio U, Koupil I, Dalman C, Micali N, Lewis G, Magnusson C. Do eating disorders in parents predict eating disorders in children? Evidence from a Swedish cohort. *Acta Psychiatrica Scandinavica*. 2015; 132:51–59. <https://doi.org/10.1111/acps.12389>. In a sample of 158,697 children born in Stockholm county 1984–1995, Sweden, the authors tested whether the diagnosis of an eating disorder in a parent was predictive of a diagnosis of an eating disorder in the offspring. Due to low rates of eating disorders in males, analyses were restricted to females who were found to be at increased risk of being diagnosed with an eating disorder. [PubMed: 25572654]
- Brown TA, Keel PK. The impact of relationships on the association between sexual orientation and disordered eating in men. *International Journal of Eating Disorders*. 2012; 45:792–799. <https://doi.org/10.1002/eat.22013>. [PubMed: 22407531]
- Bulik CM. Towards a science of eating disorders: Replacing myths with realities: The fourth Birgit Olsson lecture. *Nordic Journal of Psychiatry*. 2016; 70:224–230. <https://doi.org/10.3109/08039488.2015.1074284>. [PubMed: 26355341]
- Bulik CM, Baucom DH, Kirby JS, Pisetky E. Uniting Couples (in the treatment of) Anorexia Nervosa (UCAN). *International Journal of Eating Disorders*. 2011; 44:19–28. <https://doi.org/10.1002/eat.20790>. Uniting Couples (in the treatment of) Anorexia Nervosa (UCAN) is a couple based intervention founded on cognitive-behavioral couple therapy principles. The authors discuss the delivery of the treatment and highlight its potential to enhance both retention and treatment outcome. [PubMed: 20063308]
- Bulik CM, Von Holle A, Hamer R, Knoph Berg C, Torgersen L, Magnus P, ... Reichborn-Kjennerud T. Patterns of remission, continuation and incidence of broadly defined eating disorders during early pregnancy in the Norwegian Mother and Child Cohort Study (MoBa). *Psychological Medicine*. 2007; 37:1109–1118. <https://doi.org/10.1017/S0033291707000724>. [PubMed: 17493296]
- Campbell K, Peebles R. Eating disorders in children and adolescents: State of the art review. *Pediatrics*. 2014; 134:582–592. <https://doi.org/10.1542/peds.2014-0194>. [PubMed: 25157017]

- Carrotte ER, Vella AM, Lim MS. Predictors of “liking” three types of health and fitness-related content on social media: A cross-sectional study. *Journal of Medical Internet Research*. 2015; 17:e205. <https://doi.org/10.2196/jmir.4803>. [PubMed: 26297689]
- Carter JC, Bewell-Weiss CV. Nonfat phobic anorexia nervosa: Clinical characteristics and response to inpatient treatment. *International Journal of Eating Disorders*. 2011; 44:220–224. <https://doi.org/10.1002/eat.20820>. [PubMed: 21400560]
- Caslini M, Crocamo C, Dakanalis A, Tremolada M, Clerici M, Carrà G. Stigmatizing attitudes and beliefs about anorexia and bulimia nervosa among Italian undergraduates. *Journal of Nervous and Mental Disease*. 2016; 204:916–924. <https://doi.org/10.1097/NMD.0000000000000606>. [PubMed: 27741080]
- Chesney E, Goodwin GM, Fazel S. Risks of all-cause and suicide mortality in mental disorders: A meta-review. *World Psychiatry*. 2014; 13:153–160. <https://doi.org/10.1002/wps.20128>. [PubMed: 24890068]
- Chowdhury TG, Chen YW, Aoki C. Using the activity-based anorexia rodent model to study the neurobiological basis of anorexia nervosa. *Journal of Visualized Experiments*. 2015; 105:e52927. <https://doi.org/10.3791/52927>.
- Collantoni E, Michelon S, Tenconi E, Degortes D, Tittton F, Manara R, ... Favaro A. Functional connectivity correlates of response inhibition impairment in anorexia nervosa. *Psychiatry Research*. 2016; 247:9–16. <https://doi.org/10.1016/j.psychres.2015.11.008>. [PubMed: 26655584]
- Cook-Darzens S. The role of family meals in the treatment of eating disorders: A scoping review of the literature and implications. *Eating and Weight Disorders*. 2016; 21:383–393. <https://doi.org/10.1007/s40519-016-0263-y>. [PubMed: 26932829]
- Couturier J, Kimber M, Szatmari P. Efficacy of family-based treatment for adolescents with eating disorders: A systematic review and meta-analysis. *International Journal of Eating Disorders*. 2013; 46:3–11. <https://doi.org/10.1002/eat.22042>. [PubMed: 22821753]
- Cui H, Moore J, Ashimi SS, Mason BL, Drawbridge JN, Han S, ... Lutter M. Eating disorder predisposition is associated with ESRRA and HDAC4 mutations. *Journal of Clinical Investigation*. 2013; 123:4706–4713. <https://doi.org/10.1172/JCI71400>. [PubMed: 24216484]
- Culbert KM, Burt SA, McGue M, Iacono WG, Klump KL. Puberty and the genetic diathesis of disordered eating attitudes and behaviors. *Journal of Abnormal Psychology*. 2009; 118:788–796. <https://doi.org/10.1037/a0017207>. [PubMed: 19899848]
- Culbert KM, Racine SE, Klump KL. Research Review: What we have learned about the causes of eating disorders - a synthesis of sociocultural, psychological, and biological research. *Journal of Child Psychology and Psychiatry*. 2015; 56:1141–1164. <https://doi.org/10.1111/jcpp.12441>. [PubMed: 26095891]
- Culbert KM, Racine SE, Klump KL. Hormonal factors and disturbances in eating disorders. *Current Psychiatry Reports*. 2016; 18:65. <https://doi.org/10.1007/s11920-016-0701-6>. The role of hormonal factors influencing eating disorders is still unknown. However, the strongest evidence for etiologic effects has emerged for ovarian hormones, suggesting that estradiol reduces food intake whereas progesterone and testosterone increase food intake. [PubMed: 27222139]
- Currin L, Schmidt U, Treasure J, Jick H. Time trends in eating disorder incidence. *British Journal of Psychiatry*. 2005; 186:132–135. <https://doi.org/10.1192/bjp.186.2.132>. [PubMed: 15684236]
- Currin L, Schmidt U, Waller G. Variables that influence diagnosis and treatment of the eating disorders within primary care settings: A vignette study. *International Journal of Eating Disorders*. 2007a; 40:257–262. <https://doi.org/10.1002/eat.20355>. [PubMed: 17167756]
- Currin L, Waller G, Schmidt U. Primary care physicians' knowledge of and attitudes toward the eating disorders: Do they affect clinical actions. *International Journal of Eating Disorders*. 2009; 42:453–458. <https://doi.org/10.1002/eat.20636>. [PubMed: 19115367]
- Currin L, Waller G, Treasure J, Nodder J, Stone C, Yeomans M, Schmidt U. The use of guidelines for dissemination of “best practice” in primary care of patients with eating disorders. *International Journal of Eating Disorders*. 2007b; 40:476–479. <https://doi.org/10.1002/eat.20385>. [PubMed: 17497711]

- Dalle Grave R, Calugi S, Marchesini G. Underweight eating disorder without over-evaluation of shape and weight: Atypical anorexia nervosa. *International Journal of Eating Disorders*. 2008; 41:705–712. <https://doi.org/10.1002/eat.20555>. [PubMed: 18523958]
- Davison KM, Marshall-Fabien GL, Gondara L. Sex differences and eating disorder risk among psychiatric conditions, compulsive behaviors and substance use in a screened Canadian national sample. *General Hospital Psychiatry*. 2014; 36:411–414. <https://doi.org/10.1016/j.genhosppsych.2014.04.001>. [PubMed: 24856439]
- Diemer EW, Grant JD, Munn-Chernoff MA, Patterson DA, Duncan AE. Gender identity, sexual orientation, and eating-related pathology in a national sample of college students. *Journal of Adolescent Health*. 2015; 57:144–149. <https://doi.org/10.1016/j.jadohealth.2015.03.003>. In a sample of nearly 300,000 U.S. college students, transgender and cisgender sexual minority young adults reported a higher prevalence of past-year self-reported eating disorder diagnosis and past-month compensatory behaviors (i.e., self-induced vomiting, diet pills, and laxatives) than their cisgender heterosexual peers. [PubMed: 25937471]
- Dimitropoulos G, McCallum L, Colasanto M, Freeman VE, Gadalla T. The effects of stigma on recovery attitudes in people with anorexia nervosa in intensive treatment. *Journal of Nervous and Mental Disease*. 2016; 204:370–380. <https://doi.org/10.1097/NMD.0000000000000480>. [PubMed: 26894311]
- Duncan, AE., Ziobrowski, HN., Nicol, G. The prevalence of past 12-month and lifetime DSM-IV eating disorders by BMI category in US men and women. *European Eating Disorders Review*. 2017. <https://doi.org/10.1002/erv.2503>. In a U.S. general population sample, lifetime eating disorder prevalence was 2.22% in men and 4.93% in women. The prevalence of any lifetime and past 12-month ED, binge eating disorder and recurrent binge eating was highest among obese individuals
- Duncan L, Yilmaz Z, Gaspar H, Walters R, Goldstein J, Antilla V, Bulik-Sullivan B, ... Bulik CM. Genome-wide association study reveals first locus for anorexia nervosa and metabolic correlations. *American Journal of Psychiatry*. (in press). First identified genome-wide significant locus for anorexia nervosa on chromosome 12 (rs4622308) in a region harboring a previously reported type 1 diabetes and autoimmune disorder locus. Significant positive genetic correlations observed between anorexia nervosa and schizophrenia, neuroticism, educational attainment, and high-density lipoprotein cholesterol, and significant negative genetic correlations were observed between anorexia nervosa and body mass index, insulin, glucose, and lipid phenotypes.
- Eddy KT, Tabri N, Thomas JJ, Murray HB, Keshaviah A, Hastings ER, ... Franko DL. Recovery from anorexia nervosa and bulimia nervosa at 22-year follow-up. *Journal of Clinical Psychiatry*. 2017; 78:184–189. <https://doi.org/10.4088/JCP.15m10393>. [PubMed: 28002660]
- Edler C, Lipson SF, Keel PK. Ovarian hormones and binge eating in bulimia nervosa. *Psychological Medicine*. 2007; 37:131–141. <https://doi.org/10.1017/S0033291706008956>. [PubMed: 17038206]
- Ehrlich S, Burghardt R, Schneider N, Broecker-Preuss M, Weiss D, Merle JV, ... Hebebrand J. The role of leptin and cortisol in hyperactivity in patients with acute and weight-recovered anorexia nervosa. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2009; 33:658–662. <https://doi.org/10.1016/j.pnpbp.2009.03.007>. [PubMed: 19296912]
- Ehrlich S, Geisler D, Ritschel F, King JA, Seidel M, Boehm I, ... Kroemer NB. Elevated cognitive control over reward processing in recovered female patients with anorexia nervosa. *Journal of Psychiatry and Neuroscience*. 2015; 40:307–315. Individuals recovered from anorexia nervosa showed elevated brain activity in the dorsolateral prefrontal cortex (DLPFC) as well as greater functional coupling between the DLPFC and the orbitofrontal cortex during the anticipation phase of a monetary reward paradigm. The findings are suggestive of elevated self-regulatory processes in response to rewarding stimuli in patients recovered from anorexia nervosa. [PubMed: 26107161]
- Ehrlich S, Walton E, Roffman JL, Weiss D, Puls I, Doehler N, ... Frieling H. Smoking, but not malnutrition, influences promoter-specific DNA methylation of the proopiomelanocortin gene in patients with and without anorexia nervosa. *Canadian Journal of Psychiatry*. 2012; 57:168–176. [PubMed: 22398003]
- Ehrlich S, Weiss D, Burghardt R, Infante-Duarte C, Brockhaus S, Muschler MA, ... Frieling H. Promoter specific DNA methylation and gene expression of POMC in acutely underweight and

- recovered patients with anorexia nervosa. *Journal of Psychiatric Research*. 2010; 44:827–833. <https://doi.org/10.1016/j.jpsychires.2010.01.011>. [PubMed: 20176366]
- Eisler I. The empirical and theoretical base of family therapy and multiple family day therapy for adolescent anorexia nervosa. *Journal of Family Therapy*. 2005; 27:104–131. <https://doi.org/10.1111/j.1467-6427.2005.00303.x>.
- Emanuelli F, Waller G, Jones-Chester M, Ostuzzi R. Recovery from disordered eating: Sufferers' and clinicians' perspectives. *European Eating Disorders Review*. 2012; 20:363–372. <https://doi.org/10.1002/erv.2159>. In a checklist study, individuals with eating difficulties and clinicians ranked factors associated with recovery. Domains included psychological-emotional-social, weight-controlling behaviors, non-life-threatening and life-threatening features, and evaluation of one's own appearance. Ill individuals and clinicians agreed on the ranking of importance of these factors, but those with eating disturbances, considered 'psychological-emotional-social' and 'evaluation of one's own appearance' criteria as more important to recovery than clinicians. [PubMed: 22411482]
- Fairburn CG, Cooper Z, Doll HA, Norman P, O'Connor M. The natural course of bulimia nervosa and binge eating disorder in young women. *Archives of General Psychiatry*. 2000; 57:659–665. <https://doi.org/10.1001/pubs.Arch Gen Psychiatry-ISSN-0003-990x-57-7-yoa9404>. [PubMed: 10891036]
- Fedorowicz VJ, Falissard B, Foulon C, Dardennes R, Divac SM, Guelfi JD, Rouillon F. Factors associated with suicidal behaviors in a large French sample of inpatients with eating disorders. *International Journal of Eating Disorders*. 2007; 40:589–595. <https://doi.org/10.1002/eat.20415>. [PubMed: 17607699]
- Field AE, Austin SB, Taylor CB, Malspeis S, Rosner B, Rockett HR, ... Colditz GA. Relation between dieting and weight change among preadolescents and adolescents. *Pediatrics*. 2003; 112:900–906. <https://doi.org/10.1542/peds.112.4.900>. [PubMed: 14523184]
- Flament MF, Henderson K, Buchholz A, Obeid N, Nguyen HN, Birmingham M, Goldfield G. Weight status and DSM-5 diagnoses of eating disorders in adolescents from the community. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2015; 54:403–411. e2. <https://doi.org/10.1016/j.jaac.2015.01.020>. [PubMed: 25901777]
- Föcker M, Timmesfeld N, Scherag S, Bühren K, Langkamp M, Dempfle A, ... Hebebrand J. Screening for anorexia nervosa via measurement of serum leptin levels. *Journal of Neural Transmission*. 2011; 118:571–578. <https://doi.org/10.1007/s00702-010-0551-z>. [PubMed: 21258826]
- Foulon C, Guelfi JD, Kipman A, Adès J, Romo L, Houdeyer K, ... Gorwood P. Switching to the bingeing/purging subtype of anorexia nervosa is frequently associated with suicidal attempts. *European Psychiatry*. 2007; 22:513–519. <https://doi.org/10.1016/j.eurpsy.2007.03.004>. [PubMed: 17482799]
- Fowler SJ, Bulik CM. Family environment and psychiatric history in women with binge-eating disorder and obese controls. *Behaviour Change*. 1997; 14:106–112.
- Frank GK. Altered brain reward circuits in eating disorders: Chicken or egg. *Current Psychiatry Reports*. 2013; 15:396. <https://doi.org/10.1007/s11920-013-0396-x>. [PubMed: 23963630]
- Frank GK. Advances in the diagnosis of anorexia nervosa and bulimia nervosa using brain imaging. *Expert Opinion in Medical Diagnosis*. 2012; 6:235–244. <https://doi.org/10.1517/17530059.2012.673583>.
- Frank GK. Recent advances in neuroimaging to model eating disorder neurobiology. *Current Psychiatry Reports*. 2015a; 17:559. <https://doi.org/10.1007/s11920-015-0559-z>. [PubMed: 25749747]
- Frank GK. Advances from neuroimaging studies in eating disorders. *CNS Spectrums*. 2015b; 20:391–400. <https://doi.org/10.1017/S1092852915000012>. [PubMed: 25902917]
- French SA, Story M, Remafedi G, Resnick MD, Blum RW. Sexual orientation and prevalence of body dissatisfaction and eating disordered behaviors: A population-based study of adolescents. *International Journal of Eating Disorders*. 1996; 19:119–126. [https://doi.org/10.1002/\(SICI\)1098-108X\(199603\)19:2<119::AID-EAT2>3.0.CO;2-Q](https://doi.org/10.1002/(SICI)1098-108X(199603)19:2<119::AID-EAT2>3.0.CO;2-Q). [PubMed: 8932550]
- Friederich HC, Wu M, Simon JJ, Herzog W. Neurocircuit function in eating disorders. *International Journal of Eating Disorders*. 2013; 46:425–432. <https://doi.org/10.1002/eat.22099>. [PubMed: 23658085]

- Frieling H, Bleich S, Otten J, Römer KD, Kornhuber J, de Zwaan M, ... Hillemecher T. Epigenetic downregulation of atrial natriuretic peptide but not vasopressin mRNA expression in females with eating disorders is related to impulsivity. *Neuropsychopharmacology*. 2008; 33:2605–2609. <https://doi.org/10.1038/sj.npp.1301662>. [PubMed: 18172431]
- Frieling H, Gozner A, Römer KD, Lenz B, Bönsch D, Wilhelm J, ... Bleich S. Global DNA hypomethylation and DNA hypermethylation of the alpha synuclein promoter in females with anorexia nervosa. *Molecular Psychiatry*. 2007; 12:229–230. <https://doi.org/10.1038/sj.mp.4001931>. [PubMed: 17325715]
- Frieling H, Römer KD, Scholz S, Mittelbach F, Wilhelm J, De Zwaan M, ... Bleich S. Epigenetic dysregulation of dopaminergic genes in eating disorders. *International Journal of Eating Disorders*. 2010; 43:577–583. <https://doi.org/10.1002/eat.20745>. [PubMed: 19728374]
- Gagne DA, Von Holle A, Brownley KA, Runfola CD, Hofmeier S, Branch KE, Bulik CM. Eating disorder symptoms and weight and shape concerns in a large web-based convenience sample of women ages 50 and above: Results of the Gender and Body Image (GABI) study. *International Journal of Eating Disorders*. 2012; 45:832–844. <https://doi.org/10.1002/eat.22030>. [PubMed: 22729743]
- Galimberti E, Martoni RM, Cavallini MC, Erzegovesi S, Bellodi L. Motor inhibition and cognitive flexibility in eating disorder subtypes. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2012; 36:307–312. <https://doi.org/10.1016/j.pnpbp.2011.10.017>. [PubMed: 22079108]
- Garber AK, Mauldin K, Michihata N, Buckelew SM, Shafer MA, Moscicki AB. Higher calorie diets increase rate of weight gain and shorten hospital stay in hospitalized adolescents with anorexia nervosa. *Journal of Adolescent Health*. 2013; 53:579–584. <https://doi.org/10.1016/j.jadohealth.2013.07.014>. [PubMed: 24054812]
- Gaudiani JL, Mehler PS. Rare medical manifestations of severe restricting and purging: “Zebras,” missed diagnoses, and best practices. *International Journal of Eating Disorders*. 2016; 49:331–344. <https://doi.org/10.1002/eat.22475>. [PubMed: 26592986]
- Giel KE, Hermann-Werner A, Mayer J, Diehl K, Schneider S, Thiel A. ... GOAL study group. Eating disorder pathology in elite adolescent athletes. *International Journal of Eating Disorders*. 2016; 49:553–562. <https://doi.org/10.1002/eat.22511>. [PubMed: 26876906]
- Giel KE, Kullmann S, Preißl H, Bischoff SC, Thiel A, Schmidt U, ... Teufel M. Understanding the reward system functioning in anorexia nervosa: Crucial role of physical activity. *Biological Psychology*. 2013; 94:575–581. <https://doi.org/10.1016/j.biopsycho.2013.10.004>. [PubMed: 24161801]
- Godier LR, de Wit S, Pinto A, Steinglass JE, Greene AL, Scaife J, ... Park RJ. An investigation of habit learning in anorexia nervosa. *Psychiatry Research*. 2016; 244:214–222. <https://doi.org/10.1016/j.psychres.2016.07.051>. Exploration of the role of habits, similar to those reported in compulsive disorders, that are hypothesized to play a role in the development and maintenance of anorexia nervosa. In two parallel studies, individuals with the binge/purge subtype of anorexia nervosa, restricting subtype of anorexia nervosa, and individuals recovered from anorexia nervosa did not show reliance on habits compared to healthy controls. Intact goal-directed learning was evident across all subtypes of anorexia nervosa. [PubMed: 27497292]
- Goodman A, Heshmati A, Koupil I. Family history of education predicts eating disorders across multiple generations among 2 million Swedish males and females. *PLoS One*. 2014; 9:e106475. <https://doi.org/10.1371/journal.pone.0106475>. [PubMed: 25162402]
- Griffiths S, Mond JM, Murray SB, Touyz S. Positive beliefs about anorexia nervosa and muscle dysmorphia are associated with eating disorder symptomatology. *Australian and New Zealand Journal of Psychiatry*. 2015; 49:812–820. <https://doi.org/10.1177/0004867415572412>. [PubMed: 25690746]
- Grilo CM, Masheb RM. Onset of dieting vs binge eating in outpatients with binge eating disorder. *International Journal of Obesity and Related Metabolic Disorders*. 2000; 24:404–409. [PubMed: 10805495]
- Hadland SE, Austin SB, Goodenow CS, Calzo JP. Weight misperception and unhealthy weight control behaviors among sexual minorities in the general adolescent population. *Journal of Adolescent Health*. 2014; 54:296–303. <https://doi.org/10.1016/j.jadohealth.2013.08.021>. [PubMed: 24182939]

- Hardaway JA, Crowley NA, Bulik CM, Kash TL. Integrated circuits and molecular components for stress and feeding: Implications for eating disorders. *Genes Brain and Behavior*. 2015; 14:85–97. <https://doi.org/10.1111/gbb.12185>. Exploration of how stress modulates different forms of feeding in animal models and identification of molecularly defined brain circuits that regulate feeding. Authors discuss the potential impact of these interactions and circuits for eating disorder biology.
- Hay P, Chinn D, Forbes D, Madden S, Newton R, Surgenor L. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the treatment of eating disorders. *Australian & New Zealand Journal of Psychiatry*. 2014; 48:977–1008. <https://doi.org/10.1177/0004867414555814>. [PubMed: 25351912]
- Hebebrand J, Muller TD, Holtkamp K, Herpertz-Dahlmann B. The role of leptin in anorexia nervosa: Clinical implications. *Molecular Psychiatry*. 2007; 12:23–35. <https://doi.org/10.1038/sj.mp.4001909>. [PubMed: 17060920]
- Hege MA, Stingl KT, Kullmann S, Schag K, Giel KE, Zipfel S, Preissl H. Attentional impulsivity in binge eating disorder modulates response inhibition performance and frontal brain networks. *International Journal of Obesity*. 2015; 39:353–360. <https://doi.org/10.1038/ijo.2014.99>. [PubMed: 24909828]
- Helverskov JL, Clausen L, Mors O, Frydenberg M, Thomsen PH, Rokkedal K. Trans-diagnostic outcome of eating disorders: A 30-month follow-up study of 629 patients. *European Eating Disorders Review*. 2010; 18:453–463. <https://doi.org/10.1002/erv.1025>. Using a naturalistic design, the authors investigated the 30-months outcome (remission and relapse) of various treatments and predictors of outcome in 629 patients (adolescents and adults) with different eating disorders. Almost half the patients provided data at follow-up of which 42% attained full remission, and 30% partial remission (no longer fulfilling the criteria for an eating disorder diagnosis). A total of 22% or 35% of those obtaining full or partial remission relapsed. Adult patients with AN-like conditions had the poorest outcome, and low BMI emerged as predictor of poor outcome in AN. The frequency of bingeing and purging was a predictor of poor outcome in BN. [PubMed: 20593480]
- Hoek HW, van Hoeken D. Review of the prevalence and incidence of eating disorders. *International Journal of Eating Disorders*. 2003; 34:383–396. <https://doi.org/10.1002/eat.10222>. [PubMed: 14566926]
- Holtkamp K, Herpertz-Dahlmann B, Hebebrand K, Mika C, Kratzsch J, Hebebrand J. Physical activity and restlessness correlate with leptin levels in patients with adolescent anorexia nervosa. *Biological Psychiatry*. 2006; 60:311–313. <https://doi.org/10.1016/j.biopsych.2005.11.001>. [PubMed: 16376860]
- Huas C, Godart N, Caille A, Pham-Scottet A, Foulon C, Divac SM, ... Rouillon F. Mortality and its predictors in severe bulimia nervosa patients. *European Eating Disorders Review*. 2013; 21:15–19. <https://doi.org/10.1002/erv.2178>. The mortality risk of bulimia nervosa was estimated by following 258 individuals admitted to a hospital in France between 1988 and 2004. The mean follow-up duration for subjects was 10.5 years. A total of 10 deaths were recorded during the follow-up time period, with the majority of deaths from suicide. The results show that individuals with bulimia nervosa are at an increased risk for death, specifically suicide. [PubMed: 22528211]
- Hudson JI, Hiripi E, Pope HG, Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biological Psychiatry*. 2007; 61:348–358. <https://doi.org/10.1016/j.biopsych.2006.03.040>. [PubMed: 16815322]
- Hudson JI, Lalonde JK, Berry JM, Pindyck LJ, Bulik CM, Crow SJ, ... Pope HG. Binge-eating disorder as a distinct familial phenotype in obese individuals. *Archives of General Psychiatry*. 2006; 63:313–319. <https://doi.org/10.1001/archpsyc.63.3.313>. [PubMed: 16520437]
- Jacobi C, Hayward C, de Zwaan M, Kraemer HC, Agras WS. Coming to terms with risk factors for eating disorders: Application of risk terminology and suggestions for a general taxonomy. *Psychological Bulletin*. 2004; 130:19–65. <https://doi.org/10.1037/0033-2909.130.1.19>. [PubMed: 14717649]
- Johnson JG, Cohen P, Kasen S, Brook JS. Childhood adversities associated with risk for eating disorders or weight problems during adolescence or early adulthood. *American Journal of Psychiatry*. 2002; 159:394–400. <https://doi.org/10.1176/appi.ajp.159.3.394>. [PubMed: 11870002]

- Kally Z, Cumella EJ. 100 midlife women with eating disorders: A phenomenological analysis of etiology. *Journal of General Psychology*. 2008; 135:359–377. <https://doi.org/10.3200/GENP.135.4.359-378>. [PubMed: 18959227]
- Kaye WH. Neurobiology of anorexia and bulimia nervosa. *Physiology & Behavior*. 2008; 94:121–135. <https://doi.org/10.1016/j.physbeh.2007>. [PubMed: 18164737]
- Kaye, WH., Strober, M. *The American Psychiatric Publishing Textbook of Psychopharmacology*. Arlington, VA: American Psychiatric Association; 2009. Neurobiology of eating disorders.
- Kaye WH, Wagner A, Fudge JL, Paulus M. Neurocircuitry of eating disorders. *Current Topics in Behavioral Neuroscience*. 2011; 6:37–57. https://doi.org/10.1007/7854_2010_85.
- Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Bischoff-Grethe A. Nothing tastes as good as skinny feels: The neurobiology of anorexia nervosa. *Trends in Neuroscience*. 2013a; 36:110–120. <https://doi.org/10.1016/j.tins.2013.01.003>.
- Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Wagner A, Bischoff-Grethe A. Does a shared neurobiology for foods and drugs of abuse contribute to extremes of food ingestion in anorexia and bulimia nervosa. *Biological Psychiatry*. 2013b; 73:836–842. <https://doi.org/10.1016/j.biopsych.2013.01.002>. [PubMed: 23380716]
- Keel PK, Brown TA. Update on course and outcome in eating disorders. *International Journal of Eating Disorders*. 2010; 43:195–204. <https://doi.org/10.1002/eat.20810/abstract>. [PubMed: 20186717]
- Kendler KS, Bulik CM, Silberg J, Hettema JM, Myers J, Prescott CA. Childhood sexual abuse and adult psychiatric and substance use disorders in women: An epidemiological and cotwin control analysis. *Archives of General Psychiatry*. 2000; 57:953–959. <https://doi.org/10.1001/archpsyc.57.10.953>. [PubMed: 11015813]
- Keshaviah A, Edkins K, Hastings ER, Krishna M, Franko DL, Herzog DB, ... Eddy KT. Re-examining premature mortality in anorexia nervosa: A meta-analysis redux. *Comprehensive Psychiatry*. 2014; 55:1773–1784. <https://doi.org/10.1016/j.comppsy.2014.07.017>. [PubMed: 25214371]
- Keski-Rahkonen A, Hoek HW, Linna MS, Raevuori A, Sihvola E, Bulik CM, ... Kaprio J. Incidence and outcomes of bulimia nervosa: A nationwide population-based study. *Psychological Medicine*. 2009; 39:823–831. <https://doi.org/10.1017/S0033291708003942>. [PubMed: 18775085]
- Keski-Rahkonen A, Hoek HW, Susser ES, Linna MS, Sihvola E, Raevuori A, ... Rissanen A. Epidemiology and course of anorexia nervosa in the community. *American Journal of Psychiatry*. 2007; 164:1259–1265. <https://doi.org/10.1176/appi.ajp.2007.06081388>. [PubMed: 17671290]
- Kessler RC, Berglund PA, Chiu WT, Deitz AC, Hudson JI, Shahly V, ... Xavier M. The prevalence and correlates of binge eating disorder in the World Health Organization World Mental Health Surveys. *Biological Psychiatry*. 2013; 73:904–914. <https://doi.org/10.1016/j.biopsych.2012.11.020>. [PubMed: 23290497]
- Kessler RM, Hutson PH, Herman BK, Potenza MN. The neurobiological basis of binge-eating disorder. *Neuroscience and Biobehavioral Reviews*. 2016; 63:223–238. <https://doi.org/10.1016/j.neubiorev.2016.01.013>. [PubMed: 26850211]
- Kim YR, Kim JH, Kim MJ, Treasure J. Differential methylation of the oxytocin receptor gene in patients with anorexia nervosa: A pilot study. *PLoS One*. 2014; 9:e88673. <https://doi.org/10.1371/journal.pone.0088673>. [PubMed: 24523928]
- Kimmel MC, Ferguson EH, Zerwas S, Bulik CM, Meltzer-Brody S. Obstetric and gynecologic problems associated with eating disorders. *International Journal of Eating Disorders*. 2016; 49:260–275. <https://doi.org/10.1002/eat.22483>. [PubMed: 26711005]
- King JA, Geisler D, Bernardoni F, Ritschel F, Böhm I, Seidel M, ... Ehrlich S. Altered neural efficiency of decision making during temporal reward discounting in anorexia nervosa. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2016; 55:972–979. <https://doi.org/10.1016/j.jaac.2016.08.005>. Decreased activation in frontoparietal regions involved in decision making, but faster and more consistent choice behavior of acute patients with acute anorexia nervosa in a temporal delay discounting task, suggests that the altered efficiency of neural resource allocation might underlie an increased level of self-control. [PubMed: 27806865]

- Kirby JS, Runfola CD, Fischer MS, Baucom DH, Bulik CM. Couple-based interventions for adults with eating disorders. *Eating Disorders*. 2015; 23:356–365. <https://doi.org/10.1080/10640266.2015.1044349>. [PubMed: 26010371]
- Klein DA, Schebendach JE, Gershkovich M, Bodell LP, Foltin RW, Walsh BT. Behavioral assessment of the reinforcing effect of exercise in women with anorexia nervosa: Further paradigm development and data. *International Journal of Eating Disorders*. 2010; 43:611–618. <https://doi.org/10.1002/eat.20758>. [PubMed: 19806608]
- Klump KL, Bulik CM, Kaye WH, Treasure J, Tyson E. Academy for Eating Disorders position paper: Eating disorders are serious mental illnesses. *International Journal of Eating Disorders*. 2009; 42:97–103. <https://doi.org/10.1002/eat.20589>. [PubMed: 18951455]
- Klump KL, Burt SA, McGue M, Iacono WG. Changes in genetic and environmental influences on disordered eating across adolescence: A longitudinal twin study. *Archives of General Psychiatry*. 2007; 64:1409–1415. <https://doi.org/10.1001/archpsyc.64.12.1409>. Changes in genetic and environmental influences on disordered eating across early-, mid-, and late- adolescence were examined. Significant changes in genetic and shared environmental effects across early- to mid-adolescence were observed. Specifically, genetic factors accounted for a small proportion of variance during early adolescence, but increased in importance and accounted for approximately 50% of the variance in disordered eating at mid- and late- adolescence. Shared environmental influences decreased substantially from early- to mid- adolescence. [PubMed: 18056549]
- Klump KL, Keel PK, Culbert KM, Edler C. Ovarian hormones and binge eating: Exploring associations in community samples. *Psychological Medicine*. 2008; 38:1749–1757. <https://doi.org/10.1017/S0033291708002997>. [PubMed: 18307829]
- Klump KL, Keel PK, Sisk C, Burt SA. Preliminary evidence that estradiol moderates genetic influences on disordered eating attitudes and behaviors during puberty. *Psychological Medicine*. 2010; 40:1745–1753. <https://doi.org/10.1017/S0033291709992236>. In a sample of 198 female adolescent twins, this study explored if estradiol levels moderated disordered eating by comparing twin correlations in low vs. high estradiol groups. They found similar MZ and DZ correlations in the low estradiol group, indicating no genetic effect. In the high estradiol group the MZ twin correlation was more than double the DZ twin correlation indicating genetic effects. [PubMed: 20059800]
- Kyriacou O, Treasure J, Schmidt U. Understanding how parents cope with living with someone with anorexia nervosa: Modelling the factors that are associated with carer distress. *International Journal of Eating Disorders*. 2008; 41:233–242. <https://doi.org/10.1002/eat.20488>. [PubMed: 18008326]
- Larsen PS, Strandberg-Larsen K, Micali N, Andersen AM. Parental and child characteristics related to early-onset disordered eating: A systematic review. *Harvard Review of Psychiatry*. 2015; 23:395–412. <https://doi.org/10.1097/HRP.000000000000073>. [PubMed: 26544091]
- Lavagnino L, Arnone D, Cao B, Soares JC, Selvaraj S. Inhibitory control in obesity and binge eating disorder: A systematic review and meta-analysis of neurocognitive and neuroimaging studies. *Neuroscience and Biobehavioral Reviews*. 2016; 68:714–726. <https://doi.org/10.1016/j.neubiorev.2016.06.041>. [PubMed: 27381956]
- le Grange D, Lock J, Agras WS, Bryson SW, Jo B. Randomized clinical trial of family-based treatment and cognitive-behavioral therapy for adolescent bulimia nervosa. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2015; 54:886–94. e2. <https://doi.org/10.1016/j.jaac.2015.08.008>. [PubMed: 26506579]
- le Grange D, Lock J, Loeb K, Nicholls D. Academy for Eating Disorders position paper: the role of the family in eating disorders. *International Journal of Eating Disorders*. 2010; 43:1–5. <https://doi.org/10.1002/eat.20751>. [PubMed: 19728372]
- Lebow J, Sim LA, Kransdorf LN. Prevalence of a history of overweight and obesity in adolescents with restrictive eating disorders. *Journal of Adolescent Health*. 2015; 56:19–24. <https://doi.org/10.1016/j.jadohealth.2014.06.005>. [PubMed: 25049202]
- Lee S, Lee AM, Ngai E, Lee DT, Wing YK. Rationales for food refusal in Chinese patients with anorexia nervosa. *International Journal of Eating Disorders*. 2001; 29:224–229. [https://doi.org/10.1002/1098-108X\(200103\)29:2<224::AID-EAT1012>3.0.CO;2-R](https://doi.org/10.1002/1098-108X(200103)29:2<224::AID-EAT1012>3.0.CO;2-R). [PubMed: 11429985]

- Lilenfeld LR, Kaye WH, Greeno CG, Merikangas KR, Plotnicov K, Pollice C, ... Nagy L. A controlled family study of anorexia nervosa and bulimia nervosa: Psychiatric disorders in first-degree relatives and effects of proband comorbidity. *Archives of General Psychiatry*. 1998; 55:603–610. <https://doi.org/10.1001/archpsyc.55.7.603>. [PubMed: 9672050]
- Lindvall Dahlgren C, Wisting L. Transitioning from DSM-IV to DSM-5: A systematic review of eating disorder prevalence assessment. *International Journal of Eating Disorders*. 2016; 49:975–997. <https://doi.org/10.1002/eat.22596>. [PubMed: 27528542]
- Lock, J., le Grange, D. *Treatment Manual for Anorexia Nervosa: A Family-Based Approach*. 2. New York: Guildford Press; 2015.
- Loomba-Albrecht LA, Styne DM. Effect of puberty on body composition. *Current Opinion in Endocrinology Diabetes and Obesity*. 2009; 16:10–15.
- Lutter M, Khan MZ, Satio K, Davis KC, Kidder IJ, McDaniel L, ... Cui H. The eating-disorder associated HDAC4(A778T) mutation alters feeding behaviors in female mice. *Biological Psychiatry*. 2017; 81:770–777. <https://doi.org/10.1016/j.biopsych.2016.09.024>. [PubMed: 27884425]
- Madden S, Miskovic-Wheatley J, Clarke S, Touyz S, Hay P, Kohn MR. Outcomes of a rapid refeeding protocol in adolescent anorexia nervosa. *Journal of Eating Disorders*. 2015; 3:8. <https://doi.org/10.1186/s40337-015-0047-1>. [PubMed: 25830024]
- Mangweth-Matzek B, Hoek HW, Rupp CI, Kemmler G, Pope HG, Kinzl J. The menopausal transition--a possible window of vulnerability for eating pathology. *International Journal of Eating Disorders*. 2013; 46:609–616. <https://doi.org/10.1002/eat.22157>. [PubMed: 23847142]
- Marques L, Alegria M, Becker AE, Chen CN, Fang A, Chosak A, Diniz JB. Comparative prevalence, correlates of impairment, and service utilization for eating disorders across US ethnic groups: Implications for reducing ethnic disparities in health care access for eating disorders. *International Journal of Eating Disorders*. 2011; 44:412–420. <https://doi.org/10.1002/eat.20787>. [PubMed: 20665700]
- Marsh R, Maia TV, Peterson BS. Functional disturbances within frontostriatal circuits across multiple childhood psychopathologies. *American Journal of Psychiatry*. 2009; 166:664–674. <https://doi.org/10.1176/appi.ajp.2009.08091354>. [PubMed: 19448188]
- Marsh R, Steinglass JE, Gerber AJ, Graziano O'Leary K, Wang Z, Murphy D, ... Peterson BS. Deficient activity in the neural systems that mediate self-regulatory control in bulimia nervosa. *Archives of General Psychiatry*. 2009; 66:51–63. <https://doi.org/10.1001/archgenpsychiatry.2008.504>. [PubMed: 19124688]
- Mehler PS, Brown C. Anorexia nervosa - medical complications. *Journal of Eating Disorders*. 2015; 3:11. <https://doi.org/10.1186/s40337-015-0040-8>. [PubMed: 25834735]
- Mehler PS, Rylander M. Bulimia nervosa - medical complications. *Journal of Eating Disorders*. 2015; 3:12. <https://doi.org/10.1186/s40337-015-0044-4>. Comprehensive review of the literature on medical complications associated with purging behaviors in bulimia nervosa, focusing on self-induced vomiting and laxative abuse. While complications of laxative abuse involve mainly the gastrointestinal system and electrolyte disturbances, complications of self-induced vomiting also include cutaneous, dental, throat, cardiac, reproductive, and pulmonary domains. Effects are dependent on the mode and frequency of purging. [PubMed: 25866627]
- Micali N, Solmi F, Horton NJ, Crosby RD, Eddy KT, Calzo JP, ... Field AE. Adolescent eating disorders predict psychiatric, high-risk behaviors and weight outcomes in young adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2015; 54:652–659. e1. <https://doi.org/10.1016/j.jaac.2015.05.009>. [PubMed: 26210334]
- Mitchison D, Hay P, Slewa-Younan S, Mond J. The changing demographic profile of eating disorder behaviors in the community. *BMC Public Health*. 2014; 14:943. <https://doi.org/10.1186/1471-2458-14-943>. Using two Australian household surveys, one in 1998 (n=3010) and the other in 2008 (n=3034), the authors interrogated the stereotype that eating disorders are diseases of young, white, socioeconomically privileged women. Between the two samples, the most drastic increases in prevalence of various eating-disorder symptoms were in participants who were male, >45 years old, or from lower socioeconomic brackets. All disordered eating traits had similar negative impact on health-related quality of life regardless of demographic factors. [PubMed: 25213544]

- Mitchison D, Hay PJ. The epidemiology of eating disorders: Genetic, environmental, and societal factors. *Clinical Epidemiology*. 2014; 6:89–97. <https://doi.org/10.2147/CLEP.S40841>. [PubMed: 24728136]
- Monteleone P, Maj M. Dysfunctions of leptin, ghrelin, BDNF and endocannabinoids in eating disorders: Beyond the homeostatic control of food intake. *Psychoneuroendocrinology*. 2013; 38:312–330. <https://doi.org/10.1016/j.psychneuen.2012.10.021>. [PubMed: 23313276]
- Moore F, Keel PK. Influence of sexual orientation and age on disordered eating attitudes and behaviors in women. *International Journal of Eating Disorders*. 2003; 34:370–374. <https://doi.org/10.1002/eat.10198>. [PubMed: 12949929]
- Morrison MA, Morrison TG, Sager CL. Does body satisfaction differ between gay men and lesbian women and heterosexual men and women? A meta-analytic review. *Body Image*. 2004; 1:127–138. <https://doi.org/10.1016/j.bodyim.2004.01.002>. [PubMed: 18089146]
- Murray SM, Tulloch AJ, Chen EY, Avena NM. Insights revealed by rodent models of sugar binge eating. *CNS Spectrums*. 2015; 20:530–536. <https://doi.org/10.1017/S1092852915000656>. [PubMed: 26510689]
- Mustelin L, Raevuori A, Hoek HW, Kaprio J, Keski-Rahkonen A. Incidence and weight trajectories of binge eating disorder among young women in the community. *International Journal of Eating Disorders*. 2015; 48:1106–1112. <https://doi.org/10.1002/eat.22409>. In a community sample of young women, lifetime prevalence of DSM-5 BED was 0.7%, mean age of onset was 19 years, comorbid major depressive disorder was common, and BED was frequently preceded by relative overweight in adolescence. [PubMed: 25846672]
- National Collaborating Centre for Mental Health. Eating Disorders: Core interventions in the treatment of anorexia nervosa, bulimia nervosa and related eating disorders. 2004. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/23346610>
- Neumark-Sztainer D, Wall M, Guo J, Story M, Haines J, Eisenberg M. Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: How do dieters fare 5 years later? *Journal of the American Dietetics Association*. 2006; 106:559–568. <https://doi.org/10.1016/j.jada.2006.01.003>.
- Nicholls D, Bryant-Waugh R. Eating disorders of infancy and childhood: Definition, symptomatology, epidemiology, and comorbidity. *Child and Adolescent Psychiatric Clinics of North America*. 2009; 18:17–30. <https://doi.org/10.1016/j.chc.2008.07.008>. [PubMed: 19014855]
- Nicholls DE, Viner RM. Childhood risk factors for lifetime anorexia nervosa by age 30 years in a national birth cohort. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2009; 48:791–799. <https://doi.org/10.1097/CHI.0b013e3181ab8b75>. [PubMed: 19564797]
- Nordbø RH, Espeset EM, Gulliksen KS, Skårderud F, Geller J, Holte A. Reluctance to recover in anorexia nervosa. *European Eating Disorders Review*. 2012; 20:60–67. <https://doi.org/10.1002/erv.1097>. In depth-interviews from 36 women with anorexia nervosa were collected to explore what makes individuals with anorexia nervosa ambivalent about recovery. Core obstacles reported by patients with anorexia included: perceiving judgements, feeling stuck, feeling distressed, denying anorexia nervosa, eating, gaining weight, and perceived benefits of the illness. [PubMed: 21305676]
- Norris ML, Robinson A, Obeid N, Harrison M, Spettigue W, Henderson K. Exploring avoidant/restrictive food intake disorder in eating disordered patients: A descriptive study. *International Journal of Eating Disorders*. 2014; 47:495–499. <https://doi.org/10.1002/eat.22217>. [PubMed: 24343807]
- O'Hara CB, Campbell IC, Schmidt U. A reward-centered model of anorexia nervosa: A focused narrative review of the neurological and psychophysiological literature. *Neuroscience & Biobehavioral Reviews*. 2015; 52:131–152. <https://doi.org/10.1016/j.neubiorev.2015.02.012>. [PubMed: 25735957]
- Papadopoulos FC, Ekblom A, Brandt L, Ekselius L. Excess mortality, causes of death and prognostic factors in anorexia nervosa. *British Journal of Psychiatry*. 2009; 194:10–17. <https://doi.org/10.1192/bjp.bp.108.054742>. [PubMed: 19118319]
- Patel P, Wheatcroft R, Park RJ, Stein A. The children of mothers with eating disorders. *Clinical Child and Family Psychological Review*. 2002; 5:1–19.

- Peat CM, Peyerl NL, Muehlenkamp JJ. Body image and eating disorders in older adults: A review. *Journal of General Psychology*. 2008; 135:343–358. <https://doi.org/10.3200/GENP.135.4.343-358>. [PubMed: 18959226]
- Pike KM. Long-term course of anorexia nervosa: response, relapse, remission, and recovery. *Clinical Psychology Review*. 1998; 18:447–475. [https://doi.org/10.1016/S0272-7358\(98\)00014-2](https://doi.org/10.1016/S0272-7358(98)00014-2). [PubMed: 9638357]
- Pike KM, Dunne PE. The rise of eating disorders in Asia: A review. *Journal of Eating Disorders*. 2015; 3:33. <https://doi.org/10.1186/s40337-015-0070-2>. [PubMed: 26388993]
- Pike KM, Dunne PE, Addai E. Expanding the boundaries: reconfiguring the demographics of the “typical” eating disordered patient. *Current Psychiatry Reports*. 2013; 15:411. <https://doi.org/10.1007/s11920-013-0411-2>. [PubMed: 24122512]
- Pike KM, Hoek HW, Dunne PE. Cultural trends and eating disorders. *Current Opinions in Psychiatry*. 2014; 27:436–442. <https://doi.org/10.1097/YCO.0000000000000100>.
- Pisetsky EM, Thornton LM, Lichtenstein P, Pedersen NL, Bulik CM. Suicide attempts in women with eating disorders. *Journal of Abnormal Psychology*. 2013; 122:1042–1056. <https://doi.org/10.1037/a0034902>. [PubMed: 24364606]
- Plomin R, DeFries JC, Loehlin JC. Genotype-environment interaction and correlation in the analysis of human behavior. *Psychological Bulletin*. 1977; 84:309–322. <https://doi.org/10.1037/0033-2909.84.2.309> Examination of the effect of genotype-environment interaction and correlation in behavioral genetic studies (twin and adoption studies). [PubMed: 557211]
- Preti A, Girolamo G, Vilagut G, Alonso J, Graaf R, Bruffaerts R. ... the ESEMEd-WMH investigators. The epidemiology of eating disorders in six European countries: Results of the ESEMEd-WMH project. *Journal of Psychiatric Research*. 2009; 43:1125–1132. <https://doi.org/10.1016/j.jpsychires.2009.04.003>. In a cross-sectional survey of six European countries, lifetime estimated prevalence of AN, BN, BED, sub-threshold BED, and any binge eating were 0.48%, 0.51%, 1.12%, 0.72%, and 2.15%, respectively, and they were 3–8 times higher among women for all eating disorders. Age of onset for the majority of eating disorders was between 10 and 20 years of age. Comorbidity with other mental disorders was common. Only a minority of individuals with a lifetime eating disorder requested medical treatment. [PubMed: 19427647]
- Preti A, Rocchi MB, Sisti D, Camboni MV, Miotto P. A comprehensive meta-analysis of the risk of suicide in eating disorders. *Acta Psychiatrica Scandinavica*. 2011; 124:6–17. <https://doi.org/10.1111/j.1600-0447.2010.01641.x>. [PubMed: 21092024]
- Qian J, Hu Q, Wan Y, Li T, Wu M, Ren Z, Yu D. Prevalence of eating disorders in the general population: A systematic review. *Shanghai Archives of Psychiatry*. 2013; 25:212–223. <https://doi.org/10.3969/j.issn.1002-0829.2013.04.003>. [PubMed: 24991159]
- Racine SE, Culbert KM, Keel PK, Sisk CL, Burt SA, Klump KL. Differential associations between ovarian hormones and disordered eating symptoms across the menstrual cycle in women. *International Journal of Eating Disorders*. 2012; 45:333–344. <https://doi.org/10.1002/eat.20941>. [PubMed: 21656540]
- Redgrave GW, Coughlin JW, Schreyer CC, Martin LM, Leonpacher AK, Seide M, ... Guarda AS. Refeeding and weight restoration outcomes in anorexia nervosa: Challenging current guidelines. *International Journal of Eating Disorders*. 2015; 48:866–873. <https://doi.org/10.1002/eat.22390>. [PubMed: 25625572]
- Ricciardelli LA, McCabe MP. A biopsychosocial model of disordered eating and the pursuit of muscularity in adolescent boys. *Psychological Bulletin*. 2004; 130:179–205. <https://doi.org/10.1037/0033-2909.130.2.179>. [PubMed: 14979769]
- Richard M, Bauer S, Kordy H. Relapse in anorexia and bulimia nervosa—a 2.5-year follow-up study. *European Eating Disorders Review*. 2005; 13:180–190. <https://doi.org/10.1002/erv.638>. In Project TR-EAT, the symptomatic status of eating-disordered patients (AN, N = 233, BN, N = 422) was tracked after inpatient treatment over a 2.5-year follow-up period. The distribution of time to relapse for both disorders and possible predictors for relapse are investigated by means of discrete time survival analysis. Fifty-eight per cent of the patients with AN and 74% of those with BN achieved partial remission before end of treatment, and thus were at risk for relapse. The relapse rates within 2.5 years were 32.6% for AN and 37.4% for BN. For both eating disorders, the highest risk of relapse was within the first 6 or 7 months after achieving partial remission.

- Rousselet M, Guérineau B, Paruit MC, Guinot M, Lise S, Destrube B, ... Prétagut S. Disordered eating in French high-level athletes: Association with type of sport, doping behavior, and psychological features. *Eating and Weight Disorders*. 2017; 22:61–68. <https://doi.org/10.1007/s40519-016-0342-0>. [PubMed: 27838862]
- Runfola CD, Thornton LM, Pisetsky EM, Bulik CM, Birgegård A. Self-image and suicide in a Swedish national eating disorders clinical register. *Comprehensive Psychiatry*. 2014; 55:439–449. <https://doi.org/10.1016/j.comppsy.2013.11.007>. [PubMed: 24332388]
- Russell CJ, Keel PK. Homosexuality as a specific risk factor for eating disorders in men. *International Journal of Eating Disorders*. 2002; 31:300–306. [PubMed: 11920991]
- Saltzman JA, Liechty JM. Family correlates of childhood binge eating: A systematic review. *Eating Behaviors*. 2016; 22:62–71. <https://doi.org/10.1016/j.eatbeh.2016.03.027>. [PubMed: 27089384]
- Sanders N, Smeets PA, van Elburg AA, Danner UN, van Meer F, Hoek HW, Adan RA. Altered food-cue processing in chronically ill and recovered women with anorexia nervosa. *Frontiers in Behavioral Neuroscience*. 2015; 9:46. <https://doi.org/10.3389/fnbeh.2015.00046>. [PubMed: 25774128]
- Santonastaso P, Bosello R, Schiavone P, Tenconi E, Degortes D, Favaro A. Typical and atypical restrictive anorexia nervosa: Weight history, body image, psychiatric symptoms, and response to outpatient treatment. *International Journal of Eating Disorders*. 2009; 42:464–470. <https://doi.org/10.1002/eat.20706>. [PubMed: 19424978]
- Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature*. 2014; 511:421–427. <https://doi.org/10.1038/nature13595>. [PubMed: 25056061]
- Schmidt U, Renwick B, Lose A, Kenyon M, Dejong H, Broadbent H, ... Landau S. The MOSAIC study - comparison of the Maudsley Model of Treatment for Adults with Anorexia Nervosa (MANTRA) with Specialist Supportive Clinical Management (SSCM) in outpatients with anorexia nervosa or eating disorder not otherwise specified, anorexia nervosa type: Study protocol for a randomized controlled trial. *Trials*. 2013; 14:160. <https://doi.org/10.1186/1745-6215-14-160>. [PubMed: 23721562]
- Schroeder M, Eberlein C, de Zwaan M, Kornhuber J, Bleich S, Frieling H. Lower levels of cannabinoid 1 receptor mRNA in female eating disorder patients: Association with wrist cutting as impulsive self-injurious behavior. *Psychoneuroendocrinology*. 2012; 37:2032–2036. <https://doi.org/10.1016/j.psyneuen.2012.03.025>. [PubMed: 22542985]
- Seitz J, Bühren K, von Polier GG, Heussen N, Herpertz-Dahlmann B, Konrad K. Morphological changes in the brain of acutely ill and weight-recovered patients with anorexia nervosa. A meta-analysis and qualitative review. *Zeitschrift für Kinder- und Jugendpsychiatrie Psychotherapie*. 2014; 42:7–17. <https://doi.org/10.1024/1422-4917/a000265>.
- Seitz J, Herpertz-Dahlmann B, Konrad K. Brain morphological changes in adolescent and adult patients with anorexia nervosa. *Journal of Neural Transmission*. 2016; 123:949–959. <https://doi.org/10.1007/s00702-016-1567-9>. [PubMed: 27188331]
- Shoebri P, Gowers SG. Parental high concern and adolescent-onset anorexia nervosa. A case-control study to investigate direction of causality. *British Journal of Psychiatry*. 2000; 176:132–137. <https://doi.org/10.1192/bjp.176.2.132>. [PubMed: 10755049]
- Siervogel RM, Demerath EW, Schubert C, Remsberg KE, Chumlea WC, Sun S, ... Towne B. Puberty and body composition. *Hormone Research*. 2003; 60:36–45. <https://doi.org/71224>. [PubMed: 12955016]
- Sim LA, Homme JH, Lteif AN, Vande Voort JL, Schak KM, Ellingson J. Family functioning and maternal distress in adolescent girls with anorexia nervosa. *International Journal of Eating Disorders*. 2009; 42:531–539. <https://doi.org/10.1002/eat.20654>. [PubMed: 19189407]
- Smink FR, van Hoeken D, Hoek HW. Epidemiology of eating disorders: Incidence, prevalence and mortality rates. *Current Psychiatry Reports*. 2012; 14:406–414. <https://doi.org/10.1007/s11920-012-0282-y>. [PubMed: 22644309]
- Solomon SM, Kirby DF. The refeeding syndrome: A review. *Journal of Parenteral and Enteral Nutrition*. 1990; 14:90–97. [PubMed: 2109122]

- Spies M, Knudsen GM, Lanzenberger R, Kasper S. The serotonin transporter in psychiatric disorders: Insights from PET imaging. *Lancet Psychiatry*. 2015; 2:743–755. [https://doi.org/10.1016/S2215-0366\(15\)00232-1](https://doi.org/10.1016/S2215-0366(15)00232-1). [PubMed: 26249305]
- Steenhuis IH, Vermeer WM. Portion size: Review and framework for interventions. *International Journal of Behavioral Nutrition and Physical Activity*. 2009; 6:58. <https://doi.org/10.1186/1479-5868-6-58>. [PubMed: 19698102]
- Steinglass JE, Walsh BT. Neurobiological model of the persistence of anorexia nervosa. *Journal of Eating Disorders*. 2016; 4:19. <https://doi.org/10.1186/s40337-016-0106-2>. [PubMed: 27195123]
- Steinhausen HC. Outcome of eating disorders. *Child and Adolescent Psychiatric Clinics of North America*. 2009; 18:225–242. <https://doi.org/10.1016/j.chc.2008.07.013>. [PubMed: 19014869]
- Steinhausen HC, Jakobsen H, Helenius D, Munk-Jørgensen P, Strober M. A nation-wide study of the family aggregation and risk factors in anorexia nervosa over three generations. *International Journal of Eating Disorders*. 2015; 48:1–8. <https://doi.org/10.1002/eat.22293>. [PubMed: 24777686]
- Steinhausen HC, Weber S. The outcome of bulimia nervosa: Findings from one-quarter century of research. *American Journal of Psychiatry*. 2009; 166:1331–1341. <https://doi.org/10.1176/appi.ajp.2009.09040582>. [PubMed: 19884225]
- Sternson SM, Roth BL. Chemogenetic tools to interrogate brain functions. *Annual Review of Neuroscience*. 2014; 37:387–407. <https://doi.org/10.1146/annurev-neuro-071013-014048>.
- Striegel-Moore RH, Bulik CM. Risk factors for eating disorders. *American Psychologist*. 2007; 62:181–198. <https://doi.org/10.1037/0003-066X.62.3.181>. [PubMed: 17469897]
- Striegel-Moore RH, Dohm FA, Kraemer HC, Taylor CB, Daniels S, Crawford PB, Schreiber GB. Eating disorders in White and Black women. *American Journal of Psychiatry*. 2003; 160:1326–1331. <https://doi.org/10.1176/appi.ajp.160.7.1326>. [PubMed: 12832249]
- Striegel-Moore RH, Franko DL. Epidemiology of binge eating disorder. *International Journal of Eating Disorders*. 2003; 34:S19–S29. [PubMed: 12900983]
- Striegel RH, Bedrosian R, Wang C, Schwartz S. Why men should be included in research on binge eating: Results from a comparison of psychosocial impairment in men and women. *International Journal of Eating Disorders*. 2012; 45:233–240. <https://doi.org/10.1002/eat.20962>. [PubMed: 22031213]
- Strober M, Freeman R, Lampert C, Diamond J, Kaye W. Controlled family study of anorexia nervosa and bulimia nervosa: Evidence of shared liability and transmission of partial syndromes. *American Journal of Psychiatry*. 2000; 157:393–401. [PubMed: 10698815]
- Strober M, Humphrey LL. Familial contributions to the etiology and course of anorexia nervosa and bulimia. *Journal of Consulting and Clinical Psychology*. 1987; 55:654–659. [PubMed: 3331628]
- Suisman JL, Thompson JK, Keel PK, Burt SA, Neale M, Boker S, ... Klump KL. Genetic and environmental influences on thin-ideal internalization across puberty and preadolescent, adolescent, and young adult development. *International Journal of Eating Disorders*. 2014; 47:773–783. <https://doi.org/10.1002/eat.22321>. [PubMed: 24962440]
- Suokas JT, Suvisaari JM, Gissler M, Löfman R, Linna MS, Raevuori A, Haukka J. Mortality in eating disorders: A follow-up study of adult eating disorder patients treated in tertiary care, 1995–2010. *Psychiatry Research*. 2013; 210:1101–1106. <https://doi.org/10.1016/j.psychres.2013.07.042>. [PubMed: 23958333]
- Tau GZ, Peterson BS. Normal development of brain circuits. *Neuropsychopharmacology*. 2010; 35:147–168. <https://doi.org/10.1038/npp.2009.115>. [PubMed: 19794405]
- Thornton LM, Watson HJ, Jangmo A, Welch E, Wiklund C, von Hausswolff-Juhlin Y, ... Bulik CM. Binge-eating disorder in the Swedish national registers: Somatic comorbidity. *International Journal of Eating Disorders*. 2017; 50:58–65. <https://doi.org/10.1002/eat.22624>. Swedish register data were used to identify a sample of 850 individuals diagnosed with binge-eating disorder (BED). Associations were examined between BED and neurologic, immune, respiratory, gastrointestinal, skin, musculoskeletal, genitourinary, circulatory, and endocrine system diseases. Compared with controls matched on sex, year, month, and county of birth, BED was associated with an increased risk for most classes of disease. The strongest associations were found between BED and diabetes and cardiovascular system diseases. [PubMed: 27642179]

- Thornton LM, Welch E, Munn-Chernoff MA, Lichtenstein P, Bulik CM. Anorexia nervosa, major depression, and suicide attempts: Shared genetic factors. *Suicide and Life-Threatening Behaviors*. 2016; 46:525–534. <https://doi.org/10.1111/sltb.12235>. This is one of the first studies to investigate overlapping genetic and environmental risk factors for anorexia nervosa, major depression, and suicide attempts. Findings suggest that a portion of the genetic factors underlying anorexia nervosa also contribute to liability to major depression and suicide attempts in adult women. Individual-specific environmental factors, however, may not overlap but rather are trait specific.
- Titova OE, Hjorth OC, Schiöth HB, Brooks SJ. Anorexia nervosa is linked to reduced brain structure in reward and somatosensory regions: A meta-analysis of VBM studies. *BMC Psychiatry*. 2013; 13:110. <https://doi.org/10.1186/1471-244X-13-110>. [PubMed: 23570420]
- Trace SE, Baker JH, Peñas-Lledó E, Bulik C. The genetics of eating disorders. *Annual Review of Clinical Psychology*. 2013; 9:589–620. <https://doi.org/10.1146/annurev-clinpsy-050212-185546>.
- Treasure J, Murphy T, Szmukler G, Todd G, Gavan K, Joyce J. The experience of caregiving for severe mental illness: A comparison between anorexia nervosa and psychosis. *Social Psychiatry and Psychiatric Epidemiology*. 2001; 36:343–347. [PubMed: 11606003]
- Treasure J, Nazar BP. Interventions for the carers of patients with eating disorders. *Current Psychiatry Reports*. 2016; 18:16. <https://doi.org/10.1007/s11920-015-0652-3>. [PubMed: 26781554]
- Treasure J, Stein D, Maguire S. Has the time come for a staging model to map the course of eating disorders from high risk to severe enduring illness? An examination of the evidence. *Early Intervention in Psychiatry*. 2015; 9:173–184. <https://doi.org/10.1111/eip.12170>. [PubMed: 25263388]
- Treasure J, Zipfel S, Micali N, Wade T, Stice E, Claudino A, ... Wentz E. Anorexia nervosa. *Nature Reviews Disease Primers*. 2015; 1:15074. <https://doi.org/10.1038/nrdp.2015.74>.
- Twomey CD, Baldwin DS, Hopfe M, Cieza A. A systematic review of the predictors of health service utilisation by adults with mental disorders in the UK. *BMJ Open*. 2015; 5:e007575. <https://doi.org/10.1136/bmjopen-2015-007575>.
- van den Eynde F, Suda M, Broadbent H, Guillaume S, van den Eynde M, Steiger H, ... Schmidt U. Structural magnetic resonance imaging in eating disorders: A systematic review of voxel-based morphometry studies. *European Eating Disorders Review*. 2012; 20:94–105. <https://doi.org/10.1002/erv.1163>. [PubMed: 22052722]
- van Langenberg T, Sawyer SM, le Grange D, Hughes EK. Psychosocial well-being of siblings of adolescents with anorexia nervosa. *European Eating Disorders Review*. 2016; 24:438–445. <https://doi.org/10.1002/erv.2469>. Eighty-five parents and 55 siblings of adolescents with anorexia nervosa completed the Strengths and Difficulties Questionnaire at diagnosis. In addition, 88 parents and 46 siblings completed the Strengths and Difficulties Questionnaire after finishing treatment. Mothers and fathers reported siblings to have lower levels of conduct problems in comparison with population norms. Mothers also reported lower levels of prosocial behaviors. Siblings reported higher levels of emotional difficulties and hyperactivity in comparison with their peers. [PubMed: 27501269]
- Vandereycken W. Denial of illness in anorexia nervosa—a conceptual review: Part 1 diagnostic significance and assessment. *European Eating Disorders Review*. 2006a; 14:341–351. <https://doi.org/10.1002/erv.721>.
- Vandereycken W. Denial of illness in anorexia nervosa—a conceptual review: Part 2 different forms and meanings. *European Eating Disorders Review*. 2006b; 14:352–368. <https://doi.org/10.1002/erv.722>.
- Wade TD, Fairweather-Schmidt AK, Zhu G, Martin NG. Does shared genetic risk contribute to the co-occurrence of eating disorders and suicidality? *International Journal of Eating Disorders*. 2015; 48:684–691. <https://doi.org/10.1002/eat.22421>. Overlapping genetic and environmental risk factors for a composite measure of eating disorders (including AN, BN, BED, and purging disorder) and suicidality were examined in female twins. Results suggest that some of the genetic factors influencing vulnerability to eating disorders also influence suicidality. [PubMed: 25945699]
- Walton E, Hass J, Liu J, Roffman JL, Bernardoni F, Roessner V, ... Ehrlich S. Correspondence of DNA methylation between blood and brain tissue and its application to schizophrenia research.

Schizophrenia Bulletin. 2016; 42:406–414. <https://doi.org/10.1093/schbul/sbv074>. Blood and temporal lobe biopsy samples were obtained from twelve epilepsy patients during neurosurgical treatment. Findings indicate that most DNA methylation markers in peripheral blood do not reliably predict brain DNA methylation status, but a subset of peripheral data may proxy methylation status of brain tissue. Restricting the analysis to these markers may identify meaningful epigenetic differences in brain disorders. [PubMed: 26056378]

- Watson HJ, Bulik CM. Update on the treatment of anorexia nervosa: Review of clinical trials, practice guidelines and emerging interventions. *Psychological Medicine*. 2013; 43:2477–2500. <https://doi.org/10.1017/S0033291712002620>. [PubMed: 23217606]
- Watson, RJ., Veale, JF., Saewyc, EM. Disordered eating behaviors among transgender youth: Probability profiles from risk and protective factors. *International Journal of Eating Disorders*. 2016. <https://doi.org/10.1002/eat.22627>
- Westmoreland P, Krantz MJ, Mehler PS. Medical complications of anorexia nervosa and bulimia. *American Journal of Medicine*. 2016; 129:30–37. <https://doi.org/10.1016/j.amjmed.2015.06.031>. Review of medical complications associated with anorexia nervosa and bulimia nervosa and how the complications can be treated. *Epidemiology and psychiatric comorbidities of eating disorders are discussed*. [PubMed: 26169883]
- Whitney J, Murray J, Gavan K, Todd G, Whitaker W, Treasure J. Experience of caring for someone with anorexia nervosa: Qualitative study. *British Journal of Psychiatry*. 2005; 187:444–449. <https://doi.org/10.1192/bjp.187.5.444>. [PubMed: 16260820]
- Wierenga CE, Ely A, Bischoff-Grethe A, Bailer UF, Simmons AN, Kaye WH. Are extremes of consumption in eating disorders related to an altered balance between reward and inhibition. *Frontiers in Behavioral Neuroscience*. 2014; 8:410. <https://doi.org/10.3389/fnbeh.2014.00410>. [PubMed: 25538579]
- Witcomb GL, Bouman WP, Brewin N, Richards C, Fernandez-Aranda F, Arcelus J. Body image dissatisfaction and eating-related psychopathology in trans individuals: A matched control study. *European Eating Disorders Review*. 2015; 23:287–293. <https://doi.org/10.1002/erv.2362>. Eating disorder traits and body dissatisfaction are compared across three different groups of people: 200 trans people, 200 people with eating disorders, and 200 healthy controls. Results from the study are in line with previous literature; trans individuals do not score as highly on measures of body dissatisfaction as those with clinical eating disorders. However, compared with healthy controls, trans individuals report higher eating disorder symptoms. Particular attention is paid to gender differences, as trans males had similarly high scores on measures of body dissatisfaction as males with eating disorders. [PubMed: 25944170]
- Wu M, Hartmann M, Skunde M, Herzog W, Friederich HC. Inhibitory control in bulimic-type eating disorders: A systematic review and meta-analysis. *PLoS One*. 2013; 8:e83412. <https://doi.org/10.1371/journal.pone.0083412>. [PubMed: 24391763]
- Yager J. Family issues in the pathogenesis of anorexia nervosa. *Psychosomatic Medicine*. 1982; 44:43–60. [PubMed: 7089153]
- Yao S, Kuja-Halkola R, Thornton LM, Runfola CD, D’Onofrio BM, Almqvist C, ... Bulik CM. Familial liability for eating disorders and suicide attempts: Evidence from a population registry in Sweden. *JAMA Psychiatry*. 2016; 73:284–291. <https://doi.org/10.1001/jamapsychiatry.2015.2737>. The association between eating disorders and suicide attempts was examined using national register data from Sweden. Significantly elevated risks of suicide attempts were observed in individuals with any eating disorders, anorexia nervosa, and bulimia nervosa. Elevated risks of suicide attempts were also found in relatives (full-siblings, half-siblings, and cousins) of the individuals with the aforementioned eating disorders, suggesting familial liability for eating disorders and suicide attempts. [PubMed: 26764185]
- Yilmaz Z, Hardaway JA, Bulik CM. Genetics and epigenetics of eating disorders. *Advances in Genomics and Genetics*. 2015; 5:131–150. <https://doi.org/10.2147/AGG.S55776>. [PubMed: 27013903]
- Zerwas S, Larsen JT, Petersen L, Thornton LM, Mortensen PB, Bulik CM. The incidence of eating disorders in a Danish register study: Associations with suicide risk and mortality. *Journal of Psychiatric Research*. 2015; 65:16–22. <https://doi.org/10.1016/j.jpsychires.2015.03.003>. [PubMed: 25958083]

Zipfel S, Giel KE, Bulik CM, Hay P, Schmidt U. Anorexia nervosa: Aetiology, assessment, and treatment. *Lancet Psychiatry*. 2015; 2:1099–1111. [https://doi.org/10.1016/S2215-0366\(15\)00356-9](https://doi.org/10.1016/S2215-0366(15)00356-9). [PubMed: 26514083]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript