## Corrigendum

## Basic domain of telomere guardian TRF2 reduces D-loop unwinding whereas Rap1 restores it

Ivona Nečasová<sup>1</sup>, Eliška Janoušková<sup>1</sup>, Tomáš Klumpler<sup>2</sup> and Ctirad Hofr<sup>1,\*</sup>

<sup>1</sup>LifeB, Chromatin Molecular Complexes, CEITEC and Functional Genomics and Proteomics, National Centre for Biomolecular Research, Faculty of Science, Masaryk University, Brno CZ-62500, Czech Republic and <sup>2</sup>Structural Biology of Gene Regulation, CEITEC, Masaryk University, Brno CZ-62500, Czech Republic

Nucleic Acids Research, gkx812, https://doi.org/10.1093/nar/gkx812

The authors wish to make the following correction to their article:

Under 'Data Availability', accession number SASDC89 is incorrect and should be SASDB89.

This error has been corrected in the article.

\*To whom correspondence should be addressed. Tel: +420 549 495 952; Fax: +420 549 492 640; Email: hofr@sci.muni.cz

© The Author(s) 2017. Published by Oxford University Press on behalf of Nucleic Acids Research.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com