



Published in final edited form as:

Org Lett. 2017 August 18; 19(16): 4247–4250. doi:10.1021/acs.orglett.7b01901.

Synthesis of 1,2-Bis(trifluoromethylthio)arenes via Aryne Intermediates


Milad Mesgar and Olafs Daugulis*

Department of Chemistry, University of Houston, Houston, Texas 77204, United States

Abstract

A general method for synthesis of 1,2-bis-trifluoromethylthioarenes has been developed. Arynes generated from silyl aryl triflates or halides react with bis(trifluoromethyl)disulfide to afford 1,2-bis-trifluoromethylthioarenes. Aryl, alkyl, ester, halide, and methoxy functionalities are compatible with reaction conditions. Use of bis(perfluoroaryl)-disulfides gave moderate yields of aryne disulfenylation or cyclization to fluorinated dibenzothiophenes.

Graphical Abstract



Fluorine-containing organic substances are important in medicine, agriculture, and as materials.¹ Among fluorinated organics, aryl trifluoromethyl sulfides (ArSCF₃) are particularly interesting because they are present in many biologically active substances (Figure 1).² Furthermore, trifluoromethylthiolation of organic compounds is less studied than fluorination and trifluoromethylation.³ For many years, the trifluoromethylthio moiety was generated either by halogen–fluorine exchange in chloro- or bromomethyl sulfides or by trifluoromethylation of thiols.^{3g,h} Trifluoromethylthiolation by nucleophilic substitution usually requires harsh reaction conditions and stoichiometric amounts of transition metals.⁴ Recently, however, efficient methods utilizing milder protocols have been developed for trifluoromethylthiolation of aryl halides, boronic acids, and sp² C–H bonds under transition metal catalysis.^{5–7} Synthesis of 1,2-bis(trifluoromethylthio)arenes has been reported in a very limited number of publications, and the reactions appear to lack generality.⁸ For example, electrophilic aromatic substitution affords low yields of 1,2-bis(trifluoromethylthio)benzene.^{8a} Multistep reactions involving cyclo-addition of

*Corresponding Author: olafs@uh.edu.

ORCID

Olafs Daugulis: 0000-0003-2642-2992

Notes

The authors declare no competing financial interest.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.7b01901. Detailed experimental procedures and characterization data for new compounds (PDF)

bis(trifluoro-methylthio)acetylene have also been reported.^{8b} Perhaps the most general method involves reaction of trifluoromethylthiocopper with aromatic halides.^{8d} Unfortunately, trifluoromethylthiocopper is not commercially available. We report here a convenient method for the preparation of 1,2-bis(trifluoromethylthio)arenes from silyl aryl triflates or halides and commercially available bis(trifluoromethyl)disulfide.

A few examples of aryne reactions with trifluoromethylthio-containing nucleophiles have been reported in the literature.⁹ In 2008, Kolomeitsev reported one example of trifluoromethyl-thiobenzene synthesis in the reaction between trimethylsilyl-phenyl triflate, fluoride, and trifluoromethylsulfide anion.^{9a} In 2013, Lee and co-workers disclosed silver-mediated trifluoromethylthiolation via hexadehydro Diels–Alder reaction where a trifluoromethylthio substituent is introduced by employing excess AgSCF₃ reagent.^{9b} Recently, Hu reported the reaction of in situ generated arynes with excess AgSCF₃, affording aryl trifluoromethylsulfides in good yields.^{9c} In all these examples, the source of trifluoromethylthio substituent is not commercially available, and synthesis of bis-1,2-(trifluoromethyl-thio)-arenes was not reported.

Aryne insertion in disulfide S–S bonds, which is a potential route to 1,2-bis(trifluoromethylthio)arenes, has not been extensively studied.¹⁰ Aryne generation by employing strong bases is not compatible with electrophilic bis(trifluoromethyl)-disulfide.¹¹ Consequently, the mildest conditions of aryne generation from silyl aryl triflates and silyl aryl halides were employed.¹² Three different conditions of aryne generation from silyl aryl triflates or halides were explored based on our previous work.^{12b}

Aryne insertion in the bis(trifluoromethyl)disulfide S–S bond proceeds with best yields at relatively high temperatures. Use of cesium fluoride in dimethoxyethane afforded the highest yields. Other fluoride sources, such as tetramethylammonium fluoride, and solvents such as acetonitrile gave lower yields.¹³ The scope of the reaction with respect to silyl aryl triflates is presented in Table 1. Substrates possessing both electron-withdrawing and electron-releasing substituents are reactive. A methoxy substituent is tolerated affording products in good yields (entries 1–3). Phenyl-substituted silyl aryl triflate is reactive as well, giving bis(trifluoromethylthio)biphenyl in 78% yield (entry 4). Alkyl-substituted substrates afford products in 64–74% yields (entries 5, 8, 10, and 12).

A fluorinated substrate gives bis(trifluoromethylthio)- fluorobenzene in an acceptable yield (entry 7). Chlorinated substrates provide products in 44% and 65% yields (entries 11 and 13). An ester substituent is tolerated as well, and product is obtained in 41% yield (entry 14). Several of the products are very volatile, and their NMR yields were determined (entries 5–7, 10, 13). Silyl aryl triflate **4** derived from estrone was converted to bis(trifluoromethylthio) derivative **5** in 83% yield, showing relevance of the methodology to the modification of biologically active compounds (Scheme 1).

Unfortunately, only a few silyl aryl triflates are commercially available. In addition, their preparation requires lengthy synthetic procedures starting from *ortho*-bromophenols, which are of limited availability as well. In contrast, silyl aryl bromides and iodides can be prepared in one step from commercially available starting materials. Arynes can be

generated from these compounds under conditions similar to those employed for silyl aryl triflates.^{12b} Consequently, bis(trifluoromethyl-thiolation) of silyl aryl halides would be practical and convenient. A potential issue that could interfere with the desired reaction pathway is the trapping of arylcesium intermediate **7** by the highly electrophilic bis(trifluoromethyl)-disulfide affording monotrifluoromethylthiolation product **9** (Scheme 2). The ease of aryne formation from *ortho*-metalated species **7** is related to the strength of the leaving group (X^-) conjugate acid. Examination of hydrogen iodide and triflic acid DMSO pK_a values shows that the acidity of HI is close to that of CF_3SO_3H .¹⁴

As expected, use of silyl aryl bromides was problematic due to 2-bromoarylcium trapping with bis(trifluoromethyl)-disulfide (Scheme 3). Out of many silyl aryl bromide substrates subjected to the reaction conditions, only two gave modest yields of the desired products. Compounds **10** and **13** were reacted with bis(trifluoromethyl)disulfide to afford 61% and 53% isolated yields of **11** and **14**, respectively. Crude reaction mixture NMR spectra showed that, in addition to the desired **11** and **14**, substantial amounts of monotrifluoromethyl-sulfonylation products **12** and **15** were formed.

Considering the above results, silyl aryl iodides were used as aryne precursors. Silyl aryl iodide reactions with bis(trifluoromethyl)disulfide are summarized in Table 2. 1-Iodo-3-methoxy-2-(dimethylsilyl)benzene reacts with bis(trifluoromethyl)-disulfide to afford the product in 74% isolated yield (entry 1). 3,6-Dimethoxy-1,2-bis(trifluoromethylthio)benzene was synthesized in 71% yield (entry 2). 1,2-Bis(trifluoromethylthio)benzene was prepared in 58% yield (entry 3). Entries 4 and 5 show the formation of symmetric alkoxy-substituted products in 63% and 58% isolated yields, respectively. 1-Iodo-3,4,5-trimethoxy-2-(dimethylsilyl)benzene reacts with bis(trifluoromethyl)disulfide to afford the product in 82% isolated yield (entry 6). While the reactions appear to be more general than those of corresponding silyl aryl bromides, substantial amounts of monotrifluoromethyl-sulfonylation products are observed in all reaction mixtures. The bis-/monofunctionalized product crude ratios range from 2.4/1.0 (entry 3) to 4.5/1.0 (entry 6) showing the high electrophilicity of bis(trifluoromethyl)disulfide.

Next, silyl aryl triflate reactions with bis(perfluoroaryl)-disulfides were examined. 4-*tert*-Butyl and 3-chlorobenzynes generated from the corresponding silyl aryl triflates were reacted with bis(pentafluorophenyl)disulfide (Scheme 4). Isolated yields of the products were 26% and 21%, respectively. Attempts to improve reaction efficiency were not successful.

Interestingly, when bis(perfluoro-*p*-tolyl)disulfide was reacted with trimethylsilylphenyl triflate, a different pathway was followed (Scheme 5). Instead of aryne formal insertion into the disulfide S-S bond, a single addition of sulfide followed by intramolecular nucleophilic substitution of fluoride on the aromatic ring occurred, affording a polyfluorinated benzothiophene **23** in 67% yield. The isolation of **23** argues against a concerted aryne reaction with disulfides. Instead, initial nucleophilic attack on aryne would generate an anionic intermediate, which subsequently gives a 1,2-disulfonylation product. Alternatively, in the presence of a strongly electrophilic polyfluorinated aryl ring, internal trapping affords **23**.¹⁵

In summary, a general method for synthesis of 1,2-bis-trifluoromethylthioarenes has been developed. Arynes generated from silyl aryl triflates and silyl aryl halides react with bis(trifluoromethyl)disulfide to afford 1,2-bis(trifluoromethylthio)arenes. Aryl, alkyl, ester, halide, and methoxy functionalities are compatible with reaction conditions. Use of bis(perfluoroaryl)disulfide reagents gave moderate yields of disulfenylation products or, in the case of bis(perfluoro-*p*-tolyl)disulfide, cyclization to fluorinated dibenzothiophenes. Silyl aryl bromides and iodides gave substantial amounts of monotrifluoromethylsulfenylation products due to the high electrophilicity of bis(trifluoromethyl)disulfide.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Welch Foundation (Chair No. E-0044) and NIGMS (Grant No. R01GM077635) for supporting this research.

References

1. Wang J, Sánchez-Roselló M, Acuña JL, Del Pozo C, Sorochinsky AE, Fustero S, Soloshonok VA, Liu H. *Chem Rev.* 2014; 114:2432. [PubMed: 24299176]
2. (a) Leroux F, Jeschke P, Schlosser M. *Chem Rev.* 2005; 105:827. [PubMed: 15755078] (b) Boiko VN, Beilstein J *Org Chem.* 2010; 6:880. [PubMed: 20978611] (c) Durley RC, Grapperhaus ML, Massa MA, Mischke DA, Parnas BL, Fobian YM, Rath NP, Honda DD, Zeng M, Connolly DT, Heuvelman DM, Witherbee BJ, Glenn KC, Krul ES, Smith ME, Sikorski JA. *J Med Chem.* 2000; 43:4575. [PubMed: 11101348]
3. (a) Truong T, Klimovica K, Daugulis O. *J Am Chem Soc.* 2013; 135:9342. [PubMed: 23758609] (b) Fier PS, Hartwig JF. *Science.* 2013; 342:956. [PubMed: 24264986] (c) Ye Y, Schimler SD, Hanley PS, Sanford MS. *J Am Chem Soc.* 2013; 135:16292. [PubMed: 24160267] (d) Zhang Q, Brusoe AT, Mascitti V, Hesp KD, Blakemore DC, Kohrt JT, Hartwig JF. *Angew Chem, Int Ed.* 2016; 55:9758. (e) Dubinina GG, Furutachi H, Vicic DA. *J Am Chem Soc.* 2008; 130:8600. [PubMed: 18543912] (f) Novák P, Lishchynskiy A, Grushin VV. *J Am Chem Soc.* 2012; 134:16167. [PubMed: 22998369] (g) Nodiff EA, Lipschutz S, Craig PN, Gordon MJ. *J Org Chem.* 1960; 25:60. (h) Umemoto T, Ishihara S. *J Am Chem Soc.* 1993; 115:2156.
4. (a) Manteau B, Pazenok S, Vors JP, Leroux FR. *J Fluorine Chem.* 2010; 131:140. (b) Baert F, Colomb J, Billard T. *Angew Chem, Int Ed.* 2012; 51:10382.
5. (a) Teverovskiy G, Surry DS, Buchwald SL. *Angew Chem, Int Ed.* 2011; 50:7312. (b) Zhang CP, Vicic DA. *J Am Chem Soc.* 2012; 134:183. [PubMed: 22148268]
6. Chen C, Xie Y, Chu L, Wang RW, Zhang X, Qing FL. *Angew Chem, Int Ed.* 2012; 51:2492.
7. Tran LD, Popov I, Daugulis O. *J Am Chem Soc.* 2012; 134:18237. [PubMed: 23102009]
8. (a) Haas A, Hellwig V. *J Fluorine Chem.* 1975; 6:521. (b) Haas A, Krächter HU. *Chem Ber.* 1988; 121:1833. (c) Scribner RM. *J Org Chem.* 1966; 31:3671. (d) Kondratenko NV, Kolomeitsev AA, Popov VI, Yagupolskii LM. *Synthesis.* 1985; 1985:667.
9. (a) Kolomeitsev AA, Vorobyev M, Gillandt H. *Tetrahedron Lett.* 2008; 49:449. (b) Wang KP, Yun SY, Mamidipalli P, Lee D. *Chem Sci.* 2013; 4:3205. (c) Zeng Y, Hu J. *Org Lett.* 2016; 18:856. [PubMed: 26849240]
10. Nakayama J, Tajiri T, Hoshino M. *Bull Chem Soc Jpn.* 1986; 59:2907.
11. (a) Truong T, Daugulis O. *J Am Chem Soc.* 2011; 133:4243. [PubMed: 21384889] (b) Truong T, Daugulis O. *Chem Sci.* 2013; 4:531. [PubMed: 24077102] (c) Truong T, Mesgar M, Le KKA, Daugulis O. *J Am Chem Soc.* 2014; 136:8568. [PubMed: 24893069]

12. (a) Himeshima Y, Sonoda T, Kobayashi H. *Chem Lett.* 1983; 12:1211.(b) Mesgar M, Daugulis O. *Org Lett.* 2016; 18:3910. [PubMed: 27415183]
13. Please see Supporting Information for optimization details.
14. (a) Bordwell FG. *Acc Chem Res.* 1988; 21:456.(b) Trummal A, Lipping L, Kaljurand I, Koppel IA, Leito IJ. *J Phys Chem A.* 2016; 120:3663. [PubMed: 27115918]
15. Travieso-Puente R, Budzak S, Chen J, Stacko P, Jastrzebski JTBH, Jacquemin D, Otten E. *J Am Chem Soc.* 2017; 139:3328. [PubMed: 28218846]







Figure 1. Bioactive molecules possessing trifluoromethylthioarene functionality.




Scheme 1.
Bis(trifluoromethylthiolation) of Estrone Derivative




Scheme 2.
Reactivity of Arylmetal Intermediate



Scheme 3.
Silyl Aryl Bromide Reactions



Scheme 4.
Reactions with Bis(pentafluoroaryl)disulfides




Scheme 5.
Reaction with Bis(perfluorotolyl)disulfide

Table 1

Bis(trifluoromethylthiolation) of Silyl Aryl Triflates^a

entry	aryne precursor	product	yield (%)
1			75
2			76
3			84
4			78
5 ^b			64
6 ^b			71
7 ^b			56



entry	aryne precursor	product	yield (%)
8			74 81 ^c
9			69
10 ^b			66
11			44
12			68
13 ^b			65
14			41

^aReaction conditions: Aryne precursor (0.3 or 0.5 mmol), CsF (3 equiv), CF₃SSCF₃ (3.5 equiv), DME (2 mL), 24–48 h, 85 or 110 °C. Yields are isolated yields.

^bYield determined by NMR with PhCF₃ as an internal standard. DME is dimethoxyethane.

^cScale: 1.0 mmol. Please see Supporting Information for details.

Table 2

Silyl Aryl Iodide Reactions^a

entry	aryne precursor	16	yield (%) (16/17)
1			74 3.7:1.0
2			71 ^b 3.1:1.0
3			58 ^c 2.4:1.0
4			63 2.9:1.0
5			58 2.8:1.0
6			82 4.5:1.0
7			60 2.8:1.0

^aReaction conditions: Aryne precursor (0.3 mmol), CsF (3 equiv), CF₃SSCF₃ (3.5 equiv), DME (2 mL), 37–46 h. Yields are isolated yields.

^b3,6-Dimethoxy-2-trifluoromethylthiodobenzene (23%) also isolated.

^cYield determined by NMR with PhCF₃ as an internal standard. Please see Supporting Information for details.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript