

SCIENTIFIC REPORTS



OPEN

Effect of different organic fertilizers application on growth and environmental risk of nitrate under a vegetable field

Shuyan Li¹, Jijin Li², Bangxi Zhang^{1,3}, Danyang Li¹, Guoxue Li¹ & Yangyang Li¹

The effect of chicken manure after different disposal methods (water-logged composting, GOF; anaerobic digestion, BR; thermophilic composting, ROF) on vegetable growth and environmental risk was investigated under the tomato-celery-tomato field. Results showed that organic fertilizers significantly increased vegetable yield and quality, but with inappropriate application may cause serious environmental risk such as nitrate pollution. Maximum vegetable yield of 80.9, 68.3, 112.7 t·ha⁻¹ (first, second and third rotation crop, respectively) with best vegetable quality was obtained in ROF treatment. The highest N use efficiency with the least nitrate enrichment in soil was also found in ROF treatment. Moreover, under this fertilization way, nitrate concentration in soil leachate dropped to 6.4 mg·L⁻¹, which satisfied the threshold (<10 mg·L⁻¹) for drinking water set by the US Environmental Protection Agency. Thus, ROF was suggested to be the optimal fertilizer with the best yield, quality and the least environmental risk under the “tomato-celery” rotation system.

Although nitrogen (N) utilization has generally been optimized in agriculture, unreasonable fertilization can lead to agricultural non-point source pollution^{1–6}, and improvements are necessary to avoid adverse environmental impacts of nitrate leaching. Nitrate leaching has a significant influence on plant N supply and groundwater quality. Nitrate concentrations in soil depend on the relation between uptake by plants, soil organisms, atmospheric N₂ fixation, N mineralization (ammonification and nitrification), N deposition from the atmosphere, denitrification, and volatilization⁷.

The development of intensive agricultural areas based on irrigation with groundwater and N application in farming areas has had serious side effects on the land ecosystems including ground water depletion and nitrate leaching to ground water^{6,8}. Due to environmental pollution, high nitrate concentrations may accumulate in the edible parts of some vegetables, particularly if excessive N fertilizer has been applied⁹. Consuming these crops can harm human health.

Leaching of nitrate from soil is driven by land-use type, management (e.g., fertilization), land-use change, climate, and soil properties¹⁰. Nitrate-N leaching losses were usually less from fine-textured soils than from coarse-textured soil¹¹. The soil nitrate content may higher in spring than in autumn¹². Precipitation/irrigation can significantly increase the nitrates in the soil leachate^{13,14}. Nitrate losses decreased with the drain depth decreased¹⁵.

Organic fertilizers have been proposed as one solution to relieve environmental pressure and be a carbon-neutral alternative to liquid fossil fertilizers¹⁶. Organic matter improves soil structure, increases the water holding capacity and promotes biological transformations such as N-mineralization^{16,17}. Several researchers have examined the impact of timing of N and water applications on crop yield in field experiments^{8,18}. Behnke *et al.*¹⁹ found that N annual losses from 22.7 to 59.9 kg·ha⁻¹, and they increase with N fertilization rates increase. The soil NO₃⁻-N content under basal fertilizer was 1.65 times higher than that without fertilizer at 0–10 cm on the 36th day after sowing²⁰. Davis *et al.*²¹ found that N applications increased N leaching and N₂O emission without increasing biomass production. Liu *et al.*⁹ found that lettuce augmented with organic fertilizers had significantly

¹Beijing Key Laboratory of Farmland Soil Pollution Prevention and Remediation, College of Resources and Environmental Science, China Agricultural University, Beijing, 100193, China. ²Institute of Plant Nutrition and Resources, Beijing Academy of Agriculture and Forestry Sciences, Beijing, 100097, China. ³Guizhou Institute of Soil and Fertilizer, Guiyang, 550006, China. Correspondence and requests for materials should be addressed to G.L. (email: ligx@cau.edu.cn)




Figure 1. Effect of different organic fertilizers on vegetable yield. Mean differences in the bars are significant at $P_{0.05}$ level with different letters. Tomato 1 means the first rotation vegetable; Celery means the second rotation vegetable; Tomato 2 means the third rotation vegetable. Repeat in following figure.

longer and wider leaves, higher shoot, and lower NO_3^- -N concentrations compared with the same amount of inorganic fertilizers. Guo *et al.*²² found that N fertilizers coupled with farm yard manure resulted in 70% less NO_3^- -N accumulation in the soil profiles than that using mineral N fertilizer alone. However, some researches had found that manure applications without any pretreatment could cause serious NO_3^- -N leaching^{23,24}.

There have been many kinds of organic fertilizers, such as manure, sewage sludge, stalks, compost, biogas residues, biogas slurry and so on. An increasing body of literature has been focused on the N fertilizers for crop yield and NO_3^- -N leaching, but very little is about comparing different kinds of organic fertilizers on NO_3^- -N distribution (soil, leachate and crop), vegetable yield and quality during the agricultural process. To solve the problem of nitrate content in vegetables, soil and underground water exceeding standard caused by unreasonable fertilization, specific objectives of this study were to: (i) evaluate different organic fertilizers on vegetable yield and quality; (ii) and also determine nitrate concentrations in different soil layers and soil leachate to evaluate environmental risk.

Results and Discussion

Vegetable yield. As expected, organic fertilizers significantly increased vegetable yield by 7.6–45.2% (Fig. 1). For the first rotation, tomato yield increased by 9.2–20.1% compared with CK. Among this, ROF did best with the yield of 84.9 $\text{t}\cdot\text{ha}^{-1}$ and it was significantly higher than other treatments. However, GOF had the least effect on tomato yield and BR was similar to GOF with the increase of 9.8%. In the second rotation, all the treatments increased the celery yield by 7.6–8.4%. The maximum increase was ROF treatment with the production of 68.3 $\text{t}\cdot\text{ha}^{-1}$. Compared with tomato, the increase of celery was not obvious. After application of organic fertilizers for one year, the tomato yield in all the treatments in this rotation increased compared with the first rotation. This was mainly due to the higher N mineralization as a result of higher biological activity²⁵. For the third rotation crop, tomato yield increased by 25.8–45.2% compared with CK. In this time, ROF has the maximum yield with 112.7 $\text{t}\cdot\text{ha}^{-1}$. Among this “tomato-celery-tomato” system, organic fertilizers significantly increased the vegetable yield according to ANOVA test. This may because organic fertilizer application can increase soil organic matter and then increase yields^{26,27}.

Vegetable quality. Application of organic fertilizers can increase vegetable qualities (Fig. 2). The concentration of vitamin C (Vc) after harvest the vegetables is shown in Fig. 2a. Application organic fertilizers significantly increased the concentration of Vc by 3.0–33.5% in the first rotation crop. ROF with the Vc concentration of 122 $\text{mg}\cdot\text{kg}^{-1}$ had the best effect and GOF did worst. This may because GOF had a low humification degree without a thermophilic phase⁹. The concentration of Vc increased by 12.6–31.5% in celery planting. Like the first rotation, ROF with the concentration of 83.5 $\text{mg}\cdot\text{kg}^{-1}$ had the best effect and GOF had the least effect. After three rotations, Vc of tomato in CK treatment decreased from 91.4 (first rotation) to 79.0 (third rotation) $\text{mg}\cdot\text{kg}^{-1}$, indicating undernourishment and N depletion. Organic fertilizers application could increase the concentration of Vc by 31.6–48.1% compared with CK in the third rotation. ROF with the concentration of 117 $\text{mg}\cdot\text{kg}^{-1}$ had the best effect. This is mainly because ROF with a high stabilization and humification degree could improve soil structure, increase the water holding capacity and promote biological transformations and then improve the vegetable quality⁹.

Figure 2b gives the concentration of soluble sugar (SE) after harvest the vegetables. Application organic fertilizers can significantly increase the concentration of SE by 9.9–17.3% in the first rotation. ROF had the best effect with the concentration of 3.67%. The concentration of SE was increased by 23.6–55.6% in celery planting. ROF with the concentration of 1.12% had the best effect among all the treatments, and GOF had the least effect of all. In the third rotation, the concentration of SE was increased by 18.2–30.3%. Similar to the celery, ROF with




Figure 2. Effect of different organic fertilizers on vegetable quality Vc: vitamin C; SE: soluble sugar; TA: titratable acidity.




Figure 3. Effect of different organic fertilizers on nitrate content of tomato and celery.

the concentration of 4.30% had the best effect of all, and GOF had the least effect. The concentration of SE with the third rotation increased in all treatments including CK compared with the first rotation. This illustrates that long-term application of organic fertilizer can improve the quality of vegetables.

The concentration of titratable acidity (TA) after harvest tomatoes is shown in Fig. 2c. Organic fertilizers application has no significant influence on TA. The content of TA in the third rotation decreased compared with the first rotation, indicating that organic fertilizer can improve the vegetable taste.

Nitrate concentration in vegetable. NO_3^- -N concentration is an important quality characteristic of vegetable. NO_3^- was perceived as a purely harmful dietary component which causes infantile methaemoglobinaemia, carcinogenesis and possibly even teratogenesis²⁸. Figure 3 gives the NO_3^- -N concentration in tomato and celery. From this, NO_3^- -N concentrations of the two rotations of tomato were all less than $120 \text{ mg}\cdot\text{kg}^{-1}$ especially in the latter, which were far less than the limit of the national standard $600 \text{ mg}\cdot\text{kg}^{-1}$ (GB18406.1-2001). Celery is a crop which is easy to enrich NO_3^- and this is why the NO_3^- -N concentration in the third rotation of tomato lower than the first rotation especially in CK treatment. NO_3^- -N concentration in celery was much higher than that in tomato, but it was still less than the limit of the national standard $3000 \text{ mg}\cdot\text{kg}^{-1}$. Tomato-celery rotation system could significantly decrease the vegetable NO_3^- -N concentration under this continuous fertilization field. Through these three rotations, ROF had no significant difference with CK in terms of vegetable NO_3^- -N concentration, which indicates that ROF is a relatively safe way for fertilizing.

Nitrate content in soil. Nitrate concentration in the 0–60 cm soil layers (in time). NO_3^- -N concentration in the root zone soil of the three rotations are shown in Fig. 4. Organic fertilizers significantly affect the soil nitrate




Figure 4. Nitrate content in 0–60 cm soil profile after harvest vegetables (with time).

Physicochemical indices	Concentration
Soil texture	Clay loam
Clay (%) ^a	22.8 ± 0.8
Silt (%) ^a	43.2 ± 1.2
Sand (%) ^a	34.0 ± 0.5
pH	8.07 ± 0.07
Organic matter (g·kg ⁻¹) ^a	19.0 ± 0.2
Total N (g·kg ⁻¹) ^a	1.34 ± 0.06
Alkali-hydrolyzable N (mg·kg ⁻¹) ^a	116 ± 4
Olsen-P (mg·kg ⁻¹) ^a	105 ± 3
Rapidly available K (mg·kg ⁻¹) ^a	248 ± 8
Available Cu (mg·kg ⁻¹) ^a	3.35 ± 0.07
Available Zn (mg·kg ⁻¹) ^a	5.64 ± 0.06
Available Ca (mg·kg ⁻¹) ^a	3560 ± 42
Available Fe (mg·kg ⁻¹) ^a	31.4 ± 1.3
Available Mn (mg·kg ⁻¹) ^a	4.21 ± 0.07
Available Mg(mg·kg ⁻¹) ^a	2370 ± 35

Table 1. Characteristics of original soil. Values are given as the mean ± standard deviation (n = 3). ^aBased on dry matter (DM).

concentration in the top layers (0–60 cm). The NO_3^- -N concentration in the 0–30 cm soil layer of GOF and ROF treatments achieved the minimum value after harvest the second rotation (celery). Celery roots were mainly distributed in 0–30 cm²⁹ resulting in less absorbing of NO_3^- -N below 30 cm soil layer. This may be the reason why the NO_3^- -N concentration in 30–60 cm soil layer higher than that in 0–30 cm soil layer after harvest celery. Furthermore, for the second rotation NO_3^- -N content in the top 30 cm soil layer reduced by 60–80% compared with the first rotation, but NO_3^- -N in the 30–60 cm soil layer may have a small amount of accumulation for little absorption. However, there were no significant differences in the soil NO_3^- -N content in terms of the whole top layers (0–60 cm) compared with CK, due to NO_3^- -N absorption and enrichment in celery. The soil NO_3^- -N concentration after harvest the first rotation of tomato was significantly higher than the other rotations. This may be caused by the high nitrate content of the original soil (Table 1). Organic fertilizers application significantly increased the NO_3^- -N concentration in the 0–30 cm soil layer of all treatments after the first rotation, and especially in BR reached 138.2 g·kg⁻¹, indicating a high risk of leaching. However, NO_3^- -N in the 30–60 cm soil layer changed slightly due to root absorption. After third rotation, NO_3^- -N concentration was significantly lower than the first rotation, owing to the low nitrate background values in this rotation after harvest celery. In this time, NO_3^- -N concentration in the 0–30 cm soil layer increased slightly. Moreover, NO_3^- -N concentration in the 30–60 cm reduced significantly, which could be attributable to the higher absorption of N in the 30–60 cm soil layer by the deeper root of tomato³⁰. Thus in this tomato- celery rotation system, long time application of organic fertilizers will not affect soil nitrate content in the top layers (0–60 cm).

Nitrate content in soil profile (in depth). Organic fertilizer significantly affected soil NO_3^- -N concentration in the 0–175 cm soil layers (Fig. 5). Due to incomplete utilization of fertilizer, treatments with the organic fertilizers




Figure 5. Nitrate content in soil profile after harvest the third rotation (with depth).

Treatment	N_{initial} ($\text{kg}\cdot\text{ha}^{-1}$)	N_{input} ($\text{kg}\cdot\text{ha}^{-1}$)	N_{min} ($\text{kg}\cdot\text{ha}^{-1}$)	N_{uptake} ($\text{kg}\cdot\text{ha}^{-1}$)	N_{residual} ($\text{kg}\cdot\text{ha}^{-1}$)	$N_{\text{utilization}}$ ($\text{kg}\cdot\text{ha}^{-1}$)	NUE (%)
CK	396 ± 18	0	145 ± 12	481 ± 21	60 ± 10	—	—
GOF		1050		593 ± 34	122 ± 10	204	19.4
BR		1050		600 ± 26	120 ± 18	208	19.8
ROF		1050		726 ± 28	100 ± 15	314	30.0

Table 2. N balance and N translocation from soil and fertilizer to vegetable. Values are given as the mean ± standard deviation (n = 9).

increased the nitrate content in soil profile especially in the top layers compared with CK. Topsoil had the most obvious effect, above all GOF treatment reached 29.9 $\text{mg}\cdot\text{kg}^{-1}$, and this may be result from nitrification and mineralization for its instability²⁰. Conversely, owing to the higher humification and stability degree of ROF³¹, nitrate content in ROF treatment was lower than any other fertilization treatments. After application of BR and GOF, nitrate had a dramatic enrichment in deep soil (especially below 75 cm) indicating N surplus, and such accumulation of NO_3^- -N in soil profile posed a high risk of N leaching into groundwater. BR and GOF were all incomplete fermentation without thermophilic phase, and then they had a low humification degree with very little stabilized organic matters³². Thus a large amount of nitrogen in BR and ROF cannot be fixed like ROF, which lead to nitrate leaching seriously. In ROF treatment, nitrate content in the soil below 100 cm almost had no difference with CK. From the perspective of security, ROF is the environmentally friendly way for fertilizing.

Nitrogen balance and N translocation. Calculation of N balance is one potentially useful method for predicting the risk of nitrate leaching into groundwater¹⁰. N balance in each treatment was calculated under this tomato-celery rotation system (Table 2). Without fertilizer, the amount of N_{min} could achieve about 145 $\text{kg}\cdot\text{ha}^{-1}$. However, the N_{residual} level was lower than the N_{initial} level, indicating soil N depletion in some degree. N_{uptake} in the fertilization treatments were higher than that in CK treatment, especially in the ROF treatment, showing that fertilization can promote the absorption of N by root. After organic fertilizer application, the residual NO_3^- -N in the 0–60 cm soil layer after crop harvest accumulated to 100–122 $\text{kg}\cdot\text{ha}^{-1}$. Although this was still higher than the environmental safety standard in Europe (90–100 $\text{kg}\cdot\text{ha}^{-1}$ in the 0–100 cm soil layer)¹⁰, it resulted in 50–75% less NO_3^- -N accumulation in the soil profiles than the initial soil, and therefore the environmental risk was reduced

Treatment	Original	Tomato 1	Celery	Tomato 2
CK	38.1 ± 0.2	10.2 ± 0.2	4.2 ± 0.1	6.3 ± 0.1
GOF	38.2 ± 0.1	33.8 ± 0.3	28.5 ± 0.2	31.6 ± 0.3
BR	37.2 ± 0.3	35.1 ± 0.4	30.3 ± 0.3	32.1 ± 0.3
ROF	37.4 ± 0.3	11.6 ± 0.2	8.2 ± 0.1	6.4 ± 0.2

Table 3. NO₃⁻-N concentrations in the soil leachate after harvest the vegetables (mg·L⁻¹). Values are given as the mean ± standard deviation (n = 9).

in some degree. These results suggested that organic fertilizer application could be benefit for crop uptake, reduce the NO₃⁻-N in the soil and then alleviate the soil NO₃⁻-N leaching. NUE in these fertilization treatments were 19.4–30.0%, the ROF treatment presented the highest NUE among all the treatments due to the highest uptake by crops, implying the optimum fertilization way.

Nitrate concentration of soil leachate. Although the organic fertilizer application could be benefit for crop uptake and reduce the NO₃⁻-N in the soil, soil are still at the high risk of leaching with the high N_{residual} and low NUE in all the fertilization treatments. Then the NO₃⁻-N concentration of soil leachate at 100 cm depth below the soil surface were detected after harvest vegetables. NO₃⁻-N concentration of soil leachate varied with treatments and crop types, ranging from 6.3 to 35.1 mg·L⁻¹ for tomato and from 4.2 to 30.3 mg·L⁻¹ for celery (Table 3). Soil NO₃⁻-N leaching in tomato seasons was generally higher than in celery seasons due to higher crop N uptake and higher evaporation in celery rotation leading to less drainage into deeper layers. NO₃⁻-N leaching in all treatments decreased after application of organic fertilizers especially ROF. Fertilizer type significantly affects the NO₃⁻-N concentration in the soil leachate. The least NO₃⁻-N leaching was observed in the ROF treatment mainly due to ROF with a higher organic matter content and biological activity, stabilization and humification degree, resulting in the increase of soil aggregation, nitrogen fixation capacity and decrease of NO₃⁻-N leaching²⁵. Moreover, NO₃⁻-N concentration in CK and ROF treatments dropped below 10 mg·L⁻¹ after harvest the second rotation (celery), which satisfied the threshold (<10 mg·L⁻¹) for drinking water set by the US Environmental Protection Agency. This result suggested that application of ROF was no more likely to impair groundwater quality than the GOF, BR or even CK treatments.

Conclusions

Organic fertilizers significantly increased vegetable yield and quality, but with inappropriate application may cause serious environmental risk. Maximum vegetable yield of 80.9, 68.3, 112.7 t·ha⁻¹ (first, second and third rotation crop, respectively) with best vegetable quality was obtained in ROF treatment. The highest N use efficiency with the least nitrate enrichment in soil was also found in ROF treatment. Moreover, under this fertilization way, nitrate concentration in soil leachate satisfied the threshold for drinking water. Thus, ROF was suggested to be the optimal fertilizer with the best yield, quality and the least environmental risk under the “tomato-celery” rotation system.

Materials and methods

Site description. One and a half years of field experiments (tomato1-celery-tomato2) were conducted on clay loam soil at Liuminying Agricultural Ecological Station (39°41' N, 116°34' E) in southeast suburb area (Daxing district) of Beijing, northwest edge of North China Plain. The soil was calcareous, alkaline, and rich in phosphorus and potassium. Agriculture in the area is intensified by a double cropping system (two vegetables a year) with high-yielding cultivar and high inorganic fertilizer (more than 1000 kgN·ha⁻¹·yr⁻¹) input. Some of the characteristics of this soil were determined before this experiment (Table 1). The average air temperature during tomato planting period was about 25 °C, while in celery planting period was about 18 °C.

Crop rotation and experimental fertilizers. A typical spring tomato–autumn celery double cropping rotation was chosen, representative of the common farming practices in the area, where tomato is usually planted from March to July and celery from August to October. Tomato cultivar with Israel 1420 greenhouse grown tomato (*Lycopersicon esculentum* Mill.) was planted in the experimental plot (see Section 2.3) at a density of 36,000 plant·ha⁻¹. After tomato harvest, soil was ploughed before planting autumn celery. Celery cultivar with California celery (*Apium graveolens* L) was planted in the experimental plot at a rate of 2,300,000 plant·ha⁻¹. The selected crop varieties and planting densities is representative of that used by local farmers.

In order to evaluate of agronomic and ecological effects of soil amendment, three kinds of common organic material i.e. general organic fertilizer (GOF), biogas residue (BR) and refined organic fertilizer (ROF) were used as N fertilizer. GOF was made by chicken manure and corn stalk through water-logged composting; BR was taken from Liuminying Biogas Station, which was made by chicken manure through anaerobic digestion; and ROF was made by chicken manure and mushroom residue through a 90 days thermophilic aerobic composting. Some of the composition and characteristics of these organic fertilizers are given in Table 4.

Experimental design. The experiment was conducted in a vegetable greenhouse during the tomato and celery growing season. Four treatments with three replicates were carried out, namely CK, GOF, BR and ROF. Then the experimental area consisted of 12 plots, 5.5 m wide and 6 m long for each, and these 12 plots were arranged as split plots in a randomized complete block with a 0.5 m isolation strip in order to avoid interference. The CK was a control treatment without fertilization. GOF, BR and ROF treatments were applied with the same

Fertilizer	GOF	BR	ROF
Organic matter (g·kg ⁻¹) ^a	356 ± 3	318 ± 2	404 ± 5
Humic acid (g·kg ⁻¹) ^a	53.9 ± 1.7	61.4 ± 1.8	145 ± 7
Fulvic acid (g·kg ⁻¹) ^a	72.1 ± 2.5	59.7 ± 1.5	29.8 ± 0.9
Total N (g·kg ⁻¹) ^a	12.2 ± 0.2	13.6 ± 0.3	20.1 ± 0.2
Total P (g·kg ⁻¹) ^a	23.1 ± 0.3	51.0 ± 0.4	23.6 ± 0.2
Total K (g·kg ⁻¹) ^a	13.7 ± 0.2	9.0 ± 0.1	20.2 ± 0.2
Total Cu (mg·kg ⁻¹) ^a	128 ± 1.7	89.1 ± 1.2	75.7 ± 0.9
Total Zn (mg·kg ⁻¹) ^a	210 ± 1.4	248 ± 0.7	181 ± 1.4
Total As (mg·kg ⁻¹) ^a	4.12 ± 0.19	3.28 ± 0.29	7.85 ± 0.47
Total Cd (mg·kg ⁻¹) ^a	2.04 ± 0.42	2.33 ± 0.07	2.15 ± 0.11
Germination index (%)	83.5 ± 3.4	97.2 ± 2.8	112 ± 4.3
Escherichia coli (logCFU·g ⁻¹)	3.12 ± 0.15	2.33 ± 0.17	ND
Moisture content (%)	43.8 ± 0.2	45.2 ± 0.1	35.2 ± 0.2

Table 4. Composition and characteristics of the three organic fertilizers. Values are given as the mean ± standard deviation (n = 3). ^aBased on dry matter (DM). ND: not detected.

amount of N with 350 kgN·ha⁻¹ for each crops. Previous study has found that top dressing can increase the crop yield²⁰. Then in this experiment, 66.7% of the fertilizer was used as base fertilizer and the remaining 33.3% as top dressing in fruit swelling period and vigorous period for tomato and celery, respectively. The management practices for controlling pest, disease and weeds complied with local practices for high-yield production.

Analytical methods. Tomato and celery plant were sampled from a 5 m² area in each plot at harvest for the measurements of vegetable (tomato and celery) yield and tomato residual biomass. Samples of vegetable and tomato residual were oven-dried at 65 °C until they reached a constant weight to determine the water content and dry matter. The N content in vegetable and tomato residual of the samples were determined by the micro-Kjeldahl method by digesting the sample in H₂SO₄-H₂O₂ solution³³. N uptake by plants was estimated by multiplying the tomato, tomato residual and celery dry matter weight by their N concentrations.

Three tomatoes (or three plants of celery) per plot with similar degree of maturity and similar size and without external defects were picked for the quality indices (mainly taste quality, nutrient quality and safety quality) measurement. Tomatoes or celeries were squeezed in a blender, and then the content of vitamin C (Vc), soluble sugar (SE), and titratable acidity (TA) in the plants were detected according to³⁴. Besides, some of the squeezed vegetable was extracted with deionized water, filtered and then the concentration of NO₃⁻-N in vegetable was determined by a continuous-flow analyzer (TRAACS 2000, Bran and Luebbe, Norderstedt, Germany).

For soil N measurements, three ceramic candle extraction systems with tubes (inside diameter 50 mm) were installed in each plot at 100 cm soil depths. The amount of nitrate leached during the growing season may be minimal compared to leaching losses that occur between the harvest of one crop and the planting of the next²³. Then samples of the soil leachate were taken after each harvest and/or before sowing. Furthermore, soil samples in all plots were taken after each harvest and/or before each planting by sampling three cores per plot with an auger (3 cm inside diameter tube) to 60 cm depth in 30 cm increments. Moreover, soil samples in the depth of 15, 45, 75, 125 and 175 cm were taken after harvesting the second batch of tomato (tomato 2) to research the change of nitrate with soil depth. Soil samples obtained from the same layer and plot were thoroughly mixed. All of the soil and soil leachate samples were immediately brought to the laboratory for the measurement of NO₃⁻-N and soil moisture content.

Each fresh soil sample was extracted with CaCl₂³⁵, and the concentration of nitrate was determined by a continuous-flow analyzer (TRAACS 2000, Bran and Luebbe, Norderstedt, Germany). Soil samples were dried to a constant weight in an oven at 105 °C to determine the water content and dry matter. Bulk density of the soils was measured in the 0–60 cm soil depth with soil cores (3 cm inside diameter by 20 cm long). The NO₃⁻-N contents in soil (mg·kg⁻¹) were converted to kg·ha⁻¹ based on the bulk density of different soil layers in order to calculate the N balance. For the nitrate analysis of soil leachate, the water samples were filtered through 0.45 μm membranes and the concentration of nitrate was determined by a continuous-flow analyzer³⁶.

Nitrogen balance. Items in the N balance were estimated in each plot during the whole crop growing seasons. NO₃⁻-N below 60 cm soil depth and NH₄⁺-N throughout the soil profile will not be included in the N balance calculations because the crop roots in this experiment were mainly distributed in the 0–60 cm depth and relatively low changes in NH₄⁺-N content between seasons were found (data not presented). The N balance can be written as:

$$N_{\text{initial}} + N_{\text{input}} + N_{\text{min}} - N_{\text{uptake}} - N_{\text{residual}} = N_{\text{surplus}} \quad (1)$$

where N_{initial} is initial soil NO₃⁻-N in the 0–60 cm soil profiles; N_{input} is N application rate (350 kg N·ha⁻¹ per rotation crop plus 3 rotation crops); N_{min} is N mineralization; N_{uptake} is N uptake by plant; N_{residual} is residual NO₃⁻-N in 0–60 cm soil profiles, and N_{surplus} represent N that store in various soil fraction (mainly organic N) and N loss.

N loss is considered as mainly NO_3^- -N leaching, since other N losses via denitrification, volatilization and erosion are relatively low under such environmental conditions as reported by Fang *et al.*³⁷.

N mineralization (N_{min}) was estimated by the balance of N inputs and outputs in the control (CK) as follows:

$$N_{\text{min}} = N_{\text{uptake},0} + N_{\text{residual},0} - N_{\text{initial},0} \quad (2)$$

where $N_{\text{uptake},0}$, $N_{\text{residual},0}$ and $N_{\text{initial},0}$ are crop N uptake, residual and initial soil NO_3^- -N in the 0–60 cm soil profile of the control, respectively.

$$N_{\text{utilization}} = N_{\text{input}} - N_{\text{surplus}} \quad (3)$$

$$\text{NUE} = N_{\text{utilization}}/N_{\text{input}} \quad (4)$$

$N_{\text{utilization}}$ is the part of N_{uptake} offered by organic fertilizer. NUE is the fertilizer N use efficiency during the one and a half years of experiment period.

Statistical analyses. Analysis of variance (ANOVA) was performed with the SAS8.2 for Windows, and mean comparisons were done using the least significant difference (LSD) test at $P < 0.05$.

Data availability statement. The authors declared that none of the data in the paper had been published or was under consideration for publication elsewhere.

References

- Eriksen, J. *et al.* Nitrate leaching and residual effect in dairy crop rotations with grass–clover leys as influenced by sward age, grazing, cutting and fertilizer regimes. *Agr Ecosyst Environ* **212**, 75–84 (2015).
- Knowlton, K. F. & Cobb, T. D. Implementing waste solutions for dairy and livestock farms. *J Dairy Sci* **89**(5), 1372–83 (2006).
- Kourakos, G. *et al.* A groundwater nonpoint source pollution modeling framework to evaluate long-term dynamics of pollutant exceedance probabilities in wells and other discharge locations. *Water Resour Res* **48**(6), 2805–2814 (2012).
- Leimer, S. *et al.* Time matters for plant diversity effects on nitrate leaching from temperate grassland. *Agr Ecosyst Environ* **211**, 155–163 (2015).
- Ouyang, W. *et al.* Non-point source pollution dynamics under long-term agricultural development and relationship with landscape dynamics. *Ecol Indic* **45**(5), 579–589 (2014).
- Reichenau, T. G. *et al.* Nitrate Leaching. Regional Assessment of Global Change Impacts. 303–310 (Springer International Publishing, 2016).
- Schimel, J. P. & Bennett, J. Nitrogen mineralization: challenges of a changing paradigm. *Ecology* **85**(3), 591–602 (2004).
- Wei, Y. *et al.* Policy incentives for reducing nitrate leaching from intensive agriculture in desert oases of Alxa, Inner Mongolia, China. *Agr Water Manage* **96**(7), 1114–1119 (2009).
- Liu, C. W. *et al.* Effects of nitrogen fertilizers on the growth and nitrate content of lettuce (*Lactuca sativa* L.). *Int J Env Res Pub He* **11**(11), 4427–40 (2014).
- Ju, X. T. *et al.* Nitrogen balance and groundwater nitrate contamination: comparison among three intensive cropping systems on the north china plain. *Environ Pollut* **143**(1), 117–25 (2006).
- Fan, J., Hao, M. D. & Malhi, S. S. Accumulation of nitrate n in the soil profile and its implications for the environment under dryland agriculture in northern china: a review. *Can J Soi* **90**(3), 423–429 (2010).
- Vanik, V., Ilha, J. & Nimeèek, R. The level of soil nitrate content at different management of organic fertilizers application. *Plant Soil Environ* **49**(5), 197–202 (2003).
- Filipović, V. *et al.* Plastic mulch and nitrogen fertigation in growing vegetables modify soil temperature, water and nitrate dynamics: Experimental results and a modeling study. *Agr Water Manage* **176**, 100–110 (2016).
- Zhang, Y. *et al.* Tracing nitrate pollution sources and transformation in surface- and ground-waters using environmental isotopes. *Sci Total Environ* **490**, 213–222 (2014).
- Moriassi, D. N. *et al.* Modeling the impact of nitrogen fertilizer application and tile drain configuration on nitrate leaching using SWAT. *Agr Water Manage* **130**, 36–43 (2013).
- Evanylo, G. *et al.* Soil and water environmental effects of fertilizer-, manure-, and compost-based fertility practices in an organic vegetable cropping system. *Agr Ecosyst Environ* **127**(1–2), 50–58 (2008).
- Hassink, J., Whitmore, A. P. & Kubát, J. Size and density fractionation of soil organic matter and the physical capacity of soils to protect organic matter. *Eur J Agron* **7**(1), 189–199 (1997).
- Chaney, K. Effect of nitrogen fertilizer rate on soil nitrate nitrogen content after harvesting winter wheat. *J Agr Sci* **114**(2), 171–176 (1990).
- Behnke, G. D., David, M. B. & Voigt, T. B. Greenhouse gas emissions, nitrate leaching, and biomass yields from production of *Miscanthus × giganteus*, in illinois, usa. *Bioenerg. Res.* **5**(4), 801–813 (2012).
- Wang, X., Li, Z. & Xing, Y. Effects of mulching and nitrogen on soil temperature, water content, nitrate-N content and maize yield in the Loess Plateau of China. *Agr Water Manage* **161**, 53–64 (2015).
- Davis, M. P. *et al.* Effect of nitrogen addition on *Miscanthus × giganteus* yield, nitrogen losses, and soil organic matter across five sites. *GCB Bioenergy* **7**(6), 1222–1231 (2015).
- Guo, S. *et al.* Impacts of fertilizer practices on environmental risk of nitrate in semiarid farmlands in the loess plateau of china. *Plant Soil* **330**(1), 1–13 (2010).
- Basso, B. & Ritchie, J. T. Impact of compost, manure and inorganic fertilizer on nitrate leaching and yield for a 6-year maize–alfalfa rotation in Michigan. *Agr Ecosyst Environ* **108**(4), 329–341 (2005).
- Giola, P. *et al.* Impact of manure and slurry applications on soil nitrate in a maize–triticale rotation: Field study and long term simulation analysis. *Eur J Agron* **38**, 43–53 (2012).
- Haynes, R. J. & Naidu, R. Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. *Nutr Cycl Agroecosys* **51**(2), 123–137 (1998).
- Blair, N. *et al.* Long-term management impacts on soil c, n and physical fertility: part i: broadbalk experiment. *Soil Till Res* **91**(1–2), 30–38 (2006).
- Ding, X. *et al.* Impact of long-term application of manure, crop residue, and mineral fertilizer on organic carbon pools and crop yields in a mollisol. *J Soil Sediment* **14**(5), 854–859 (2014).
- Santamaria, P. Nitrate in vegetables: toxicity, content, intake and EC regulation. *J Sci Food Agr* **86**(1), 10–17 (2006).

29. Santamaria, P. *et al.* Comparison between nitrate and ammonium nutrition in fennel, celery, and Swiss chard. *J Plant Nutr* **22**(7), 1091–1106 (1999).
30. Mohammad, M. *et al.* Tomato root and shoot responses to salt stress under different levels of phosphorus nutrition. *J Plant Nutr* **21**(8), 1667–1680 (1998).
31. Zhou, Y., Selvam, A. & Wong, J. W. Evaluation of humic substances during co-composting of food waste, sawdust and Chinese medicinal herbal residues. *Bioresour Technol* **168**, 229–234 (2014).
32. Ko, H. J. *et al.* Evaluation of maturity parameters and heavy metal contents in composts made from animal manure. *Waste Manag* **28**(5), 813–820 (2008).
33. Bremner, J. M. Nitrogen-total. In: *Methods of soil analysis part 3. Chemical methods*. SSSA Book Ser. No. 5. (eds Sparks, D. L. *et al.*) 1058–1121 (SSSA, Madison, 1996).
34. Bao S.D. Soil agro-chemistry analysis method. 25–38 (China agriculture pres, Beijing, 2000).
35. Wang, H. *et al.* Impact of tillage practices on nitrogen accumulation and translocation in wheat and soil nitrate-nitrogen leaching in drylands. *Soil Till Res* **153**, 20–27 (2015).
36. Shi, W. M., Yao, J. & Yan, F. Vegetable cultivation under greenhouse conditions leads to rapid accumulation of nutrients, acidification and salinity of soils and groundwater contamination in South-Eastern China. *Nutr Cycl Agroecosys* **83**(1), 73–84 (2009).
37. Fang, Q. X. *et al.* Soil nitrate accumulation, leaching and crop nitrogen use as influenced by fertilization and irrigation in an intensive wheat–maize double cropping system in the north china plain. *Plant Soil* **284**(1), 335–350 (2006).

Acknowledgements

This research was financially supported by the High Effective Intelligent Composting Process and Key Technology Research (2016YFD0800600-01) and the Research Project for Technical System Construction in Cashmere Goat Industry of Ministry of Agriculture of the People's Republic of China (CARS-39-19).

Author Contributions

J.L., G.L. and S.L. designed the experiment, S.L., B.Z., D.L., and Y.L. performed the experiment and did the statistical analysis, S.L. wrote the first draft of the manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017