

## Identification of a 4-fluorobenzyl L-valinate amide benzoxaborole (AN11736) as a potential development candidate for the treatment of Animal African Trypanosomiasis (AAT)

Tsutomu Akama<sup>a</sup>, Yong-Kang Zhang<sup>a</sup>, Yvonne R. Freund<sup>a</sup>, Pamela Berry<sup>a</sup>, Joanne Lee<sup>a</sup>, Eric E. Easom<sup>a</sup>, Robert T. Jacobs<sup>a</sup>, Jacob J. Plattner<sup>a</sup>, Michael J. Witty<sup>b,\*</sup>, Rosemary Peter<sup>b</sup>, Tim G. Rowan<sup>b</sup>, Kirsten Gillingwater<sup>c,d</sup>, Reto Brun<sup>c,d</sup>, Bakela Nare<sup>e</sup>, Luke Mercer<sup>e</sup>, Musheng Xu<sup>f</sup>, Jiangong Wang<sup>f</sup>, Hao Liang<sup>f</sup>

<sup>a</sup>Anacor Pharmaceuticals, Inc., 1020 E. Meadow Circle, Palo Alto, CA 94303, USA

<sup>b</sup>Global Alliance for Livestock and Veterinary Medicine, Doherty Building, Pentlands Science Park, Penicuik, Edinburgh EH26 0PZ, UK

<sup>c</sup>Swiss Tropical and Public Health Institute, Socinstrasse 57, 4051 Basel, Switzerland

<sup>d</sup>University of Basel, Petersplatz 1, 4003 Basel, Switzerland

<sup>e</sup>Avista Pharma Solutions, 350 Tricenter Boulevard, Suite C, Durham, NC 27713, USA

<sup>f</sup>Wuxi AppTec (Tianjin) Co. Ltd., No. 168 NanHai Road, 10th Avenue, TEDA, Tianjin 300457, PR China

### ARTICLE INFO

#### Article history:

Received 4 September 2017

Accepted 14 November 2017

Available online 20 November 2017

#### Keywords:

Benzoxaborole  
Trypanosomiasis  
Cattle  
SAR  
Lead optimisation  
Protozoan

### ABSTRACT

Novel L-valinate amide benzoxaboroles and analogues were designed and synthesized for a structure-activity-relationship (SAR) investigation to optimize the growth inhibitory activity against *Trypanosoma congolense* (*T. congolense*) and *Trypanosoma vivax* (*T. vivax*) parasites. The study identified 4-fluorobenzyl (1-hydroxy-7-methyl-1,3-dihydrobenzo[c][1,2]oxaborole-6-carbonyl)-L-valinate (**5**, AN11736), which showed IC<sub>50</sub> values of 0.15 nM against *T. congolense* and 1.3 nM against *T. vivax*, and demonstrated 100% efficacy with a single dose of 10 mg/kg against both *T. congolense* and *T. vivax* in mouse models of infection (IP dosing) and in the target animal, cattle, dosed intramuscularly. AN11736 has been advanced to early development studies.

© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Animal African Trypanosomiasis (AAT) is a fatal, parasitic wasting disease of livestock and wild animals in Sub-Saharan Africa.<sup>1</sup> It is caused primarily by the two protozoan parasites *Trypanosoma congolense* (*T. congolense*) and *Trypanosoma vivax* (*T. vivax*), which are transmitted by tsetse flies.<sup>1</sup> AAT is responsible for 3 million cattle deaths annually and costs African livestock farmers approximately US\$ 1–5 billion per year.<sup>2</sup> The current standard-of-care drugs, such as diminazene aceturate, isometamidium and homidium chloride, have been in use for several decades and are often ineffective with drug resistance becoming an increasing concern.<sup>1</sup> No new trypanocides have been approved for use in cattle for many years. Initial screening of the Anacor Pharmaceuticals library of novel boron-containing compounds identified an active compound (**1**, Fig. 1), which had an IC<sub>50</sub> = 5 nM against *T. congolense* and 69 nM against *T. vivax* while its enantiomer was much less

\* Corresponding author.

E-mail address: [michael.witty@galvmed.org](mailto:michael.witty@galvmed.org) (M.J. Witty).

<https://doi.org/10.1016/j.bmcl.2017.11.028>

0960-894X/© 2017 The Authors. Published by Elsevier Ltd.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

active. A quick and simple modification on the amino acid side chain with an isopropyl group generated **2** (Fig. 1) with improved *in vitro* potency (IC<sub>50</sub> = 2 nM against both *T. congolense* and *T. vivax*). This encouraging result prompted us to investigate this chemical series further. We designed and synthesized a series of novel benzoxaboroles (**3–71**, Figs. 2–8) to optimize anti-parasitic activity, physicochemical and *in vitro* ADME properties, and the pharmacokinetic profile. Specifically, these molecules were designed to examine the effects of oxaborole 3-substituent variation (**3** vs **2**, Figs. 1 and 2), oxaborole 7-substituent variation (**4** vs **2**, Figs. 1 and 2; **5** vs **20–27**, Fig. 4), substituent changes on the benzyl group (**5–19**, Fig. 3), modification of the amino acid (**28–32**, Fig. 5), heteroaromatic methyl esters (**33–48**, Fig. 6), introduction of water-solubilizing scaffolds to the benzyl group (**49–54**, Fig. 7) and aliphatic esters (**55–71**, Fig. 8). Herein, we report the synthesis and antiparasitic activity against *T. congolense* and *T. vivax* of these novel compounds.

Compounds **1–71** were convergently synthesized from three building blocks: the left side alcohols (**72**), amino acid linkers




Fig. 1. Chemical structures of early hits (1 and 2).



Fig. 2. Structures of benzoxaboroles with additional 3,3-Me<sub>2</sub> (3) or 7-Me (4) modification as compared to analog 2.

Fig. 3. Structures of benzoxaboroles with variation of substituents on the benzyl ring (5–19) as compared to analog 4.




Fig. 4. Structures of benzoxaboroles with variation of 7-substituents on the benzene ring (20–27) as compared to analog 5.






Fig. 5. Structures of benzoxaboroles with variation on the amino acid side chain (28–32) as compared to analog 5.



	R	R
33		41
34		42
35		43
36		44
37		45
38		46
39		47
40		48

Fig. 6. Structures of benzoxaboroles with variation of the left side arylmethyl groups (33–48) as compared to analog 5.



	49	50	51	52	53	54
R						


Fig. 7. Structures of benzoxaboroles with water-solubilizing scaffolds on the benzyl ring (49–54) as compared to analog 5.

(73) and benzoxaborole 6-carboxylic acids (76).<sup>3,4</sup> The general synthetic route is shown in Scheme 1. Reaction of alcohols 72 with *N*-Boc protected amino acids 73 gave ester intermediates 74, which were treated with dry hydrogen chloride to generate ester amine salts 75. Condensation of these amine salts 75 with benzoxaborole 6-carboxylic acids 76 provided the final compounds 1–71.


Scheme 2 illustrates the synthesis of 1-hydroxy-7-methyl-1,3-dihydrobenzo[*c*][1,2]oxaborole-6-carboxylic acid (83) as an example of key boron intermediates. Esterification of the acid 77 produced the ester 78, which was formylated to yield 79. Treatment of 79 with trifluoromethyl sulfonyl anhydride afforded the triflate compound 80, which was converted to the pinacol boron intermediate 81. Reduction of 81 and subsequent cyclization under aqueous acidic conditions generated the benzoxaborole ester 82. Hydrolysis of the ester group in 82 afforded the acid 83. The experimental procedures for the synthesis of 5 are described in the reference and note section.<sup>5</sup>

Activity of compounds 1–71 against *T. congolense* and *T. vivax* was determined using the whole cell assays as described<sup>6</sup> and their IC<sub>50</sub> values are summarized in Table 1.

Lead compound 2 exhibited an IC<sub>50</sub> of 2 nM against both *T. congolense* and *T. vivax*. The 3,3-dimethyl analog 3 was essentially inactive (IC<sub>50</sub> = 2580 nM against *T. c.* and 9190 nM against *T. v.*) but better activity was observed for the 7-methyl analog 4 (IC<sub>50</sub>




**Fig. 8.** Structures of benzoxaboroles with variation of the left side aliphatic ester groups (55–71) as compared to analog 5.



**Scheme 1.** General route for syntheses of 1–71. Reagents and conditions: (a) DCC, DMAP, DCM, 0–20 °C, 16 h; (b) HCl/EtOAc, 20 °C, 5 h; (c) HOBt, EDCI, TEA, DCM, 0–20 °C, 15 h.

= 0.46 nM against *T. c.* and 0.79 nM against *T. v.*). We focused future SAR development on compounds incorporating the 7-methyl group, as *in vivo* activity of **4** was superior to that observed for **2** (*vide infra*). A wide range of substituents, such as halogens, trifluoromethyl, trifluoromethoxy, cyano and methylsulfonyl (**5–19** in Fig. 3) on the benzyl ring were introduced to examine their effects on the antiparasitic activity. The majority of these fifteen compounds, with exception of **11** and **12**, were very potent showing  $IC_{50}$  values around 1 nM (see Table 1). We next explored variation of the substituent at the 7-position of benzoxaborole (**20–27** in Fig. 4). The difluoromethyl (**22**) and ethyl (**24**) analogs had similar activity to that of **5**, but the electron-withdrawing fluoro (**21**) and trifluoromethyl (**23**) analogs were of significantly reduced potency. The amino acid linker was also modified (Fig. 5), with the cyclopropyl (**28**), 2-hydroxyisopropyl (**29**), 2-fluoroisopropyl (**30**) and spirocyclobutyl (**32**) analogs exhibiting potency similar to **5**, but the spirocyclopropyl analog (**31**) exhibited decreased activity against *T. v.* parasite. Replacement of the 4-fluorophenyl in **5** with various heteroaryl groups (**33–48**, Fig. 6) resulted in the excellent activity in all cases except the NH-imidazole analog **44**. Introduction of basic nitrogen-containing groups on the benzyl ester (**49–54**, Fig. 7) provided compounds **50–54** that were generally similar to **5**. Lastly, aliphatic and heterocyclic esters (**55–71**, Fig. 8) were



**Scheme 2.** Synthetic route for preparation of **83**. Reagents and conditions: (a)  $H_2SO_4$ , EtOH, reflux, 24 h; (b)  $MgCl_2$ ,  $(CH_2O)_n$ , TEA, THF, reflux, 14 h; (c)  $Tf_2O$ , pyridine, DMAP, DCM, 0–15 °C, 1 h; (d)  $Pin_2B_2$ , KOAc,  $Pd(dppf)Cl_2$ , 1,4-dioxane,  $N_2$ , 85 °C, 15 h; (e)  $NaBH_4$ , MeOH, THF, 0–15 °C, 1 h, then HCl,  $H_2O$ ; (f) NaOH,  $H_2O$ , 40 °C, 3 h, then HCl for acidification.

**Table 1**

Activity of compounds 1–71 against *T. congolense* (*T. c.*) and *T. vivax* (*T. v.*).<sup>a</sup>

Compound	$IC_{50}$ (nM)		Compound	$IC_{50}$ (nM)	
	<i>T. c.</i>	<i>T. v.</i>		<i>T. c.</i>	<i>T. v.</i>
<b>1</b>	4.9	69	<b>37</b>	0.78	0.50
<b>2</b>	2.0	2.0	<b>38</b>	0.68	0.31
<b>3</b>	2580	9190	<b>39</b>	0.57	0.24
<b>4</b>	0.46	0.79	<b>40</b>	0.27	0.50
<b>5</b>	0.14	1.3	<b>41</b>	0.78	0.11
<b>6</b>	0.47	2.9	<b>42</b>	0.20	0.19
<b>7</b>	0.59	0.10	<b>43</b>	0.062	NT <sup>b</sup>
<b>8</b>	0.28	0.07	<b>44</b>	5000	980
<b>9</b>	0.18	0.10	<b>45</b>	0.37	0.33
<b>10</b>	0.22	2.7	<b>46</b>	0.42	0.24
<b>11</b>	0.16	24	<b>47</b>	0.20	0.081
<b>12</b>	0.31	19	<b>48</b>	<0.005	0.71
<b>13</b>	0.23	0.04	<b>49</b>	0.39	68
<b>14</b>	0.10	0.06	<b>50</b>	0.32	0.48
<b>15</b>	0.15	0.05	<b>51</b>	0.25	0.21
<b>16</b>	0.08	0.07	<b>52</b>	1.3	0.29
<b>17</b>	0.21	0.04	<b>53</b>	0.51	0.35
<b>18</b>	0.20	0.06	<b>54</b>	1.0	0.37
<b>19</b>	0.61	0.44	<b>55</b>	4.2	3.3
<b>20</b>	3.0	1.0	<b>56</b>	5.2	14
<b>21</b>	28	NT <sup>b</sup>	<b>57</b>	0.46	0.26
<b>22</b>	0.67	0.92	<b>58</b>	0.66	0.69
<b>23</b>	37	51	<b>59</b>	0.70	0.52
<b>24</b>	0.11	0.05	<b>60</b>	<0.005	0.78
<b>25</b>	3.0	4.7	<b>61</b>	0.36	0.21
<b>26</b>	2.3	0.05	<b>62</b>	0.38	0.16
<b>27</b>	3.9	0.71	<b>63</b>	0.39	0.09
<b>28</b>	0.12	0.38	<b>64</b>	2.3	0.78
<b>29</b>	0.26	0.10	<b>65</b>	9.4	1.2
<b>30</b>	0.09	0.25	<b>66</b>	5.9	4.1
<b>31</b>	0.13	18	<b>67</b>	0.43	0.79
<b>32</b>	0.45	2.5	<b>68</b>	1.8	37
<b>33</b>	0.28	26	<b>69</b>	0.47	1.6
<b>34</b>	0.14	0.07	<b>70</b>	0.34	2.28
<b>35</b>	0.15	0.09	<b>71</b>	0.2	0.14
<b>36</b>	0.26	0.14			

<sup>a</sup> Experimental procedures are described in the reference and note section.<sup>6</sup>

<sup>b</sup> NT = Not tested.

synthesized and many of these had  $IC_{50}$  values less than 1 nM as shown in Table 1.

Selected compounds were screened in both mouse and bovine *in vitro* metabolic stability assays (mouse S9 and bovine S9), as summarized in Table 2. These two species were chosen because the primary *in vivo* assays were conducted in mice, and the target animal of this research program is cattle. As shown in Table 2, out of 36 compounds tested, 27 compounds had  $Cl_{int} < 10 \mu\text{L}/\text{min}/\text{mg}$  protein in both mouse and bovine S9 assays suggesting moderate to excellent *in vitro* metabolic stability. We evaluated the efficacy of selected compounds in two *in vivo* mouse models of infection, against *T. congolense* and *T. vivax*, respectively. Mice were infected with either  $1 \times 10^5$  *T. c.* parasites or  $1 \times 10^4$  *T. v.* parasites, and then treated with a test compound via intraperitoneal administration for 1, 2 or 4 consecutive days. The mice were then monitored for the presence of parasitemia for up to 60 days post treatment.<sup>7</sup> We tested in *T. c.* model first, then followed up with *T. v.* for interesting compounds. As shown in Table 2, the 7-methyl analog **4** was superior to the 7-unsubstituted analog **2** in both *T. c.* and *T. v.* mouse models of infection. Of the 38 compounds tested with the

*in vivo* mouse models, seven compounds (**5**, **8**, **33**, **34**, **49**, **62** and **71**) demonstrated  $\geq 50\%$  curative efficacy in the *T. c.*-infected mouse model and 100% curative efficacy in the *T. v.*-infected mouse model, when tested as a single dose of 10 mg/kg. To select further from these seven compounds, four (**5**, **8**, **33** and **71**) had  $\geq 75\%$  curative efficacy in the *T. c.*-infected mouse model at a single dose of 10 mg/kg, and two (**5** and **8**) showed 100% curative efficacy. These two compounds were further tested at a single 5 mg/kg dose, but were unable to cure the *T. c.*-infected mice. We selected compound **5** (**AN11736**) to progress to exploratory studies to determine the efficacy and safety in a preliminary formulation against induced infections of *T. vivax* and *T. congolense* in cattle.<sup>8</sup> **AN11736** demonstrated 100% curative efficacy with a single intramuscular injection of 10 mg/kg against both *T. congolense* and *T. vivax* in cattle.

In summary, a novel series of *L*-valinate amide benzoxaboroles was discovered to be active against *T. congolense* and *T. vivax*, which are the main causative agents of Animal African Trypanosomiasis (AAT) in cattle. Two compounds (**5** and **8**) showed 100% curative efficacy in both *T. c.*- and *T. v.*-infected mice with a single dose of 10 mg/kg. Compound **5** (**AN11736**) demonstrated 100% curative efficacy with a single IM dose of 10 mg/kg against both *T. congolense* and *T. vivax* in cattle for a duration of 100 days. **AN11736**, as a novel chemical entity, was selected as a potential developmental candidate for the treatment of AAT.

## Acknowledgement

The research was conducted with a grant from the Global Alliance for Livestock Veterinary Medicines (GALVmed) and funded by the Bill & Melinda Gates Foundation [OPP1093639] and the UK Government. The findings and conclusions contained are those of the authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation or the UK Government.

## References

- Giordani F, Morrison LJ, Rowan TG, de Koning HP, Barrett MP. *Parasitology*. 2016;143:1862.
- Shaw AP, Cecchi G, Wint GR, Mattioli RC, Robinson TP. *Prev Vet Med*. 2014;113:197.
- Li X, Zhang S, Zhang Y-K, et al. *Bioorg Med Chem Lett*. 2011;21:2048.
- Ding CZ, Zhang Y-K, Li X, et al. *Bioorg Med Chem Lett*. 2010;20:7317.
- Experimental procedures for the synthesis of 4-fluorobenzyl (1-hydroxy-7-methyl-1,3-dihydrobenzo[c][1,2]oxaborole-6-carbonyl)-L-valinate (5, AN11736)*: To a solution of **77** (1.65 kg, 10.8 mol) in EtOH (6.50 L) was added conc.  $\text{H}_2\text{SO}_4$  (326 g, 3.25 mol). The reaction mixture was stirred at 105 °C for 24 h. TLC (thin layer chromatography) showed the starting material was consumed completely. The mixture was cooled to 15 °C and concentrated. The residue was poured into aqueous 2 M  $\text{NaHCO}_3$  (3 L) and the solid was filtered off. The filtrate was concentrated to give **78** (1.75 kg, yield 90%) as a brown solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.41 (d,  $J = 7.9$  Hz, 1H), 7.11 (t,  $J = 7.9$  Hz, 1H), 6.94 (d,  $J = 7.9$  Hz, 1H), 4.58 (br s, 1H), 4.37 (q,  $J = 7.4$  Hz, 2H), 2.46 (s, 3H), 1.40 (t,  $J = 7.1$  Hz, 3H). To a solution of **78** (800 g, 4.44 mol) in THF (6.50 L) were added  $\text{MgCl}_2$  (634 g, 6.66 mol), TEA (1.80 kg, 17.8 mol) and  $(\text{HCHO})_n$  (600 g, 6.66 mol). The mixture was immediately heated to 90 °C and was stirred for 14 h. The reaction mixture was cooled to 15 °C. To it were added ice  $\text{H}_2\text{O}$  (3 L) and then 12 N HCl (1.5 L) slowly. The mixture was stirred for 0.5 h and then extracted with EtOAc (2 L). The combined organic layer was washed with sat.  $\text{NaHCO}_3$  to neutral, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under reduced pressure to give crude **79** (880 g) as a brown oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.40 (s, 1H), 9.93 (s, 1H), 7.46 (d,  $J = 7.6$  Hz, 1H), 7.37 (d,  $J = 8.0$  Hz, 1H), 4.40 (q,  $J = 7.4$  Hz, 2H), 2.44 (s, 3H), 1.41 (t,  $J = 7.1$  Hz, 3H). To a solution of **79** (900 g, 4.32 mol) in DCM (7.56 L) were added pyridine (1.02 kg, 12.9 mol) and DMAP (27 g, 221 mmol). The mixture was cooled to 0 °C and  $\text{TiF}_2\text{O}$  (1.60 kg, 5.66 mol) was added dropwise. The reaction mixture was warmed to 15 °C and stirred for 1 h. The mixture was quenched with water (7.65 L) and then extracted with DCM (2  $\times$  7.65 L). The combined organic layer was washed with water (2 L), dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under reduced pressure to give **80** (685 g, 47%) as a light yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.27 (s, 1H), 7.99 (d,  $J = 8.0$  Hz, 1H), 7.91–7.87 (m, 1H), 4.43 (q,  $J = 7.0$  Hz, 2H), 2.64 (s, 3H), 1.43 (t,  $J = 7.3$  Hz, 3H). To a solution of **80** (1.00 kg, 2.94 mol), bis(pinacolato)diboron (1.12 kg, 4.41 mol) and KOAc (573 g, 5.84 mol) in 1,4-dioxane (6.50 L) was added  $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$  (150 g, 184 mmol). The mixture was stirred at 85 °C for 15 h under  $\text{N}_2$ . The mixture was

**Table 2**  
*In vitro* metabolic stability and *in vivo* mouse efficacy of selected compounds.<sup>a</sup>

Compound	$Cl_{int}$ ( $\mu\text{L}/\text{min}/\text{mg}$ protein)		Efficacy in mouse model <sup>b</sup>	
	Mouse S9	Bovine S9	<i>T. c.</i>	<i>T. v.</i>
<b>2</b>	NT <sup>c</sup>	NT <sup>c</sup>	0/5 (2 $\times$ 10)	5/5 (4 $\times$ 10) 1/5 (1 $\times$ 10)
<b>4</b>	NT <sup>c</sup>	NT <sup>c</sup>	4/5 (2 $\times$ 10)	5/5 (2 $\times$ 10) 4/4 (1 $\times$ 10)
<b>5</b>	5.4	9.3	5/5 (2 $\times$ 10) 4/4 (1 $\times$ 10) 0/4 (1 $\times$ 5)	5/5 (2 $\times$ 10) 5/5 (1 $\times$ 10) 4/4 (1 $\times$ 10)
<b>6</b>	17	9.1	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>8</b>	7.9	5.6	4/4 (1 $\times$ 10) 0/4 (1 $\times$ 5)	4/4 (1 $\times$ 10)
<b>14</b>	9.1	2.7	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>16</b>	15	8.9	4/4 (2 $\times$ 10) 1/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>17</b>	14	10	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>18</b>	9.7	3.1	1/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>19</b>	2.6	1.3	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>20</b>	1.5	1.3	0/5 (4 $\times$ 10)	NT <sup>c</sup>
<b>21</b>	<1	<1	1/4 (2 $\times$ 10)	4/4 (2 $\times$ 10)
<b>22</b>	1.4	0.9	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>24</b>	21	23	1/4 (2 $\times$ 10)	NT <sup>c</sup>
<b>26</b>	32	32	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>29</b>	<1	<1	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>30</b>	3.7	3.1	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>31</b>	<1	<1	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>33</b>	6.3	2.0	3/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>34</b>	11.8	4.8	2/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>35</b>	5.2	7.8	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>39</b>	1.6	3.9	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>41</b>	1.3	1.8	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>46</b>	7.4	7.0	1/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>47</b>	4.8	10	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>48</b>	4.0	23	1/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>49</b>	1.5	4.0	2/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>50</b>	8.6	12.7	1/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>51</b>	1.8	5.0	0/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>52</b>	1.8	6.0	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>53</b>	5.4	6.8	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>54</b>	5.5	3.8	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>55</b>	<1	3.1	0/4 (1 $\times$ 10)	NT <sup>c</sup>
<b>62</b>	10	20	2/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>63</b>	0.9	1.5	1/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>69</b>	0.7	0.6	1/4 (1 $\times$ 10)	1/4 (1 $\times$ 10)
<b>70</b>	1.5	0.3	1/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)
<b>71</b>	6.4	8.1	3/4 (1 $\times$ 10)	4/4 (1 $\times$ 10)

<sup>a</sup> Methods for testing compound efficacy in mouse models are described in the reference and note section.<sup>7</sup>

<sup>b</sup> The efficacy data in the table is presented as n/m (q  $\times$  10), where n = number of mice survived, m = total number of mice in the study group, q = how many time dosed, and 10 = 10 mg/kg.

<sup>c</sup> NT = Not tested.

cooled to 15 °C, filtered and concentrated to give the crude product. The residue was purified by column chromatography (SiO<sub>2</sub>, petroleum ether/ethyl acetate = 40/1 to 4:1) to give crude **81** (942 g) as a yellow oil. To a solution of **81** (1.20 kg, 3.77 mol) in MeOH (300 mL) and THF (6 L) was added NaBH<sub>4</sub> (80 g, 2.11 mol) in portions at 0 °C. Then the reaction mixture was stirred at 15 °C for 1 h. HPLC showed **81** was consumed completely. The reaction solution was adjusted to pH = 4 with 2 M HCl. The organic layer was removed in vacuum and the mixture was filtered. The cake was washed with petroleum ether (5 L) and dried in vacuum to give **82** (665 g, 80%) as a white solid. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 9.18 (s, 1H), 7.89 (d, *J* = 8.0 Hz, 1H), 7.32 (d, *J* = 8.0 Hz, 1H), 5.00 (s, 2H), 4.30 (q, *J* = 7.0 Hz, 2H), 2.68 (s, 3H), 1.33 (t, *J* = 7.0 Hz, 3H). To a mixture of **82** (867 g, 3.94 mol) in H<sub>2</sub>O (5 L) was added NaOH (394 g, 9.85 mol) in one portion. The solution was heated at 40 °C for 3 h. HPLC showed **82** was consumed completely. This batch was worked-up together with the other batches and acidified with 2 N HCl to pH = 2. The solid was filtered and washed with H<sub>2</sub>O (10 L). The cake was dried to give **83** (2.00 kg, yield 87%) as a white solid. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 12.75 (s, 1H), 9.13 (s, 1H), 7.89 (d, *J* = 8.0 Hz, 1H), 7.28 (d, *J* = 8.0 Hz, 1H), 4.98 (s, 2H), 2.68 (s, 3H); HPLC purity: 100% at both 220 nm and 254 nm; MS (ESI+): *m/z* = 193 (M+1). To a solution of 4-fluorobenzylalcohol (**72**, R<sup>1</sup> = 4-fluorobenzyl, 290 g, 2.30 mol, 248.10 mL) and *N*-Boc-(*S*)-valine (**73**, R<sup>2</sup> = isopropyl, R<sup>3</sup> = H, 500 g, 2.30 mol) in dry DCM (6.0 L) were added DCC (854 g, 4.14 mol, 838 mL) and DMAP (39.36 g, 322.19 mmol). The reaction mixture was stirred at 25 °C for 15 h. The mixture was filtered and washed with DCM (2 L) and concentrated to give the crude product. The residue was purified by column chromatography (SiO<sub>2</sub>, petroleum ether/ethyl acetate = 50:1 to 10:1) to give 4-fluorobenzyl (*tert*-butoxycarbonyl)-*L*-valinate (**74**, R<sup>1</sup> = 4-fluorobenzyl, R<sup>2</sup> = isopropyl, R<sup>3</sup> = H, 708 g, 95% yield) as a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.35 (dd, *J* = 8.2 & 5.5 Hz, 2H), 7.05 (t, *J* = 8.6 Hz, 2H), 5.19–5.08 (m, 2H), 5.01 (d, *J* = 8.4 Hz, 1H), 4.25 (dd, *J* = 8.4 & 4.4 Hz, 1H), 2.13 (dd, *J* = 11.9 & 6.2 Hz, 1H), 1.44 (s, 9H), 0.93 (d, *J* = 7.1 Hz, 3H), 0.84 (d, *J* = 7.1 Hz, 3H). The mixture of 4-fluorobenzyl (*tert*-butoxycarbonyl)-*L*-valinate (**74**, R<sup>1</sup> = 4-fluorobenzyl, R<sup>2</sup> = isopropyl, R<sup>3</sup> = H, 1.06 kg, 3.26 mol) in HCl/EtOAc (6.0 L) was stirred at 25 °C for 14 h. The solvent was removed under reduced pressure to give 4-fluorobenzyl *L*-valinate hydrochloride (**75**, R<sup>1</sup> = 4-fluorobenzyl, R<sup>2</sup> = isopropyl, R<sup>3</sup> = H, 780 g, 91% yield) was obtained as a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.90 (br, s, 3H), 7.37 (dd, *J* = 8.2 & 5.5 Hz, 2H), 7.03 (t, *J* = 8.4 Hz, 2H), 5.29–5.10 (m, 2H), 3.95 (br, s, 1H), 2.44 (dd, *J* = 11.0 & 6.6 Hz, 1H), 1.08 (dd, *J* = 10.1 & 7.1 Hz, 6H). To the mixture of **83** (150 mg, 0.77 mmol), **75** (203 mg, 0.77 mmol) and DIEA (0.4 mL, 2.33 mmol) in DMF was added HATU (325 mg, 0.86 mmol). The mixture was stirred at rt for 3 h. The crude product was purified by preparative TLC and preparative HPLC to get the final product **5** (**AN11736**, 125 mg, 40% yield). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 9.03 (s, 1H), 8.57 (d, *J* = 7.2 Hz, 1H), 7.47–7.43 (m, 2H), 7.33 (d, *J* = 7.6 Hz, 1H), 7.23–7.18 (m, 3H), 5.21 (d, *J* = 12.4 Hz, 1H), 5.11 (d, *J* = 12.4 Hz, 1H), 4.96 (s, 2H), 4.33 (t, *J* = 7.2 Hz, 1H), 2.42 (s, 3H), 2.19–2.10 (m, 1H), 0.94 (d, *J* = 6.8 Hz, 3H), 0.92 (d, *J* = 6.8 Hz, 3H); HPLC purity: 100% at both 220 nm and 254 nm; MS (ESI+): *m/z* = 400 (M+1).

6. *Assay methods for determination of T. c. and T. v. IC<sub>50</sub> values:* Bloodstream-form *Trypanosoma congolense* (*T. congolense* IL-3000 strain) were cultured in 24-well

plates at 34 °C and 5% CO<sub>2</sub>. Culture media consisted of complete HMI-9 (IMDM) with 20% bovine serum. To ensure log growth phase, trypanosomes were sub-cultured at appropriate dilutions every 2–3 days. Bloodstream form *Trypanosoma vivax* (*T. vivax*) cannot be cultured axenically, therefore trypanosomes (*T. vivax* STIB719/ILRAD 560 strain) were harvested from a highly parasitemia mouse via cardiac puncture and used directly in the *ex vivo* drug sensitivity assay on the same day. IC<sub>50</sub> determination was carried out using the Alamar Blue assay for *T. congolense* (72 h *in vitro* assay) and modified slightly for *T. vivax* (48 h *ex vivo* assay). Parasite starting concentrations of 2 × 10<sup>5</sup> (*T. c.*) and 4 × 10<sup>5</sup> (*T. v.*) were calculated respectively, using a cell analyzer system or haemo-cytometer, followed by quantification at 536 nm excitation and 588 nm emission wavelengths using a flow cytometer reader. Test compounds were prepared as 10 mg/mL DMSO stocks for each assay run. Compounds were assayed in at least three separate, independent test runs and an 11-point dilution curve was used to determine the IC<sub>50</sub> values. Data points were averaged to generate sigmoidal dose-response curves and IC<sub>50</sub> values were determined using Softmax Pro 5.2 software.

7. *Methods for testing compound efficacy in mouse models:* *In vivo* mouse efficacy studies were performed at the Swiss TPH, using established mouse models of infection for *T. c.* and *T. v.* NMRI female mice were independently infected either with 1 × 10<sup>5</sup> *T. c.* parasites/mouse (STIB736/IL1180 strain) or 1 × 10<sup>4</sup> *T. v.* parasites/mouse (STIB719/ILRAD560 strain) using an intraperitoneal route. Parasitemia was allowed to develop over 7 days (*T. c.*) or 3 days (*T. v.*), respectively, before treatment was administered. Compounds were administered via intraperitoneal injection in 10% DMSO/water. Four mice were used per treatment group. Mice were monitored for the presence of trypanosomes, via tail blood examination microscopically with twice-per-week parasitemia checks for 60 days post treatment. Parasitemia was graded on a scale of 0–3, with 0 indicating no trypanosomes seen in 20 fields of view, 1 indicating the presence of 1–5 trypanosomes per field, 2 indicating the presence of 6–20 trypanosomes per field and 3 indicating greater than 20 trypanosomes per field. Mice scoring 2 or 3 were immediately euthanized. After 60 days, aparasitemic mice were considered cured. Untreated control mice survive on average for 10 and 6 days post infection for *T. c.* and *T. v.*, respectively. All *in vivo* mouse experiments were conducted in accordance with the strict guidelines set out by the Swiss Federal Veterinary Office, under the ethical approval of license number #2813.

8. *Method for testing compound efficacy in cattle:* The cattle studies were conducted in accordance with the method described for cattle by Eisler et al. (2001) in studies conducted in fly-proof facilities and using *T. congolense* and *T. vivax* isolates that had previously been confirmed resistant in cattle to diminazine (7 mg/kg live weight) and/or isometamidium (1 mg/kg live weight). Cattle studies included negative (saline) controls, all assessments were made for 100 days post treatment (unless animals relapsed sooner) and were conducted by staff blinded (masked) to allocation of animals to treatment groups and in accordance with the principles of veterinary good clinical practice (VICH, 2000).