

Topical administration of Esculetin as a potential therapy for experimental dry eye syndrome

D Jiang¹, X Liu¹ and J Hu

Abstract

Purpose In this study, we investigated the therapeutic effects of topical Esculetin for dry eye rabbits through the ocular tests, inflammatory factor levels and specific phosphorylated protein expressions of ERK1/2 signal pathway.

Patients and methods Thirty-two healthy adult male New Zealand white rabbits were chosen for the study. DES models were established after removing of the main lacrimal gland, Harderian gland and nictitating membrane in the left eyes and randomly divided into group DES control, group CsA, group Esculetin and group Esculetin combined with CsA (C&E), meanwhile the right eyes served as group Normal control. Schirmer's I tests, fluorescein scores, goblet cell densities, inflammatory cytokines IL-1 α , IL-1 β , TNF- α levels were observed by slit-lamp microscope, conjunctival impression cytology and ELISA essay at week 0, 1, 2, 4, 8. Phosphorylated-ERK1/2 expressions were detected in Western blot analysis at week 8.

Results After induction of DES, aqueous tear production and goblet cell density were decreased, FL score was much higher in group DES control throughout the study ($P < 0.05$). Both topical Esculetin and Esculetin combined with CsA increased the SIT values (10 ± 1 mm, 14 ± 3 mm, $P < 0.05$) and goblet cell densities (77 ± 12 /HP, 92 ± 12 /HP, $P < 0.05$), decreased FL scores (7.48 ± 0.33 , 5.09 ± 0.24 , $P < 0.05$) at week 8. Alterations of IL-1 α , IL-1 β , TNF- α levels had similar trend. In Western blot analysis, downregulations of p-ERK1/2 were observed in therapy groups when compared with group DES control and the most decreasing was found in group C&E ($P < 0.05$).

Conclusion Topical Esculetin improved DES symptoms, downregulated the inflammatory cytokine expressions, suppressed the ERK1/2

pathway and enhanced the therapeutic effect of CsA.

Eye (2017) 31, 1724–1732; doi:10.1038/eye.2017.117; published online 23 June 2017

Introduction

Dry eye syndrome (DES) is a multifactorial disease of tears and ocular surface that results in symptoms of discomfort, visual disturbance and tear film instability. It is accompanied by increased osmolarity of the tear film and inflammation of the ocular surface.¹ There are many etiological factors of DES including chronic diseases, infectious, aging, radiation and environmental factors like smoking or video display.² According to the Dry Eye Workshop (DEWS) report in 2007, prevalence of dry eye ranges from 5 to 30% in people aged 50 years and older.¹ Cyclosporine A (CsA) is one kind of drug proved therapeutic by FDA in curing DES through inhibiting the immune reaction and inflammation of the ocular surface. However, topical CsA has potential side effects, for example, ocular burning, conjunctival hyperemia, epiphora, eye pain, foreign body sensation,³ especially when used chronically. Appropriately administration of this agent and monitoring for side effects mentioned above remained tough problems. In addition, topical CsA eyedrops were not easily available in our country and many patients cannot afford the high price of it. These illuminated us to seek for lower cost, more effective or less side effect drugs for DES.

Esculetin, which was extracted from Chinese herb drug 'Qinpi', was known for inducing apoptosis in human cancer cells,⁴ attenuated the expression of pro-inflammatory mediators in the inflamed colon⁵ and inhibiting VEGF-induced cell proliferation and DNA synthesis with no cell toxicity both *in vitro* and *in vivo*⁶ in recent studies. Researches on Esculetin were systemic

Department of Ophthalmology, Longhua Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China

Correspondence: X Liu, Department of Ophthalmology, Longhua Hospital, Shanghai University of Traditional Chinese Medicine, 725 South Wanping Road, Shanghai 200032, China
Tel: +86 21 64385700; Fax: +86 21 64875390.
E-mail: drliuxinquan@hotmail.com

¹These authors contributed equally to this work.

Received: 11 August 2016
Accepted in revised form: 27 April 2017
Published online: 23 June 2017

administration in mice or rats, while topical Esculetin in curing ocular diseases had not ever been reported before. In this study, we aimed to investigate whether topical Esculetin can improve the DES symptoms and prevent the damage on ocular surface via cytokine expressions and signal pathways. We also intended to observe whether Esculetin is capable of enhance the therapeutic effects of CsA, which may give a chance of reducing the dosage and attenuating side effects of CsA, and contribute to the treatment strategies of DES.

Materials and methods

Experimental animals and ethics statement

Thirty-two adult male New Zealand white rabbits with no signs of ocular inflammatory or gross abnormalities, weighing 2.0–2.5 kg, were obtained from the animal facility of Fudan University, China. Rabbits were reared under standard laboratory conditions as follows: room temperature 22 ± 2 °C, relative humidity $60 \pm 10\%$, and a 12-h light-dark cycle (0800 to 2000). All rabbits were quarantined and acclimatized a week before the experiment and free to food and water throughout the experiment. This study conformed to the standards and procedures for the proper care and use of animals, as described in the US Public Health Service Policy on Humane Care and Use of Laboratory Animals as well as the Association for Research in Vision and Ophthalmology Statement for the use of animals in ophthalmic and vision research. The protocol was approved by the committee on the ethics of animal experiments of Fudan Experimental Center. All operations and tests were performed under general anesthesia by an intramuscular injection of 40 mg/kg ketamine hydrochloride to keep the rabbits immobile and given an overdose of 80 mg/kg nembutal for euthanasia, with every effort to minimize suffering. No blinding research method was executed in the experiments.

Induction of the DE models

The left eyes of all the 32 rabbits were chosen for the induction of DE models and posterior experiments. Eye drops containing 0.3% tobramycin (Alcon Inc., Fort Worth, TX, USA) were applied three times daily for 3 days preoperatively and all operations were performed under sterile conditions in the operating room for animals in Fudan Experimental Center after washing the conjunctival sac and ocular surface with povidone iodine. After instilling 0.5% proparacaine hydrochloride (Alcon Inc.) eye drops, an incision was made in the bulbar conjunctiva, the main lacrimal gland, Harderian gland and nictitating membrane were removed and each

incision was sutured. Eye drops containing 0.3% tobramycin and 0.1% dexamethasone (Alcon Inc.) four times daily, as well as an eye ointment of 0.3% tobramycin and 0.1% dexamethasone (Alcon Inc.) once a night for 7 days to prevent infection of eyes. Esculetin (St Louis, MO, USA) and/or 0.05% CsA (Restasis; Allergan Inc., Irvine, CA, USA) eyedrops were administered 1 week after surgery and DES models were confirmed by a Schirmer I test (SIT) result of ≤ 10 mm/5 min and/or tear-film break-up time (TBUT) ≤ 10 s without anesthetic eye drops and throughout the treatment period (8 weeks) for DES maintenance.

Groups

The 32 rabbits were randomly divided into four groups and the timing of the treatment eye drop instillation was 0800, 0000, 1400 and 1800: Group DES control ($n=8$; 100 μ l of phosphate buffer saline(PBS) topically applied quarter in die(QID) in left eyes); Group CsA ($n=8$; 100 μ l of 0.05% CsA topically applied QID in left eyes); Group Esculetin ($n=8$; 100 μ l of 0.05% Esculetin, which was dissolved in PBS, topically applied QID in left eyes); Group CsA & Esculetin (gourp C&E, $n=8$; 50 μ l of 0.05% CsA combined with 50 μ l of 0.05% Esculetin topically applied QID in left eyes, with a 10 min interval between the instillations of different eyedrops); all the right eyes served as normal controls exposed to no solutions).

Ocular tests

The following ocular tests were performed pre-operation (week 0) and week 1, 2, 4, and 8 after induction of DES, at 1000 in the morning and repeated under the same conditions at 1600 in the afternoon, the average values of each test were calculated and recorded.

Aqueous tear production

The Schirmer's I test was carried out to determine the quantitative of aqueous tear production without anesthetic eye drops. The lower eyelid was slightly pulled down and the Schirmer's test strip (Tianjin Jingming New Technological Development Co., Ltd, China) was placed on the palpebral conjunctival vesica, located near the junction of the middle and outer third of the lower eyelid. After 5 min, the wetting length of the strip was read and data were expressed in millimeters.

Fluorescein on ocular surface

After instilling one drop of 1% fluorescein (FL) solution (Tianjin Jingming New Technological Development Co., Ltd, China), eyes of all rabbits were examined under the

slit-lamp microscope (Topcon SL-D7, Tokyo, Japan) with a cobalt blue filter at a magnification of 16. The cornea was divided into four quadrants and the staining intensity in each quadrant was scored in a scale of 0–3 and for a maximum score of 12. Scoring was achieved using the following guidelines: no punctate staining on the cornea was assigned 0 point; punctate staining of 1–10 was assigned 1 point; punctate staining 11–30 was assigned 2 points; either punctate staining >30 or clumped staining was assigned 3 points.¹

Conjunctival impression cytology

Conjunctival impression cytology (CIC) specimens were collected on week 0, 1, 2, 4, and 8. Two 3.5 × 3.5 mm circular disks of nitrocellulose filter paper (Pall Co., NY, USA) were placed separately on the nasal and temporal bulbar conjunctiva with the filter paper dull-side down. After the filter paper was pressed for 10 s with constant pressure, it was gently lifted and fixed with 95% alcohol. Hematoxylin and periodic acid-Schiff (PAS) reagents were used to stain the specimens. After staining, the number of goblet cells was counted under a microscope (Olympus, Tokyo, Japan) with a 400 objective. Three different sections of each specimen were selected randomly for counting, and the average value was calculated as cells/high-power (HP) visual field with × 400).

Tear fluid collection and determination of IL-1 α , IL-1 β , TNF- α by ELISA

To analyze inflammatory cytokine levels, tear samples were collected according to the method described previously.^{7,8} Total of 50 μ l of PBS was instilled onto the ocular surface, while the animal was alert. The eyelid was manually blinked for several times and the fluids on the ocular surface were collected by sterile capillary tube. All of the tear samples were obtained from the lateral canthus, which parallel to the ocular surface, without stimulating reflex tearing, followed by immediate transfer to a 0.5 ml Eppendorf tube and centrifugation at 1000 r.p.m. for 3 min at 4 °C. The supernatants then were stored at –80°C. The amount of tear IL-1 α , IL-1 β , and TNF- α were measured using an enzyme-linked immunosorbent assay (ELISA) kit (eBioscience, San Diego, CA, USA) according to the manufacturer's instructions. The optical density of each well was determined at a wavelength of 450 nm. Samples were considered positive when the signal was higher than the background signal (modified Krebs solution), and was within the range of the standard curve. The experiment was repeated thrice.

Western blot analysis to detect pERK1/2 protein expression

Western blotting was performed to analyze phosphorylated-ERK1/2 (p-ERK1/2) protein expression. All the rabbits were given an overdose of 80 mg/kg nembutal for euthanasia at week 8. The lower bulbar conjunctivas were blunt desected and cut into small pieces for the determination. Specimens were stored at –80 °C and grindied into powder. Radio immunoprecipitation assay buffer 1 ml and 10 μ l phenylmethyl sulfonyl fluoride were added to each 100 mg tissues on the 60 Hz tissue homogenate machine for 5 min. The supernatants were obtained by centrifugation at 10,000g for 15 min at 4 °C. The protein concentrations in the lysate were quantified by the BCA protocols. The proteins were resolved by 10% SDS–polyacrylamide gel electrophoresis (SDS–PAGE) and transferred onto a nitrocellulose membrane. The membranes were blocked with 5% nonfat milk in TBST (pH 7.6) for 1 hr, incubated overnight at 4°C with anti-p-ERK1/2 antibody (mouse anti-rabbit IgG, NO: 4370S; Cell Signaling Technology Inc., Boston, MA, USA), diluted 1 : 1000, washed extensively and then incubated with secondary antibody (1:5000, goat anti-mouse IgG, No: 7076; Cell Signaling Technology Inc.) for 2 h. The bands were visualized by chemiluminescence and observed at an optimal timepoint. Protein bands were quantified by densitometric analysis using GENE TOOL (UVP). The Western blot was triplicated.

Analysis

Statistical analysis was performed using SPSS17.0 software (SPSS Inc., Chicago, IL, USA). All data were presented as means \pm SD. Values were compared among the groups and time points using analysis of one-way ANOVA, and $P < 0.05$ was considered statistically significant.

Results

Schirmer's I test

No significant difference in aqueous tear production was identified in each group at week 0 ($P > 0.05$). After induction of DES, the wetting length of Schirmer's test strip at week 1, 2, 4, and 8 (8 ± 3 mm, 9 ± 2 mm, 9 ± 2 mm, 8 ± 2 mm) was shorter than week 0 ($P < 0.01$). Schirmer's I test (SIT) values of group CsA (11 ± 2 mm), group Esculetin (9 ± 3 mm), and group C&E (12 ± 2 mm) were obviously higher than group DES control since week 1 ($P < 0.05$). Meanwhile, SIT of group C&E was increased the most at any timespot when compared with groups CsA and group Esculetin ($P < 0.05$), and the highest SIT

values of the three therapy groups (12 ± 1 mm, 10 ± 2 mm, 14 ± 1 mm) were observed at week 4 (Figure 1).

Cornea fluorescein scores

Photographs were obtained at week 0 and 8 in each group (Figures 2a–j). There was no significant difference in minimal corneal punctuate staining in each group at baseline (Figures 2a–e, $P > 0.05$). In the normal control group, no apparent changes in scores were observed throughout the study, while FL scores were significantly increased after operation in other four groups (Figure 2k, $P < 0.01$). FL scores in group DES (9.33 ± 0.33) were higher than group CsA (7.92 ± 0.56), group Esculetin (8.62 ± 0.32) and group C&E (6.71 ± 0.26) since week 1 (Figure 2k, $P < 0.05$). Clumped FL staining was still shown on cornea in group DES (Figure 2g) and punctuate staining in each quadrant of cornea in group Esculetin (Figure 2i). Scores in three therapy groups decreased obviously after instilling different drugs and the lowest results were found in group C&E at week 1, 2, 4, and 8 (Figure 2k, $P < 0.05$). FL scores in group CsA were lower than group Esculetin from week 1 to week 8 (Figure 2k, $P < 0.05$).

Goblet cell density

Assessment of the CIC focused on goblet cell density (Table 1). There was no difference in goblet cell number in

each group preoperative at baseline ($P < 0.05$). Goblet cell density in group DES decreased to 72 ± 21 cells/HP at week 1 and 50 ± 14 cells/HP at week 8. Meanwhile, both group CsA and group Esculetin alleviate decreasing percentage of goblet cell density, respectively $54.83 \pm 6.72\%$ and $52.75 \pm 8.14\%$ at week 8, which showed no significant difference ($P > 0.05$) between the two groups. However, obvious increase of goblet cell density were observed at any time spot in group C&E when compared with group CsA and group Esculetin ($P < 0.05$).

Concentrations of IL-1 α , IL-1 β , and TNF- α

The concentrations of IL-1 α , IL-1 β , and TNF- α were detected by ELISA to estimate the chronic inflammation on ocular surface (Table 2). No significant difference was found of IL-1 α , IL-1 β , and TNF- α concentrations in each group at week 0 ($P > 0.05$). As shown in the tables, DE resulted in higher levels of IL-1 α , IL-1 β , and TNF- α . IL-1 α and IL-1 β concentrations increased gradually during the study ($P < 0.05$), TNF- α increased from week 1 to week 4, and then decreased at week 8 ($P < 0.05$). Upregulations of IL-1 α , IL-1 β , and TNF- α were inhibited in group CsA, group Esculetin and group C&E ($P < 0.05$). The maximum inhibiting effect was noted in group C&E ($P < 0.05$).

Expression of pERK1/2 protein

Immunoblotting analysis was performed to determine the levels of p-ERK1/2 protein in each group before and after therapy. DE activated the ERK1/2 signal pathway and the bands in group DES were much more stronger (gray value was 2.35 fold than GAPDH) than other groups (Figure 3a). Upregulations of p-ERK1/2 in group CsA and group Esculetin were observed and the bands in group CsA were stronger than that in group Esculetin, and gray values were 1.87 fold and 1.44 fold than GAPDH respectively and reduced to 1.12 fold in group C&E (Figure 3b, $P < 0.05$). The difference were obviously revealed among the three therapy groups and the most decreasing of p-ERK1/2 expression was observed in group C&E ($P < 0.05$).

Discussion

Human tear film coats the anterior ocular surface and is composed of three distinct layers: an inner mucin coating, a middle aqueous component, a lipid overlay and the aqueous component is derived from the lacrimal gland.² The lacrimal gland fluid contributes 58–79% of the total volume of tears in rabbits.⁹ The secretory cells of the lacrimal gland produce a highly complex product of water, ions, and proteins,¹⁰ which are the main components of the aqueous layer. In this study, accessory




Figure 1 Comparison of the wetting length of Schirmer I test at different time points. Wetting length of Schirmer's test strip was shorter than week 0 after induction of DES and increasing of SIT was observed after therapy from week 1 to week 8 in group CsA, group Esculetin and group C&E. (* $P < 0.05$ compared with group Normal control at week 0, * $P < 0.05$ compared with group DES at week 1–8 respectively, $\Delta P < 0.05$ compared with group C&E at week 1–8, respectively).




Figure 2 Cornea fluorescein staining observed under the slit-lamp microscope with cobalt blue filter at a magnification of 16. (a–e) were figures observed at week 0 (a) group normal control; (b) group DES; (c) group CsA; (d) group Esculetin; (e) group (c and e). No significant difference on the ocular surface was shown among groups. (f–j) were figures observed at week 8 (f) group normal control; (g) group DES; (h) group CsA; (i) group Esculetin; (j) group (c and e). Clumped staining was observed in (g), punctate staining diminished in (i) and nearly vanished in (h and j). Cornea fluorescein scores of groups at week 0 to week 8 were shown in (k). There was no significant difference in each group preoperatively and the scores increased after establishing of DE models and were persistent. The decreasing of FL scores were observed after therapy with CsA or/and Esculetin eyedrops instilling. (* $P < 0.05$ compared with group Normal control at week 0, # $P < 0.05$ compared with group DES at week 1–8 respectively, $\Delta P < 0.05$ compared with group C&E at week 1–8, respectively).

Table 1 Goblet cell density calculated with microscope at magnification of 400

	Week 0	Week 1	Week 2	Week 4	Week 8
Group Normal	140 ± 23	145 ± 19	138 ± 17	142 ± 11	145 ± 13
Group DES	141 ± 25	72 ± 21*	55 ± 16*	59 ± 15*	50 ± 14*
Group CsA	138 ± 18*	108 ± 20 [#] Δ	86 ± 10 [#] Δ	81 ± 10 [#] Δ	77 ± 12 [#] Δ
Group Esculetin	143 ± 19*	110 ± 17 [#] Δ *	88 ± 13 [#] Δ *	83 ± 13 [#] Δ *	74 ± 12 [#] Δ *
Group C&E	137 ± 17*	126 ± 25 [#]	109 ± 15 [#]	101 ± 14 [#]	92 ± 12 [#]

* $P < 0.05$ compared with group Normal control at week 0–8.

[#] $P < 0.05$ compared with group DES at week 1–8.

$\Delta P < 0.05$ compared with group C&E.

* $P < 0.05$ compared with group CsA at week 1–8.

lacrimal glands in the superior palpebral conjunctiva of rabbits are reserved, which acinar systems have cellular features that resemble of the main lacrimal glands.¹¹ Sjögren’s syndrome (SS) is a chronic and progressive systemic autoimmune disease that primarily involves immune-mediated damage to the lacrimal and salivary glands.^{1,12} This pathology translates into dry eyes (kerato

conjunctivitis sicca) and dry mouth. SS-induced dry eye is one of the primary type of DES. We established the DE models, which were extremely imitating the ocular surface condition of SS-induced dry eye. Objective ocular tests can help diagnosing or monitoring of dry eye.⁴ In group DES control, obvious alternations of the SIT and FL scores and symptoms after establishing the model

Table 2 IL-1 α , IL-1 β and TNF- α expressions detected by ELISA

	Week 0	Week 1	Week 2	Week 4	Week 8
<i>IL-1α (pg/ml)</i>					
Group Normal	5.13 \pm 0.54	5.01 \pm 0.34	4.98 \pm 0.32	5.19 \pm 0.59	5.07 \pm 0.47
Group DES	5.08 \pm 0.62*	7.87 \pm 0.96*	9.57 \pm 1.37*	11.56 \pm 2.33*	13.55 \pm 2.89*
Group CsA	5.21 \pm 0.34*	5.98 \pm 0.86 [#] Δ	6.73 \pm 0.99 [#] Δ	7.37 \pm 1.01 [#] Δ	8.56 \pm 1.67 [#] Δ
Group Esculetin	5.28 \pm 0.69*	6.92 \pm 1.01 [#] Δ	7.85 \pm 1.17 [#] Δ	9.33 \pm 1.58 [#] Δ	8.92 \pm 1.04 [#] Δ
Group C&E	5.15 \pm 0.57*	5.45 \pm 0.57 [#]	5.94 \pm 0.60 [#]	6.29 \pm 0.77 [#]	6.81 \pm 0.69 [#]
<i>IL-1β (pg/ml)</i>					
Group Normal	4.17 \pm 0.32	4.05 \pm 0.19	4.21 \pm 0.41	4.37 \pm 0.39	4.19 \pm 0.30
Group DES	4.26 \pm 0.24*	6.28 \pm 0.51*	8.53 \pm 0.87*	10.11 \pm 1.07*	12.53 \pm 2.19*
Group CsA	4.01 \pm 0.13*	5.46 \pm 0.49 [#] Δ	6.35 \pm 0.67 [#] Δ	7.02 \pm 0.71 [#] Δ	7.95 \pm 0.91 [#] Δ
Group Esculetin	4.23 \pm 0.38*	5.98 \pm 0.54 [#] Δ	6.71 \pm 0.59 [#] Δ	8.82 \pm 0.76 [#] Δ	9.37 \pm 1.25 [#] Δ
Group C&E	4.14 \pm 0.37*	5.15 \pm 0.56 [#]	5.88 \pm 0.62 [#]	6.51 \pm 0.67 [#]	6.72 \pm 0.58 [#]
<i>TNF-α (pg/ml)</i>					
Group Normal	13.81 \pm 2.01	14.11 \pm 2.23	13.75 \pm 1.94	13.92 \pm 2.00	14.25 \pm 2.31
Group DES	13.42 \pm 1.93*	19.87 \pm 3.16*	25.32 \pm 4.01*	37.59 \pm 4.31*	35.92 \pm 3.57*
Group CsA	14.02 \pm 1.57*	16.17 \pm 1.89 [#] Δ	18.97 \pm 2.03 [#] Δ	19.54 \pm 2.12 [#] Δ	19.63 \pm 1.78 [#] Δ
Group Esculetin	13.94 \pm 1.26*	19.13 \pm 2.91 [#] Δ	21.67 \pm 3.11 [#] Δ	20.09 \pm 2.34 [#] Δ	20.98 \pm 2.02 [#] Δ
Group C&E	13.28 \pm 0.96*	14.29 \pm 1.54 [#]	16.12 \pm 1.79 [#]	16.78 \pm 1.59 [#]	16.85 \pm 1.46 [#]

* $P < 0.05$ compared with group Normal control.

[#] $P < 0.05$ compared with group DES.

$\Delta P < 0.05$ compared with group C&E.




Figure 3 Western blot analysis of p-ERK1/2 protein in group Normal, DES, CsA, Esculetin and C&E. (a) Representative Western blots. (b) Quantification of p-ERK1/2 in each group using densitometric analysis of the bands shown in (a). Expression levels values normalized to GAPDH levels. (* $P < 0.05$ compared with group DES, [#] $P < 0.05$ compared with group CsA, $\Delta P < 0.05$ compared with group C&E).

continued during the whole experimental period (Figure 1). SIT valve was 15.75 \pm 2 mm before surgery and reduced to 8 \pm 3 mm in week 1. Tear secretion and tear film may be related to either compensation of neurally regulated conjunctival goblet cell mucin secretion¹³ or secretion of the accessory lacrimal gland on the conjunctiva.

Cyclosporine A (CsA) is a neutral, hydrophobic, cyclic peptide of amino acids which was isolated from several species of fungi¹⁴ and used for corneal graft rejection, autoimmune uveitis, and dry eye syndrome¹⁵ through suppressing cell-mediated immune responses.^{16,17} The mechanism based on its effects on subconjunctival and lacrimal gland inflammation, resulting in an increase in tear production, and conjunctival goblet cell density.^{3,18} Ophthalmic CsA has been available for a few years and beneficial in DES, which recommended concentration is 0.05%, could slow down or prevent the progression of DES.¹⁹ It is important to consider both the benefits and risks of dry eye therapy and select the treatment appropriately. Esculetin is a derivative of coumarin that is present in Chinese herbal drug 'Qinpi', used as folk medicines, such as *Fraxinus rhynchophylla*, *Rehmannia glutinosa* and *Artemisia capillaries*.^{20–23} Previously, the inhibitory effect of Esculetin on cell proliferation in human cell lines had been reported.^{24,25} Esculetin reduced CCl₄-induced hepatic apoptosis in rats probably against hepatocellular apoptosis with its free-radical scavenging ability.²⁶ It was also suggested to be a potent cancer therapeutic and preventive agent. Lee *et al* identified Esculetin as a potent inhibitor of Wnt signaling, decreased viability, inhibited anchorage-independent growth of colon cancer cells and tumor growth in a xenograft mouse model.²⁷ Esculetin was capable of inhibiting growth of hepatocellular carcinoma both *in vivo* and *in vitro*, it also resulted in Bax/Bcl-2 ratio, activation of the caspase cascade and loss of MMP.²⁸ Topical Esculetin on ocular

surface had never been reported ever before. We administrated 0.05% Esculetin solution which concentration was identical to the CsA to the DE rabbits and found that Esculetin have effectively increased SIT values (9 ± 3 mm), goblet cell densities (110 ± 17 cells/HP) and decreased FL scores (9.12 ± 0.32) since week 1. Effects of Esculetin continuously maintained till week 8, while the SIT values were 9 ± 1 mm, goblet cell densities were 74 ± 12 cells/HP and FL scores were 7.48 ± 0.33 . These results were superior to that in group DES control, even though the extent was moderate than CsA or Esculetin combining with CsA (Figures 1, 2 and Table 1). We can also estimate that Esculetin improve DE symptoms through stimulating aqueous secretion of accessory lacrimal gland or neurally regulated conjunctival goblet cell mucin secretion.

There are many cellular and molecular process involved in the pathogenesis of dry eye, including increased inflammatory cytokines, tear film osmolarity, metalloproteinases, chemokines and their receptors, inflammatory cascades, and activation of immune cells.²⁹ The major pro-inflammatory cytokines found to be important in pSS are interferons, IL-12, IL-18, TNF α , IL-1 β , IL-6, IL-17, IL-23 and B-cell activating factor (BAFF).³⁰ There was profound increment of IL-4 and IL-5, which were produced by T helper2 (Th2) cells, in the salivary proteome of patients with pSS.³¹ Cai *et al*³² found that Esculetin inhibit IL-6 expression and it could also inhibit NO, TNF- α , IL-1 β effectively when combined with curcumin and hesperetin. IL-6 and IL-8 production were prevented with Esculetin in LPS-induced rat intestinal microvascular endothelial cells.³³ We investigated the IL-1 α , IL-1 β , TNF α concentrations of tear samples in ELISA assay and gradual increasing of all the three kinds of cytokines were shown from week 1 to week 8 in group DES control. IL-1 α , IL-1 β , TNF α concentrations were gradually increased in three therapy groups at relatively low amplitude (Table 2). Therefore, effects of topical Esculetin and/or CsA were detected obviously in this essay and the most expecting results appeared in group C&E. This was not only proving Esculetin inhibit expressions of some interleukins, but also reveal that Esculetin enhance the anti-inflammatory function of CsA.

ERK cascade plays a crucial role in multiple cellular processes such as cell proliferation, differentiation, adhesion, migration, survival and apoptosis.^{34,35} ERK1/2 pathway participated in the pro-inflammatory responses mounted by pSS salivary gland epithelial cells.³⁶ Experimental induced dry eye in mice stimulated expression and production of IL-1 β , TNF- α , MMP-9, and activated ERK1/2 signaling pathway on the ocular surface,³⁷ some growth factor besides EGF stimulated conjunctival goblet cell proliferation by ERK1/2 pathway.³⁸ Dryness also increased expressions of IL-1 α ,

IL-1 β and TNF- α mRNA and stimulated phosphorylation of JNK1/2, ERK1/2 and p38 MAPKs in the corneal epithelium³⁹ and even activation of several signaling pathway, including ERK in RPE-choroid AMD phenotypes and MAPK inhibitors could provide a novel therapeutic strategy for prevention or treatment of AMD.^{40,41} In normal control group, gray value of p-ERK1/2 was 0.46 fold than GAPDH, while it was 2.31 fold in group DES. This once again verified the correlation between chronic ocular surface inflammation and ERK1/2 signal pathway. When refer to the therapy groups, gray values were respectively 1.87 fold in group CsA, 1.44 fold in group Esculetin and 1.12 fold in group C&E (Figure 3). Results indicated suppression of ERK1/2 pathway after different therapy, consistent with observations in the former parts of the study and confirmed the effect of Esculetin combining with CsA. We found that p-ERK1/2 expression in group Esculetin was nearly 30% lower than group CsA. It was maybe because of ERK1/2 pathway play a more important role in DES treating with Esculetin, rather than with CsA.

In conclusion, we topically applied Esculetin firstly to search for a new potential drug for the patient suffering from DES. The results of ocular tests, cytokine levels and specific phosphorylated protein of ERK1/2 signal pathway proved that Esculetin has resemble therapeutic effect on curing DES when compared with CsA. Esculetin was capable of enhancing the effect of CsA and meanwhile it was more available and in lower price in eastern countries. We only observed the ERK1/2 pathway in this article and whether there exist other pathways crosstalking with it remains uncertain. All these encourage us to approaching for the concrete mechanism and most effective concentration of topical Esculetin in anti-inflammatory of DES.

Summary

What was known before

- High prevalence of dry eye syndrome in aged people was noticed and its clinical manifestation is chronic inflammation on the ocular surface. CsA can improve the symptoms and signs of dry eye, however, it has side effects.

What this study adds

- Topical Esculetin improves DE symptoms, downregulates the inflammatory cytokine expressions, suppresses the ERK1/2 signal pathway and enhances the therapeutic effect of CsA.
-

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

We gratefully acknowledge Fudan Animal Experimental Center for supporting sterile experimental operation room, technical assistance in animal rearing and sample testing, Xiaofei Song (Department of Urology, Minhang Affiliated Hospital of Fudan University) for data statistical analysis, Zhaoyang Wang (Department of Ophthalmology, Shanghai No. 9 Hospital) for native English instructions. This work was supported by the Natural Science Foundation of China (No. 81574028), Shanghai Health Bureau of Traditional Chinese Medicine Scientific Fund (No. 2012J005A) and Longhua Medical Scholar (LYTD-50).

References

- 1 The definition and classification of dry eye disease: report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop. *Ocul Surf* 2007; **5**(2): 75–92.
- 2 Conrady CD, Joos ZP, Patel BC. Review: the lacrimal gland and its role in dry eye. *J Ophthalmol* 2016; **2016**: 7542929.
- 3 Sall K, Stevenson OD, Mundorf TK, Reis BL. Two multicenter, randomized studies of the efficacy and safety of cyclosporine ophthalmic emulsion in moderate to severe dry eye disease. CsA Phase 3 Study Group. *Ophthalmology* 2000; **107**(4): 631–639.
- 4 Kim AD, Madduma Hewage SR, Piao MJ, Kang KA, Cho SJ, Hyun JW. Esculetin induces apoptosis in human colon cancer cells by inducing endoplasmic reticulum stress. *Cell Biochem Funct* 2015; **33**(7): 487–494.
- 5 Yum S, Jeong S, Lee S, Kim W, Nam J, Jung Y. HIF-prolyl hydroxylase is a potential molecular target for esculetin-mediated anti-colitic effects. *Fitoterapia* 2015; **103**(6): 55–62.
- 6 Park SL, Won SY, Song JH, Lee SY, Kim WJ, Moon SK. Esculetin inhibits VEGF-induced angiogenesis both *in vitro* and *in vivo*. *Am J Chin Med* 2016; **44**(1): 61–76.
- 7 Lee SY, Han SJ, Nam SM, Yoon SC, Ahn JM, Kim TI et al. Analysis of tear cytokines and clinical correlations in Sjögren syndrome dry eye patients and non-Sjögren syndrome dry eye patients. *Am J Ophthalmol* 2013; **156**(2): 247–253.e1.
- 8 Zhou L, Wei R, Zhao P, Koh SK, Beuerman RW, Ding C. Proteomic analysis revealed the altered tear protein profile in a rabbit model of Sjögren's syndrome-associated dry eye. *Proteomics* 2013; **13**(16): 2469–2481.
- 9 Thörig L, van Agtmaal EJ, Glasius E, Tan KL, van Haeringen NJ. Comparison of tears and lacrimal gland fluid in the rabbit and guinea pig. *Curr Eye Res* 1985; **4**(8): 913–920.
- 10 Walcott B. The lacrimal gland and its veil of tears. *News Physiol Sci* 1998; **13**: 97–103.
- 11 Bergmanson JP, Doughty MJ, Blocker Y. The acinar and ductal organization of the tarsal accessory lacrimal gland of Wolfring in rabbit eyelid. *Exp Eye Res* 1999; **68**(4): 411–421.
- 12 Tincani A, Andreoli L, Cavazzana I, Doria A, Favero M, Franceschini F et al. Novel aspects of Sjögren's syndrome in 2012. *BMC Med* 2013; **4**(11): 93.
- 13 Dartt DA. Regulation of mucin and fluid secretion by conjunctival epithelial cells. *Prog Retin Eye Res* 2002; **21**(6): 555–576.
- 14 Petcher TJ, Weber H, Rueegger A. Crystal and molecular structure of an iodo derivative of the cyclic undecapeptide cyclosporin A. *Helv Chim Acta* 1976; **59**(5): 1480–1489.
- 15 Nussenblatt RB, Palestine AG, Chan CC, Stevens Jr G, Mellow SD, Green SB. Randomized, double-masked study of cyclosporine compared to prednisolone in the treatment of endogenous uveitis. *Am J Ophthalmol* 1991; **112**(2): 138–146.
- 16 Noble S, Markham A. Cyclosporin. A review of the pharmacokinetic properties, clinical efficacy and tolerability of a microemulsion-based formulation (Neoral). *Drugs* 1995; **50**(5): 924–941.
- 17 Donnenfeld E, Pflugfelder SC. Topical ophthalmic cyclosporine: pharmacology and clinical uses. *Surv Ophthalmol* 2009; **54**(3): 321–338.
- 18 Wilson SE, Perry HD. Long-term resolution of chronic dry eye symptoms and signs after topical cyclosporine treatment. *Ophthalmology* 2007; **114**(1): 76–79.
- 19 Yavuz B, Bozdag Pehlivan S, Unlu N. An overview on dry eye treatment: approaches for cyclosporine a delivery. *ScientificWorldJournal* 2012; **2012**: 194848.
- 20 Hong GE, Park HS, Kim JA, Nagappan A, Zhang J, Kang SR et al. Anti-oxidant and anti-inflammatory effects of Fraxinus rhynchophylla on lipopolysaccharide (LPS)-induced murine Raw 264.7 cells. *J Biomed Res* 2012; **13**(4): 331–338.
- 21 Wu ZB, Liu Y, Tian SS, Wen C. Chemical constituents of the stem bark of Fraxinus rhynchophylla. *Chem Nat Compd* 2014; **49**(6): 1162–1163.
- 22 Si CL, Liu Z, Su YF, Kim JK, Bae YS. Coumarins and secoiridoid glucosides from bark of Fraxinus rhynchophylla Hance. *Holzforchung* 2008; **62**(5): 553–555.
- 23 Prabakaran D, Ashokkumar N. Protective effect of esculetin on hyperglycemia-mediated oxidative damage in the hepatic and renal tissues of experimental diabetic rats. *Biochimie* 2013; **95**(2): 366–373.
- 24 Ng TB, Liu F, Wang ZT. Antioxidative activity of natural products from plants. *Life Sci* 2000; **66**(8): 709–723.
- 25 Adfa M, Yoshimura T, Komura K, Koketsu M. Antitermite activities of coumarin derivatives and scopoletin from Protium javanicum Burm. *J Chem Ecol* 2010; **36**(7): 720–726.
- 26 Tien YC, Liao JC, Chiu CS, Huang TH, Huang CY, Chang WT et al. Esculetin ameliorates carbon tetrachloride-mediated hepatic apoptosis in rats. *Int J Mol Sci* 2011; **12**(6): 4053–4067.
- 27 Lee SY, Lim TG, Chen H, Jung SK, Lee HJ, Lee MH et al. Esculetin suppresses proliferation of human colon cancer cells by directly targeting β -catenin. *Cancer Prev Res (Phila)* 2013; **6**(12): 1356–1364.
- 28 Wang J, Lu ML, Dai HL, Zhang SP, Wang HX, Wei N. Esculetin, a coumarin derivative, exerts *in vitro* and *in vivo* anti proliferative activity against hepatocellular carcinoma by initiating a mitochondrial-dependent apoptosis pathway. *Braz J Med Biol Res* 2015; **48**(3): 245–253.
- 29 Coursey TG, de Paiva CS. Managing Sjögren's syndrome and non-Sjögren's syndrome dry eye with anti-inflammatory therapy. *Clin Ophthalmol* 2014; **8**: 1447–1458.
- 30 Roescher N, Tak PP, IIIei GG. Cytokines in Sjögren's syndrome. *Oral Dis* 2009; **15**(8): 519–526.
- 31 Delaleu N, Mydel P, Kwee I, Brun JG, Jonsson MV, Jonsson R. High fidelity between saliva proteomics and the biologic state of salivary glands defines biomarker signatures for primary Sjögren's syndrome. *Arthritis Rheumatol* 2015; **67**(4): 1084–1095.
- 32 Cai C, Chen Y, Zhong S, Zhang Y, Jiang J, Xu H et al. Synergistic effect of compounds from a Chinese herb:

- compatibility and dose optimization of compounds from N-Butanol extract of *Ipomoea stolonifera*. *Sci Rep* 2016; **6**: 27014.
- 33 Hu Y, Chen X, Duan H, Hu Y, Mu Y. Chinese herbal medicinal ingredients inhibit secretion of IL-6, IL-8, E-selectin and TXB2 in LPS-induced rat intestinal microvascular endothelial cells. *Immunopharmacol Immunotoxicol* 2009; **31**(4): 550–555.
- 34 Yoon S, Seger R. The extracellular signal-regulated kinase: multiple substrates regulate diverse cellular functions. *Growth Factors* 2006; **24**(1): 21–44.
- 35 Busca R, Pouyssegur J, Lenormand P. ERK1 and ERK2 map kinases: specific roles or functional redundancy. *Front Cell Dev Biol* 2016; **4**: 53.
- 36 Sisto M, Lisi S, D'Amore M, Lofrumento DD. The metalloproteinase ADAM17 and the epidermal growth factor receptor (EGFR) signaling drive the inflammatory epithelial response in Sjögren's syndrome. *Clin Exp Med* 2015; **15**(2): 215–225.
- 37 Luo L, Li DQ, Doshi A, Farley W, Corrales RM, Pflugfelder SC. Experimental dry eye stimulates production of inflammatory cytokines and MMP-9 and activates MAPK signaling pathways on the ocular surface. *Invest Ophthalmol Vis Sci* 2004; **45**(12): 4293–4301.
- 38 Li D, Shatos MA, Hodges RR, Dartt DA. Role of PKC α activation of Src, PI-3K/AKT, and ERK in EGF-stimulated proliferation of rat and human conjunctival goblet cells. *Invest Ophthalmol Vis Sci* 2013; **54**(8): 5661–5674.
- 39 De Paiva CS, Corrales RM, Villarreal AL, Farley WJ, Li DQ, Stern ME et al. Corticosteroid and doxycycline suppress MMP-9 and inflammatory cytokine expression, MAPK activation in the corneal epithelium in experimental dry eye. *Exp Eye Res* 2006; **83**(3): 526–535.
- 40 Makarev E, Cantor C, Zhavoronkov A, Buzdin A, Aliper A, Csoka AB. Pathway activation profiling reveals new insights into age-related macular degeneration and provides avenues for therapeutic interventions. *Aging (Albany NY)* 2014; **6**(12): 1064–1075.
- 41 SanGiovanni JP, Lee PH. AMD-associated genes encoding stress-activated MAPK pathway constituents are identified by interval-based enrichment analysis. *PLoS One* 2013; **8**(8): e71239.