

PROSTHETIC MANAGEMENT OF PATIENTS WITH ORO-MAXILLO-FACIAL DEFECTS: A LONG-TERM FOLLOW-UP RETROSPECTIVE STUDY

G. GASTALDI, L. PALUMBO, C. MORESCHI, E.F. GHERLONE, P. CAPPARÉ

Dental School, Vita-Salute University, Milan, Italy (Dean: Prof. Enrico Felice Gherlone)

SUMMARY

Introduction. The aim of this study is to determine the outcome of maxillofacial prosthetic rehabilitation after oncological resections, including both intra- and extra-oral prosthetic devices.

Methods. In this study were included 72 patients, who have undergone an intra or extra-oral maxillofacial prosthetic rehabilitation after an oncologic resection.

Tumors on the head and neck were analyzed and the defects of these resections have been divided in two different groups: intra and extra-oral defects.

Results. 72 participants were treated with maxillofacial prosthesis, 3 of which with post-traumatic wounds and 69 with resections of tumors on the head and neck. Of the 69 treated for neoplastic disease, 43 received an intraoral prosthesis (palatal obturator) and 29 with an extraoral epithesis (18 with nasal prostheses, 8 with orbital implants and 3 with ear implants). The group included patients with different types of tumors. All the patients were evaluated in terms of aesthetic appearance after the construction of the prostheses and the results were satisfactory.

Conclusion. Within the limitations of this study, after the use of maxillofacial prostheses patients feel more confident and self-assured. Maxillofacial prostheses are a good solution in order to improve the life's quality in patients with tumors resections: prostheses are easy to handle and provide a satisfying social interaction for the patients.

Key words: maxillofacial, cancer, defect, prosthodontics.

Introduction

In the last years, an increase of oral-pharynx cancer has been registered. Tumors of oral-cranial-facial area, with 6% of prevalence, are placed at 6th position of malignant tumors ranking. The survival rate of a localized tumor at 5 years is around 82% (1, 2).

Males are more affected than women, 2:1. The therapy, based on the onset place and stage, can be divided in surgery, chemotherapy, radiotherapy or a combination of these actions (3-5). The resections of these tumors can produce defects of the oral and nasal cavities, nasopharynx, oropharynx

and extra-oral defects.

The defects can be divided in two different groups: intra and extra-oral results. The most frequent intra-oral defects are related to the loss of palatal portion.

Optimal aesthetical and functional reconstruction in the head and neck area is important for the social integration and the quality of life of patients (6).

Extended craniofacial defects can led to wide functional and psychosocial impairment in patients. Functional limitations can affect vision, speech, mastication and swallowing (7).

The postsurgical defects are critical in many ways, leading to these patients retracting from their fam-

ily and society and living a life of exclusion and depression (7).

After the surgical resection, a rehabilitation is needed. Reconstruction of the resulting defects can be achieved by means of reconstructive plastic surgery and/or maxillofacial prostheses.

The restoration of defects (8) can be traditionally achieved with the aid of the conventional surgery; disadvantage, however, is the necessity for multiple procedures (9, 10). In addition, surgical reconstruction may be limited by general medical condition, insufficient residual tissue, vascular compromise subsequent to radiation, age, inadequacy of the donor sites, or patient preference. It is not always possible to reconstruct the defect with a surgical approach (11).

In these cases, prosthetic rehabilitation become the first choice treatment (12). The rehabilitation with maxillofacial prosthesis aims to restore an effective division between oral, nasal or orbital cavities and gives faster reconstructive possibilities, simplifying the post-surgery period and trying to recover an adequate patient lifestyle.

A collaboration between surgeon, prosthodontist and technician is required to realize an obturator prosthesis immediately after surgery, in order to improve the wound healing and the integration with the patient tissue (13).

Nevertheless, more protocols are needed in order to improve the predictability of results and guidelines are necessary.

The aim of this study is to determine the successful of maxillofacial prosthetic rehabilitations after oncological resections or post traumatic results, including both intra- and extra-oral devices.

Materials and methods

In this retrospective study were included 72 patients, who have undergone an intra or extra-oral maxillofacial prosthetic rehabilitation after an oncological resection or post traumatic results, treated within the San Donato Hospital Group, Italy.

Tumors on the head and neck were analyzed and

the defects of these resections can be divided in two different groups: intra and extra-oral.

Patients with extra-oral lesions have been treated immediately after the end of the healing process with an epithesis: an impression in alginate (Hydrogum 5, Zhermack, Rovigo, Italy) was performed, using a wet gauze to prevent material from infiltrating into the cavities. In many cases, it was considered necessary to ensure completion of a temporary prosthesis; despite the aesthetic limitations, this solution could be helpful in improving the patient's psychological aspect.

Even for the intra-oral restorations, the palatal obturator has been realized after an alginate (Hydrogum 5, Zhermack Rovigo, Italy) impression for the realization of temporary prosthesis: the palatal obturator has been made of an acrylic resin on the patient casts; it was later refined with methyl methacrylate (Ivocron, Ivoclar Vivadent, Bolzano, Italy) and refined with soft materials directly in the oral cavity in order to adapt the prosthesis to the defect. Patients were included in a very detailed follow-up program (every 7 days) for 30-45 days. When the healing process was complete, more accurate impressions were taken with polysulfide using the palatal obturator as a guide; the final restoration can also contain the teeth.

Results

Seventy-two participants were treated with maxillofacial prosthesis, 3 of which with post-traumatic wounds and 69 with resections of tumors on the head and neck.

Of the 69 treated for neoplastic disease, 43 received an intraoral prosthesis (palatal obturator) and 29 with an extraoral epithesis (18 with nasal prostheses, 8 with orbital implants and 3 with ear implants).

The group included patients with different types of tumors (Chart 1, Table 1), respectively:

- 36% squamous cancer
- 20% basal cell carcinoma
- 16% adenoid cystic carcinoma

- 11% mucoepidermoid carcinoma
- 17% others.

In total 30 patients were treated with radiotherapy, 4 with radio- and chemotherapy, 10 were not treated with radiation and 28 are not available for the series.

The radiation dose was calculated around 12-70 Gy.

No case of osteoradionecrosis was documented in our study group; no complications like oral lesions, sensibility impairment, or wound infections were observed at medium-term follow-up.

Any implant failure or soft tissue reactions have been observed in our study group.

All the patients were evaluated in terms of aesthetic appearance after the construction of the prostheses and the results were satisfactory (Figures 1-5).

The patients had to receive a new epithesis 1-2 years after anchorage of the initial epithesis, especially due to deterioration of the colour and quality.

The follow-up ranged from 2.5 to 7.8 years (mean 5.6 years \pm 2.8).

The patient survival rate was 93.06% (5 patients died).



Figure 1
Frontal view of the patient after surgical resection.

Discussion

In this study 72 patients were treated, 3 of them with post-traumatic results and 69 with resec-

tions of head and neck tumors. According to the literature, studies have shown that an unrepaired maxillary defect can result in high incidence of hypernasality (14, 15) and low speech intelligi-




Figure 2
Detail of defects: a) intraoral view of oro-nasal defect; b) frontal view of nasal defect.



Figure 3
A lateral view of palatal obturator prosthesis with extension for epithesis attachment.




Figure 4
The epithesis allows both prompt inspection of the resection site and makes daily care easier.



Figure 5
Frontal view of the patient after superior intraoral prosthesis and nasal epithesis delivery.




Table 1 - The percentage of cancer's type.

Squamous Cancer	36
Basel Cell Cancer	20
Adenoid Cystic Carcinoma	16
Mucoepidermoid Carcinoma	11
Adenocarcinoma	4
Chondrosarcoma	2
Osteosarcoma	4
Giant Cell Tumor	2
Other	5

bility (14, 16). This is due to inadequate separation of the oral and nasal cavities. After prosthetic treatment, speech were improved. However, maxillary obturators have been cited as being uncomfortable to wear and some patients find them inconvenient to remove and clean (17, 18). The success of a prosthesis is related to the extent on the size and location of the maxillary defect, as well as the presence of remaining dentition (10). These factors can affect the stability of the prosthesis influencing its effectiveness. Maxillary obturators require that a maxillofacial prosthodontist be available for construction and maintenance of the prosthesis (15, 18).

With regard to functional and aesthetic aspects,

combination of dentures linked to extraoral facial prostheses, enabled optimal orofacial rehabilitation. Oral functions such as chewing, swallowing and speaking were facilitated by the dentures which at the same time worked as a stabilisation element for the prostheses.

The facial prostheses were individually adapted in order to resemble the familiar preoperative appearance of the patient.

The available ways of prostheses anchorage are four (19, 20): the anatomical (to already existing structures), the mechanical (to spectacle frames), the chemical (using adhesive) and the surgical anchorage (by osseointegrated titanium implants with magnets).

The use of fixed facial prostheses is based on a close cooperation between prosthodontic and surgeon to provide the optimal aesthetic and functional outcome.

Maxillofacial prostheses is appropriate (21) in the cases of large midfacial defect after a disfiguring cancer surgery, since it is very difficult if not impossible to reconstruct these defects by the grafting techniques using autogenous tissue and to achieve satisfactory results.

Orofacial rehabilitation of patients with maxillofacial defects using obturator prostheses is an appropriate treatment modality. To improve the situation of patients prior to and after maxillectomy sufficient information about the treatment, adequate psychological care and speech therapy should be

provided.

With the use of obturator protheses and/or epithesis, patients regain self-confidence and assurance. The patients recounted a high level of satisfaction and a positive impact on daily life. According to Sullivan et al., Wondergem et al. and Rieger et al., the goal of any intervention, whether surgical or prosthetic, is to limit the impact of the oncologic treatment on these aspects of patients' lives.

One of the main pros reported by patients is the fact that they are submit to surgery only-once reaching the same result as the surgery. On the other hand, disadvantages and limitations of these protheses include discoloration (as regards epithesis) and protheses deterioration and skin reactions (19, 20).

According to our experience, an intra- or extra oral protheses rehabilitation should be favored over a plastic reconstruction in the cases of the previous multiple necessary plastic operations, due to prior huge surgical resection which makes the surgical reconstruction technically impossible, or in cases, the patients prefer the prosthetic solution.

Nevertheless, further studies are needed in order to draft new guidelines. It is expected that future improvements in techniques of implantation and virtual technique for optical 3d acquisition (22, 23) will further improve the overall satisfaction and wellbeing of the patients, but this requires a well-considered approach and close collaboration between the surgeon and the prosthodontics (15).

Conclusion

The clinical use of fixed facial protheses is based on a close cooperation between surgeon and prosthodontist to provide the optimal aesthetic and functional outcome.

Within the limitations of this study, after the use of maxillofacial protheses patients feel more confident and self-assured. Maxillofacial protheses are a good solution in order to improve the life's quality in patients with tumors resections: protheses are easy to handle and provide a satisfying social interaction for the patients.

Authors' contributions

Giorgio Gastaldi: clinical procedures, conception and design, drafting the article.

Luca Palumbo: drafting the article, data analysis.

Chiara Moreschi: data acquisition and analysis.

Enrico F. Gherlone: critical revision, final approval.

Paolo Capparé: drafting the article, critical revision, clinical procedures.

Funding

No funding were requested for this study. The Authors had no conflict of interest in connection to this study.

References

1. Cancer incidence by site. Age-standardized rate per 100,000. Statistics Canada and the Canadian Council of Cancer Registries, Health Protection Branch - Laboratory Centre for Disease Control, 1999.
2. WHO Mortality Database. Age-standardized rate per 100,000. WHO Databank, 1999.
3. Murrah VA, Batzakis LG. Proliferative verrucous leukoplakia and verrucous hyperplasia. *Ann Otol Rhinol Laryngol.* 1994;103:660-3.
4. Buglione M, Cavagnini R, Di Rosario F, Sottocornola L, Maddalo M, Vassalli L, Grisanti S, Salgarello S, Orlandi E, Paganelli C, Majorana A, Gastaldi G, Bossi P, Berruti A, Pavanato G, Nicolai P, Maroldi R, Barasch A, Russi EG, Raber-Durlacher J, Murphy B, Magrini SM. Oral toxicity management in head and neck cancer patients treated with chemotherapy and radiation: Dental pathologies and osteoradionecrosis (Part 1) literature review and consensus statement. *Crit Rev Oncol Hematol.* 2016. Jan;97:131-42. doi: 10.1016/j.critrevonc.2015.08.010.
5. Buglione M, Cavagnini R, Di Rosario F, Maddalo M, Vassalli L, Grisanti S, Salgarello S, Orlandi E, Bossi P, Majorana A, Gastaldi G, Berruti A, Trippa F, Nicolai P, Barasch A, Russi EG, Raber-Durlacher J, Murphy B, Magrini SM. Oral toxicity management in head and neck cancer patients treated with chemotherapy and radiation: Xerostomia and trismus (Part 2). Literature review and consensus statement. *Crit Rev Oncol Hema-*

- tol. 2016 Jun;102:47-54. doi: 10.1016/j.critrevonc.2016.03.012.
6. Gherlone EF, Capparé P, Tecco S, Polizzi E, Pantaleo G, Gastaldi G, Grusovin MG. Implant Prosthetic Rehabilitation in Controlled HIV-Positive Patients: A Prospective Longitudinal Study with 1-Year Follow-Up. *Clin Implant Dent Relat Res*. 2016 Aug;18(4):725-34. doi: 10.1111/cid.12353.
 7. Suarez-Cunqueiro MM, Schramm A, Schoen R, et al. Speech and swallowing impairment after treatment for oral and oropharyngeal cancer. *Arch Otolaryngol Head Neck Surg*. 2008;134:1299-1304.
 8. Park SS. Reconstruction of nasal defects larger than 1.5 centimeters in diameter. *Laryngoscope*. 2000 Aug;110(8):1241-50. PubMed PMID: 10942120.
 9. Korfage A, Raghoobar GM, Noorda WD, Plaat BE, Vissink A, Visser A. Recommendations for implant-retained nasal prostheses after ablative tumor surgery: Minimal surgical aftercare, high implant survival, and satisfied patients. *Head Neck*. 2016 Apr;38(Suppl 1):E619-24. doi: 10.1002/hed.24053. Epub 2015 Jul 18. PubMed PMID: 25784187.
 10. Rohrich RJ, Griffin JR, Ansari M, Beran SJ, Potter JK. Nasal reconstruction-beyond aesthetic subunits: a 15-year review of 1334 cases. *Plast Reconstr Surg*. 2004 Nov;114(6):1405-16; discussion 1417-9. Review. PubMed PMID 15509926.
 11. Kumar TP, Azhagarasan NS, Shankar KC, Rajan M. Prosthetic rehabilitation of orofacial donor site fistula following surgical reconstruction: a clinical report. *J Prosthodont*. 2008;17:336-339.
 12. Sakuraba M, Kimata Y, Ota Y, et al. Simple maxillary reconstruction using free tissue transfer and prostheses. *Plast Reconstr Surg*. 2003;111:594-8. doi:10.1097/01.PRS.0000041941.98504.B6.
 13. Davis BK, Roumanas ED, Nishimura RD. Prosthetic-surgical collaborations in the rehabilitation of patients with head and neck defects. *Otolaryngol Clin North Am*. 1997 Aug;30(4):631-45. Review. PubMed PMID: 9233862.
 14. Arigbede AO, Dosumu OO, Shaba OP, Esan TA. Evaluation of speech in patients with partial surgically acquired defects: pre and post prosthetic obturation. *J Contemp Dent Pract*. 2006;7:89-96.
 15. Keyf F. Obturator prostheses for hemimaxillectomy patients. *J Oral Rehabil*. 2001;28:821-9. doi:10.1046/j.1365-2842.2001.00754.x.
 16. Umino S, Masuda G, Ono S, Fujita K. Speech intelligibility following maxillectomy with and without a prosthesis: an analysis of 54 cases. *J Oral Rehabil*. 1998;25:153-8. doi:10.1046/j.1365-2842.1998.00238.x.
 17. Triana RJ, Uglesic V, Virag M, et al. Microovascular free flap reconstructive options in patients with partial and total maxillectomy defects. *Arch Facial Plast Surg*. 2000;2:91-101. doi:10.1001/archfaci.2.2.91.
 18. Browne JD, Burke AJC. Benefits of routine maxillectomy and orbital reconstruction with the rectus abdominis free flap. *Otolaryngol Head Neck Surg*. 1999;121:203-9. doi:10.1016/S0194-5998(99)70172-5.
 19. Federspil PA. Auricular prostheses. *Adv Otorhinolaryngol*. 2010;68:65-80. doi 10.1159/000314563. Epub 2010 May 3. PubMed PMID: 20442562.
 20. Federspil PA. Implant-retained craniofacial prostheses for facial defects. *GMS Curr Top Otorhinolaryngol Head Neck Surg*. 2009;8:Doc03. doi: 10.3205/cto000055. Epub 2011 Mar 10. PubMed PMID: 22073096; PubMed Central PMCID: PMC3199820.
 21. Scolozzi P, Jaques B. Treatment of midfacial defects using prostheses supported by ITI dental implants. *Plast Reconstr Surg*. 2004 Nov;114(6):1395-404. Review. PubMed PMID: 15509925.
 22. Sansoni G, Trebeschi M, Docchio F, Gastaldi G. State-of-The-Art and Applications of 3D Imaging Sensors in Industry, Cultural Heritage, Medicine, and Criminal Investigation. *Sensors (Basel)*. 2009;9(1):568-601. doi: 10.3390/s90100568. Epub 2009 Jan 20. PubMed PMID: 22389618; PubMed Central PMCID: PMC3280764.
 23. Sansoni G, Cavagnini G, Docchio F, Gastaldi G. Virtual and physical prototyping by means of a 3D optical digitizer: Application to facial prosthetic reconstruction. *Virtual Physical Prototyping*. 2009;4(4):217-226.

Correspondence to:

Dr. Paolo Capparé
 Department of Dentistry
 IRCCS San Raffaele Hospital
 Via Olgettina 48
 20132 Milan, Italy
 Phone: 0039 0226433619; fax 0039 0226432953;
 E-mail: cappare.paolo@hsr.it