

# Correction: Integration of flux measurements to resolve changes in anabolic and catabolic metabolism in cardiac myocytes


Andrew A. Gibb, Pawel K. Lorkiewicz, Yu-Ting Zheng, Xiang Zhang, Aruni Bhatnagar, Steven P. Jones and Bradford G. Hill



*Biochem. J.* 2017, volume 474, issue 16, pages, 2785–2801; <https://doi.org/10.1042/BCJ20170474>

In the isotopologue analysis of glycerol(phospho)lipids (Figure 5), the authors mistakenly included glycerolipids with only the glycerol moiety labeled (i.e.,  $m+3$ ) in the Glycerol + Fatty Acid labeling groups (Glycl+FA). Fractional enrichment of  $^{13}\text{C}$  into Glycl+FA of panels B–F should include only those glycerolipids having both the glycerol and fatty acyl chains labeled (e.g. this would include  $m+5$ ,  $m+7$ ,  $m+9$ ,  $m+11$ ,  $m+13$ ,  $m+15$ , etc.). The corrected figure is presented here. These changes do not change the conclusions drawn from the study and do not require modifications to the text of the manuscript.

Correction published:  
14 December 2017



**Figure 5. Phosphofructokinase regulates glycerolipid biosynthesis.**

Stable isotope tracing of phospholipids and triacylglycerols in cardiomyocytes incubated with media containing  $^{13}\text{C}_6$ -glucose for 18 h: (A) atom-resolved map illustrating the biological and biochemical history of  $^{13}\text{C}$  incorporation into glycerolipids; fractional enrichment values of  $^{13}\text{C}$  into: (B) PC; (C) PI; (D) PE; (E) PS; and (F) TAG. Graph represents three replicates per group from one isolation. \*\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\*\* $P < 0.0001$ .