

Genomic and Transcriptomic Analysis of *Escherichia coli* Strains Associated with Persistent and Transient Bovine Mastitis and the Role of Colanic Acid

John D. Lippolis,^a  Devin B. Holman,^b Brian W. Brunelle,^b  Tyler C. Thacker,^c Bradley L. Bearson,^d Timothy A. Reinhardt,^a Randy E. Sacco,^a Thomas A. Casey^b

^aRuminant Diseases and Immunology Research Unit, National Animal Disease Center, Agricultural Research Service, U.S. Department of Agriculture, Ames, Iowa, USA

^bFood Safety and Enteric Pathogens Research Unit, National Animal Disease Center, Agricultural Research Service, U.S. Department of Agriculture, Ames, Iowa, USA

^cBacterial Diseases of Livestock Research Unit, National Animal Disease Center, Agricultural Research Service, U.S. Department of Agriculture, Ames, Iowa, USA

^dAgroecosystems Management Research Unit, National Laboratory for Agriculture and the Environment, Agricultural Research Service, U.S. Department of Agriculture, Ames, Iowa, USA

ABSTRACT *Escherichia coli* is a leading cause of bacterial mastitis in dairy cattle. It is most often transient in nature, causing an infection that lasts 2 to 3 days. However, *E. coli* has been shown to cause a persistent infection in a minority of cases. Mechanisms that allow for a persistent *E. coli* infection are not fully understood. The goal of this work was to determine differences between *E. coli* strains originally isolated from dairy cattle with transient and persistent mastitis. Using RNA sequencing, we show gene expression differences in nearly 200 genes when bacteria from the two clinical phenotypes are compared. We sequenced the genomes of the *E. coli* strains and report genes unique to the two phenotypes. Differences in the *wca* operon, which encodes colanic acid, were identified by DNA as well as RNA sequencing and differentiated the two phenotypes. Previous work demonstrated that *E. coli* strains that cause persistent infections were more motile than those that cause transient infections. Deletion of genes in the *wca* operon from a persistent-infection strain resulted in a reduction of motility as measured in swimming and swarming assays. Furthermore, colanic acid has been shown to protect bacteria from complement-mediated killing. We show that transient-infection *E. coli* strains were more sensitive to complement-mediated killing. The deletion of genes from the *wca* operon caused a persistent-infection *E. coli* strain to become sensitive to complement-mediated killing. This work identifies important differences between *E. coli* strains that cause persistent and transient mammary infections in dairy cattle.

KEYWORDS colanic acid, *Escherichia coli*, mastitis, pathogenesis

Mastitis is the most common health problem in the dairy industry. The approximate economic impact of mastitis in dairy cattle to the U.S. economy is \$2 billion of lost revenue per year (1). Numerous microorganisms can cause mastitis, and manifestation of the disease can range from completely subclinical to severe and life threatening.

Escherichia coli is a common cause of mastitis in dairy cows and is a leading cause of acute mastitis (2). Typically, an *E. coli* mastitis infection is transient in duration, but persistent intramammary infections can occur (3, 4). Research suggests that 5 to 20% of mammary-pathogenic *E. coli* (MPEC) infections are persistent (4–6). Strains of MPEC that were isolated from persistent intramammary infections have been distinguished from MPEC strains that were isolated from transient infections in several ways. First,

Received 4 August 2017 Returned for modification 1 September 2017 Accepted 13 October 2017

Accepted manuscript posted online 23 October 2017

Citation Lippolis JD, Holman DB, Brunelle BW, Thacker TC, Bearson BL, Reinhardt TA, Sacco RE, Casey TA. 2018. Genomic and transcriptomic analysis of *Escherichia coli* strains associated with persistent and transient bovine mastitis and the role of colanic acid. Infect Immun 86:e00566-17. <https://doi.org/10.1128/IAI.00566-17>.

Editor Guy H. Palmer, Washington State University

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply. Address correspondence to John D. Lippolis, john.lippolis@ars.usda.gov.

MPEC strains that were associated with persistent infections have been shown to invade cultured mammary epithelial cells more effectively than strains that cause transient infections (7). Second, MPEC strains that were associated with persistent infections were found to have greater motility *in vitro*, as demonstrated by increased rates of swimming and swarming, than MPEC strains that cause transient mastitis (8).

Previously, we have shown that growth of *E. coli* in milk compared to that in Luria-Bertani (LB) broth caused massive changes in protein expression levels (9). In fact, over 50% of the 1,000 proteins identified had significant expression changes between the *E. coli* grown in milk and that grown in bacterial growth medium. Some of the proteins downregulated in milk were important for bacterial motility. Because previous swimming and swarming assays were performed in LB broth-based medium, we sought to determine if these phenotypes were affected by milk.

Analysis of protein expression differences between the persistent- and transient-infection strains (8) identified 28 proteins whose change in expression levels correlated with disease phenotype (e.g., persistent versus transient). One group of these proteins consisted of flagellar and chemotaxis proteins. The goal of the transcriptomics experiment in this paper was to identify additional differences between the persistent- and transient-infection strains grown in milk to help explain these pathogenic differences.

The objective of this work is to identify mechanisms that allow a subset of MPEC strains to establish a persistent infection and therefore improve our understanding of bacterial pathogenesis in the mammary gland. Utilizing transcriptomics, genomics, and phenotypic assays, we report that the *wca* operon plays a critical role in determining whether the *E. coli* strain will produce a transient or persistent infection.

RESULTS

Motility in milk plates. Our previous work showed that three persistent-infection MPEC strains (strains P1 to P3) were more motile in standard LB broth-based swimming and swarming plates than transient-infection MPEC strains (strains T1 to T3) (10). Because milk affects the expression of motility proteins (9), milk-based agar plates were developed to assay swimming and swarming phenotypes among the MPEC isolates. Commercially available milk agar plates are made with tryptone, yeast extract, dextrose, skim milk powder, and agar, providing nutrients not typically found in milk. Therefore, for this swimming assay we developed milk-based agar plates that contained 90% fresh skim milk and a 10× agar solution (final agar concentration, 0.35%). Our plates did not contain typical bacterial growth medium components (e.g., tryptone or yeast extract) in order to generate growth conditions more similar to *in vivo* conditions. Figure 1A shows a representative photograph of a less motile MPEC strain, and Fig. 1B shows a highly motile MPEC strain. The persistent-infection strains, as a group, were significantly more motile in this assay than the transient-infection strains ($P < 0.0001$) (Fig. 1C). Additionally, all three persistent-infection strains were significantly more motile than any of the three transient-infection strains (Fig. 1C). These results are consistent with our results previously reported using normal LB swimming plates (10). In the milk and LB swimming assays the time required for the bacteria to swim the same distance was greater for the milk-based plates than for the LB swimming plates (15 versus 5 h).

Gene expression differences between transient- and persistent-infection strains. Each of the six MPEC strains was grown in fresh milk to mid-log phase as previously described (8). This experiment was repeated three times, and RNA from each strain was independently isolated and sequenced. Transcriptome sequencing (RNA-seq) data between replicate experiments for individual strains were consistent, as shown by a principal-component analysis (PCA) plot (Fig. 2). The variation between replicates for each strain was highly consistent ($r = 0.98$ or greater for all 6 strains).

Gene expression analysis identified 285 genes that were differentially expressed between the two phenotypic groups (false discovery rate [FDR], $P < 0.05$) (see Table S2 in the supplemental material). Of the 285 genes, 132 were expressed at higher levels in persistent-infection strains, and 153 were expressed at higher levels in transient-infection strains. To categorize the functions associated with the differentially ex-




FIG 1 Milk swimming plates. Shown are examples of two *E. coli* strains on milk swimming plates. The first is strain T2 (A), which is a slower swimmer, and the second is P1 (B), a faster swimmer. The relative swimming abilities of these two strains on the milk swimming plates are consistent with data on the traditional swimming plate with the exception that growth is much slower. (C) Combined results from three swimming assays on skim milk plates. There is a significant difference between the persistent-versus transient-infection groups ($P < 0.001$). In addition, all three P strains are significantly different from all three T strains. Different lowercase letters indicate significantly different means. Error bars represent the standard errors of the means.

pressed genes, functional enrichment analysis was performed using DAVID (11, 12). The biological process gene ontology (GO) terms associated with higher expression in persistent-infection strains include flagellum-dependent motility, assembly and organization, lipopolysaccharide and slime layer biosynthetic process, iron ion homeostasis and response to iron, and cell adhesion (Table 1). The biological process terms associated with higher gene expression in transient-infection strains involved the molybdopterin cofactor biosynthetic process (Table 1). In agreement with the pheno-




FIG 2 Principal-component analysis (PCA) plot of RNA-seq data. The PCA plot shows the variance of the three experimental replicates of each of the three transient-infection strains (T1, T2, and T3) and the three persistent-infection strains (P1, P2, and P3). The percentages on each axis represent the percentages of variation explained by the principal components.

TABLE 1 Enriched gene ontology biological processes in persistent- versus transient-infection MPEC strains based on differentially expressed genes

Gene group and GO term	No. of genes	P value	Gene names
Genes with higher expression in persistent-infection isolates			
Bacterial-type flagellum-dependent cell motility	15	1.10E-13	<i>fliF, fliE, flgK, flgL, fliK, fliG, flhM, fliN, fliS, fliJ, flgJ, flhI, fliH, flgI, flhT</i>
Lipopolysaccharide biosynthetic process	15	2.10E-10	<i>arnT, wzzE, wcaH, wecA, wcaK, wcaM, arnA, eptA, wcaI, wcaL, wcaN, wzzB, arnC, arnD, arnB</i>
Bacterial-type flagellum assembly	9	7.10E-09	<i>flgN, fliQ, flhD, flgK, fliK, fliG, fliS, flgJ, flhI</i>
Bacterial-type flagellum organization	7	2.50E-07	<i>flgM, fliO, flhP, flhC, fliJ, fliH, flhT</i>
Iron ion homeostasis	7	1.10E-04	<i>fecI, fecA, fecR, fecE, fecB, fecC, fecD</i>
Chemotaxis	6	6.80E-04	<i>dppA, flhO, fliG, fliN, fliJ, trg</i>
Response to iron(III) ion	4	7.00E-04	<i>arnT, basR, basS, arnD</i>
Lipid A biosynthetic process	6	8.40E-04	<i>arnT, arnA, eptA, arnC, arnD, arnB</i>
Slime layer polysaccharide biosynthetic process	4	3.80E-03	<i>wcaK, wcaM, wcaI, wcaL</i>
Ion transport	6	3.90E-03	<i>fecI, fecR, fecE, fecB, fecC, fecD</i>
Gamma-aminobutyric acid catabolic process	3	1.10E-02	<i>gabT, gabP, gabD</i>
Cell adhesion	5	2.40E-02	<i>fimI, fimD, fimF</i>
Genes with higher expression in transient-infection isolates			
Mo-molybdopterin cofactor biosynthetic process	4	8.20E-03	<i>moaB, moaC, moaE, moaD</i>
Molybdopterin cofactor biosynthetic process	3	1.50E-02	<i>moaB, moaC, moaE</i>

typic motility results, 15 flagellar genes were upregulated in the persistent-infection strains compared to their levels in the transient-infection isolates. Additionally, we show that 11 genes involved in iron regulation have increased transcription levels in the persistent-infection isolates. This observation is consistent with previously published data (8, 10). Unique to this data set are the differentially expressed genes within the *wca* operon. The *wcaGHIKLM* genes all had greater expression levels in the persistent-infection strains.

Genetic differences between transient- and persistent-infection strains. We reasoned that the difference in gene expression levels between these strains could be due to differential transcription or a difference in gene content at the genome level. To address this second possibility, we sequenced all six strains. Four strains have been previously sequenced (13), and we added two previously unsequenced genomes to the NCBI database for ECA-B and ECC-M. Sequence information was aligned to the *E. coli* strain MG1655 (GenBank accession number [NC_000913](https://www.ncbi.nlm.nih.gov/nuccore/NC_000913)). There were 3,782 genes in MG1655 that were found in at least one of the three persistent- or three transient-infection strains (Table S3). There were 3,788 genes in all six strains (using a minimal read number of 75). In the MG1655 matched data, there were 14 genes that were present in all three persistent-infection strains but absent in the transient-infection strains. There were three genes that were in the transient-infection strains but not in the persistent-infection strains. *De novo*-assembled sequences from the six strains that did not match MG1655 were subjected to a BLAST search against the NCBI nonredundant (nr) database. An additional 16 genes from the transient-infection strains were identified that did not exist in the persistent-infection strains (Table S4). Additionally, 90 genes were identified in the persistent-infection strains that did not exist in the transient-infection strains (Table S5). The genes identified by the BLAST search are likely plasmid genes as they included conjugative genes on several contigs. Of all the genetic differences between the strains, the most striking were the 14 genes matched to MG1655 that were present only in the three persistent-infection strains. All 14 of these genes are in the same location on the chromosome, which includes part of the *wca* operon (Table 2). The *wca* operon encodes part of the colanic acid metabolic pathway. The absence of the genes within the *wca* operon in the transient-infection strains is consistent with the transcriptomics data showing differential expression of the colanic acid metabolic process (Table 1).

Effect of gene deletions in the *wca* operon. The most motile MPEC strain, P1 (strain ECC-Z), was used to make gene deletion mutants to test the hypothesis that the

TABLE 2 MiSeq read data of six MPEC strains

Gene name ^a	No. of reads by strain ^b					
	T1	T2	T3	P1	P2	P3
<i>hisL</i>	1139.9	1022.0	792.8	1269.1	1197.0	1207.2
<i>hisG</i>	257.5	12.0	222.1	258.6	277.2	235.1
<i>hisD</i>	236.2	9.5	186.7	242.4	273.7	243.3
<i>hisC</i>	275.4	11.1	203.3	258.9	285.5	252.6
<i>hisB</i>	235.4	20.9	198.0	224.5	246.5	233.5
<i>hisH</i>	276.2	12.1	233.5	286.9	313.2	268.2
<i>hisA</i>	269.8	15.7	246.3	288.0	274.0	250.8
<i>hisF</i>	269.2	6.0	223.4	285.7	246.7	248.6
<i>hisI</i>	235.5	9.5	188.5	277.9	84.9	256.5
wzzB	0.0	0.0	0.0	136.8	101.7	58.8
<i>ugd</i>	10.2	0.0	0.0	20.0	188.2	22.8
gnd	16.4	28.2	0.0	179.9	194.8	173.1
<i>wbbL</i> (pseudogene)	NA	NA	NA	NA	NA	NA
<i>insH1</i>	0.0	229.4	0.0	0.0	0.0	0.8
<i>wbbK</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>wbbJ</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>wbbI</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>wbbH</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>glf</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>wzxB</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>rfbC</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>rfbA</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>rfbD</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>rfbB</i>	0.0	0.0	0.0	157.3	0.0	0.0
wcaN	27.6	5.2	0.4	202.3	177.7	163.6
wcaM	0.0	0.0	0.0	200.4	207.9	200.5
wcaL	0.0	0.0	0.0	195.3	200.2	237.3
wcaK	0.0	0.4	0.0	200.1	188.3	243.6
wzxC	2.3	0.0	0.0	245.4	250.5	209.4
wcaJ	15.4	0.0	0.0	238.9	230.3	228.7
cpsG	1.7	0.0	0.3	249.2	252.7	215.5
cpsB	0.0	2.3	0.0	204.8	228.6	193.6
wcaI	0.0	0.0	0.0	269.5	290.5	254.3
wcaH	0.0	0.0	0.0	360.3	389.0	317.4
wcaG	0.0	0.0	0.0	276.4	313.1	270.1
gmd	42.6	0.7	0.0	235.7	265.2	237.3
<i>wcaF</i>	243.7	0.5	0.0	294.5	304.9	288.1
<i>wcaE</i>	269.7	0.7	0.0	264.9	278.1	268.4
<i>wcaD</i>	266.3	57.1	0.0	249.8	260.4	239.5
<i>wcaC</i>	265.2	242.0	0.0	255.6	292.8	237.3
<i>wcaB</i>	337.1	321.5	0.0	339.7	320.3	288.8
<i>wcaA</i>	266.7	215.7	0.0	221.2	233.8	247.4
<i>wzc</i>	231.2	231.3	0.2	223.5	239.3	211.0
<i>wzb</i>	387.4	367.1	0.0	354.7	373.9	313.7
<i>wza</i>	263.8	231.0	0.0	239.8	277.9	224.8
<i>yegH</i>	247.4	235.7	155.5	232.3	239.3	247.1
<i>asmA</i>	225.6	223.4	190.1	225.9	231.2	221.0
<i>dcd</i>	326.1	321.7	263.2	330.9	333.2	291.6

^aGenes in boldface were absent from all three transient strains (T1 to T3) but present in all three persistent strains (P1 to P3).

^bData are reads per kilobase of transcript per million mapped reads. Shading indicates genes with insufficient read numbers to be counted as present. NA, not available.

missing genes are responsible for differences in swimming and swarming between the persistent- and transient-infection MPEC strains. The first deletion removed the genes from *gmd* through *wcaN*, making P1 similar to strain T1 in that region of the chromosome (Table 2). The second mutation deleted the *motAB* genes, which encode proteins that are part of the flagellar motor, thereby allowing the bacteria to make structurally complete but nonfunctional flagella.

Figure 3 shows the data from three independent experiments for the motility assays of swimming (Fig. 3A) and swarming (Fig. 3B). As expected, deletion of the *motAB* genes caused a loss of both swimming and swarming. The deletion of part of the *wca* operon




FIG 3 Motility assays with deletion mutants. Results of three independent assays with three replicate plates per assay were averaged. The relative growth for each strain was calculated using the average P1 growth as a reference. (A) Swimming assay. (B) Swarming assay. Error bars represent the standard errors of the means.

that approximates the genes missing from the transient-infection strains caused a 20% reduction in swimming and a 40% reduction in swarming compared to levels of the most motile strain, P1.

A recent paper has shown that colanic acid has a protective role against complement-mediated killing in serum (14). To test whether MPEC bacteria that cause persistent and transient infections have different susceptibilities to complement in serum, the three persistent-infection strains, the three transient-infection strains, and the Δwca mutant were grown in 50% fresh bovine serum and 50% LB broth or in 50% heat-inactivated serum and 50% LB broth (Fig. 4). Growth of strain T2 and the Δwca mutant was completely inhibited in normal serum. Strains T1 and T3 showed growth curves that were repressed in normal serum compared to growth of the persistent-infection strains. Growth of all three persistent-infection strains was unaffected by culture with normal serum. Importantly, heat inactivation of the serum (56°C for 30 min) removed any growth inhibition and allowed all strains to grow at roughly similar rates.

To test whether preexposure to an MPEC mammary infection affected the efficacy of serum to inhibit bacterial growth, sera from four cows that were part of an MPEC mastitis study within the last year, as well as sera from four heifers (never lactated), were compared for their ability to inhibit growth of the persistent- and transient-infection MPEC strains (Fig. 5). The growth of the three transient-infection strains was more inhibited in normal serum than that of the persistent-infection strains, consistent with data presented in Fig. 4. There was no difference in the abilities to inhibit growth




FIG 4 Serum inhibition assay. Persistent- and transient-infection strains and the P1 Δwca mutant were grown in 50% LB broth and either 50% fresh normal bovine serum (A) or heat-inactivated serum (B), as indicated. Optical density (at 600 nm) was monitored in five replicate wells for each bacterial strain, and the experiment was repeated three times. Groups of strains that were statistically different ($P < 0.05$) are indicated by different lowercase letters.

of the transient-infection strain between the sera from cows with recent known MPEC mastitis and heifers without any known mastitis.

DISCUSSION

Our research goal is to better understand the pathogenic mechanisms that allow a subset of MPEC strains to persist in the mammary gland of dairy cattle. Our previous work demonstrated that growth of bacteria in whole fresh milk significantly altered the proteome of an MPEC strain compared to that of a strain grown in bacterial culture medium (9). The differentially expressed proteins that were categorized as chemotaxis and motility were all downregulated in the MPEC grown in milk, including the flagellin protein *FliC* which is the major subunit of flagella. We have shown a correlation between motility and protein expression in persistent-infection MPEC strains, and a greater motility phenotype was determined in swimming and swarming assays of persistent-infection strains than in transient-infection strains (10). We previously used bacterial culture medium-based swimming and swarming plates, and because MPEC growth in milk had a negative effect on motility protein expression, we developed a milk-based motility plate assay. Using semisolid agar motility plates that were 90% fresh




FIG 5 Sera from four previously MPEC-infected cows (Challenged) were compared to sera from four heifers (Naive) for the ability to inhibit growth of various MPEC strains. Normal serum (A) and serum heat inactivated at 56°C for 30 min (B) were used. Error bars represent standard errors of the means.

skim milk, we have shown that persistent-infection MPEC strains were more motile than the transient-infection strains, consistent with our previous results in standard medium (LB broth). The change in medium did lead to a proportional increase in motility assay time for all the strains (approximately three times as long).

To better understand the differences between the persistent- and transient-infection MPEC strains, we performed RNA sequencing on each strain grown in whole fresh milk. Confirmatory to our previous proteomics data (10), our gene expression data showed higher expression of motility genes in the persistent-infection MPEC strains than in the transient-infection strains. Our RNA sequencing data also demonstrated higher expression levels of genes important for iron acquisition and transport in the persistent-infection strains. It has been shown that iron acquisition and transport genes are important for motility in these MPEC strains (8). However, the largest group of genes with significant expression changes consisted of those involved in colanic acid biosynthesis.

Our subsequent genome sequencing determined that a significant portion of the operon (*wca*) responsible for colanic acid synthesis is missing in the transient-infection strains. Previous work has shown a connection between motility and colanic acid synthesis (15). In that work, the Rcs signaling system was found to positively regulate colanic acid synthesis and negatively regulate motility. We have deleted part of the *wca* operon that is absent in all three transient-infection strains from the persistent-infection strain P1 (P1 Δ*wca*). We observed that the deletion of a significant segment of the *wca*

operon (P1 Δwca) inhibited swimming by approximately 20% and swarming by 40% compared to the levels of the wild-type strain (P1).

Colanic acid is induced by culturing bacteria under suboptimal conditions, such as osmotic and oxidative stress (16) and adverse temperature (17). Recently, colanic acid has been shown to have a protective role against complement-mediated killing in serum (14). The three persistent-infection MPEC strains showed little difference in growth rates in either normal bovine serum or heat-inactivated bovine serum. The three transient-infection MPEC strains were inhibited for growth in normal serum. Growth of the transient-infection strain T2 and the deletion mutant P1 Δwca was completely inhibited in normal bovine serum. Growth of the three transient-infection strains and P1 Δwca was comparable to that of the persistent-infection strains when the bovine serum was heat inactivated. Together, these data highlight the dual role of colanic acid synthesis in motility and resistance to serum for the persistent-infection MPEC isolates.

We also have shown that previous exposure to MPEC mastitis does not affect the serum inhibition of the transient-infection strains. Sera from heifers, adult female cattle that have not calved, were equal to sera from cows that were part of a recent MPEC challenge experiment in their ability to inhibit the transient-infection strains. This indicates that an innate immune response that potentially helps to control MPEC infections is inhibited by MPEC that can express colanic acid.

Understanding of the role of complement in mastitis is incomplete. While levels of the major complement proteins are significantly lower in milk than in serum (18), complement proteins can rise dramatically in clinical mastitis in cows (18, 19). MPEC infections have been shown to increase permeability between the vascular system and the mammary gland, as shown by increased bovine serum albumin (BSA) levels in milk in the acute phase of an infection (20, 21). Disruption of the physiological barrier between blood and milk may be how complement can play a role in intramammary infections. The ability of certain MPEC strains to synthesize colanic acid may provide these strains with protection against complement during the acute phase of the infection.

E. coli mastitis is caused by a highly diverse group of MPEC strains (22, 23). No specific virulence factor has been identified that is necessary for infection by *E. coli* of the mammary gland. The great diversity of clinical outcomes of *E. coli* mastitis is thought to be due largely to host factors (24). However, prior to our work, there have been no genetic differences identified between *E. coli* strains that cause persistent infections and those that cause transient infections (22). Our observations of the differences between MPEC strains that cause persistent versus transient infection indicate that persistent-infection strains are more motile, have increased gene expression of the iron acquisition system, and are more resistant to growth inhibition caused by complement. A relationship between iron acquisition and motility has been shown in these strains (8), and a relationship between colanic acid synthesis and motility has been described previously (15). Generation of specific deletion mutants that will test the role of motility, iron acquisition, and colanic acid synthesis *in vivo* will be necessary to further determine the mechanisms necessary for persistence of *E. coli* in mammary gland infections.

The use of antibiotics in agriculture is of concern to many consumers, policy makers, and farmers. Current treatment recommendations for dairy cattle with mild to moderate *E. coli* mastitis include use of anti-inflammatory therapy, frequent milking, and fluid treatment. Antibiotic treatment of *E. coli* mastitis is not recommended except in severe cases (25). Currently, there are no recommendations for treatment of persistent MPEC infections. Identification of *E. coli* strains that might progress to persistent infection will lead to better decisions and options for mastitis treatment.

MATERIALS AND METHODS

Bacteria. *E. coli* strains used for these experiments were described previously (7). Abbreviations for the bacterial strains used in the manuscript correspond with the published names as follows: T1,

ECA-727; T2, ECA-O157; T3, ECA-B; P1, ECC-Z; P2, ECC-M; P3, ECC-1470 (kindly provided by Ynte Schukken).

E. coli deletion mutants were derivatives of the P1 (ECC-Z) strain. Recombineering was performed using a *neo* cassette encoding kanamycin resistance as previously described (26). Briefly, the oBBI 92/93 *neo* fragment was amplified using a set of knockout primers containing nucleotide sequences corresponding to the *E. coli* genomic region targeted for deletion. Following PCR purification, the amplification product was transformed into strain EBB 667 containing pKD46, and knockout mutants were selected on LB medium containing kanamycin. The *neo* gene was deleted by transforming EBB 671 (*motAB::neo*) and EBB 683 [*wcaNMLK wzcW wcaJ cpsGB wcaIHG gmd::neo*] with pCP20 and screening the resulting transformants for sensitivity to kanamycin. PCR primers were used confirm gene deletions in EBB 673 (Δ *motAB*) and EBB 684 [Δ (*wcaNMLK wzcW wcaJ cpsGB wcaIHG gmd*)]; referred to as the Δ *wca* strain in the manuscript for simplicity. A list of strains and primers is found in Table S1 in the supplemental material.

Animals. Blood samples from eight Holstein cows were obtained from our herd at the National Animal Disease Center. Milk used in transcriptomic experiments was fresh unprocessed bovine milk obtained from multiple cows and pooled. Milk was plated on bacterial growth plates and incubated overnight at 37°C to ensure clean milk samples prior to use for bacterial assays. The National Animal Disease Center Animal Care and Use Committee approved all animal-related procedures used in this study.

Milk swimming assay. A modified swimming assay was developed using skim milk as a medium. Fresh unprocessed bovine milk from multiple cows was pooled and then centrifuged at $1,500 \times g$ for 45 min. Fat was scraped off, and milk was transferred to a new bottle that was centrifuged at $10,000 \times g$ for 45 min. The skim milk and the 3.5% (10 \times) agar solution were equilibrated to 80°C and then mixed for a final agar concentration of 0.35%. Plates were poured and left at room temperature for 1 h and then transferred to 4°C for 1 h. Plates were inoculated by spotting 5 μ l of a bacterial overnight culture to the center of the plate, and the bacteria were allowed to grow at 37°C in a humidified incubator for approximately 15 h. In this assay, motile bacteria will swim from the center toward the outside edge of the plates. Bacterial motility zones were visualized by crystal violet staining. Briefly, 10 ml of 100% ethanol was gently overlaid on the surface of the plate for 5 min. After the ethanol was removed, the plates were then stained with 0.002% crystal violet in 10% ethanol for 1 h. Plates were destained with water in three washes. The diameter of the bacterial motility zones was measured from photographs of the plates using ImageJ (NIH). There were four plates per strain used in each experiment, and the diameter for each strain was averaged. The experiment was repeated four times. Statistical analysis of swimming on milk plates comparing all persistent-infection strains to all transient-infection strains was accomplished with an unpaired *t* test and one-way analysis of variance (ANOVA) with Tukey's multiple-comparison test. The statistical software used was Prism, version 7 (GraphPad, San Diego, CA).

RNA sequencing. (i) Bacterial growth and isolation. Clean fresh milk was pooled from three cows and inoculated with bacterial stocks for overnight cultures grown at 37°C with shaking at 180 rpm. Overnight cultures were diluted into 100 ml of fresh clean milk and grown to mid-log phase (8). Milk cultures were then poured into 300 ml of ice-cold Hanks balanced salt solution with 10 mM EDTA, and the solutions were cooled in a salted ice bath for 5 min. Diluted milk cultures were centrifuged at $10,000 \times g$ for 30 min at 4°C. The pellets were resuspended in 2 ml of RNA Protect and 1 ml of phosphate-buffered saline (PBS) and held at room temperature for 5 to 30 min. The resuspended pellet was passed through a 5- μ m-pore-size syringe filter to remove somatic cells (27). The filtrates were centrifuged at $10,000 \times g$ for 2 min in microcentrifuge tubes, and the supernatants were removed. Cell pellets were stored at -80°C .

(ii) RNA isolation and sequencing. An RNeasy minikit (Qiagen, Germantown, MD) was used to extract RNA from the bacterial cells, and genomic DNA was subsequently removed using Turbo DNA-free DNase (Ambion, Austin, TX) as per the manufacturer's directions. Quantitation of total RNA was performed using a NanoDrop ND-1000 spectrophotometer (Thermo Scientific, Wilmington, DE). RNA quality was assessed with a 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). A quantitative PCR (qPCR) assay was performed on all samples using primers for both *E. coli gapA* and bovine *BRPS9* housekeeping genes to identify potential *E. coli* DNA and bovine RNA contamination. rRNA was depleted using a Ribo-Zero rRNA Removal kit (Bacteria) according to the manufacturer's instructions (Illumina, Inc., San Diego, CA), and rRNA removal was verified on the 2100 Bioanalyzer. The cDNA libraries were constructed using TruSeq Stranded Total RNA Library Prep kits and sequenced on a HiSeq 3000 using a 100-cycle single-end run (Illumina, Inc., San Diego, CA) at the Iowa State University DNA core facility. As per the recommendations of Haas et al. (28), we targeted 10 to 50 million reads per sample; our sequencing resulted in an average of 38.2 ± 5.8 million reads per sample.

RNA-seq analysis. FastQC, version 0.11.5, was used to assess the quality of all RNA-seq reads. Bowtie2, version 2.2.9 (29), was then used to align the reads to the *E. coli* K-12 substrain MG1655 genome (GenBank accession number [NC_000913](https://ncbi.nlm.nih.gov/nuccore/NC_000913)) using the default parameters for the very-sensitive-local setting. The *E. coli* strain MG1655 was used as a reference genome because of its level of gene functional annotation and coverage. The number of reads that aligned to each gene from the *E. coli* K-12 substrain MG1655 genome was quantified using HTSeq, version 0.6.1 (30). Gene counts were then normalized, and differentially expressed genes (false discovery rate [FDR] of <0.05) between the persistent- and transient-infection strains were identified using the DESeq2 package, version 1.12.4 (31) in R, version 3.2.5. The regularized logarithm-transformed counts for each *E. coli* strain were used to generate a principal-component analysis (PCA) plot. To assess the efficiency of the removal of cattle RNA, we aligned the raw RNA-seq reads against the *Bos taurus* reference genome (GenBank accession number [GCA_000003055.3](https://ncbi.nlm.nih.gov/nuccore/GCA_000003055.3)).

It was determined that $0.17\% \pm 0.03\%$ of reads aligned to the *Bos taurus* genome, indicating that host cell contamination was not a significant issue for downstream analysis.

DNA sequencing and analysis. DNA was isolated and purified using CsCl gradients as described by Hull et al. (32), with the exceptions that bacteria were harvested from 6 ml of overnight LB cultures and that bacterial lysates were centrifuged at $372,000 \times g$ for 4 h in a Beckman VTI 65 rotor. Isolated DNA was sent to the Iowa State University DNA core facility. DNA was fragmented using a Bioruptor followed by TruSeq (Illumina, Inc.) library preparation with indexing. Insert size was around 700 bp. Samples were run on a MiSeq using a MiSeq reagent kit, version 3 (Illumina, Inc.) (600 cycles). Sequencing resulted in over 3.7×10^6 reads per genome. Sequences were aligned to the reference sequence of *E. coli* strain MG1655 (GenBank accession number [NC_000913](#)) using DNASTar LaserGene NGen, version 14 (Madison, WI). Average read depth was $>150\times$. Genes with few or no reads were considered absent. In addition to the alignment of the MG1655 reference genome, all reads were *de novo* assembled using Abyss (version 1.9) (33). Sequence reads that were not aligned to the reference strain (MG1655) were *de novo* assembled using NGen. The resulting contigs were subjected to a BLAST search against the NCBI nr database and were identified as plasmids.

Bacterial swimming and swarming assays. A 5- μ l aliquot of a fresh overnight culture of each *E. coli* strain was plated on 0.3% swimming agar plates (10 g of Bacto tryptone, 5 g of Bacto yeast extract, 5 g of NaCl, 3 g of Bacto agar in 1 liter of deionized water) or 0.5% swarming agar plates (10 g of Bacto tryptone, 5 g of Bacto yeast extract, 5 g of NaCl, 5 g of Bacto agar, 0.5% [wt/vol] glucose in 1 liter of deionized water) (34, 35). Each *E. coli* strain was inoculated onto four plates and then incubated for approximately 5 h for swimming and 12 h for swarming in a humidified 37°C incubator. Photographs of all the plates were taken with a fixed distance between the plate and the camera. Swimming and swarming experiments had four replicate plates for each strain, and results were averaged for each experiment. The swimming and swarming experiments were performed three times. ImageJ (NIH) was used to determine the diameter of the swimming plates and area of the outer bacterial growth circle of swarming plates. Differences were assessed among the strains within each motility condition by one-way ANOVA with Tukey's multiple-comparison test using Prism, version 7 (GraphPad).

Serum inhibition test. Serum was isolated from two cows and was mixed in equal volumes. Half the serum was heat inactivated at 56°C for 30 min. Bacterial strains were grown in Luria-Bertani (LB) broth overnight at 37°C with shaking at 200 rpm. Aliquots of the overnight culture (10 μ l) were added to a 1:1 mixture of LB broth and normal cow serum or to a 1:1 mixture of LB broth and heat-inactivated cow serum (190 μ l) into a 100-well microtiter plate. Each condition was replicated in five wells. Plates were loaded onto a Bioscreen C growth curve instrument (Growth Curves, Ltd., Raisio, Finland). Plates were incubated at 37°C with continuous shaking, and the optical density at 600 nm (OD_{600}) was read every 15 min. Growth data from five replicates from each plate were averaged, and the complete assay was repeated three times. The area under the curve (AUC) was determined for each growth curve. The replicate AUCs for strains P1 to P3 were compared to the AUCs for strains T1 to T3 using an unpaired *t* test. Comparisons of results with P1 to P3, T1 to T3, and P1 Δwca were accomplished with one-way ANOVA with Tukey's multiple-comparison test. The AUC and statistics were done using Prism, version 7 (GraphPad).

Accession number(s). The genome sequences of ECA-B and ECC-M were deposited in the GenBank under accession numbers [NHTF00000000](#) and [NHZH00000000](#), respectively. All RNA-seq files were deposited in the Short Read Archive (SRA) under BioProject accession number [PRJNA326931](#) and BioSample accession numbers [SAMN07125434](#) to [SAMN07125451](#). The genomic sequences were deposited in the SRA under BioSample numbers [SAMN07125105](#) and [SAMN07125106](#).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/IAI.00566-17>.

SUPPLEMENTAL FILE 1, XLSX file, 0.1 MB.

SUPPLEMENTAL FILE 2, PDF file, 0.1 MB.

SUPPLEMENTAL FILE 3, XLSX file, 1.7 MB.

SUPPLEMENTAL FILE 4, XLSX file, 0.3 MB.

SUPPLEMENTAL FILE 5, XLSX file, 0.1 MB.

SUPPLEMENTAL FILE 6, XLSX file, 0.1 MB.

ACKNOWLEDGMENT

We thank Duane Zimmerman, Briony Atkinson, Tera Nyholm, and Kellie Winter for their technical expertise.

Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendations or endorsement by the U.S. Department of Agriculture.

USDA is an equal opportunity provider and employer.

REFERENCES

1. USDA Animal and Plant Health Inspection Service. 2007. Highlights of Dairy 2007. Part I: reference of dairy health and management in the United States, 2007. USDA Animal and Plant Health Inspection Service, Fort Collins, CO. https://www.aphis.usda.gov/animal_health/nahms/dairy/downloads/dairy07/Dairy07_is_PartI_Highlights.pdf.
2. Hogan JS, Smith KL. 2003. Coliform mastitis. *Vet Res* 34:507–519. <https://doi.org/10.1015/vetres:2003022>.
3. Roberson JR. 2012. Treatment of clinical mastitis. *Vet Clin North Am Food Anim Pract* 28:271–288. <https://doi.org/10.1016/j.cvfa.2012.03.011>.
4. Döpfer D, Barkema HW, Lam TJ, Schukken YH, Gaastra W. 1999. Recurrent clinical mastitis caused by *Escherichia coli* in dairy cows. *J Dairy Sci* 82:80–85. [https://doi.org/10.3168/jds.S0022-0302\(99\)75211-2](https://doi.org/10.3168/jds.S0022-0302(99)75211-2).
5. Bradley AJ, Green MJ. 2001. Adaptation of *Escherichia coli* to the bovine mammary gland. *J Clin Microbiol* 39:1845–1849. <https://doi.org/10.1128/JCM.39.5.1845-1849.2001>.
6. Lam TJ, Lipman LJ, Schukken YH, Gaastra W, Brand A. 1996. Epidemiological characteristics of bovine clinical mastitis caused by *Staphylococcus aureus* and *Escherichia coli* studied by DNA fingerprinting. *Am J Vet Res* 57:39–42.
7. Dogan B, Klaessig S, Rishniw M, Almeida RA, Oliver SP, Simpson K, Schukken YH. 2006. Adherent and invasive *Escherichia coli* are associated with persistent bovine mastitis. *Vet Microbiol* 116:270–282. <https://doi.org/10.1016/j.vetmic.2006.04.023>.
8. Lippolis JD, Brunelle BW, Reinhardt TA, Sacco RE, Thacker TC, Looft TP, Casey TA. 2016. Differential gene expression of three mastitis-causing *Escherichia coli* strains grown under planktonic, swimming, and swarming culture conditions. *mSystems* 1:e00064-16. <https://doi.org/10.1128/mSystems.00064-16>.
9. Lippolis JD, Bayles DO, Reinhardt TA. 2009. Proteomic changes in *Escherichia coli* when grown in fresh milk versus laboratory media. *J Proteome Res* 8:149–158. <https://doi.org/10.1021/pr800458v>.
10. Lippolis JD, Brunelle BW, Reinhardt TA, Sacco RE, Nonnecke BJ, Dogan B, Simpson K, Schukken YH. 2014. Proteomic analysis reveals protein expression differences in *Escherichia coli* strains associated with persistent versus transient mastitis. *J Proteomics* 108:373–381. <https://doi.org/10.1016/j.jprot.2014.06.008>.
11. Huang DW, Sherman BT, Lempicki RA. 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4:44–57. <https://doi.org/10.1038/nprot.2008.211>.
12. Seo SW, Kim D, Latif H, O'Brien EJ, Szubin R, Palsson BO. 2014. Deciphering Fur transcriptional regulatory network highlights its complex role beyond iron metabolism in *Escherichia coli*. *Nat Commun* 5:4910. <https://doi.org/10.1038/ncomms5910>.
13. Richards VP, Lefébure T, Bitar PD, Dogan B, Simpson KW, Schukken YH, Stanhope MJ. 2015. Genome-based phylogeny and comparative genomic analysis of intra-mammary pathogenic *Escherichia coli*. *PLoS One* 10:e0119799. <https://doi.org/10.1371/journal.pone.0119799>.
14. Miajlovic H, Cooke NM, Moran GP, Rogers TR, Smith SG. 2014. Response of extraintestinal pathogenic *Escherichia coli* to human serum reveals a protective role for Rcs-regulated exopolysaccharide colanic acid. *Infect Immun* 82:298–305. <https://doi.org/10.1128/IAI.00800-13>.
15. Wang S, Shi H, Li Y, Shi Z, Zhang X, Baek CH, Mothershead T, Curtiss R. 2013. A colanic acid operon deletion mutation enhances induction of early antibody responses by live attenuated *Salmonella* vaccine strains. *Infect Immun* 81:3148–3162. <https://doi.org/10.1128/IAI.00097-13>.
16. Chen J, Lee SM, Mao Y. 2004. Protective effect of exopolysaccharide colanic acid of *Escherichia coli* O157:H7 to osmotic and oxidative stress. *Int J Food Microbiol* 93:281–286. <https://doi.org/10.1016/j.ijfoodmicro.2003.12.004>.
17. Navasa N, Rodríguez-Aparicio L, Martínez-Blanco H, Arcos M, Ferrero MÁ. 2009. Temperature has reciprocal effects on colanic acid and polysialic acid biosynthesis in *E. coli* K92. *Appl Microbiol Biotechnol* 82:721–729. <https://doi.org/10.1007/s00253-008-1840-4>.
18. Rainard P. 2003. The complement in milk and defense of the bovine mammary gland against infections. *Vet Res* 34:647–670. <https://doi.org/10.1051/vetres:2003025>.
19. Reinhardt TA, Sacco RE, Nonnecke BJ, Lippolis JD. 2013. Bovine milk proteome: quantitative changes in normal milk exosomes, milk fat globule membranes and whey proteomes resulting from *Staphylococcus aureus* mastitis. *J Proteomics* 82:141–154. <https://doi.org/10.1016/j.jprot.2013.02.013>.
20. Harmon RJ, Schanbacher FL, Ferguson LC, Smith KL. 1976. Changes in lactoferrin, immunoglobulin G, bovine serum albumin, and α -lactalbumin during acute experimental and natural coliform mastitis in cows. *Infect Immun* 13:533–542.
21. Bannerman DD, Paape MJ, Lee J-W, Zhao X, Hope JC, Rainard P. 2004. *Escherichia coli* and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection. *Clin Diagn Lab Immunol* 11:463–472. <https://doi.org/10.1128/CDLI.11.3.463-472.2004>.
22. Zadoks RN, Middleton JR, McDougall S, Katholm J, Schukken YH. 2011. Molecular epidemiology of mastitis pathogens of dairy cattle and comparative relevance to humans. *J Mammary Gland Biol Neoplasia* 16:357. <https://doi.org/10.1007/s10911-011-9236-y>.
23. Leimbach A, Poehlein A, Vollmers J, Görlich D, Daniel R, Dobrindt U. 2017. No evidence for a bovine mastitis *Escherichia coli* pathotype. *BMC Genomics* 18:359. <https://doi.org/10.1186/s12864-017-3739-x>.
24. Burvenich C, van Merris V, Mehrzad J, Diez-Fraile A, Duchateau L. 2003. Severity of *E. coli* mastitis is mainly determined by cow factors. *Vet Res* 34:521–564. <https://doi.org/10.1051/vetres:2003023>.
25. Suojala L, Kaartinen L, Pyörälä S. 2013. Treatment for bovine *Escherichia coli* mastitis—an evidence-based approach. *J Vet Pharmacol Ther* 36:521–531. <https://doi.org/10.1111/jvp.12057>.
26. Bearson BL, Bearson SM, Uthe JJ, Dowd SE, Houghton JO, Lee I, Toscano MJ, Lay DC, Jr. 2008. Iron regulated genes of *Salmonella enterica* serovar Typhimurium in response to norepinephrine and the requirement of *fepDGC* for norepinephrine-enhanced growth. *Microbes Infect* 10:807–816. <https://doi.org/10.1016/j.micinf.2008.04.011>.
27. Richards VP, Choi SC, Pavinski Bitar PD, Gurjar AA, Stanhope MJ. 2013. Transcriptomic and genomic evidence for *Streptococcus agalactiae* adaptation to the bovine environment. *BMC Genomics* 14:920. <https://doi.org/10.1186/1471-2164-14-920>.
28. Haas BJ, Chin M, Nusbaum C, Birren BW, Livny J. 2012. How deep is deep enough for RNA-seq profiling of bacterial transcriptomes? *BMC Genomics* 13:734. <https://doi.org/10.1186/1471-2164-13-734>.
29. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
30. Anders S, Pyl PT, Huber W. 2015. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169. <https://doi.org/10.1093/bioinformatics/btu638>.
31. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550. <https://doi.org/10.1186/s13059-014-0550-8>.
32. Hull RA, Gill RE, Hsu P, Minshew BH, Falkow S. 1981. Construction and expression of recombinant plasmids encoding type 1 or α -mannose-resistant pili from a urinary tract infection *Escherichia coli* isolate. *Infect Immun* 33:933–938.
33. Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJM, Birol I. 2009. ABySS: a parallel assembler for short read sequence data. *Genome Res* 19:1117–1123. <https://doi.org/10.1101/gr.089532.108>.
34. Gómez-Gómez J-M, Manfredi C, Alonso J-C, Blázquez J. 2007. A novel role for RecA under non-stress: promotion of swarming motility in *Escherichia coli* K-12. *BMC Biol* 5:14. <https://doi.org/10.1186/1741-7007-5-14>.
35. Wozniak CE, Chevance FFV, Hughes KT. 2010. Multiple promoters contribute to swarming and the coordination of transcription with flagellar assembly in *Salmonella*. *J Bacteriol* 192:4752–4762. <https://doi.org/10.1128/JB.00093-10>.