

Responses of lichen communities to 18 years of natural and experimental warming

Juha M. Alatalo^{1,*}, Annika K. Jägerbrand², Shengbin Chen³ and Ulf Molau⁴

¹*Department of Biological and Environmental Sciences, College of Arts and Sciences, Qatar University, PO Box 2713, Doha, Qatar*, ²*Calluna AB, Hästholmsvägen 28, 131 30 Nacka, Sweden*, ³*College of Environment and Civil Engineering, Chengdu University of Technology, Chengdu 610059, China* and ⁴*Department of Biological and Environmental Sciences, University of Gothenburg, PO Box 461, SE-405 30 Gothenburg, Sweden*

*For correspondence. E-mail jalatalo@qu.edu.qa

Received: 9 January 2017 Returned for revision: 21 February 2017 Editorial decision: 20 March 2017 Accepted: 10 April 2017

- **Background and Aims** Climate change is expected to have major impacts on high alpine and arctic ecosystems in the future, but empirical data on the impact of long-term warming on lichen diversity and richness are sparse. This study reports the effects of 18 years of ambient and experimental warming on lichens and vascular plant cover in two alpine plant communities, a dry heath with sparse canopy cover (54 %) and a mesic meadow with a more developed (67 %) canopy cover, in sub-arctic Sweden.
- **Methods** The effects of long-term passive experimental warming using open top chambers (OTCs) on lichens and total vascular plant cover, and the impact of plant cover on lichen community parameters, were analysed.
- **Key Results** Between 1993 and 2013, mean annual temperature increased about 2 °C. Both site and experimental warming had a significant effect on cover, species richness, effective number of species evenness of lichens, and total plant canopy cover. Lichen cover increased in the heath under ambient conditions, and remained more stable under experimental warming. The negative effect on species richness and effective number of species was driven by a decrease in lichens under experimental warming in the meadow. Lichen cover, species richness, effective number of species evenness were negatively correlated with plant canopy cover. There was a significant negative impact on one species and a non-significant tendency of lower abundance of the most common species in response to experimental warming.
- **Conclusions** The results from the long-term warming study imply that arctic and high alpine lichen communities are likely to be negatively affected by climate change and an increase in plant canopy cover. Both biotic and abiotic factors are thus important for future impacts of climate change on lichens.

Key words: Arctic, climate change, effective number of species, global warming, plant–climate interactions, species richness, tundra.

INTRODUCTION

Climate change is predicted to have a large impact on a wide range of ecosystems and ecosystem services (Shen and Ma, 2014; Wu *et al.*, 2014; Zhang *et al.*, 2014; Hao *et al.*, 2017). Polar and high-elevation alpine ecosystems are likely to experience rapid climate change (Chapin *et al.*, 1995; Mack *et al.*, 2004; Stocker *et al.*, 2013). Changes in species composition in alpine and arctic plant communities have already been recorded (Capers and Stone, 2011; Erschbamer *et al.*, 2011; Callaghan *et al.*, 2013), and a meta-analysis incorporating 1367 species responses provided evidence of a rapid latitudinal and elevational range shift in species across a large geographical range (Chen *et al.*, 2011). However, it is not always possible to identify the cause of range shifts, and they may sometimes be due to other changes brought about by human activities (Groom, 2013). The loss of habitat due to climate change is predicted to increase the extinction risks on mountain ranges worldwide (Colwell *et al.*, 2008; Raxworthy *et al.*, 2008; Dirnböck *et al.*, 2011; Engler *et al.*, 2011). However, as glaciers retreat, this may uncover new habitats to be colonized, e.g. a number of studies have shown that lichen diversity and abundance is correlated with

increasing time since glacier retreat (Bilovitz *et al.*, 2014a, b, 2015a). However, the ice-free glacial tills exposed sometimes require several hundred years to reach the climax stage of alpine grassland (Raffl and Erschbamer, 2004). In polar regions and high alpine areas, lichens tend to be more important in terms of cover and biomass for N₂ fixation and as a food source for herbivores as vascular plants become smaller (Longton, 1984; Heggberget *et al.*, 2002; Wirtz *et al.*, 2003; Nash, 2008; Rees *et al.*, 2008; Rai *et al.*, 2014). Thus, lichens are a very important part of high-altitude/latitude ecosystems but, despite this, the majority of climate change studies to date focus on vascular plants (Alatalo and Totland, 1997; Arft *et al.*, 1999; Dumais *et al.*, 2014; Wheeler *et al.*, 2016; Zhang and Wang, 2016). Long-term studies have shown that vascular plants have increased in abundance in response to warming (Sturm *et al.*, 2001; Myers-Smith *et al.*, 2011; Hobbie *et al.*, 2017). Most studies on lichens lack information about species-level responses, and only a few incorporate species-level data to study the impact on different species or on lichen diversity and richness (Molau and Alatalo, 1998; Klanderud and Totland, 2005;

Lang *et al.*, 2012; Alatalo *et al.*, 2014a, 2015a). However, studies on lichens are currently being initiated worldwide to study potential impacts of climate change and other anthropogenic disturbances on lichen communities (Maphangwa *et al.*, 2012; Rai *et al.*, 2012a, b; Darnajoux *et al.*, 2015; Shukla *et al.*, 2015; Upreti *et al.*, 2015; Piercey-Normore *et al.*, 2016). Existing long-term studies (9–20 years) show that lichen biomass and/or cover is sensitive to long-term warming at alpine and arctic sites (Chapin *et al.*, 1995; Van Wijk *et al.*, 2003; Elmendorf *et al.*, 2012; Lang *et al.*, 2012; Sistla *et al.*, 2013), while shorter term studies (2–7 years) report contrasting results (Press *et al.*, 1998; Klanderud and Totland, 2005; Biasi *et al.*, 2008; Alatalo *et al.*, 2014a, 2015a). Lichen diversity has also been shown to decrease due to warming in arctic Alaska (16 years of warming) and sub-arctic Sweden (9 years of warming) (Lang *et al.*, 2012). Furthermore, the response may be context dependent, depending on potential competition with vascular plants (Alatalo, 1998; Cornelissen *et al.*, 2001). Modelling studies on the potential impact of climate change on lichens suggest that many lichen species are potentially threatened (Allen and Lendemer, 2016; Nascimbene *et al.*, 2016; Rubio-Salcedo *et al.*, 2017).

Here we examined lichen communities following 18 years of experimental warming in two contrasting alpine sub-arctic plant communities (mesic meadow and dry, poor heath) in Sweden. The hypotheses tested were that (1) lichen cover and diversity are negatively affected (decreasing) by long-term warming; (2) the negative impacts of warming are greater for a meadow community with a more developed vascular plant community (67 % canopy cover) than a poor heath with a less developed vascular plant community (54 % canopy cover); and (3) more species are lost than gained owing to long-term warming.

MATERIALS AND METHODS

Study area

Fieldwork took place at Latnjajaure field station, which is located in the Latnjavagge valley (68°21'N, 18°29'E; 1000 m above sea level) in northern Sweden. Climate parameters were measured daily from early spring 1992 onwards (by data loggers that collect hourly data on temperature, and by a manned climate station during summers). The climate at the site is classified as sub-arctic (Polunin, 1951), with snow cover for most of the year, cool summers and relatively mild, snow-rich winters. The growing season starts in late May and ends in early September (Molau *et al.*, 2005). Mean annual air temperature ranged from -0.76 to -2.92 °C between 1993 and 2013 (Fig. 1; Supplementary Data Fig. S1). The mean temperature was highest in July, with mean temperature ranging from 5.9 °C in 1995 to 13.1 °C in 2013 (Fig. S1). Mean annual precipitation during that period was 846 mm, but in individual years it ranged from a low of 607 mm (1996) to a high of 1091 mm (2003) (Fig. S1). Climate data are collected throughout the year at the weather station at Latnjajaure field station, with hourly means, maxima and minima. Physical conditions in the soils in the valley vary from dry to wet and poor and acidic to base rich, with an associated variation in plant communities (Molau and Alatalo, 1998; Lindblad *et al.*, 2006; Björk *et al.*, 2007; Alatalo *et al.*, 2014b). The mesic meadow community is dominated by *Carex vaginata*, *C. bigelowii*, *Festuca ovina*, *Salix reticulata*, *S. polaris*,

Cassiope tetragona, *Polygonum viviparum* and *Thalictrum alpinum* (Molau and Alatalo, 1998; Alatalo *et al.*, 2014b). The more sparsely vegetated poor heath community is dominated by *Betula nana*, *S. herbacea* and *Calamagrostis lapponica* (Molau and Alatalo, 1998; Alatalo *et al.*, 2015b).

Experimental design and measurements

In July 1995, 1 × 1 m plots with homogeneous vegetation cover were marked out in an alpine mesic meadow plant community and in a heath plant community, and randomly assigned to treatments (control and warming) in a factorial design. At the start of the experiment there were eight control (CTR) plots and four plots with warming in each plant community (a total of 12 in each plant community). However, as we could not identify all initial control plots in 2013, in that year we only made measurements in four control and four warming plots in each community. Warming was applied by open top chambers (OTCs), and we monitored the temperature in control and plots with OTCs in the initial years with Delta™ and Tinytag™ loggers (Molau and Alatalo, 1998). As found in other studies (Marion *et al.*, 1997; Molau and Alatalo, 1998; Hollister and Webber, 2000), OTCs increased the air temperature by 1.5–3 °C compared with control plots with ambient temperature. OTCs have also been shown to decrease canopy moisture (Hollister and Webber, 2000), causing earlier snow melt and prolonging the growing season (Molau and Alatalo, 1998; Hollister and Webber, 2000). The OTCs were then left on plots with warming treatment all year around. The majority of lichens in the plots were identified to species level. When necessary, we collected a specimen of the same species outside the experimental plots to be determined in the laboratory. In the case of *Cladonia*, when we were not able to determine the specimen to species level, we labelled it *Cladonia* spp. Coverage of each species was assessed by point-intercept using a 1 × 1 m frame with 100 grid points (Walker, 1996) in the peak of the 1995, 1999, 2001 and 2013 growing seasons. Due to their hexagonal shape, the OTCs reduced the number of points per plot to 77–87, and thus warmed plots had fewer pin-point intercepts than control plots.

Statistical analyses

The following community parameters were calculated for comparison of the lichen assemblages: cover, species richness, Shannon's evenness and effective number of species (expH = exponential of Shannon entropy), which is the number of equally abundant species needed for the average proportional abundance of the species to equal that observed in the data set (where all species may not be equally abundant) (Jost, 2006). For vascular plants, we calculated total canopy cover. All data were checked for normality assumptions and homogeneity of variance by the Kolmogorov–Smirnov test and Levene's test of equality of error variances, respectively. We then applied a univariate analysis of variance (ANOVA) with treatment (control or warming) and vegetation type (meadow or heath) as fixed factors, and the ratio of the value in 2013 to the value in 1995 (relative change of the community parameters mentioned above) on each plot as the response variable. The use of the




Fig. 1. Mean annual temperature (°C), 1993–2013, at Latnjaure, Northern Sweden. Solid line, mean annual temperature; dotted line, trend line.

ratio as the response variable was due to the fact that the number of hits per plot differed between treatments, and that plots differed in their starting values of cover, richness and species composition. Thus we opted to analyse relative changes between 1995 and 2013 instead of actual numbers. The ANOVA included the initial value in 1995 of each variable as covariate, which may potentially have affected the response variable. To check for the effect of time on cover, species richness, effective number of species and Shannon's evenness, we applied a repeated measurement ANOVA on relative changes between 1995 and 1999, 1995 and 2001, and 1995 and 2013. As the species-level data did not meet the assumption of normal distribution after transformation, we used the conservative non-parametric Mann–Whitney U-test to analyse the effect of treatment on the relative change between 1995 and 2013 of the most common species. Only species with >100 pin-point intercept hits in total from the years 1995, 1999, 2001 and 2013 were included (Alatalo *et al.*, 2015a). Two-tailed Pearson correlation was used to analyse correlations between lichen community parameters (total lichen cover, species richness, Shannon's evenness and effective number of species) and vascular plant canopy cover. All analyses were performed in IBM® SPSS® Statistics Version 23.0.0.2.

RESULTS

Initial lichen cover in the heath and meadow control plots was 35.75 % (s.d. ± 5.44) and 11.75 % (± 5.56), respectively. There was a significant effect of time on lichen total cover ($P < 0.001$), species richness ($P = 0.009$) and effective number of species ($P = 0.004$), but not on Shannon's evenness ($P = 0.3160$). Cover of lichens increased by 87 % over the 18 year study period in the heath control plots. Over the same period, lichen cover in control plots in the meadow increased by 21 %. Eighteen years of experimental warming and site both had significant effects on cover, species richness, effective number of species and evenness of lichens (Fig. 2; Table 1). Lichen cover decreased significantly in response to the long-term warming in both heath and meadow communities ($P < 0.001$); there was an 8 % decline in lichen cover in the heath and a

31 % decline in the meadow compared with the starting year. A total of 22 lichen species were recorded in the study plots during the course of the study, with 19 in 1995 and 16 in 2013 (three new species recorded and six species lost in 2013) (Table 2). Lichen richness and, effective number of species, and evenness declined significantly ($P < 0.001$, $P < 0.014$, $P < 0.035$, respectively) in response to long-term warming, an effect mainly driven by a decline in the meadow, while species richness and effective number of species both remained more stable in the heath community (Fig. 2). Total lichen cover (-0.542 , $P < 0.001$), species richness (-0.370 , $P < 0.001$), effective number of species (-0.350 , $P = 0.001$), but not Shannon's evenness (0.070 , $P = 0.545$), were negatively correlated with vascular plant canopy cover (Fig. 3). Eighteen years of experimental warming and site both had significant effects on plant canopy cover (Table 3); warming having a positive impact on plant cover, with the largest increase found in the heath (Fig. 4).

Of the eight most common lichen species, there was a significant negative effect of long-term warming on cover of one species, *Flavocetraria cucullata* ($P = 0.029$), in the heath, but not in the meadow ($P = 1$). For the other seven species, *Cetrariella delisei* ($P = 0.114$, $P = 1$), *Flavocetraria nivalis* ($P = 0.114$, $P = 0.686$), *Cladina arbuscula* ($P = 0.343$, $P = 0.486$), *Cladonia unicalis* ($P = 0.114$, $P = 0.114$), *Ochrolechia frigida* ($P = 1$, $P = 1$), *Sphaerophorus globosus* ($P = 0.20$, $P = 0.114$) and *Stereocaulon alpinum* ($P = 0.114$, $P = 1$), there were no significant effects on cover in either the heath or the meadow (Fig. 5).

DISCUSSION

The results from this long-term experiment confirm predictions that lichens may be less vulnerable in plant communities with less developed plant cover than in communities with more developed plant cover (Alatalo, 1998; Cornelissen *et al.*, 2001). As hypothesized, lichens were more vulnerable in the meadow with its more developed plant canopy cover. We found that all lichen community parameters (total lichen cover, species richness, effective number of species and Shannon's evenness) were negatively correlated with plant canopy cover. Similar to




FIG. 2. Boxplots of changes in lichens during an 18 year period (1995–2013) of experimental warming in an alpine heath and an alpine meadow community at Latnjajaure, sub-arctic Sweden. Total cover of lichens (percentage), species richness of lichens, effective number of lichen species (exponential of Shannon's entropy) and Shannon evenness. Boxplots show the 10th–90th percentile of the data. Treatments: CTR, control; T, temperature warming. Number of plots: $n = 4$ for CTR and T at each site. Asterisks (*) indicate significant differences in relative changes between treatments.

TABLE 1. Result of univariate ANOVAs testing the effects of treatment (18 years of experimental warming) and site (alpine meadow and alpine heath) on relative change in: cover, species richness, effective number of species and evenness for lichens

Source	Type III sum of squares	df.	Mean square	F	P-value
Relative change in cover					
Initial cover	1.900	1	1.900	9.815	0.010
Treatment	4.050	1	4.050	20.926	<0.001
Site	2.652	1	2.652	13.703	0.003
Treatment × Site	0.269	1	0.269	1.388	0.264
Total	29.178	16			
$R^2 = 0.704$ (Adjusted $R^2 = 0.596$)					
Relative change in species richness					
Initial richness	0.885	1	0.885	8.919	0.012
Treatment	2.142	1	2.142	21.580	<0.001
Site	1.088	1	1.088	10.963	0.007
Treatment × Site	0.259	1	0.259	2.613	0.134
Total	17.714	16			
$R^2 = 0.716$ (Adjusted $R^2 = 0.613$)					
Relative change in effective number of species					
Initial effective no. sp.	0.839	1	0.839	5.737	0.036
Treatment	1.256	1	1.256	8.594	0.014
Site	1.196	1	1.196	8.180	0.016
Treatment × Site	0.206	1	0.206	1.411	0.260
Total	17.035	16			
$R^2 = 0.579$ (Adjusted $R^2 = 0.425$)					
Relative change in evenness					
Initial evenness	0.004	1	0.004	0.060	0.811
Treatment	0.397	1	0.397	5.750	0.035
Site	0.677	1	0.6777	9.807	0.010
Treatment × Site	0.660	1	0.660	9.566	0.010
Total	12.352	16			
$R^2 = 0.700$ (Adjusted $R^2 = 0.591$)					

D.f., degrees of freedom; F, F-statistics; P-value, significance level.

TABLE 2. List of lichens recorded in the heath (H) and meadow (M) alpine vegetation communities at Latnjajaure, sub-arctic Sweden

Lichen species	1995	1999	2001	2013
<i>Alectoria nigricans</i>	H	H	H	Not recorded
<i>Alectoria ochroleuca</i>	H	H	H	H
<i>Cetrariella delisei</i>	H, M	H, M	H, M	H, M
<i>Cladina arbuscula</i>	H, M	H, M	H, M	H, M
<i>Cladina rangiferina</i>	H	Not recorded	Not recorded	Not recorded
<i>Cladonia</i> spp.	H, M	H, M	H, M	H
<i>Cladonia furcata</i>	H	H, M	H	Not recorded
<i>Cladonia gracilis</i>	Not recorded	H	H	H, M
<i>Cladonia pyxidata</i>	Not recorded	Not recorded	Not recorded	H
<i>Cladonia uncialis</i>	H, M	H, M	H, M	H, M
<i>Cornicularia divergens</i>	H	H	H	Not recorded
<i>Flavocetraria cucullata</i>	H, M	H, M	H, M	H, M
<i>Flavocetraria nivalis</i>	H, M	H, M	H, M	H, M
<i>Nephroma arctica</i>	H, M	H, M	H, M	H
<i>Ochrolechia frigida</i>	H, M	H, M	M	Not recorded
<i>Peltigera aphthosa</i>	H, M	H, M	H, M	H, M
<i>Peltigera scabrosa</i>	H	H	H	H, M
<i>Pertusaria dactylina</i>	Not recorded	Not recorded	Not recorded	H
<i>Solorina crocea</i>	H	H	H	Not recorded
<i>Sphaerophorus globosus</i>	H, M	H, M	H, M	H, M
<i>Stereocaulon alpinum</i>	H	H	H	H
<i>Thamnomia vermicularis</i>	H, M	H, M	H, M	H, M

Species in bold are the eight most common lichen species found.

other studies (Sturm *et al.*, 2001; Myers-Smith *et al.*, 2011; Harte *et al.*, 2015; Hobbie *et al.*, 2017), we found that long-term warming caused a significant increase in total plant canopy cover, with canopy expanding most in the heath. As both lichens (this study) and bryophytes (Jägerbrand *et al.*, 2012) have been shown to be negatively correlated with vascular plant canopy, a future increase in vascular plant canopy is therefore likely to have a detrimental effect on these groups. The differences in canopy cover of vascular plant communities may thus help to explain the significant site effect on all response variables. The negative impact on diversity is in line with previous long-term studies in Alaska showing a decrease in lichen diversity under 16 years of warming (Lang *et al.*, 2012). This highlights the importance of long-term studies, as shorter term studies have tended to find contrasting effects on lichens (Klanderud and Totland, 2005; Alatalo *et al.*, 2014a), while longer term studies have typically found negative effects (Chapin *et al.*, 1995; Van Wijk *et al.*, 2003; Elmendorf *et al.*, 2012; Lang *et al.*, 2012; Sistla *et al.*, 2013).

The responses to long-term warming differed between the two plant communities studied, with the meadow experiencing a larger decline in lichen cover relative to the initial year of the study than the heath community (Fig. 2). A similar pattern was found for richness and effective number of species, which declined in the meadow but not in the heath. This shows that both abiotic and biotic interactions may play important roles in lichen responses and reveals the necessity of including different




FIG. 3. Relationship between lichens and vascular plant canopy cover at Latnjajaure, sub-arctic Sweden: total cover of lichens, species richness of lichens, effective number of lichen species (exponential of Shannon's entropy) and Shannon evenness. Number of plots: $n = 88$.

TABLE 3. Result of univariate ANOVAs testing the effects of treatment (18 years of experimental warming) and site (alpine meadow and alpine heath) on relative change in vascular plant canopy cover

Source	Type III sum of squares	d.f.	Mean square	F	P-value
Relative change in cover					
Initial cover	2.247	1	2.247	54.156	<0.001
Treatment	0.698	1	0.698	16.831	0.002
Site	0.194	1	0.194	4.664	0.054
Treatment × Site	0.456	1	1.018	24.542	<0.001
Total	36.623	16			
$R^2 = 0.906$ (adjusted $R^2 = 0.872$)					

plant communities in experimental studies. In our study, of the eight most common species, we only found a significant negative impact on one species, *F. cucullata*, in the heath, but not in the meadow. However, while there were no significant effects on the other species, there was a general tendency for lower abundance of the most common species in the plots experiencing long-term warming. Thus, it is likely that the increase in

evenness observed in our long-term study may have been caused by an overall decline in dominant lichen species over the study period.

The long-term data from our control plots also indicate that bare ground in high alpine areas that has been ice free for a long period (since the retreat of glaciers) can continue to be colonized by lichens, similarly to the plant progression that occurs when glaciers retreat (Raffl and Erschbamer, 2004; Bilovitz et al., 2015a, 2015b).

In fact, Latnjajaure experienced a natural increase in mean annual temperature of about 2 °C between 1995 and 2013. Thus, the lichens in control plots were also exposed to natural climate warming, which may have contributed to the positive effect on lichen cover seen in control plots. It is likely that the main driver for the significant difference between treatments was the increase in lichen cover in the control plots, not the decrease in lichen cover in long-term warming plots. Thus, while the lichens within the warming plots declined less over the study period in the heath compared with the meadow community, the difference compared with the control plots was larger in the heath than in the meadow community. Another potential




FIG. 4. Boxplots of changes in vascular plant canopy cover during an 18 year period (1995–2013) of experimental warming in an alpine heath and an alpine meadow community at Latnjajaure, sub-arctic Sweden. Boxplots show the 10th–90th percentile of the data. Treatments: CTR, control; T, temperature warming. Number of plots: $n = 4$ for CTR and T at each site.

explanation for the large difference in increase in cover may be differences in initial levels of bare ground and vascular plant cover between the plant communities, with the meadow having less bare ground and a more developed initial plant cover (Molau and Alatalo, 1998). Thus, the heath both offered more bare ground to be colonized by lichens and experienced less competition from vascular plants. The fact that lichen cover, species richness, and effective number of species were all negatively correlated to plant canopy cover indicates that lichens were most probably outcompeted, or overgrown, by vascular plants.

It should be noted that a constant level of warming is not the most realistic scenario for future climate change, which will most probably increase both the variability and magnitude of climate events (Stocker *et al.*, 2013). For example, increased precipitation during winter could potentially increase snow accumulation, and this in turn may reduce survival and growth of dominant arctic–alpine lichens (Bidussi *et al.*, 2016). However, if an increase in precipitation during winter is accompanied by an earlier onset of spring due to warmer climate, this may be counterbalanced. While we did not measure precipitation or snow accumulation in our study, the experimental long-term warming caused a loss of soil carbon, nitrogen, C/N ratio and soil moisture in the mineral soil layer of the meadow (Alatalo *et al.*, 2017). However, the warming did not have any effect on the soil parameters in the thin organic soil layers in the meadow or heath (Alatalo *et al.*, 2017). As lichens do not have a well-developed root system, drying of deeper soil layers is therefore unlikely to have had a direct effect on lichens. However, OTCs have been shown to decrease moisture in the canopy layer (Hollister and Webber, 2000), which may potentially have had a negative effect on lichens that depend on moisture levels for their photosynthesis and growth. There are very few experimental studies that examine different warming scenarios across years (Jonasson *et al.*, 1999; Alatalo *et al.*, 2014a, 2016), and none is long term. The one existing short-term study (3 years)

applying different climate change scenarios to lichens in an alpine meadow found the lichens to be highly resistant to a constant level of warming, a stepwise increase in warming and a single season of high-level pulse warming (Alatalo *et al.*, 2014a). However, as the short-term and longer term responses of lichens have been shown to differ (Alatalo *et al.*, 2015a), there is a need to initiate long-term experiments that incorporate different warming and precipitation scenarios.

Overall, 18 years of experimental warming brought a significant decline in lichen cover, species richness and, effective number of species, and evenness. The results showed that lichen responses are most probably dependent on both biotic and abiotic interactions and that responses differ among communities. Specifically, lichens increased more under ambient conditions in heath with a less developed plant community and decreased more under experimental warming in meadow with a more developed plant community. Species richness and effective number of species remained stable in the heath, but decreased under experimental warming in the meadow. Lichen cover, species richness, effective number of species and evenness were all negatively correlated with plant canopy cover.

SUPPLEMENTARY DATA

Supplementary data are available online at <https://academic.oup.com/aob> and consist of Figure S1: mean, minimum and maximum monthly temperatures ($^{\circ}\text{C}$), and monthly precipitation (mm) January 1993–December 2013, at Latnjajaure, Northern Sweden.

ACKNOWLEDGEMENTS

The authors thank the staff of Abisko Scientific Research Station for their help and hospitality, and Matthias Molau for assistance in the field. This study was supported by Carl




FIG. 5. Boxplots of species-specific responses of lichen cover (percentage) during an 18 year period (1995–2013) of experimental warming in a poor alpine heath and a rich alpine meadow community at Latnjajaure, sub-arctic Sweden. Boxplots show the 10th–90th percentile of the data. Treatments: CTR, control, T, temperature warming. Number of plots: $n = 4$ for CTR and T at each site.




FIG. 5. Continued




FIG. 5. Continued

Tryggers stiftelse för vetenskaplig forskning through a grant to J.M.A. The authors thank J. S. Heslop-Harrison, Anne Brysting and two anonymous reviewers whose comments improved the manuscript.

AUTHOR CONTRIBUTIONS

J.M.A and U.M. designed the experiment, J.M.A. and U.M. carried out fieldwork. J.M.A., A.K.J. and S.C. carried out data analyses, J.M.A. and A.K.J. made the figures, and J.M.A. made the tables. J.M.A. drafted the manuscript. All authors read, commented on and approved the final manuscript. The authors declare no competing financial interests.

LITERATURE CITED

- Alatalo J. 1998. *Climate change: impacts on structure and biodiversity of sub-arctic plant communities*. PhD Thesis, Göteborg University, Sweden.
- Alatalo JM, Totland Ø. 1997. Response to simulated climatic change in an alpine and subarctic pollen-risk strategist, *Silene acaulis*. *Global Change Biology* 3: 74–79.
- Alatalo JM, Jägerbrand AK, Molau U. 2014a. Climate change and climatic events: community-, functional- and species-level responses of bryophytes and lichens to constant, stepwise, and pulse experimental warming in an alpine tundra. *Alpine Botany* 124: 81–91.
- Alatalo JM, Little CJ, Jägerbrand AK, Molau U. 2014b. Dominance hierarchies, diversity and species richness of vascular plants in an alpine meadow: contrasting short and medium term responses to simulated global change. *PeerJ* 2: e406. doi: 10.7717/peerj.406.
- Alatalo JM, Jägerbrand AK, Molau U. 2015a. Testing reliability of short-term responses to predict longer-term responses of bryophytes and lichens to environmental change. *Ecological Indicators* 58: 77–85.
- Alatalo JM, Little CJ, Jägerbrand AK, Molau U. 2015b. Vascular plant abundance and diversity in an alpine heath under observed and simulated global change. *Scientific Reports* 5: 10197. doi: 10.1038/srep10197.
- Alatalo JM, Jägerbrand AK, Molau U. 2016. Impacts of different climate change regimes and extreme climatic events on an alpine meadow community. *Scientific Reports* 6: 21720. doi: 10.1038/srep21720.
- Alatalo JM, Jägerbrand AK, Juhanson J, Michelsen A, Luptáček P. 2017. Impacts of twenty years of experimental warming on soil carbon, nitrogen, moisture and soil mites across alpine/subarctic tundra communities. *Scientific Reports* 7: 44489. doi: 10.1038/srep44489.
- Allen JL, Lendemer JC. 2016. Climate change impacts on endemic, high-elevation lichens in a biodiversity hotspot. *Biodiversity and Conservation* 25: 555–568.
- Arft AM, Walker MDM, Gurevitch J, et al. 1999. Responses of tundra plants to experimental warming: meta-analysis of the international tundra experiment. *Ecological Monographs* 69: 491–511.
- Biasi C, Meyer H, Rusalimova O, et al. 2008. Initial effects of experimental warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant and Soil* 307: 191–205.
- Bidussi M, Solhaug KA, Gauslaa Y. 2016. Increased snow accumulation reduces survival and growth in dominant mat-forming arctic-alpine lichens. *The Lichenologist* 48: 237–247.
- Bilovitz PO, Nascimbene J, Tutzer V, et al. 2014a. Terricolous lichens in the glacier forefield of the Rötkees (Eastern Alps, South Tyrol, Italy). *Phyton; annales rei botanicae* 54: 245–250.

- Bilovitz PO, Wallner A, Tutzer V, et al. 2014b.** Terricolous lichens in the glacier forefield of the Gaisbergferner (Eastern Alps, Tyrol, Austria). *Phyton; annales rei botanicae* **54**: 235–243.
- Bilovitz PO, Nascimbene J, Mayrhofer H. 2015a.** Terricolous lichens in the glacier forefield of the Morteratsch Glacier (Eastern Alps, Graubünden, Switzerland). *Phyton; annales rei botanicae* **55**: 193–199.
- Bilovitz PO, Wallner A, Tutzer V, Nascimbene J, Mayrhofer H. 2015b.** Terricolous lichens in the glacier forefield of the Pasterze (Eastern Alps, Carinthia, Austria). *Phyton; annales rei botanicae* **55**: 201–214.
- Björk RG, Klemedtsson L, Molau U, Harndorf J, Ödman A, Giesler R. 2007.** Linkages between N turnover and plant community structure in a tundra landscape. *Plant and Soil* **294**: 247–261.
- Callaghan T, Jonasson C, Thierfelder T, et al. 2013.** Ecosystem change and stability over multiple decades in the Swedish subarctic: complex processes and multiple drivers. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368**: 20120488. doi: 10.1098/rstb.2012.0488.
- Capers R, Stone A. 2011.** After 33 years, trees more frequent and shrubs more abundant in northeast US alpine community. *Arctic, Antarctic, and Alpine Research* **43**: 495–502.
- Chapin F III, Shaver G, Giblin A, Nadelhoffer K, Laundre J. 1995.** Responses of arctic tundra to experimental and observed changes in climate. *Ecology* **76**: 694–711.
- Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD. 2011.** Rapid range shifts of species associated with high levels of climate warming. *Science* **333**: 1024–1026.
- Colwell RK, Brehm G, Cardelús CL, Gilman AC, Longino JT. 2008.** Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science* **322**: 258–261.
- Cornelissen JHC, Callaghan TV, Alatalo JM, et al. 2001.** Global change and arctic ecosystems: is lichen decline a function of increases in vascular plant biomass? *Journal of Ecology* **89**: 984–994.
- Darnajoux R, Lutzoni F, Miadlikowska J, Bellenger J-P. 2015.** Determination of elemental baseline using peltiligeralean lichens from Northeastern Canada (Québec): Initial data collection for long term monitoring of the impact of global climate change on boreal and subarctic area in Canada. *Science of the Total Environment* **533**: 1–7.
- Dirnböck T, Essl F, Rabitsch W. 2011.** Disproportional risk for habitat loss of high-altitude endemic species under climate change. *Global Change Biology* **17**: 990–996.
- Dumais C, Ropars P, Denis M-P, Dufour-Tremblay G, Boudreau S. 2014.** Are low altitude alpine tundra ecosystems under threat? A case study from the Parc National de la Gaspésie, Québec. *Environmental Research Letters* **9**: 94001.
- Elmendorf S, Henry G, Hollister R, et al. 2012.** Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time. *Ecology Letters* **15**: 164–175.
- Engler R, Randin CF, Thuiller W, et al. 2011.** 21st century climate change threatens mountain flora unequally across Europe. *Global Change Biology* **17**: 2330–2341.
- Erschbamer B, Unterluggauer P, Winkler E, Mallaun M. 2011.** Changes in plant species diversity revealed by long-term monitoring on mountain summits in the Dolomites (northern Italy). *Preslia* **83**: 387–401.
- Groom QJ. 2013.** Some poleward movement of British native vascular plants is occurring, but the fingerprint of climate change is not evident. *PeerJ* **1**: e77. doi: 10.7717/peerj.77.
- Hao R, Yu D, Liu Y, et al. 2017.** Impacts of changes in climate and landscape pattern on ecosystem services. *Science of the Total Environment* **579**: 718–728.
- Harte J, Saleska SR, Levy C. 2015.** Convergent ecosystem responses to 23-year ambient and manipulated warming link advancing snowmelt and shrub encroachment to transient and long-term climate–soil carbon feedback. *Global Change Biology* **21**: 2349–2356.
- Hegberger TM, Gaare E, Ball JP. 2002.** Reindeer (*Rangifer tarandus*) and climate change: importance of winter forage. *Rangifer* **22**: 13–31.
- Hobbie JE, Shaver GR, Rastetter EB, et al. 2017.** Ecosystem responses to climate change at a Low Arctic and a High Arctic long-term research site. *Ambio* **46**: 160–173.
- Hollister RD, Webber PJ. 2000.** Biotic validation of small open-top chambers in a tundra ecosystem. *Global Change Biology* **6**: 835–842.
- Jägerbrand AK, Kudo G, Alatalo JM, Molau U. 2012.** Effects of neighboring vascular plants on the abundance of bryophytes in different vegetation types. *Polar Science* **6**: 200–208.
- Jonasson S, Michelsen A, Schmidt I, Nielsen E. 1999.** Responses in microbes and plants to changed temperature, nutrient, and light regimes in the Arctic. *Ecology* **80**: 1828–1843.
- Jost L. 2006.** Entropy and diversity. *Oikos* **113**: 363–375.
- Klanderud K, Totland Ø. 2005.** Simulated climate change altered dominance hierarchies and diversity of an alpine biodiversity hotspot. *Ecology* **86**: 2047–2054.
- Lang SI, Cornelissen JHC, Shaver GR, et al. 2012.** Arctic warming on two continents has consistent negative effects on lichen diversity and mixed effects on bryophyte diversity. *Global Change Biology* **18**: 1096–1107.
- Lindblad KEM, Nyberg R, Molau U. 2006.** Generalization of heterogeneous alpine vegetation in air photo-based image classification, Latnjajaure catchment, northern Sweden. *Pirineos* **161**: 74–79.
- Longton R. 1984.** The role of bryophytes in terrestrial ecosystems. *Journal of the Hattori Botanical Laboratory* **55**: 147–163.
- Mack MC, Schuur EAG, Bret-Harte MS, Shaver GR, Chapin FS. 2004.** Ecosystem carbon storage in arctic tundra reduced by long-term nutrient fertilization. *Nature* **431**: 440–443.
- Maphangwa KW, Musil CF, Raitt L, Zedda L. 2012.** Experimental climate warming decreases photosynthetic efficiency of lichens in an arid South African ecosystem. *Oecologia* **169**: 257–268.
- Marion G, Henry GHR, Freckman DW, et al. 1997.** Open-top designs for manipulating field temperature in high-latitude ecosystems. *Global Change Biology* **3**: 20–32.
- Molau U, Alatalo JM. 1998.** Responses of subarctic–alpine plant communities to simulated environmental change: biodiversity of bryophytes, lichens, and vascular plants. *Ambio* **27**: 322–329.
- Molau U, Nordenhäll U, Eriksen B. 2005.** Onset of flowering and climate variability in an alpine landscape: a 10-year study from Swedish Lapland. *American Journal of Botany* **92**: 422–31.
- Myers-Smith IH, Forbes BC, Wilmsing M, et al. 2011.** Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environmental Research Letters* **6**: 45509. doi: 10.1088/1748–9326/6/4/045509.
- Nascimbene J, Casazza G, Benesperi R, et al. 2016.** Climate change fosters the decline of epiphytic *Lobaria* species in Italy. *Biological Conservation* **201**: 377–384.
- Nash TH III. 2008.** Nitrogen, its metabolism and potential contribution to ecosystems. In: *Lichen biology*. Cambridge University Press: Cambridge, 216–233.
- Piercey-Normore MD, Brodo IM, Deduke C. 2016.** Lichens on the Hudson Bay Lowlands: a long-term survey in Wapusk National Park, Manitoba. *The Lichenologist* **48**: 581–592.
- Polunin N. 1951.** The real arctic: suggestions for its delimitation, subdivision, and characterization. *Journal of Ecology* **39**: 308–315.
- Press M, Potter J, Burke M, Callaghan T, Lee J. 1998.** Responses of a subarctic dwarf shrub heath community to simulated environmental change. *Journal of Ecology* **86**: 315–327.
- Raffl C, Erschbamer B. 2004.** Comparative vegetation analyses of two transects crossing a characteristic glacier valley in the Central Alps. *Phytocoenologia* **34**: 225–240.
- Rai H, Khare R, Gupta RK, Upreti DK. 2012a.** Terricolous lichens as indicator of anthropogenic disturbances in a high altitude grassland in Garhwal (Western Himalaya), India. *Botanica Orientalis: Journal of Plant Science* **8**: 16–23.
- Rai H, Upreti DK, Gupta RK. 2012b.** Diversity and distribution of terricolous lichens as indicator of habitat heterogeneity and grazing induced trampling in a temperate-alpine shrub and meadow. *Biodiversity and Conservation* **21**: 97–113.
- Rai H, Khare R, Upreti DK. 2014.** Lichenological studies in India with reference to terricolous lichens. In: Rai H, Upreti FK, eds. *Terricolous lichens in India*. Berlin: Springer; 1–20.
- Raxworthy CJ, Pearson RG, Rabibisoa N, et al. 2008.** Extinction vulnerability of tropical montane endemism from warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar. *Global Change Biology* **14**: 1703–1720.
- Rees WG, Stammler FM, Danks FS, Vitebsky P. 2008.** Vulnerability of European reindeer husbandry to global change. *Climatic Change* **87**: 199–217.
- Rubio-Salcedo M, Psomas A, Prieto M, Zimmermann NE, Martínez I. 2017.** Case study of the implications of climate change for lichen diversity and distributions. *Biodiversity and Conservation* **26**: 1121. doi:10.1007/s10531-016-1289-1.

- Shen Z, Ma K. 2014. Effects of climate change on biodiversity. *Chinese Science Bulletin* **59**: 4637–4638.
- Shukla P, Bajpai R, Singh CP, Sharma N, Upreti DK. 2015. Lichen diversity in alpine regions of eastern Sikkim with respect to long term monitoring programme of Indian Space Research Organization. *Geophytology* **45**: 57–62.
- Sistla SA, Moore JC, Simpson RT, Gough L, Shaver GR, Schimel JP. 2013. Long-term warming restructures Arctic tundra without changing net soil carbon storage. *Nature* **497**: 615–618.
- Stocker TF, Qin D, Plattner GK. 2013. *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge.
- Sturm M, Racine C, Tape K. 2001. Climate change: increasing shrub abundance in the Arctic. *Nature* **411**: 546–547.
- Upreti DK, Bajpai R, Nayaka S. 2015. Lichenology: current research in India. In: Bahadur B, Rajam MV, Shijram L, Krishnamurthy KV, eds. *Plant biology and biotechnology*. Berlin: Springer; 263–280.
- Van Wijk MT, Clemmensen KE, Shaver GR, et al. 2003. Long-term ecosystem level experiments at Toolik Lake, Alaska, and at Abisko, Northern Sweden: generalizations and differences in ecosystem and plant type responses to global change. *Global Change Biology* **10**: 105–123.
- Walker MD. 1996. Community baseline measurements for ITEX studies. In: Molau U, Mielgaard P, eds. *ITEX Manual*, 2nd edn. Copenhagen, Denmark: Danish Polar Centre, 39–41.
- Wheeler JA, Cortés AJ, Sedlacek J, et al. 2016. The snow and the willows: earlier spring snowmelt reduces performance in the low-lying alpine shrub *Salix herbacea*. *Journal of Ecology* **104**: 1041–1050.
- Wirtz N, Lumbsch HT, Green TG, et al. 2003. Lichen fungi have low cyanobiont selectivity in maritime Antarctica. *New Phytologist* **160**: 177–183.
- Wu X, Lin X, Zhang Y, Gao J, Guo L, Li J. 2014. Impacts of climate change on ecosystem in Priority Areas of Biodiversity Conservation in China. *Chinese Science Bulletin* **59**: 4668–4680.
- Zhang Y, Wang W. 2016. Interactions between warming and soil moisture increase overlap in reproductive phenology among species in an alpine meadow. *Biology Letters* **12**: 20150749. doi: 10.1098/rsbl.2015.0749.
- Zhang Y, Wang Y, Zhang M, Ma K. 2014. Climate change threats to protected plants of China: an evaluation based on species distribution modeling. *Chinese Science Bulletin* **59**: 4652–4659.