



OPEN ACCESS

Correlates of self-reported exposure to advertising of tobacco products and electronic cigarettes across 28 European Union member states

Filippos T Filippidis,^{1,2} Anthony A Laverty,² Esteve Fernandez,^{3,4,5} Ute Mons,⁶ Olena Tigova,^{3,4} Constantine I Vardavas^{7,8}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2016-053479>).

For numbered affiliations see end of article.

Correspondence to

Dr Filippos T Filippidis, Department of Primary Care and Public Health, School of Public Health, Imperial College London, 310 Reynolds Building, St. Dunstan's Road, London W6 8RP, UK; f.filippidis@imperial.ac.uk

Received 12 October 2016

Revised 19 April 2017

Accepted 21 April 2017

Published Online First

12 June 2017

ABSTRACT

Background Despite advertising bans in most European Union (EU) member states, outlets for promotion of tobacco products and especially e-cigarettes still exist. This study aimed to assess the correlates of self-reported exposure to tobacco products and e-cigarette advertising in the EU.

Methods We analysed data from wave 82.4 of the Eurobarometer survey (November–December 2014), collected through interviews in 28 EU member states (n=27 801 aged ≥15 years) and data on bans of tobacco advertising extracted from the Tobacco Control Scale (TCS, 2013). We used multilevel logistic regression to assess sociodemographic correlates of self-reported exposure to any tobacco and e-cigarette advertisements.

Results 40% and 41.5% of the respondents reported having seen any e-cigarette and tobacco product advertisement respectively within the past year. Current smokers, males, younger respondents, those with financial difficulties, people who had tried e-cigarettes and daily internet users were more likely to report having seen an e-cigarette and a tobacco product advertisement. Respondents in countries with more comprehensive advertising bans were less likely to self-report exposure to any tobacco advertisements (OR 0.87; 95% CI 0.79 to 0.96 for one-unit increase in TCS advertising score), but not e-cigarette advertisements (OR 1.08; 95% CI 0.95 to 1.22).

Conclusion Ten years after ratification of the Framework Convention for Tobacco Control, self-reported exposure to tobacco and e-cigarette advertising in the EU is higher in e-cigarette and tobacco users, as well as those with internet access. The implementation of the Tobacco Products Directive may result in significant changes in e-cigarette advertising, therefore improved monitoring of advertising exposure is required in the coming years.

INTRODUCTION

Comprehensive bans on tobacco advertising, promotion and sponsorship are effective measures to decrease smoking rates.¹ Article 13 of the WHO Framework Convention for Tobacco Control² calls for comprehensive bans on all available media. Despite this, legislation to regulate tobacco product advertising across the globe is heterogeneous.³ In response to the increase in e-cigarette advertising and use,^{4–6} the Tobacco Products Directive (TPD) passed by the European Union (EU) in 2014⁷ addresses issues pertinent to e-cigarette advertising in print and in audiovisual avenues. There are no

specific provisions in the TPD, but ‘a restrictive approach to advertising electronic cigarettes and refill containers’ is suggested.⁷ However, the speed of transposition of the TPD into national jurisdictions potentially varies, and hence discrepancies in exposure to e-cigarette advertising may exist across the EU.

As transposition of the TPD has commenced, it will be of interest to monitor changes in exposure to tobacco and e-cigarette advertising across the EU that may be potentially attributable to differing adoption of TPD articles or inherent population differences. The objective of this study was to explore factors associated with self-reported exposure of the EU population to tobacco products and electronic cigarettes advertising. For this purpose, we used a Eurobarometer survey, which was conducted before TPD implementation had commenced; hence our findings on e-cigarette advertising may act as a pre-TPD assessment of self-reported exposure to e-cigarette advertising in the past year.

METHODS

Data source

We conducted a secondary analysis of data from wave 82.4 of the Eurobarometer survey,⁸ collected in 28 EU member states in November–December 2014. A multistage sampling design was employed to collect samples representative of the population aged ≥15 years. Face-to-face interviews were used to collect self-reported data from n=27 801 individuals from all 28 EU member states.

Measures

Exposure to tobacco products advertising was assessed with the question ‘in the past 12 months, have you seen advertisements or promotions for tobacco products in (OUR COUNTRY)? Advertisement and promotion for electronic cigarettes or similar devices should not be reported here’. A similar question on advertising of ‘electronic cigarettes or any similar devices (eg, e-shisha, e-pipe)’ was also asked. Exposure to advertising was analysed as a binary variable (‘often’; ‘from time to time’ and ‘rarely’ vs ‘never’) in order to reduce potential misclassification due to inaccurate recall of exposure frequency.

The survey also collected data on participants’ smoking (current, former and never smokers), e-cigarette ever use, age (15–24, 25–39, 40–54 and ≥55 years), gender (male, female), age at which



CrossMark

To cite: Filippidis FT, Laverty AA, Fernandez E, et al. *Tob Control* 2017;**26**:e130–e133.

they stopped full-time education (≤ 15 , 16–19 and ≥ 20 years), their difficulties to pay bills during the last 12 months (almost never/never and from time to time/most of the time) and internet use (everyday, occasionally, never). More details on Eurobarometer have been given elsewhere.^{9 10}

Data on existing bans on tobacco advertising across the EU were extracted from the 2013 version of the Tobacco Control Scale (TCS).¹¹ We used each country's score on the domain 'bans of tobacco advertising', in which countries are assigned a score ranging from 0 to 12, based on the extent of advertising and promotion bans. Higher scores correspond to more comprehensive smoking bans.

Statistical analysis

We explored factors associated with reporting having seen any advertisements of tobacco products and—separately—e-cigarettes with a multilevel logistic regression model, with country being the higher level of analysis. The independent variables included in the model were sex, age, area of residence, education, difficulty to pay bills, internet use, tobacco smoking, e-cigarette ever use and TCS advertising score (at a country level). Sensitivity analyses were performed categorising only those reporting that they had seen advertisements often or from time to time as exposed. Analyses were performed with Stata V.14.0, incorporating the weights provided in the Eurobarometer data set to account for the complex survey design.

RESULTS

A total of 41.5% (95% CI 40.5% to 42.5%) and 40.0% (95% CI 39.0% to 41.0%) of the respondents reported having been exposed to a tobacco product and e-cigarette advertising or promotion in the past 12 months, respectively. Detailed results by country have been presented in the official Eurobarometer report.⁹

Current smokers were significantly more likely to report exposure to tobacco advertising (OR 1.26; 95% CI 1.17 to 1.36) and e-cigarette advertising (OR 1.11; 95% CI 1.04 to 1.19) in the past 12 months, compared with never-smokers (table 1). Former smokers were also more likely to report exposure to e-cigarette advertising compared with never smokers (OR 1.30; 95% CI 1.20 to 1.40), but not to tobacco advertising (OR 1.03; 95% CI 0.96 to 1.11). Those who had ever tried e-cigarette, occasional and daily internet users, younger respondents, men, people with financial difficulties and those with higher education were also significantly more likely to report having seen, read or heard tobacco and e-cigarette advertisements or promotions in the past 12 months. Results from the sensitivity analyses were very similar (supplementary table 1).

Respondents living in countries with more comprehensive bans in tobacco advertising were less likely to report exposure to tobacco advertisements (OR 0.87; 95% CI 0.79 to 0.96, for one-point increase in the TCS advertising score); however, this was not the case with e-cigarette advertising (OR 1.08; 95% CI 0.95 to 1.22) (table 1).

DISCUSSION

Our analysis showed that sociodemographic factors were associated with the self-reported exposure to any tobacco and e-cigarette promotions in the past year, with those who had tried e-cigarettes and had frequent access to internet being more likely to report having seen an advertisement. Respondents living in member states with more comprehensive advertising bans were less likely to report exposure to tobacco advertisements, but did not differ in self-reported exposure to e-cigarette advertising.

Table 1 Association of exposure to tobacco and e-cigarette advertisements or promotions with tobacco smoking, e-cigarette experimentation, internet use and sociodemographic factors in 28 European Union member states, Eurobarometer 2014

| | Exposure to tobacco advertisements (n=25 454) OR (95% CI) | Exposure to e-cigarette advertisements (n=25 528) OR (95% CI) |
|---|---|---|
| Age (years) | | |
| ≥ 55 (ref.) | 1.00 | 1.00 |
| 40–54 | 1.17 (1.09 to 1.26) | 1.17 (1.08 to 1.27) |
| 25–39 | 1.28 (1.18 to 1.38) | 1.28 (1.18 to 1.40) |
| 15–24 | 1.60 (1.44 to 1.78) | 1.46 (1.31 to 1.63) |
| Gender | | |
| Female (ref.) | 1.00 | 1.00 |
| Male | 1.32 (1.25 to 1.40) | 1.14 (1.07 to 1.20) |
| Difficulties paying bills | | |
| Never/almost never (ref.) | 1.00 | 1.00 |
| From time to time/most of the time | 1.10 (1.04 to 1.18) | 1.09 (1.02 to 1.16) |
| Age when stopped education (years) | | |
| Up to 15 (ref.) | 1.00 | 1.00 |
| 16–19 | 1.22 (1.12 to 1.33) | 1.29 (1.17 to 1.42) |
| ≥ 20 | 1.40 (1.27 to 1.54) | 1.51 (1.36 to 1.67) |
| Area of residence | | |
| Rural (ref.) | 1.00 | 1.00 |
| Urban | 1.09 (1.03 to 1.16) | 1.11 (1.04 to 1.18) |
| Tobacco smoking | | |
| Never smokers (ref.) | 1.00 | 1.00 |
| Current smoker | 1.11 (1.04 to 1.19) | 1.26 (1.17 to 1.36) |
| Former smoker | 1.03 (0.96 to 1.11) | 1.30 (1.20 to 1.40) |
| Internet use | | |
| Never/no access (ref.) | 1.00 | 1.00 |
| Often/sometimes | 1.63 (1.47 to 1.80) | 1.82 (1.64 to 2.03) |
| Everyday | 1.71 (1.57 to 1.86) | 2.21 (2.01 to 2.42) |
| E-cigarette experimentation | | |
| No (ref.) | 1.00 | 1.00 |
| Yes | 1.24 (1.13 to 1.37) | 1.76 (1.60 to 1.94) |
| Tobacco Control Scale tobacco advertising score | 0.87 (0.79 to 0.96) | 1.08 (0.95 to 1.22) |

Results from a multilevel regression model adjusted for all the variables in the table; country was the higher level of analysis. Statistically significant associations are shown in bold.

The main outcome measure in Eurobarometer is self-reported frequency of exposure to advertising over the past 12 months, which was dichotomised to represent any exposure. While self-report is the most commonly used measure of media exposure, its validity and reliability may vary and it can be subject to recall bias,¹² especially in the current context of multiple and overlapping exposures to digital media.¹³ Self-report is considered an adequate measure of the relative levels of media exposure,¹² but the timeframe (ie, 12 months) in the Eurobarometer question is considerably larger than the one typically used in tobacco surveillance; for example, the Global Adult Tobacco Survey assesses exposure in the past 30 days.¹⁴ Additionally, we analysed the exposure measure as a binary variable which could not reflect the frequency of exposure. Nonetheless, sensitivity analyses using different level of exposure gave similar results. Therefore, our results should be interpreted with caution and may not fully reflect actual levels of exposure.

The association between TCS advertising scores and self-reported exposure to tobacco advertising suggests that advertising bans may be effective in reducing exposure to marketing activities for tobacco products, as exemplified by the contrasting cases of Finland (maximum TCS advertising score and low self-reported exposure) and Germany (lowest TCS score and high self-reported exposure).^{9,11} The lack of an association between TCS advertising scores and self-reported exposure to e-cigarette advertising could imply that other factors than the tobacco control environment, such as the market development, also play a role.

Having daily or frequent access to the internet was strongly associated with increased odds of having seen tobacco, but especially e-cigarette advertisements. E-cigarettes have a strong presence within social media^{15–18} and online advertising is generally cheaper than in traditional media, allowing smaller companies to promote their products at relatively low cost. Online promotions are also harder to monitor and regulate and may attract younger users,¹⁹ who are more likely to experiment with e-cigarettes.²⁰ Currently, the TPD does not address issues of online advertising of e-cigarettes, which should be further considered by policymakers, as it is a main source of e-cigarette promotions in the EU.⁹ E-cigarette advertising is further complicated by the direct involvement of vapour store owners, who seem to be following strategies typical of the tobacco industry, including pricing discounts, loyalty programmes and use of print and social media²¹ and of current e-cigarette users, who serve as an important source of information about e-cigarettes¹⁹ and can disseminate information to a large number of people through social media.

The Eurobarometer survey is cross-sectional, thus precluding any conclusion on potential causal relationships, but the large and representative sample of the EU population allows for analyses that control for several confounding factors and yield results that are generalisable at an EU level.

CONCLUSIONS

The likelihood of self-reported exposure to any tobacco or e-cigarettes advertising in the past year was higher among smokers, male, young, literate, urban dwellers and users of the internet. Those living in countries with more comprehensive advertising bans were less likely to report exposure to tobacco advertising but not to e-cigarette advertising. Further longitudinal research with more sensitive measures of advertising exposure is needed to support these results and monitor potential changes on the characteristics of tobacco and e-cigarette advertising that may arise from national transposition and implementation of the TPD.

What this paper adds

- ▶ 41.5% of people aged ≥ 15 years self-reported past year exposure to tobacco products advertising and 40% to e-cigarette advertising in the European Union in 2014.
- ▶ E-cigarette ever-users, current smokers and those with frequent access to the internet reported higher past year exposure to tobacco and e-cigarette advertising.
- ▶ In member states with more comprehensive advertising bans, self-reported past year exposure to tobacco products advertisements was lower, but that was not the case with e-cigarettes.

Author affiliations

¹Center for Health Services Research, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece

²Public Health Policy Evaluation Unit, School of Public Health, Imperial College London, London, UK

³Tobacco Control Unit, WHO Collaborating Center for Tobacco Control, Institut Català d'Oncologia-ICO, L'Hospitalet de Llobregat, Barcelona, Catalonia, Spain
⁴Cancer Prevention and Control Group, Institut d'Investigació Biomèdica de Bellvitge-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Catalonia, Spain
⁵Department of Clinical Sciences, School of Medicine, Campus de Bellvitge, Universitat de Barcelona, L'Hospitalet del Llobregat Barcelona, Catalonia, Spain
⁶Cancer Prevention Unit and WHO Collaborating Center for Tobacco Control, German Cancer Research Center, Heidelberg, Germany
⁷Clinic of Social and Family Medicine, School of Medicine, University of Crete, Heraklion, Greece
⁸European Network for Smoking and Tobacco Prevention (ENSP), Brussels, Belgium

Correction notice This paper has been amended since it was published Online First. Owing to a scripting error, some of the publisher names in the references were replaced with 'BMJ Publishing Group'. This only affected the full text version, not the PDF. We have since corrected these errors and the correct publishers have been inserted into the references. The Funding statement has been added back into the paper.

Contributors FTF had the main role in data analysis. FTF, AAL and EF had the main role in manuscript preparation. All authors contributed to data interpretation and manuscript preparation. FTF is responsible for the overall content.

Funding This work was supported by a grant from the European Commission (Horizon2020 HCO-6-2015; EUREST-PLUS: 681109; Vardavas). EF was partly supported by the Instituto de Salud Carlos III of the Government of Spain and the European Regional Development Fund, ERDF (RTICC RD12/0036/0053) and the Department of Universities and Research, Government of Catalonia (2014SGR999).

Competing interests None declared.

Ethics approval Data were freely available and de-identified, thus no ethical approval was required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data set supporting the conclusions of this article is freely available online, doi:10.4232/1.12265 in <https://dbk.gesis.org/dbksearch/sdesc.asp?no=5933&db=e&doi=10.4232/1.12264>.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Levy DT, Chaloupka F, Gitchell J. The effects of tobacco control policies on smoking rates: a tobacco control scorecard. *J Public Health Manag Pract* 2004;10:338–53.
- 2 World Health Organization. *WHO Framework Convention on tobacco control*. Geneva: World Health Organization, 2003.
- 3 World Health Organization. *WHO Report on the global tobacco epidemic 2013*, 2013.
- 4 Filippidis FT, Laverty AA, Gerovasili V, et al. Two-year trends and predictors of e-cigarette use in 27 European Union member states. *Tob Control* 2017;26:98–104.
- 5 Kornfield R, Huang J, Vera L, et al. Rapidly increasing promotional expenditures for e-cigarettes. *Tob Control* 2015;24:110–1.
- 6 Duke JC, Lee YO, Kim AE, et al. Exposure to electronic cigarette television advertisements among youth and young adults. *Pediatrics* 2014;134:e29–e36.
- 7 European Commission. Directive of the European Union on the approximation of the laws, regulations and administrative provisions of the Member States concerning the manufacture, presentation and sale of tobacco and related products and repealing directive 2001/37/EC. http://ec.europa.eu/health/tobacco/docs/dir_201440_en.pdf (accessed 25 Sep 2015).
- 8 European Commission. Eurobarometer 82.4, November–December 2014. GESIS Data Archive: za5933, dataset version 5.0.0 (2014). Brussels: TNS OPINION & SOCIAL, 2014.
- 9 European Commission. Special eurobarometer 429. *Attitudes of Europeans towards tobacco* 2015.
- 10 Filippidis FT, Laverty AA, Vardavas CI. Experimentation with e-cigarettes as a smoking cessation aid: a cross-sectional study in 28 European Union member states. *BMJ Open* 2016;6:e012084.
- 11 Joossens LRM. *The tobacco control Scale 2013 in Europe*, 2013.
- 12 de Vreese CH, Neijens P. Measuring media exposure in a changing communications environment. *Commun Methods Meas* 2016;10:69–80.
- 13 Niederdeppe J. Meeting the challenge of Measuring Communication exposure in the Digital Age. *Commun Methods Meas* 2016;10:170–2.

- 14 World Health Organization. Tobacco questions for surveys.
- 15 Jo CL, Kornfield R, Kim Y, *et al.* Price-related promotions for tobacco products on Twitter. *Tob Control* 2016;25:476–9.
- 16 Huang J, Kornfield R, Szczyka G, *et al.* A cross-sectional examination of marketing of electronic cigarettes on Twitter. *Tob Control* 2014;23(Suppl 3):iii26–iii30.
- 17 Romito LM, Hurwich RA, Eckert GJ. A snapshot of the depiction of electronic cigarettes in YouTube Videos. *Am J Health Behav* 2015;39:823–31.
- 18 Luo C, Zheng X, Zeng DD, *et al.* Portrayal of electronic cigarettes on YouTube. *BMC Public Health* 2014;14:1028.
- 19 Pepper JK, Emery SL, Ribisl KM, *et al.* How U.S. adults find out about electronic cigarettes: implications for public health messages. *Nicotine Tob Res* 2014;16:1140–4.
- 20 Vardavas CI, Filippidis FT, Agaku IT. Determinants and prevalence of e-cigarette use throughout the European Union: a secondary analysis of 26 566 youth and adults from 27 Countries. *Tob Control* 2015;24:442–8.
- 21 Cheney M, Gowin M, Wann TF. Marketing practices of vapor store owners. *Am J Public Health* 2015;105:e16–e21.