

Assessing Exposure to Household Air Pollution: A Systematic Review and Pooled Analysis of Carbon Monoxide as a Surrogate Measure of Particulate Matter

Ellison Carter,^{1*} Christina Norris,² Kathie L. Dionisio,³ Kalpana Balakrishnan,⁴ William Checkley,^{5,6} Maggie L. Clark,⁷ Santu Ghosh,⁴ Darby W. Jack,⁸ Patrick L. Kinney,⁸ Julian D. Marshall,⁹ Luke P. Naeher,¹⁰ Jennifer L. Peel,⁷ Sankar Sambandam,⁴ James J. Schauer,^{11,12} Kirk R. Smith,¹³ Blair J. Wylie,¹⁴ and Jill Baumgartner^{1,2,15}

¹Institute on the Environment, University of Minnesota, St. Paul, Minnesota, USA

²Department of Epidemiology, Biostatistics & Occupational Health, McGill University, Montreal, Quebec, Canada

³National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, USA

⁴Department Environmental Health Engineering, Sri Ramachandra University, Porur, Chennai, India

⁵Division of Pulmonary and Critical Care, School of Medicine, Johns Hopkins University, Baltimore, Maryland, USA

⁶Program in Global Disease Epidemiology and Control, Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

⁷Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, Colorado, USA

⁸Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, New York, USA

⁹Department of Civil and Environmental Engineering, University of Washington, Seattle, Washington, USA

¹⁰Department of Environmental Health Science, College of Public Health, The University of Georgia, Athens, Georgia, USA

¹¹Environmental Chemistry & Technology Program, University of Wisconsin-Madison, Madison, Wisconsin, USA

¹²Department of Civil & Environmental Engineering, University of Wisconsin-Madison, Madison, Wisconsin, USA

¹³Division of Environmental Health Sciences, School of Public Health, University of California, Berkeley, Berkeley, California, USA

¹⁴Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

¹⁵Institute for Health and Social Policy, McGill University, Montreal Quebec, Canada

BACKGROUND: Household air pollution from solid fuel burning is a leading contributor to disease burden globally. Fine particulate matter (PM_{2.5}) is thought to be responsible for many of these health impacts. A co-pollutant, carbon monoxide (CO) has been widely used as a surrogate measure of PM_{2.5} in studies of household air pollution.

OBJECTIVE: The goal was to evaluate the validity of exposure to CO as a surrogate of exposure to PM_{2.5} in studies of household air pollution and the consistency of the PM_{2.5}–CO relationship across different study settings and conditions.

METHODS: We conducted a systematic review of studies with exposure and/or cooking area PM_{2.5} and CO measurements and assembled 2,048 PM_{2.5} and CO measurements from a subset of studies (18 cooking area studies and 9 personal exposure studies) retained in the systematic review. We conducted pooled multivariate analyses of PM_{2.5}–CO associations, evaluating fuels, urbanicity, season, study, and CO methods as covariates and effect modifiers.

RESULTS: We retained 61 of 70 studies for review, representing 27 countries. Reported PM_{2.5}–CO correlations (*r*) were lower for personal exposure (range: 0.22–0.97; median = 0.57) than for cooking areas (range: 0.10–0.96; median = 0.71). In the pooled analyses of personal exposure and cooking area concentrations, the variation in ln(CO) explained 13% and 48% of the variation in ln(PM_{2.5}), respectively.

CONCLUSIONS: Our results suggest that exposure to CO is not a consistently valid surrogate measure of exposure to PM_{2.5}. Studies measuring CO exposure as a surrogate measure of PM exposure should conduct local validation studies for different stove/fuel types and seasons. <https://doi.org/10.1289/EHP767>

Introduction

Over 2.8 billion people are exposed to household air pollution from cooking and heating with solid fuels, which include biomass (e.g., wood, crop residues, animal dung, charcoal) and coal (Bonjour et al. 2013). Household air pollution comprises many pollutants (Zhang and Smith 2007; Naeher et al. 2007) and is a leading health risk factor, annually responsible for an estimated

2.9 million premature deaths (GBD 2013 Risk Factors Collaborators et al. 2015). Two widely studied air pollutants from solid fuel combustion are particulate matter (PM) and carbon monoxide (CO). Strong epidemiologic and experimental evidence point to the mass of PM with a diameter $\leq 2.5 \mu\text{m}$ (PM_{2.5}) as a pollutant that is causally associated with many health outcomes (Pope and Dockery 2006; U.S. EPA 2009) and is likely a strong driver of many health effects associated with household air pollution (Brook et al. 2010; WHO 2014). Evidence for adverse health outcomes related to low-to-moderate CO exposure is sparse and less consistent, with associations between infant low birth weight and women's CO exposure during pregnancy demonstrated in some studies (Ritz and Yu 1999; Ha et al. 2001; Gouveia et al. 2004; Salam et al. 2005), but not in others (Alderman et al. 1987; Koren et al. 1991; Chen et al. 2002; Parker et al. 2005; Wylie et al. 2016). In epidemiologic and exposure studies of household air pollution, including those evaluating maternal exposure and birth outcomes (Thompson et al. 2011; Dix-Cooper et al. 2012), CO exposure is usually measured as a surrogate of PM_{2.5} exposure (Balakrishnan et al. 2011; Clark et al. 2013).

Accurate exposure assessment is the basis for evaluating exposure–response relationships (Armstrong 1998, 2004), and in the context of household air pollution, critical to interpreting the effectiveness of stove-fuel interventions (Peel et al. 2015). Direct

Address correspondence to J. Baumgartner, Institute for Health and Social Policy and Department of Epidemiology, Biostatistics and Occupational Health, 1130 des Pins Ave. Ouest, Montréal, Québec H3A 1A3 Canada. Telephone: (514) 398-6688. Email: jill.baumgartner@mcgill.ca

*Current affiliation: Department of Civil and Environmental Engineering, Colorado State University, 400 Isotope Dr, Fort Collins, CO 80521 USA.

Supplemental Material is available online (<https://doi.org/10.1289/EHP767>).

The authors declare they have no actual or potential competing financial interests.

Received 6 July 2016; Revised 19 December 2016; Accepted 20 December 2016; Published 28 July 2017.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

measurement of personal exposure to PM_{2.5} mass is considered the “gold standard” in epidemiologic studies (Smith 1993; Northcross et al. 2015), but is challenging to measure in large populations (Northcross et al. 2015) and in infants (Naeher et al. 2001; Dionisio et al. 2008). Questionnaires and cooking area PM_{2.5} have been used alone or in combination as surrogates but were poorly associated with personal PM_{2.5} exposure in validation studies (Ezzati et al. 2000; Bruce et al. 2004; Cynthia et al. 2008; Baumgartner et al. 2011; Ni et al. 2016). As an alternative, many health and intervention studies have measured personal exposure to CO as a surrogate for PM_{2.5} given that it is also a major component of household air pollution but is easier and less costly to measure than PM_{2.5} (Naeher et al. 2001; Dionisio et al. 2008; Smith et al. 2010).

The empirical evidence supporting the validity of personal CO exposure as a surrogate of personal PM_{2.5} exposure is limited and inconclusive, despite its common use. Direct measurements of personal PM_{2.5} and CO exposure were not correlated (Pearson $r = -0.04$) in children living in homes cooking with wood in The Gambia (Dionisio et al. 2012) and were only moderately correlated in women in Peru (Spearman $r = 0.41$; Commodore et al. 2013), Tanzania (Spearman $r = 0.34$; Wylie et al. 2016), and China (Spearman $r = 0.60$; Ni et al. 2016). In rural Guatemala, however, variation in personal CO exposures explained 78% of the variation in personal PM_{2.5} exposures among women (McCracken et al. 2013). It is not known whether the strength of a PM_{2.5}–CO relationship in one setting is transportable to other settings. Here, we use definitions for a validation study and transportability adapted from Spiegelman (2010):

Validation study: a study in which data are simultaneously collected on the exposure surrogate (CO) and the gold standard method of exposure assessment (PM_{2.5}). This study may be external to the main epidemiologic study, or be a subsample internal to the main study.

Transportability: the extent to which the PM_{2.5}–CO relationship in a validation study is similar to the one that generates the surrogate exposure in the main study (PM_{2.5}).

Studies in Bolivia, Peru, Ghana, Kenya, Mexico, South Africa, the Philippines, and Burkina Faso (Röllin et al. 2004; Saksena et al. 2007; Riojas-Rodriguez et al. 2011; Ochieng et al. 2013; Commodore et al. 2013; Alexander et al. 2014; Thorsson et al. 2014; Jack et al. 2015; Yip et al. 2017) measured CO exposure as a surrogate for PM_{2.5} without prior validation; one reason given was the strong PM_{2.5}–CO exposure relationship observed in Guatemala (McCracken et al. 2013; Naeher et al. 2000a). Similarly, it is unknown whether the PM_{2.5}–CO exposure relationship within a single study setting and population under one set of study conditions is transportable to other study conditions (e.g., pre- vs. postintervention; heating- vs. nonheating season) in the same setting and population, which is an approach taken in some studies (Smith-Sivertsen et al. 2009; Smith et al. 2011; Guarnieri et al. 2014; Pope et al. 2015). Finally, it is unclear whether the PM_{2.5}–CO correlation in cooking areas can be extrapolated to personal exposures in the same setting, which several studies have done (Bruce et al. 2004; Northcross et al. 2010; Dionisio et al. 2012; Alnes et al. 2014). Of these, only one (Dionisio et al. 2012) directly compared actual versus predicted PM_{2.5} exposure, finding no relationship (Pearson $r = 0.01$).

Both PM and CO are products of incomplete combustion and co-emitted during solid fuel burning. The amount and relative proportion of these pollutants emitted from stoves can vary by factors including fuel type and moisture content; combustion

efficiency and power throughout burn cycles; stove ventilation; and the behavior of energy users (Roden et al. 2009; Chen et al. 2012; Jetter et al. 2012; Carter et al. 2014). For personal exposures, the presence of other community or regional air pollution sources with different pollutant composition (Huang et al. 2015) could further impact the strength and consistency of a personal PM–CO association.

We systematically reviewed the methods and correlation coefficients reported in studies with paired measurements of PM_{2.5} and CO personal exposures and/or cooking area concentrations in settings where biomass is the primary household fuel. We also obtained 2,048 paired PM_{2.5} and CO measurements from previously completed studies along with relevant information on season, level of urbanicity, fuel type, and other energy use behaviors to conduct pooled analyses of the PM_{2.5}–CO relationship for personal exposures and cooking area concentrations. For the pooled analysis, our first objective was to evaluate the validity of exposure to CO as a surrogate of exposure to PM_{2.5} in epidemiologic and intervention studies of household air pollution. Because most health studies aim to evaluate daily or “usual” exposure, we limited our pooled analysis to studies of PM_{2.5} and CO concentration and/or exposure relationships for at least 24-hr in settings where biomass was the dominant household fuel. Our second objective was to evaluate whether the PM_{2.5}–CO relationships estimated under one set of conditions are transportable to other conditions.

We provide a timely assessment of CO exposure as a surrogate of PM_{2.5} exposure, as a systematic review has been lacking but is critical to exposure measurement method selection for ongoing (Rosa et al. 2014; Klasen et al. 2013; Tielsch et al. 2014; Jack et al. 2015; NIH 2015) randomized controlled trials and other epidemiologic studies.

Methods

Search Strategy and Selection Criteria for the Systematic Review

We searched publications included in the electronic database PubMed (from 1966 to present; <https://www.ncbi.nlm.nih.gov/pubmed/>) and the Science Citation index, as well as the electronic databases Ovid MEDLINE® In-Process & Other NonIndexed Citations, Ovid MEDLINE® Daily, Ovid OLDMEDLINE®, (1946 to present, <http://ospguides.ovid.com/OSPguides/medline.htm>) and Embase Classic+ Embase (1947 to present, <http://www.elsevier.com/embase>). We searched for combinations of the key words “carbon monoxide” or “CO” and “indoor air pollution” or “indoor*” or “house*” or “home*” or “personal exposure and particulate*” or “PM*” and “biomass or coal” or “fuel*” or “wood*” or “dung” or “crop” or “agricultural residue*.” The search was restricted to articles available in English, French, Spanish, or Chinese. We retained articles for which cooking area or personal measurements of PM and CO were done concurrently in a setting where biomass was burned for cooking and/or heating. Two researchers independently extracted information from these articles and hand-searched their reference lists to identify additional publications for retrieval. Finally, we contacted 15 researchers to directly obtain data from published and unpublished studies with paired PM and CO measurements. These studies were identified from the literature review and recent conference proceedings or academic meetings. We adhered to systematic review guidelines from PRISMA-P guidelines and the Cochrane Collaboration (Van Tulder et al. 2003; Moher et al. 2015).

We classified studies retained for this review into two groups (Figure 1): studies with paired measurements of personal PM and




Figure 1. Flow diagram of systematic search of literature for review.

CO exposures or stationary PM and CO concentrations in cooking areas. Though personal exposures were our primary interest, we reviewed studies with paired cooking area measurements because they are more common than studies with personal exposure (Balakrishnan et al. 2011; Clark et al. 2013) and may shed additional light on the $PM_{2.5}$ –CO relationship for personal exposures. For every study, the following information was extracted: authors, year of publication, year(s) of data collection, location, season(s), description of setting, elevation, description of study population (see Table S1), stove types, cooking location, cooking area ventilation, fuel types, other local air pollution sources, number of paired PM and CO measurements, pollutant measurement methods (i.e., protocols, instrumentation, quality assurance, quality control measures), and reported $PM_{2.5}$ –CO correlation coefficients. When information was not reported, we requested it from corresponding authors.

Compiling Paired $PM_{2.5}$ and CO Measurements for Pooled Data Analysis

We contacted the corresponding authors of studies identified in our systematic review to obtain the paired $PM_{2.5}$ and CO data. If authors did not respond but the data were available in published studies, we downloaded those data. We also requested information on the stoves and fuels used, stove ventilation, other local air pollution sources, season of data collection, level of urbanicity (rural vs. peri-urban/urban), and $PM_{2.5}$ and CO measurement methods. If these variables were unavailable for individual observations, we assigned them at the study level based on the information reported in the manuscript or communication with authors.

We obtained 2,048 paired $PM_{2.5}$ and CO measurements and covariate data from 9 studies of personal $PM_{2.5}$ and CO exposures ($n = 714$ pairs) and 18 studies of cooking area $PM_{2.5}$ and CO concentrations ($n = 1,334$ pairs). Personal exposure data were obtained from authors for 6 studies and extracted from tables or figures from 3 studies (Fitzgerald et al. 2012; McCracken et al. 2013; Naeher et al. 2000b). For paired cooking area measurements, data were obtained from authors of 9 studies, and we extracted data from 9 studies (Naeher et al. 2000a; Naeher et al. 2001; Park and Lee 2003; Chengappa et al. 2007; Dutta et al. 2007; Henkle et al.

2010; Fitzgerald et al. 2012; Chowdhury et al. 2013; Huboyo et al. 2014). We created dichotomous variables for covariates (see Table S2), including fuel use (exclusive vs. nonexclusive biomass use), whether other local sources of air pollution were reported, level of urbanicity, season (nonheating vs. heating), and CO measurement method (colorimetric-based vs. sensor-based). PM measurement type was almost exclusively gravimetric for personal exposures and a mix of gravimetric and optical measurements in cooking areas. We summarized the protocols and quality assurance/control procedures for personal PM and CO in Tables S3 and S4.

Statistical Analysis of the Pooled Data

We conducted a series of univariate and multivariate regression models to evaluate the coefficient of determination (R^2) and the slope between $PM_{2.5}$ and CO, with separate models for personal exposure and cooking area measurements. Natural cubic spline functions with 2–5 degrees of freedom were used to evaluate whether the pollutant relationships were linear functions. Covariates including fuel use, other local sources of air pollution, urbanicity, season, and CO measurement method were added to the models to determine the extent to which their inclusion improved the R^2 . We incorporated a random intercept for study into the linear regression models to account for clustering of data by study. The R^2 values were compared to quantify the proportion of variation in $\ln(PM_{2.5})$ explained by $\ln(CO)$ alone and after including other covariates in the models. Differences in the slope of $\ln(PM_{2.5})$ on $\ln(CO)$ by fuel use, urbanicity, season, and CO measurement method were also compared to evaluate transportability (i.e., similarity) of the $PM_{2.5}$ –CO relationships between study conditions. Finally, in studies for which personal exposure and cooking area measurements of $PM_{2.5}$ and CO were concurrent, we graphically compared the slopes of the personal exposure versus cooking area $PM_{2.5}$ –CO relationships to assess within-study transportability of the cooking area $PM_{2.5}$ –CO relationship to personal exposures.

As sensitivity analyses, we conducted the same models with untransformed CO. We also conducted the analyses without nonwood biomass (e.g., dung, charcoal), which may differ from wood in its proportional contribution of PM and CO to

overall emissions (Jetter et al. 2012), and excluding studies not meeting the U.S. EPA (2016) Quality Assurance Guidelines for gravimetric PM analysis ($n=2$). Finally, we compared the R^2 values for univariate and multivariate models within studies to investigate the extent to which individual-level covariate heterogeneity improved explanation of variation in $\ln(\text{PM}_{2.5})$ by $\ln(\text{CO})$. All model assumptions were verified by routine diagnostic analysis of the residuals. The statistical analysis was conducted using Stata 13.1 (Stata Corporation, College Station, Texas, USA).

Results

Systematic Review of the Literature

Our search criteria yielded 70 studies, including 2 unpublished studies that were eligible for review, representing measurements in 27 countries. Of these, we retained 61 studies for review after excluding 5 studies of outdoor wildfires, 3 studies of emissions measurements, and 1 urban outdoor air pollution study. Publication year ranged from 1968 to 2016, though most studies (92%) were published after 2000. Studies were conducted in Sub-Saharan Africa ($n=12$), Latin America ($n=23$), South and East Asia ($n=16$), Eastern Mediterranean ($n=1$), and Western Pacific ($n=9$).

Studies with paired measurements of personal exposures to $\text{PM}_{2.5}$ and CO in adults and/or children accounted for 23% ($n=14$) of all studies reviewed (Table 1). Sample sizes ranged from 10 to 268 paired measurements (median=80 pairs). Twelve of the 14 studies enrolled women who were the primary household cooks; in 2 studies, all enrolled participants were pregnant (St. Helen et al. 2015; Wylie et al. 2016). One study enrolled children 15–61 months of age (Dionisio et al. 2012), and another (Naeher et al. 2000a) enrolled mother–child pairs in which both mother and child (<15 months) wore the CO and $\text{PM}_{2.5}$ monitors.

Personal $\text{PM}_{2.5}$ and CO exposures were integrated over 22-, 24-, or 48-hr periods to represent “usual” daily exposure. Most studies ($n=12$ of 14) measured personal $\text{PM}_{2.5}$ exposures with

gravimetric instruments. Nine studies used a sensor-based method for CO measurement, and five used a colorimetric dosimeter.

Studies of paired cooking area $\text{PM}_{2.5}$ and CO concentrations comprised 93% ($n=57$) of those identified in this systematic review (see Table S5). Sample sizes ranged from 9 to 350 paired measurements (median=60 pairs). Most stationary $\text{PM}_{2.5}$ and CO concentrations were measured in kitchens and cooking areas located in the same building as the living quarters, although some were conducted in rooms adjacent to the kitchen or in separate cookhouses. In five studies (Fischer and Koshland 2007; Pearce et al. 2009; Leavey et al. 2015; Muralidharan et al. 2015; Saksena et al. 1992), $\text{PM}_{2.5}$ and CO measurements were limited to cooking events, but the rest were integrated over 22-, 24-, or 48-hr periods. Light-scattering, optical techniques ($n=27$) and integrated, gravimetric techniques ($n=30$) were used for $\text{PM}_{2.5}$ measurements. Of the studies with optical $\text{PM}_{2.5}$ measurements, 85% measured CO with an electrochemical or optical sensor. Of studies with gravimetric measurements of $\text{PM}_{2.5}$, CO was measured with a sensor in 67% of studies ($n=20$) and with a colorimetric dosimeter in 33% of studies ($n=10$).

Correlations between paired $\text{PM}_{2.5}$ and CO personal exposure measurements. Correlation coefficients (Spearman r) were reported or calculated for 11 of the 14 studies measuring personal exposures (Table 1). The highest correlation ($r=0.97$) was observed in the study with the smallest sample size, namely 12 observations from mother–child pairs using biomass in open fires and traditional stoves in Guatemala (Naeher et al. 2000a). In the remaining studies, the correlations ranged from $r=0.22$ to $r=0.71$ [$n=10$ studies; median=0.53; interquartile range (IQR)=0.34–0.68]. Personal $\text{PM}_{2.5}$ –CO correlations were generally higher for studies reporting exclusive use of biomass fuel ($n=9$ studies; median=0.60; IQR=0.49–0.71), conducted in rural settings ($n=10$ studies; median=0.64; IQR=0.53–0.77), and using sensor-based CO measurements ($n=9$ studies; median=0.57; IQR=0.41–0.71).

Correlations between paired measurements of cooking area $\text{PM}_{2.5}$ and CO concentrations. Correlation coefficients were reported or calculated in 45 of the 57 studies with paired $\text{PM}_{2.5}$

Table 1. Characteristics of studies with paired measurements of personal exposure to $\text{PM}_{2.5}$ and CO.

Author/year (country)	Fuel(s) ^a	Other local air pollution sources	CO/PM method		$\text{PM}_{2.5}$ –CO correlation ^f (correlation coeff; r)
			CO S ^b /D ^c	PM G ^d /LS ^e	
Cynthia et al. 2008 (Mexico)	Wood	ETS ^g	S	LS	0.82 ($n=45$) preintervention 0.84 ($n=45$) postintervention
Balakrishnan et al. 2015 (India)	Wood, dung		S	G	0.49 ($n=45$)
Commodore et al. 2013 (Peru)	Wood		S	LS	0.41 ($n=19^h$)
Dionisio et al. 2012 (The Gambia)	Wood		D	G	0.22 ($n=29$)
Ellegård and Egnéus 1993 (Zambia)	Wood, charcoal, electricity	ETS	D	G	NR ⁱ ($n=268$)
Fitzgerald et al. 2012 (Peru)	Wood		S	G	0.68 ($n=80$)
Hartinger et al. 2013 (Peru)	Wood	ETS	S	G	NR ($n=79$)
McCracken et al. 2013 (Guatemala)	Wood		S	G	0.70 ($n=216$)
Mukhopadhyay et al. 2012 (India)	Wood, dung, LPG ^j		S	G	NR ($n=10$)
Naeher et al. 2000a (Guatemala)	Wood		D	G	0.97 ($n=12$)
Ni et al. 2016 (China)	Wood	ETS	D	G	0.60 ($n=22$)
Peel JL, written and oral communication, 2016 (Honduras)	Wood		S	G	0.57 ($n=105$)
St. Helen et al. 2015 (Peru)	Wood, coal, LPG, kerosene	ETS	S	G	0.33 ($n=93$)
Wylie et al. 2016 (Tanzania)	Wood, charcoal, kerosene	ETS; major road ≤ 200 m	D	G	0.34 ($n=118$)

^aBiomass (e.g., wood, crop residue, dung) and non-biomass fuels.

^bSensor-based.

^cColorimetric/diffusion-based.

^dGravimetric.

^eLight-scattering.

^fSpearman correlation.

^gEnvironmental tobacco smoke.

^h4-Hr mean CO and $\text{PM}_{2.5}$ concentrations.

ⁱNot reported.

^jLiquefied petroleum gas.

and CO measurements in cooking areas (see Table S5) and ranged from $r=0.10$ to $r=0.96$ (median = 0.71; IQR = 0.54–0.80). Overall, the $PM_{2.5}$ –CO correlations were higher for studies with exclusive biomass use ($n=18$ studies; median = 0.74; IQR = 0.65–0.86) than use of multiple fuels ($n=26$ studies; median = 0.64; IQR = 0.50–0.79) but the same in rural ($n=37$ studies, median = 0.71, IQR = 0.53–0.80) and peri-urban settings ($n=7$ studies; median = 0.72; IQR = 0.54–0.80). The $PM_{2.5}$ –CO correlations were similar for all combinations of $PM_{2.5}$ and CO measurement techniques and in homes with or without a tobacco or pipe smoker. In one-third of studies reviewed, the authors reported $PM_{2.5}$ –CO correlation coefficients for subgroup analyses (see Table S5). Within studies, the $PM_{2.5}$ –CO correlation was often higher for observations in rural settings or where wood was burned in open fires or traditional stoves.

Results from Pooled Data Analyses of Paired $PM_{2.5}$ and CO Measurements

Paired personal exposures to $PM_{2.5}$ and CO. The $PM_{2.5}$ and CO personal exposure means, ranges (see Table S6), and correlations varied between studies (Figure 2). The overall $PM_{2.5}$ –CO correlation was $r=0.36$ [95% confidence intervals (CI): 0.30, 0.42; $n=714$ pairs] (Figure 2a). The majority of study participants lived in rural settings (68%) and used biomass fuels exclusively (80%). Participants who did not use biomass exclusively also used other fuels including liquefied petroleum gas (LPG; 9%), charcoal (6%), kerosene (3%), coal (2%), and electricity (0.1%). The $PM_{2.5}$ –CO correlation among only those living with a tobacco or pipe smoker ($n=46$ pairs) was low ($r=0.12$, 95% CI: -0.17 , 0.40). All $PM_{2.5}$ exposure measurements were gravimetric. Most (75%) CO observations were sensor-based, and the rest were colorimetric-based.




Figure 2. Paired personal $PM_{2.5}$ and personal CO exposure measurements for (a) all observations combined from nine studies and for (b–i) individual studies. One outlying data point for Tanzania (CO: 25.2 ppm, $PM_{2.5}$: 42.9 $\mu\text{g}/\text{m}^3$), one for Peru (CO: 25.2 ppm, $PM_{2.5}$: 42.9 $\mu\text{g}/\text{m}^3$), two for Guatemala (CO: 18.5 ppm, $PM_{2.5}$: 284 $\mu\text{g}/\text{m}^3$; CO: 23.6 ppm, $PM_{2.5}$: 1,843 $\mu\text{g}/\text{m}^3$), and two for India (CO: 14.7 ppm, $PM_{2.5}$: 1,226 $\mu\text{g}/\text{m}^3$; CO: 9.5 ppm, $PM_{2.5}$: 1,243 $\mu\text{g}/\text{m}^3$) are not pictured to improve data visualization. 2h has an expanded CO concentration range along the horizontal axis.




Figure 3. Natural log-transformed PM_{2.5} personal exposures versus natural log-transformed CO personal exposures plotted for nine unique studies. The Spearman correlation (\pm 95% confidence intervals) for all observations ($n = 714$ pairs) is presented at the bottom left of the figure.

Associations between personal exposures to PM_{2.5} and CO in the pooled analysis. Pollutant concentrations for personal exposure to PM_{2.5} and CO were not normally distributed (right-skewed), and were natural log-transformed prior to evaluating their relationship using scatter plots (Figure 3) and locally weighted scatter plot smoothing and natural cubic spline models (see Figure S1a,b). Visual inspection of these plots indicated that the personal $\ln(\text{PM}_{2.5})$ – $\ln(\text{CO})$ relationship was approximately linear.

None of the univariate or multivariate linear regression models explained more than 50% of the variance in $\ln(\text{PM}_{2.5})$ exposure. The proportion of variation in $\ln(\text{PM}_{2.5})$ exposure explained

by $\ln(\text{CO})$ exposure was 13% with CO alone in the model and 19% in the model including fuel use, urbanicity, season, and CO method (Figure 4). Restricting the multivariate analysis to observations conducted in rural settings ($n = 478$) or during the heating season ($n = 453$) resulted in the highest explanation of variation in $\ln(\text{PM}_{2.5})$, namely 42% and 47%, respectively (Figure 4). Excluding two studies not meeting the U.S. EPA Quality Assurance Guidelines for gravimetric analysis did not substantially change our results ($n = 376$ pairs; $R^2 = 0.16$).

We observed significant differences in the $\ln(\text{PM}_{2.5})$ – $\ln(\text{CO})$ slope by fuel use, level of urbanicity, and season (all interaction




Figure 4. Comparison of estimates of the slope of $\ln(\text{PM}_{2.5})$ on $\ln(\text{CO})$ (\pm 95% confidence intervals) for personal exposures using univariate and multivariate linear regression models for the full data set and stratified by subsets of the data. The R^2 values and root mean squared error (RMSE) for each model are reported to the right of the plotted $\ln(\text{CO})$ slope. Note: CI, confidence interval; RMSE, root mean squared error.

p -values < 0.02). The slope was three to five times greater for measurements with exclusive use of biomass fuel, in rural settings, and during the heating season (Figure 4). In one study, the $PM_{2.5}$ –CO relationship also varied by whether the measurements were conducted pre- versus postintervention. In rural Peru (Fitzgerald et al. 2012), the slope of $\ln(PM_{2.5})$ on $\ln(CO)$ was 0.50 (95% CI: 0.30, 0.70; $n = 41$) among participants cooking with open fires, which was twice that of the slope among participants using chimney stoves (0.22, 95% CI: 0.03, 0.42; $n = 36$) in the same setting (interaction p -value < 0.05).

Paired stationary concentrations of $PM_{2.5}$ and CO in cooking areas. Cooking area $PM_{2.5}$ and CO concentration means and ranges (see Table S7) and the strength of the $PM_{2.5}$ –CO correlation varied by study ($n = 18$ studies; median = 0.80; range = 0.10–0.92; IQR = 0.57–0.82). After combining the paired cooking area $PM_{2.5}$ and CO concentrations from these studies, the $PM_{2.5}$ –CO correlation was $r = 0.46$ ($n = 1,336$; 95% CI: 0.42–0.50) but improved to $r = 0.74$ ($n = 981$; 95% CI: 0.71–0.76) after removing 350 observations (26% of all observations) from a study in India with a $PM_{2.5}$ –CO correlation of $r = 0.10$ (Balakrishnan et al. 2013). The CO measurements in the India data set had minimal variability (range: 0.2–11.0 ppm; IQR: 0.3–3.0 ppm), whereas the PM measurements ranged from 25 to 8,820 $\mu g/m^3$. As the low CO variability could be attributable to instrument failure, these data were excluded from subsequent analyses. Of the remaining 981 observations, biomass was the primary cooking fuel for 82% ($n = 807$) of observations, followed by LPG (12%), dung (4%), kerosene (0.5%), coal (1.4%), and electricity (0.1%). Over 86% ($n = 847$) of observations were conducted in rural settings, and 64% ($n = 625$) took place during the nonheating season.

Associations between cooking area concentrations of $PM_{2.5}$ and CO in the pooled analysis. A natural cubic spline model of $\ln(PM_{2.5})$ and $\ln(CO)$ with three knots was consistent with a linear relationship (see Figure S2). The proportion of variation in $\ln(PM_{2.5})$ concentrations explained by $\ln(CO)$ concentrations was 48% in both the univariate and the multivariate models, which included fuel use, setting, season, and CO method (Figure 5). The $\ln(PM_{2.5})$ – $\ln(CO)$ slope for cooking area measurements was twice as large in homes exclusively using biomass fuels compared with homes using multiple fuels (interaction p -value < 0.001) and in rural compared with peri-urban settings (interaction p -value < 0.001) (see Figure S3). The slope for cooking area measurements did not significantly differ by season (interaction p -value = 0.18) or CO method (interaction p -value = 0.73).

Comparison of personal exposure and cooking area $PM_{2.5}$ –CO correlations and slopes. For five studies (Dionisio et al. 2012; Fitzgerald et al. 2012; Ni et al. 2016; Peel JL, written and oral communication, 2016; St. Helen et al. 2015), personal exposure and cooking area $PM_{2.5}$ and CO measurements were collected concurrently (Figure 5). With the exception of a study in China (Ni et al. 2016), the R^2 value was considerably higher for cooking area measurements than for personal exposures in the same study, suggesting that studies planning to use personal CO exposures as a surrogate for personal $PM_{2.5}$ exposures would benefit from prior validation studies of personal exposure measurements, rather than cooking area measurements alone. In four of the five studies shown in Figure 5, the slope of $\ln(PM_{2.5})$ on $\ln(CO)$ concentrations in cooking areas was two to eight times steeper than the slope of $\ln(PM_{2.5})$ on $\ln(CO)$ exposures in the same study, which further suggests that use of cooking area concentrations to develop a model to estimate personal $PM_{2.5}$ from measurements of personal CO exposure could lead to biased estimates.

Conducting our multivariate models with untransformed CO concentrations and exposures did not change our overall results (see Tables S8 and S9). In a subset of studies, adding covariates at the individual level in the models led to modest changes (3–19%) in the explanation of variation in $\ln(PM)$ explained by $\ln(CO)$ in the fully adjusted model relative to the univariate model (see Table S10). Removing observations where dung ($n = 49$ kitchens) or wood-charcoal ($n = 42$ exposures) or coal ($n = 13$ exposures) was used as the primary fuel with biomass did not appreciably change our results (data not shown). The variance inflation factors for our independent variables did not exceed 2.5, indicating a lack of multicollinearity.

Discussion

Our results suggest that exposure to CO is not a consistently valid surrogate of exposure to $PM_{2.5}$ in settings with household air pollution, as indicated by low-to-moderate personal $PM_{2.5}$ –CO correlations [range: 0.22 ($n = 29$)–0.97 ($n = 12$); median = 0.57]. None of the multivariate regression models explained > 50% of the variation in personal $PM_{2.5}$ exposures. Further, the personal $PM_{2.5}$ –CO relationship was not transportable across different energy-use and environmental settings, suggesting that, if personal CO exposure is pursued as a surrogate measure of personal $PM_{2.5}$ exposure, a separate $PM_{2.5}$ –CO validation may be needed for each unique study setting and, within studies, potentially each season or pre-versus post-stove/fuel intervention.

We found a stronger correlation between personal $PM_{2.5}$ and CO exposures among exclusive biomass users relative to mixed fuel users ($R^2 = 0.29$ vs. 0.18), supporting previous studies (Naeher et al. 2000a; Naeher et al. 2000b; McCracken et al. 2013). This finding is consistent with results from stove emission tests in laboratory and field settings. For example, Jetter et al. (2012) observed a higher coefficient of determination for $PM_{2.5}$ versus CO emissions from biomass stoves compared with nonbiomass stoves during Water Boiling Tests (see Figure S4). This study and others describe fundamental sources of variability in combustion conditions and energy-use behaviors that limit the strength and consistency of the correlation we may expect for $PM_{2.5}$ and CO exposures and concentrations in real-world settings with biomass combustion (Zhang et al. 2000; Roden et al. 2009; Shen et al. 2010; Chen et al. 2012; Shen et al. 2013). With widespread use of multiple stoves and fuels (i.e., stove stacking), exclusive biomass use is increasingly less common (Masera and Navia 1997; Masera et al. 2000; Ruiz-Mercado et al. 2011; Rehfuess et al. 2014; Ni et al. 2016); and may reduce the number of settings in which validation studies will demonstrate CO to be a valid PM surrogate.

We observed a stronger personal $PM_{2.5}$ –CO relationship for measurements conducted in rural versus peri-urban settings ($R^2 = 0.42$ vs. 0.25; interaction p -value < 0.001), likely because densely populated peri-urban neighborhoods may have more community (i.e., solid waste burning) and regional pollution. At the same time, a stronger $PM_{2.5}$ –CO relationship for personal exposure measurements conducted in the heating season relative to the nonheating season ($R^2 = 0.47$ vs. 0.13; interaction p -value < 0.001) may reflect the greater proportion of time people spend indoors next to the fire, which is also where stationary indoor monitors are usually located. Notably, the R^2 for the personal $PM_{2.5}$ –CO exposure relationship in the heating season is almost identical to the cooking area relationship (0.47 vs. 0.46). Separately, we found that the personal $PM_{2.5}$ –CO relationship was modified significantly by season (interaction p -value = 0.02), but the cooking area $PM_{2.5}$ –CO relationship was not (interaction p -value = 0.34). This finding supports recent studies showing that personal exposures are impacted by other (i.e. noncooking) air pollution sources (Baumgartner et al. 2014;

○ Personal Exposure △ Household Concentration




Figure 5. Paired personal and cooking area $PM_{2.5}$ and CO (24- or 48-hr integrated concentrations) for (a) China (Ni et al. 2016), (b) Honduras (Peel JL, written and oral communication, spring 2016), (c) The Gambia (Dionisio et al. 2012), (d) Peru (St. Helen et al. 2015), and Peru (e) pre- and (f) postintervention (Fitzgerald et al. 2012). The R^2 and slope of the $\ln(PM_{2.5})$ - $\ln(CO)$ relationship is shown for cooking area measurements (blue) and personal exposures (black).

Huang et al. 2015; Secrest et al. 2017). These other air pollution sources impact noncooking area measurements and weaken the basis for transportability of the cooking area $PM_{2.5}$ -CO relationship to personal exposures. Using a cooking area $PM_{2.5}$ -CO relationship to estimate personal $PM_{2.5}$ from measurements of personal CO exposure may yield inaccurate results, as our graphical comparison of these two relationships from the same studies suggests (Figure 5).

The $PM_{2.5}$ -CO exposure relationships may also vary by age or gender. Although 24-hr CO and $PM_{2.5}$ exposures were measured in participants ranging from 18 months to 90 years of age, most exclusively measured adult women's exposures, highlighting the limited data on exposures in infants and young children and the need to reduce technological barriers to measuring their exposure. No studies evaluated this relationship in men, though it is unlikely that the $PM_{2.5}$ -CO relationship would be stronger in

men who, in many settings, are less likely to be the primary cooks and more likely to spend time outside of the home.

The type and range of pollutants of interest may vary depending on the health endpoint. Though the exact PM components responsible for its health impacts are unclear, there is strong and consistent evidence that both short- and long-term exposures to PM_{2.5} are associated with a range of clinical health outcomes in adults and children (WHO 2007; Chen et al. 2008; U.S. EPA 2009; Brook et al. 2010), including a number of PM exposure-response studies conducted in settings of biomass burning (Ezzati and Kammen 2001; Smith et al. 2011; Baumgartner et al. 2011; Norris et al. 2016). The evidence base for direct health impacts of CO exposure, beyond acute poisoning, is less strong. Animal studies indicate that fetal carboxyhemoglobin levels equilibrate with maternal levels (Longo 1977), and that very high maternal CO exposures are associated with adverse pregnancy outcomes, including pregnancy loss and low birth weight (Astrup et al. 1972; Garvey and Longo 1978). In epidemiologic studies, exposure to low-to-moderate CO concentrations in pregnant women was associated with reduced fetal growth in some studies (Ritz and Yu 1999; Ha et al. 2001; Gouveia et al. 2004; Salam et al. 2005) but not others (Alderman et al. 1987; Koren et al. 1991; Chen et al. 2002). Notably, a recent study that measured both personal PM_{2.5} and personal CO exposure in pregnant Tanzanian women cooking with biomass stoves found that only PM_{2.5} exposure was associated with adverse birth outcomes (Wylie et al. 2016), supporting a similar finding among pregnant women in the urban United States (Parker et al. 2005).

A strength of our pooled analysis is the inclusion of multiple independent variables that have been shown to influence the PM_{2.5}-CO relationship. Still, it is possible that inclusion of other individual- or study-level variables could further improve the ability of CO to predict PM exposures or concentrations. For example, we did not have access to detailed socio-demographic data for study participants and could not include variables for altitude, monitor placement, or stove type because these variables were collinear with study, measurement type, and fuel type, respectively, and were thus excluded from the models.

Though our systematic review revealed inconsistencies in the reporting of quality assurance and quality control protocols for PM_{2.5} and CO measurements, which are subject to systematic error and may introduce bias, we recognize that investigators have to balance the scientific, logistical, technical, and cost trade-offs in selecting an exposure metric for their study. Standardized and transparent reporting could improve the comparability of pollution measurements collected across diverse settings. Such reporting would include, for example, filter handling, collection, and transport; field and lab blank correction; duplicate precision estimates, if possible; method detection limits and instrument sensitivity; instrument flow rate precision estimates; and traceability and calibration levels of gas standards used to calibrate CO sensors. Then, the trade-offs in pollutant selection, study design, and measurement precision, stability, practicality, and number— inherent to efforts to reduce exposure misclassification for long-term air pollution exposures (McCracken et al. 2009; Pillarisetti 2016)—could be evaluated more consistently across settings. Further, improving our collective assessment of these trade-offs would bring to light more suitable and effective approaches and technologies to measure exposure, especially among young children and infants, for whom we have the least information on PM exposure (Balakrishnan et al. 2011; Clark et al. 2013).

Conclusions

Our systematic review and pooled analysis suggest that personal CO exposures are a poor surrogate of measured personal PM

exposures, even when biomass is exclusively burned. Our conclusions support those reached in recent studies reporting low PM_{2.5}-CO correlations for cooking area concentrations (Klassen et al. 2015; Bartington et al. 2016). The relationship between cooking area PM_{2.5} and CO concentrations in this review was stronger than for personal exposures, potentially due to the closer proximity of stationary monitors to the solid fuel emission source, but still the variation in ln(CO) did not explain more than 48% of the variation in ln(PM_{2.5}). Based on the evidence presented in this analysis, to use CO exposure as a surrogate for PM exposure would require repeated validation studies, especially if study conditions change over time. Lowering the barriers to PM_{2.5} exposure assessment, particularly for infants and young children, is an important direction for future research. Recent developments in portable lightweight PM_{2.5} monitors that are virtually silent and low-profile (Birch et al. 2015; Volckens et al. 2017) could expand the feasibility of PM exposure assessment to different populations and settings. Given that PM_{2.5} is likely one of the important drivers of the health effects associated with air pollution exposure, further research and development is needed to minimize PM_{2.5} measurement error, to reduce the logistical and technological challenges of large-scale PM exposure assessment, and to identify better surrogate measures of PM_{2.5} exposure and dose, potentially including internal biomarkers, for epidemiologic and intervention studies involving household air pollution.

Acknowledgments

We thank K. Ni, S. Pollard, O. Adetona, and M. Bechle for assistance with data compilation. This publication was made possible by U.S. EPA grant 83542201. Its contents are solely the responsibility of the grantee and do not necessarily represent the official views of the U.S. EPA. Further, the U.S. EPA does not endorse the purchase of any commercial products or services mentioned in the publication. J.B. was supported by Canadian Institutes of Health Research New Investigator Award 141959.

References

- Alderman BW, Baron AE, Savitz DA. 1987. Maternal exposure to neighborhood carbon monoxide and risk of low infant birth weight. *Public Health Rep* 102(4):410–414, PMID: 3112852.
- Alexander D, Linnes JC, Bolton S, Larson T. 2014. Ventilated cookstoves associated with improvements in respiratory health-related quality of life in rural Bolivia. *J Public Health* 36(3):460–466, PMID: 23965639, <https://doi.org/10.1093/pubmed/ftd086>.
- Ales LW, Mestl HE, Berger J, Zhang H, Wang S, Dong Z, et al. 2014. Indoor PM and CO concentrations in rural Guizhou, China. *Energy Sustain Dev* 21:51–59, <https://doi.org/10.1016/j.esd.2014.05.004>.
- Armstrong BG. 1998. Effect of measurement error on epidemiological studies of environmental and occupational exposures. *Occup Environ Med* 55(10):651–656, PMID: 9930084, <https://doi.org/10.1136/oem.55.10.651>.
- Armstrong BG. 2004. Exposure measurement error: consequences and design issues. *Exposure Assessment in Occupational and Environmental Epidemiology*. Nieuwenhuijsen MJ, ed. New York, NY:Oxford University Press.
- Astrup P, Trolle D, Olsen HM, Kjeldsen K. 1972. Effect of moderate carbon monoxide exposure on fetal development. *Lancet* 2(7789):1220–1222, PMID: 4117709, [https://doi.org/10.1016/S0140-6736\(72\)92270-2](https://doi.org/10.1016/S0140-6736(72)92270-2).
- Balakrishnan K, Ghosh S, Ganguli B, Sambandam S, Bruce N, Barnes DF, et al. 2013. State and national household concentrations of PM_{2.5} from solid cookfuel use: results from measurements and modeling in India for estimation of the global burden of disease. *Environ Health* 12(1):77, <https://doi.org/10.1186/1476-069X-12-77>.
- Balakrishnan K, Sambandam S, Ghosh S, Mukhopadhyay K, Vaswani M, Arora NK, et al. 2015. Household air pollution exposures of pregnant women receiving advanced combustion cookstoves in India: implications for intervention. *Ann Glob Health* 81(3):375–385, PMID: 26615072, <https://doi.org/10.1016/j.aogh.2015.08.009>.
- Balakrishnan K, Thangavel G, Ghosh S, Sambandam S, Mukhopadhyay K, Adair-Rohani H, et al., 2011. The Global Household Air Pollution Database 2011 (Version 3.0).

- Bartington SE, Bakolis I, Devakumar D, Kurmi OP, Gulliver J, Chaube G, et al. 2016. Patterns of domestic exposure to carbon monoxide and particulate matter in households using biomass fuel in Janakpur, Nepal. *Environ Pollut* 220(pt A):38–45, PMID: 27707597, <https://doi.org/10.1016/j.envpol.2016.08.074>.
- Baumgartner J, Schauer JJ, Ezzati M, Lu L, Cheng C, Patz J, et al. 2011. Patterns and predictors of personal exposure to indoor air pollution from biomass combustion among women and children in rural China. *Indoor Air* 21(6):479–488, PMID: 21692855, <https://doi.org/10.1111/j.1600-0668.2011.00730.x>.
- Baumgartner J, Zhang Y, Schauer JJ, Huang W, Wang Y, Ezzati M. 2014. Highway proximity and black carbon from cookstoves as a risk factor for higher blood pressure in rural China. *Proc Natl Acad Sci USA* 111(36):13229–13234, PMID: 25157159, <https://doi.org/10.1073/pnas.1317176111>.
- Birch A, Hystad P, Arku R, Volkens J, Miller-Lionberg D, Quinn C, et al. 2015. Household and personal air pollution monitoring in two communities in rural India using established and prototype devices. We-P-30. 34th Annual Conference of the International Society of Exposure Science, Nevada, USA.
- Bonjour S, Adair-Rohani H, Wolf J, Bruce NG, Mehta S, Prüss-Ustün A, et al. 2013. Solid fuel use for household cooking: country and regional estimates for 1980–2010. *Environ Health Perspect* 121(7):784–790, PMID: 23674502, <https://doi.org/10.1289/ehp.1205987>.
- Brook RD, Rajagopalan S, Pope CA III, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121(21):2331–2378, <https://doi.org/10.1161/CIR.0b013e3181d8e1>.
- Bruce N, McCracken J, Albalak R, Schei MA, Smith KR, Lopez V, et al. 2004. Impact of improved stoves, house construction and child location on levels of indoor air pollution exposure in young Guatemalan children. *J Expo Anal Environ Epidemiol* 14:S26–S33, PMID: 15118742, <https://doi.org/10.1038/sj.jea.7500355>.
- Carter EM, Shan M, Yang X, Li J, Baumgartner J. 2014. Pollutant emissions and energy efficiency of Chinese gasifier cooking stoves and implications for future intervention studies. *Environ Sci Technol* 48(1):6461–6467, PMID: 24784418, <https://doi.org/10.1021/es405723w>.
- Chen H, Goldberg MS, Villeneuve PJ. 2008. A systematic review of relation between long-term exposure to ambient air pollution and chronic disease. *Rev Environ Health* 23(4):243–296, PMID: 19235364.
- Chen L, Yang W, Jennison BL, Goodrich A, Omaye ST. 2002. Air pollution and birth weight in northern Nevada, 1991–1999. *Inhal Toxicol* 14(2):141–157, PMID: 12122577, <https://doi.org/10.1080/089583701753403962>.
- Chen Y, Roden CA, Bond TC. 2012. Characterizing biofuel combustion with Patterns of Real-Time Emission Data (PaRTED). *Environ Sci Technol* 46(11):6110–6117, PMID: 22533493, <https://doi.org/10.1021/es3003348>.
- Chengappa C, Edwards R, Bajpai R, Shields KN, Smith KR. 2007. Impact of improved cookstoves on indoor air quality in the Bundelkhand region in India. *Energy Sustain Dev* 11(2):33–44, [https://doi.org/10.1016/S0973-0826\(08\)60398-1](https://doi.org/10.1016/S0973-0826(08)60398-1).
- Chowdhury Z, Campanella L, Gray C, Al Masud A, Mater-Kenyon J, Pennise D, et al. 2013. Measurement and modeling of indoor air pollution in rural households with multiple stove interventions in Yunnan, China. *Atmos Environ* 67:161–169, <https://doi.org/10.1016/j.atmosenv.2012.10.041>.
- Clark ML, Peel JL, Balakrishnan K, Breyse PN, Chillrud SN, Naeher LP, et al. 2013. Health and household air pollution from solid fuel use: the need for improved exposure assessment. *Environ Health Perspect* 121(10):1120–1128, PMID: 23872398, <https://doi.org/10.1289/ehp.1206429>.
- Commodore AA, Hartinger SM, Lanata CF, Mäusezahl D, Gil AI, Hall DB, et al. 2013. A pilot study characterizing real time exposures to particulate matter and carbon monoxide from cookstove related woodsmoke in rural Peru. *Atmos Environ* 79:380–384, PMID: 24288452, <https://doi.org/10.1016/j.atmosenv.2013.06.047>.
- Cynthia AA, Edwards RD, Johnson M, Zuk M, Rojas L, Jiménez RD, et al. 2008. Reduction in personal exposures to particulate matter and carbon monoxide as a result of the installation of a Patsari improved cook stove in Michoacan, Mexico. *Indoor Air* 18(2):93–105, PMID: 18333989, <https://doi.org/10.1111/j.1600-0668.2007.00509.x>.
- Dionisio KL, Howie SR, Dominici F, Fornace KM, Spengler JD, Adegbola RA, et al. 2012. Household concentrations and exposure of children to particulate matter from biomass fuels in The Gambia. *Environ Sci Technol* 46(6):3519–3527, PMID: 22304223, <https://doi.org/10.1021/es203047e>.
- Dionisio KL, Howie S, Fornace KM, Chimah O, Adegbola RA, Ezzati M. 2008. Measuring the exposure of infants and children to indoor air pollution from biomass fuels in The Gambia. *Indoor Air* 18(4):317–327, PMID: 18422570, <https://doi.org/10.1111/j.1600-0668.2008.00533.x>.
- Dix-Cooper L, Eskenazi B, Romero C, Balmes J, Smith KR. 2012. Neurodevelopmental performance among school age children in rural Guatemala is associated with prenatal and postnatal exposure to carbon monoxide, a marker for exposure to woodsmoke. *Neurotoxicology* 33(2):246–254, PMID: 21963523, <https://doi.org/10.1016/j.neuro.2011.09.004>.
- Dutta K, Shields KN, Edwards R, Smith KR. 2007. Impact of improved biomass cookstoves on indoor air quality near Pune, India. *Energy Sustain Dev* 11(2):19–32, [https://doi.org/10.1016/S0973-0826\(08\)60397-X](https://doi.org/10.1016/S0973-0826(08)60397-X).
- Ellegård A, Egnéus H. 1993. Urban energy: exposure to biomass fuel pollution in Lusaka. *Energy Policy* 21(5):615–622, [https://doi.org/10.1016/0301-4215\(93\)90044-G](https://doi.org/10.1016/0301-4215(93)90044-G).
- Ezzati M, Kammen DM. 2001. Indoor air pollution from biomass combustion and acute respiratory infections in Kenya: an exposure-response study. *Lancet* 358(9282):619–624, PMID: 11530148, [https://doi.org/10.1016/S0140-6736\(01\)05777-4](https://doi.org/10.1016/S0140-6736(01)05777-4).
- Ezzati M, Mbinda BM, Kammen DM. 2000. Comparison of emissions and residential exposure from traditional and improved cookstoves in Kenya. *Environ Sci Technol* 34(4):578–583, <https://doi.org/10.1021/es9905795>.
- Fischer SL, Koshland CP. 2007. Daily and peak 1-h indoor air pollution and driving factors in a rural Chinese village. *Environ Sci Technol* 41(9):3121–3126, PMID: 17539514, <https://doi.org/10.1021/es060564o>.
- Fitzgerald C, Aguilar-Villalobos M, Eppler AR, Dorner SC, Rathbun SL, Naeher LP. 2012. Testing the effectiveness of two improved cookstove interventions in the Santiago de Chuco Province of Peru. *Sci Total Environ* 420:54–64, PMID: 22309740, <https://doi.org/10.1016/j.scitotenv.2011.10.059>.
- Garvey DJ, Longo LD. 1978. Chronic low level maternal carbon monoxide exposure and fetal growth and development. *Biol Reprod* 19(1):8–14, PMID: 687711, <https://doi.org/10.1095/biolreprod19.1.8>.
- GBD 2013 Risk Factors Collaborators, Forouzanfar MH, Alexander L, Anderson HR, Bachman VF, Biryukov S, et al. 2015. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 386(10010):2287–2323, PMID: 27733284, [https://doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/10.1016/S0140-6736(16)31679-8).
- Gouveia N, Bremner SA, Novaes HMD. 2004. Association between ambient air pollution and birth weight in São Paulo, Brazil. *J Epidemiol Community Health* 58:11–17, PMID: 14684720, <https://doi.org/10.1136/jech.58.1.11>.
- Guarnieri MJ, Diaz JV, Basu C, Diaz A, Pope D, Smith KR, et al. 2014. Effects of woodsmoke exposure on airway inflammation in rural Guatemalan women. *PLoS One* 9:e88455, PMID: 24625755, <https://doi.org/10.1371/journal.pone.0088455>.
- Ha EH, Hong YC, Lee BE, Woo BH, Schwartz J, Christiani DC. 2001. Is air pollution a risk factor for low birth weight in Seoul? *Epidemiology* 12(6):643–648, PMID: 11679791, <https://doi.org/10.1097/00001648-200111000-00011>.
- Hartinger SM, Commodore AA, Hattendorf J, Lanata CF, Gil AI, Verastegui H, et al. 2013. Chimney stoves modestly improved indoor air quality measurements compared with traditional open fire stoves: results from a small-scale intervention study in rural Peru. *Indoor Air* 23(4):342–352, PMID: 23311877, <https://doi.org/10.1111/jia.12027>.
- Henkle J, Mandzuk C, Emery E, Schrowe L, Sevilla-Martir J. 2010. Global health and international medicine: Honduras Stove Project. *Hisp Health Care Int* 8(1):36–46, <https://doi.org/10.1891/1540-4153.8.1.36>.
- Huang W, Baumgartner J, Zhang Y, Wang Y, Schauer JJ. 2015. Source apportionment of air pollution exposures of rural Chinese women cooking with biomass fuels. *Atmos Environ* 104:79–87, <https://doi.org/10.1016/j.atmosenv.2014.12.066>.
- Huboyo HS, Tohno S, Lestari P, Mizohata A, Okumura M. 2014. Characteristics of indoor air pollution in rural mountainous and rural coastal communities in Indonesia. *Atmos Environ* 82:343–350, <https://doi.org/10.1016/j.atmosenv.2013.10.044>.
- Jack DW, Asante KP, Wylie BJ, Chillrud SN, Whyatt RM, Ae-Ngibise KA, et al. 2015. Ghana Randomized Air Pollution and Health Study (GRAPHS): study protocol for a randomized controlled trial. *Trials* 16(1):420, PMID: 26395578, <https://doi.org/10.1186/s13063-015-0930-8>.
- Jetter J, Zhao Y, Smith KR, Khan B, Yelverton T, Decarlo P, et al. 2012. Pollutant emissions and energy efficiency under controlled conditions for household biomass cookstoves and implications for metrics useful in setting international test standards. *Environ Sci Technol* 46(19):10827–10834, PMID: 22924525, <https://doi.org/10.1021/es301693f>.
- Klasen E, Miranda JJ, Khatry S, Menya D, Gilman RH, Tielsch JM, et al. 2013. Feasibility intervention trial of two types of improved cookstoves in three resource-limited settings: study protocol for a randomized controlled trial. *Trials* 14:327, PMID: 24112419, <https://doi.org/10.1186/1745-6215-14-327>.
- Klasen EM, Wills B, Naithani N, Gilman RH, Tielsch JM, Chiang M, et al. 2015. Low correlation between household carbon monoxide and particulate matter concentrations from biomass-related pollution in three resource-poor settings. *Environ Res* 142:424–431, PMID: 26245367, <https://doi.org/10.1016/j.envres.2015.07.012>.
- Koren G, Sharav T, Pastuszak A, Garrettson LK, Hill K, Samson I, et al. 1991. A multicenter, prospective study of fetal outcome following accidental carbon monoxide poisoning in pregnancy. *Reprod toxicol* 5:397–403, PMID: 1806148, [https://doi.org/10.1016/0890-6238\(91\)90002-VV](https://doi.org/10.1016/0890-6238(91)90002-VV).
- Leavey A, Londeree J, Priyadarshini P, Puppala J, Schechtman KB, Yadama G, et al. 2015. Real-time particulate and CO concentrations from cookstoves in rural households in Udaipur, India. *Environ Sci Technol* 49(12):7423–7431, PMID: 25985217, <https://doi.org/10.1021/acs.est.5b02139>.

- Longo LD. 1977. The biological effects of carbon monoxide on the pregnant woman, fetus and newborn infant. *Am J Obstet Gynecol* 129(1):69–103, PMID: 561541, [https://doi.org/10.1016/0002-9378\(77\)90824-9](https://doi.org/10.1016/0002-9378(77)90824-9).
- Masera OR, Navia J. 1997. Fuel switching or multiple cooking fuels? Understanding inter-fuel substitution patterns in rural Mexican households. *Biomass Bioenergy* 12(5):347–361, [https://doi.org/10.1016/S0961-9534\(96\)00075-X](https://doi.org/10.1016/S0961-9534(96)00075-X).
- Masera OR, Saatkamp BD, Kammen DM. 2000. From linear fuel switching to multiple cooking strategies: a critique and alternative to the energy ladder model. *World Dev* 28(12):2083–2103, [https://doi.org/10.1016/S0305-750X\(00\)00076-0](https://doi.org/10.1016/S0305-750X(00)00076-0).
- McCracken JP, Schwartz J, Bruce N, Mittleman M, Ryan LM, Smith KR. 2009. Combining individual-and group-level exposure information: child carbon monoxide in the Guatemala woodstove randomized control trial. *Epidemiology* 20:127–136, <https://doi.org/10.1097/EDE.0b013e31818ef327>.
- McCracken JP, Schwartz J, Diaz A, Bruce N, Smith KR. 2013. Longitudinal relationship between personal CO and personal PM_{2.5} among women cooking with wood-fired cookstoves in Guatemala. *PLoS One* 8(2):e55670, PMID: 23468847, <https://doi.org/10.1371/journal.pone.0055670>.
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. 2015. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 4:1, PMID: 25554246, <https://doi.org/10.1186/2046-4053-4-1>.
- Mukhopadhyay R, Sambandam S, Pillariseti A, Jack D, Mukhopadhyay K, Balakrishnan K, et al. 2012. Cooking practices, air quality, and the acceptability of advanced cookstoves in Haryana, India: an exploratory study to inform large-scale interventions. *Glob Health Action* 5(1):1–13, PMID: 22989509, <https://doi.org/10.3402/gha.v5i0.19016>.
- Muralidharan V, Sussan TE, Limaye S, Koehler K, Williams DL, Rule AM, et al. 2015. Field testing of alternative cookstove performance in a rural setting of Western India. *Int J Environ Res Public Health* 12(2):1773–1787, PMID: 25654775, <https://doi.org/10.3390/ijerph120201773>.
- Naeher LP, Brauer M, Lipsett M, Zelikoff JT, Simpson CD, Koenig JQ, et al. 2007. Woodsmoke health effects: a review. *Inhal Toxicol* 19(1):67–106, PMID: 17127644, <https://doi.org/10.1080/08958370600985875>.
- Naeher LP, Leaderer BP, Smith KR. 2000a. Particulate matter and carbon monoxide in highland Guatemala: indoor and outdoor levels from traditional and improved wood stoves and gas stoves. *Indoor Air* 10:200–205, PMID: 10979201.
- Naeher LP, Smith KR, Leaderer BP, Mage D, Grajeda R. 2000b. Indoor and outdoor PM_{2.5} and CO in high-and low-density Guatemalan villages. *J Expo Anal Environ Epidemiol* 10(3):544–551, PMID: 11140438, <https://doi.org/10.1034/j.1600-0668.2000.010003200.x>.
- Naeher LP, Smith KR, Leaderer BP, Neufeld L, Mage DT. 2001. Carbon monoxide as a tracer for assessing exposures to particulate matter in wood and gas cookstove households of highland Guatemala. *Environ Sci Technol* 35(3):575–581, PMID: 11351731, <https://doi.org/10.1021/es991225g>.
- Ni K, Carter EM, Schauer JJ, Ezzati M, Zhang Y, Niu H, et al. 2016. Seasonal variation in ambient, household, and personal air pollution exposures of women using traditional wood stoves in the Tibetan Plateau: baseline assessment for an energy intervention study. *Environ Int* 94:449–457, <https://doi.org/10.1016/j.envint.2016.05.029>.
- NIH (National Institutes of Health). 2015. “Household Air Pollution (HAP) Health Outcomes Trial (UM1).” RFA-HL-16-012, <https://grants.nih.gov/grants/guide/rfa-files/RFA-HL-16-012.html> [accessed 22 June 2017].
- Norris C, Goldberg MS, Marshall JD, Valois MF, Pradeep T, Narayanswamy M, et al. 2016. A panel study of the acute effects of personal exposure to household air pollution on ambulatory blood pressure in rural Indian women. *Environ Res* 147:331–342, PMID: 26928412, <https://doi.org/10.1016/j.envres.2016.02.024>.
- Northcross A, Chowdhury Z, McCracken J, Canuz E, Smith KR. 2010. Estimating personal PM_{2.5} exposures using CO measurements in Guatemalan households cooking with wood fuel. *J Environ Monit* 12(14):873–878, PMID: 20383368, <https://doi.org/10.1039/b916068j>.
- Northcross AL, Hwang N, Balakrishnan K, Mehta S. 2015. Assessing exposures to household air pollution in public health research and program evaluation. *Ecohealth* 12(1):57–67, PMID: 25380652, <https://doi.org/10.1007/s10393-014-0990-3>.
- Ochieng CA, Vardoulakis S, Tonne C. 2013. Are rocket mud stoves associated with lower indoor carbon monoxide and personal exposure in rural Kenya? *Indoor Air* 23(1):14–24, PMID: 22563898, <https://doi.org/10.1111/j.1600-0668.2012.00786.x>.
- Park E, Lee K. 2003. Particulate exposure and size distribution from wood burning stoves in Costa Rica. *Indoor Air* 13(3):253–259, PMID: 12950588.
- Parker JD, Woodruff TJ, Basu R, Schoendorf KC. 2005. Air pollution and birth weight among term infants in California. *Pediatrics* 115(1):121–128, PMID: 15629991, <https://doi.org/10.1542/peds.2004-0889>.
- Pearce JL, Aguilar-Villalobos M, Rathbun SL, Naeher LP. 2009. Residential exposures to PM_{2.5} and CO in Cusco, a high-altitude city in the Peruvian Andes: a pilot study. *Arch Environ Occup Health* 64(4):278–282, PMID: 20007125, <https://doi.org/10.1080/19338240903338205>.
- Peel JL, Baumgartner J, Wellenius GA, Clark ML, Smith KR. 2015. Are randomized trials necessary to advance epidemiologic research on household air pollution? *Curr Epidemiol Rep* 2(4):263–270, <https://doi.org/10.1007/s40471-015-0054-4>.
- Pillariseti A. 2016. Long-term PM_{2.5} monitoring in kitchens cooking with wood: implications for measurement strategies. In “Inspecting What you Expect: Applying Modern Tools and Techniques to Evaluate the Effectiveness of Household Energy Interventions.” PhD Dissertation. Accessible via University of California, Berkeley library http://guides.lib.berkeley.edu/dissertations_theses.
- Pope CA III, Dockery DW. 2006. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 56(6):709–742, <https://doi.org/10.1080/10473289.2006.10464485>.
- Pope D, Diaz E, Smith-Sivertsen T, Lie RT, Bakke P, Balmes JR, et al. 2015. Exposure to household air pollution from wood combustion and association with respiratory symptoms and lung function in nonsmoking women: results from the RESPIRE trial, Guatemala. *Environ Health Perspect* 123:285–292, PMID: 25398189, <https://doi.org/10.1289/ehp.1408200>.
- Rehfuess EA, Puzolo E, Stanistreet D, Pope D, Bruce NG. 2014. Enablers and barriers to large-scale uptake of improved solid fuel stoves: a systematic review. *Environ Health Perspect* 122(2):120–130, PMID: 24300100, <https://doi.org/10.1289/ehp.1306639>.
- Riojas-Rodriguez R, Schilman A, Marron-Mares AT, Masera O, Li Z, Romanoff L, et al. 2011. Impact of the improved Patsari biomass stove on urinary polycyclic aromatic hydrocarbon biomarkers and carbon monoxide exposures in rural Mexican women. *Environ Health Perspect* 119(9):1301–1307, PMID: 21622083, <https://doi.org/10.1289/ehp.1002927>.
- Ritz B, Yu F. 1999. The effect of ambient carbon monoxide on low birth weight among children born in southern California between 1989 and 1993. *Environ Health Perspect* 107(1):17–25, PMID: 9872713, <https://doi.org/10.1289/ehp.9910717>.
- Roden CA, Bond TC, Conway S, Pinel ABO, MacCarty N, Still D. 2009. Laboratory and field investigations of particulate and carbon monoxide emissions from traditional and improved cookstoves. *Atmos Environ* 43(6):1170–1181, <https://doi.org/10.1016/j.atmosenv.2008.05.041>.
- Röllin HB, Mathee A, Bruce N, Levin J, Von Schirnding YER. 2004. Comparison of indoor air quality in electrified and un-electrified dwellings in rural South African villages. *Indoor Air* 14(3):208–216, PMID: 15104789, <https://doi.org/10.1111/j.1600-0668.2004.00238.x>.
- Rosa G, Majorin F, Boisson S, Barstow C, Johnson M, Kirby M, et al. 2014. Assessing the impact of water filters and improved cook stoves on drinking water quality and household air pollution: a randomised controlled trial in Rwanda. *PLoS One* 9(3):e91011, <https://doi.org/10.1371/journal.pone.0091011>.
- Ruiz-Mercado I, Masera O, Zamora H, Smith KR. 2011. Adoption and sustained use of improved cookstoves. *Energy Policy* 39(12):7557–7566, <https://doi.org/10.1016/j.enpol.2011.03.028>.
- Saksena S, Prasad R, Pal RC, Joshi V. 1992. Patterns of daily exposure to TSP and CO in the Garhwal Himalaya. *Atmos Environ. Part A. General Topics* 26(11):2125–2134, [https://doi.org/10.1016/0960-1686\(92\)90096-4](https://doi.org/10.1016/0960-1686(92)90096-4).
- Saksena S, Subida R, Büttner L, Ahmed L. 2007. Indoor air pollution in coastal houses of southern Philippines. *Indoor Built Environ* 16(2):159–168, <https://doi.org/10.1177/1420326X07076783>.
- Salam MT, Millstein J, Li YF, Lurmann FW, Margolis HG, Gilliland FD. 2005. Birth outcomes and prenatal exposure to ozone, carbon monoxide, and particulate matter: results from the Children’s Health Study. *Environ Health Perspect* 113(11):1638–1644, PMID: 16263524, <https://doi.org/10.1289/ehp.8111>.
- Secret MH, Schauer JJ, Carter E, Baumgartner J. 2017. Particulate matter chemical component concentrations and sources in settings of household solid fuel use. *Indoor Air*, PMID: 28401994, <https://doi.org/10.1111/ina.12389>.
- Shen G, Yang Y, Wang W, Tao S, Zhu C, Min Y, et al. 2010. Emission factors of particulate matter and elemental carbon for crop residues and coals burned in typical household stoves in China. *Environ Sci Technol* 44(18):7157–7162, PMID: 20735038, <https://doi.org/10.1021/es101313y>.
- Smith KR. 1993. Fuel combustion, air pollution exposure, and health: the situation in developing countries. *Annu Rev Energy Environ* 18:529–566, <https://doi.org/10.1146/annurev.energy.18.1.529>.
- Smith KR, McCracken JP, Thompson L, Edwards R, Shields KN, Canuz E, et al. 2010. Personal child and mother carbon monoxide exposures and kitchen levels: methods and results from a randomized trial of woodfired chimney cookstoves in Guatemala (RESPIRE). *J Expo Sci Environ Epidemiol* 20(5):406–416, PMID: 19536077, <https://doi.org/10.1038/jes.2009.30>.
- Smith KR, McCracken JP, Weber MW, Hubbard A, Jenny A, Thompson LM, et al. 2011. Effect of reduction in household air pollution on childhood pneumonia in Guatemala (RESPIRE): a randomised controlled trial. *Lancet* 378(9804):1717–1726, PMID: 22078686, [https://doi.org/10.1016/S0140-6736\(11\)60921-5](https://doi.org/10.1016/S0140-6736(11)60921-5).
- Smith-Sivertsen T, Diaz E, Pope D, Lie RT, Diaz A, McCracken J. 2009. Effect of reducing indoor air pollution on women’s respiratory symptoms and lung function: the RESPIRE Randomized Trial, Guatemala. *Am J Epidemiol* 170(2):211–220, PMID: 19443665, <https://doi.org/10.1093/aje/kwp100>.

- Spiegelman D. 2010. Approaches to uncertainty in exposure assessment in environmental epidemiology. *Annu Rev Public Health* 31:149–163, PMID: 20070202, <https://doi.org/10.1146/annurev.publhealth.012809.103720>.
- St. Helen G, Aguilar-Villalobos M, Adetona O, Cassidy B, Bayer CW, Hendry R, et al. 2015. Exposure of pregnant women to cookstove-related household air pollution in urban and periurban Trujillo, Peru. *Arch Environ Occup Health* 70(1):10–18, PMID: 24215174, <https://doi.org/10.1080/19338244.2013.807761>.
- Thompson LM, Bruce N, Eskenazi B, Diaz A, Pope D, Smith KR. 2011. Impact of reduced maternal exposures to wood smoke from an introduced chimney stove on newborn birth weight in rural Guatemala. *Environ Health Perspect* 119(10):1489–1494, PMID: 21652290, <https://doi.org/10.1289/ehp.1002928>.
- Thorsson S, Holmer B, Andjelic A, Lindén J, Cimerman S, Barregard L. 2014. Carbon monoxide concentrations in outdoor wood-fired kitchens in Ouagadougou, Burkina Faso—implications for women’s and children’s health. *Environ Monit Assess* 186(7):4479–4492, PMID: 24652378, <https://doi.org/10.1007/s10661-014-3712-y>.
- Tielsch JM, Katz J, Zeger SL, Khattry SK, Shrestha L, Breyse P, et al. 2014. Designs of two randomized, community-based trials to assess the impact of alternative cookstove installation on respiratory illness among young children and reproductive outcomes in rural Nepal. *BMC Public Health* 14:1271–1281, PMID: 25511324, <https://doi.org/10.1186/1471-2458-14-1271>.
- U.S. EPA (U.S. Environmental Protection Agency). 2009. Integrated Science Assessment for Particulate Matter (Final Report). EPA/600/R-08/139F. Washington, DC:U.S. EPA.
- U.S. EPA. 2016. Quality Assurance Guidance Document 2.12. Monitoring PM_{2.5} in Ambient Air Using Designated Reference or Class I Equivalent Methods. EPA-454/B-16-001. Washington, DC:U.S. EPA.
- Van Tulder M, Furlan A, Bombardier C, Bouter L; Editorial Board of the Cochrane Collaboration Back Review Group, 2003. Updated method guidelines for systematic reviews in the Cochrane Collaboration Back Review Group. *Spine (Phila Pa 1976)* 28(12):1290–1299, PMID: 12811274, <https://doi.org/10.1097/01.BRS.0000065484.95996.AF>.
- Volckens J, Quinn C, Leith D, Mehaffy J, Henry CS, Miller-Lionberg D. 2017. Development and evaluation of an ultrasonic personal aerosol sampler (UPAS). *Indoor Air* 27(2):409–416, PMID: 27354176, <https://doi.org/10.1111/ina.12318>.
- WHO (World Health Organization). 2007. Health Relevance of Particulate Matter from Various Sources. Copenhagen, Denmark:WHO, Regional Office for Europe. http://www.euro.who.int/__data/assets/pdf_file/0007/78658/E90672.pdf [accessed May 2016].
- WHO. 2014. WHO Indoor Air Quality Guidelines: Household Fuel Combustion. Geneva, Switzerland:WHO.
- Wylie BJ, Kishashu Y, Matechi E, Zhou Z, Coull B, Abioye AI, et al. 2016. Maternal exposure to carbon monoxide and fine particulate matter during pregnancy in an urban Tanzanian cohort. *Indoor Air* 27(1):136–146, <https://doi.org/10.1111/ina.12289>.
- Yip F, Christensen B, Sircar K, Naeher L, Bruce N, Pennise D, et al. 2017. Assessment of traditional and improved stove use on household air pollution and personal exposures in rural western Kenya. *Environ Int* 99:185–191, PMID: 27923586, <https://doi.org/10.1016/j.envint.2016.11.015>.
- Zhang JJ, Smith KR. 2007. Household air pollution from coal and biomass fuels in China: measurements, health impacts, and interventions. *Environ Health Perspect* 115(6):848–855, PMID: 17589590, <https://doi.org/10.1289/ehp.9479>.
- Zhang J, Smith KR, Ma Y, Jiang F, Qi W, Khalil MAK, et al. 2000. Greenhouse gases and other airborne pollutants from household stoves in China: a database for emission factors. *Atmos Environ* 34(26):4537–4549, [https://doi.org/10.1016/S1352-2310\(99\)00450-1](https://doi.org/10.1016/S1352-2310(99)00450-1).