

RESEARCH

Open Access



Tissue transglutaminase in astrocytes is enhanced by inflammatory mediators and is involved in the formation of fibronectin fibril-like structures

Nathaly Espitia Pinzón¹, John J. P. Brevé¹, John G. J. M. Bol¹, Benjamin Drukarch¹, Wia Baron^{2†} and Anne-Marie van Dam^{1*†}

Abstract

Background: During multiple sclerosis (MS) lesion formation, inflammatory mediators are produced by microglial cells and invading leukocytes. Subsequently, hypertrophic astrocytes fill the lesion and produce extracellular matrix (ECM) proteins that together form the astroglial scar. This is beneficial because it seals off the site of central nervous system (CNS) damage. However, astroglial scarring also forms an obstacle that inhibits remyelination of brain lesions. This is possibly an important cause for incomplete remyelination of the CNS in early stage MS patients and for failure of remyelination when the disease progresses. Tissue transglutaminase (TG2), a Ca²⁺-dependent enzyme that can cross-link proteins, appears in astrocytes in inflammatory MS lesions and may contribute to the rearrangement of ECM protein deposition and aggregation.

Methods: The effect of different inflammatory mediators on TG2 and fibronectin, an ECM protein, protein levels was examined in primary rat microglia and astrocytes by western blotting. Also, TG2 activity was analyzed in primary rat astrocytes by a TG activity assay. To determine the role of TG2 in the deposition and cross-linking of fibronectin, a TG2 inhibitor and TG2 knockdown astrocytes were used.

Results: Our data show that under inflammatory conditions in vitro, TG2 production is enhanced in astrocytes and microglia. We observed that in particular, astrocytes produce fibronectin that can be cross-linked and aggregated by exogenous TG2. Moreover, inflammatory stimulus-induced endogenously produced TG2 is involved in the appearance of morphological fibril-like fibronectin deposits but does not lead to cross-linked fibronectin aggregates.

Conclusions: Our in vitro observations suggest that during MS lesion formation, when inflammatory mediators are produced, astrocyte-derived TG2 may contribute to ECM rearrangement, and subsequent astroglial scarring.

Keywords: Tissue transglutaminase, Neuroinflammation, Astrocytes, Astroglial scarring, Extracellular matrix, Fibronectin

Background

In multiple sclerosis (MS), a chronic inflammatory, demyelinating disease of the central nervous system (CNS), various classes of inflammatory white matter lesions can be identified [1, 2]. Active white matter lesions appear with a massive influx of leukocytes and ongoing demyelination. In

these lesions, e.g., inflammatory cytokines and chemokines are produced by activated microglial cells and invading leukocytes [1, 3–6]. In chronic MS lesions, infiltrating cells, particularly macrophages, remain present at the rim of the lesion where demyelination is still ongoing [7]. Simultaneously, astrogliosis appears in the lesion which mainly consists of astrocytes with a hypertrophic phenotype [8–11]. The formed astroglial scar is beneficial because it seals off inflammation in the CNS into focal areas [12]. However, the scar also impedes remyelination by inhibiting the migration and differentiation of

* Correspondence: amw.vandam@vumc.nl

†Equal contributors

¹Department Anatomy and Neurosciences, VU University Medical Center, Amsterdam Neuroscience, De Boelelaan 1108, 1081, HZ, Amsterdam, The Netherlands

Full list of author information is available at the end of the article



oligodendrocyte precursor cells [13]. Remyelination in MS lesions is therefore not very effective [14–16], which might contribute to the chronic neurodegenerative character of this disease with progressive loss of motor, sensory, and cognitive functions [17].

During astrogliosis, the extracellular matrix (ECM) is largely modified, which is reflected by an enhanced production of ECM proteins by astrocytes [18–22]. Previous studies, under in vitro conditions, in MS animal models and in brain material from patients already showed changes in the production and deposition of chondroitin sulfate proteoglycans, a family of ECM proteoglycans [23, 24]. Furthermore, the ECM proteins fibronectin and laminin have been shown to be more expressed by astrocytes [25–28] and aggregated in chronically demyelinated multiple sclerosis lesions [29], possibly contributing to the non-regenerative nature of these lesions [30, 31].

Tissue transglutaminase (TG2) is a Ca^{2+} -dependent enzyme with various catalytic functions, including protein deamidation, transamidation, and cross-linking [32–38]. TG2 is localized intracellularly [39], on the cell surface and in the ECM [40, 41]. In the ECM, TG2 is capable of cross-linking a wide range of proteins, which are important in ECM deposition and stabilization [42, 43]. TG2 can also directly, i.e., non-enzymatically, interact with fibronectin, an important ECM protein in tissue repair processes, and various beta-integrins, thereby mediating cell-ECM interactions [44–46]. TG2 is therefore thought to play an important role in various physiological and pathological situations including inflammation and fibrosis [36, 47] and can possibly contribute to the process of astroglial scarring in the CNS of MS patients and other brain injuries.

We have previously shown the appearance of TG2 immunoreactivity in astrocytes in active and chronic active MS lesions [48] and in demyelinating areas in the mouse brain after cuprizone treatment [49]. Moreover, TG2 expression in astrocytes is regulated by some inflammatory mediators as are present in MS lesions [50]. In the present study, we questioned whether a wide array of inflammatory mediators can affect TG2 production in astrocytes and possibly also in microglia which can be a source of ECM proteins during the formation of the astroglial scar [51–56]. We also questioned whether TG2 contributes to the production, deposition, and cross-linking of the ECM protein fibronectin under inflammatory conditions. A better understanding of the inflammatory regulation of TG2 and its role in the production, deposition, and cross-linking of ECM proteins by astrocytes and possibly microglia could be of therapeutic interest to overcome astroglial scarring and promote a remyelinating milieu.

Methods

Primary astrocytes and microglia

Primary rat glial cells were isolated from cerebral cortices of 2-day-old Wistar rats (Harlan CPB, Zeist, The Netherlands), as described previously [57], and approved by the Animal Experiment Committee of the VU University Medical Center (ID FGA 11-03). The meninges and blood vessels were removed from the cortices. Cortices were mechanically homogenized in Dulbecco's Modified Eagle's Medium-F10 (Gibco, Life Technologies, Breda, The Netherlands), supplemented with 10% *v/v* heat-inactivated fetal calf serum (Gibco), 2 mM L-glutamine (Sigma-Aldrich, St. Louis, MO, USA), 50 U/ml penicillin (Sigma-Aldrich), and 50 $\mu\text{g}/\text{ml}$ streptomycin (Gibco) to make a cell suspension. Dissociated cells from 2 to 3 pups were plated in poly-L-lysine (PLL, 15 $\mu\text{g}/\text{ml}$ (2 $\mu\text{g}/\text{cm}^2$; Sigma-Aldrich) coated T75 culture flasks (Nunc, Hamstrup, Denmark) and incubated at 37 °C in humidified air containing 5% CO_2 . The medium was changed at days 1, 6, and 8 after seeding. After 10 days in culture, microglia and astrocytes were separated by shaking the flasks on a rotary platform (Heidolph Unimax 2010) at 230 rpm for 16 h. Astrocytes were further purified by treatment with 5 mM leucine methyl ester (Sigma-Aldrich) in serum-free medium, for 24 h at 37 °C. Microglia were cultured in fresh medium mixed with conditioned medium (1:1 ratio) collected from the mixed cell culture before separation. The purity of astrocyte cultures was between 80 and 90%, as described previously [58], determined by immunofluorescent staining using glial fibrillary acidic protein (GFAP) antibody (1:6000; DAKO, Glostrup, Denmark). Purity of primary microglial cultures was between 90 and 94% as determined by immunofluorescent staining using Iba1 antibody (1:1000; WAKO Chemicals USA).

Treatments

Astrocytes and microglia cultured on 2 $\mu\text{g}/\text{cm}^2$ PLL-coated 6-well plates (Nunc) were incubated for 48 h in medium alone (control) or with lipopolysaccharide (LPS, *E. coli* 055-B5, Difco, 100 ng/ml), human recombinant (rec) transforming growth factor (TGF)- β 1 (BioLegend), human rec TGF- β 2 (BioLegend), rat rec interleukin (IL)-1 β (Glaxo), rat rec tumor necrosis factor (TNF)- α (Biolegend), rat rec TNF- α +IL-1 β , rat rec interleukin (IL)-4 (BioVision), rat rec interleukin (IL)-6 (gift from Steve Poole), or rat rec interleukin (IL)-10 (Pharmingen) (all 20 ng/ml). In addition, astrocytes and ECM deposited by astrocytes were treated for 48 h with TNF+IL-1 β in the absence or presence of 10 μM of the bona fide dihydroisoxazole TG2 activity inhibitor ERW1041E (Quinolin-3-ylmethyl(S)-2-(((S)-3-bromo-4,5-dihydroisoxazol-5-yl)methyl)carbamoyl)pyrrolidine-1-carboxylate) diluted in 0.1% (*v/v*) dimethyl sulfoxide (DMSO); kind gift from C. Khoshla, Stanford University, USA) [59–61]. Astrocytes were alternatively treated with

exogenous recombinant guinea pig TG2 (Sigma-Aldrich) in pathophysiological concentrations (0.13 and 1.3 μM) according to previously published studies [62, 63] for 48 h. Additionally, deposited ECM, after removal of astrocytes, was treated with 0.64 μM exogenous recombinant guinea pig TG2 (Sigma-Aldrich) for 16 h combined with a pre-incubation with a selective inhibitor of TG2 (Z-DON, 1 μM , Zedira) diluted in 0.001% DMSO or only with DMSO for 30 min (min) at 37 °C. In some experiments, astrocytes were cultured in 8-well Lab-Tek Permanox chamber slides (Nunc) or 6-well plates (Nunc) coated with 2 $\mu\text{g}/\text{cm}^2$ laminin (mixture of laminin-1 and laminin-2, from Sigma-Aldrich) instead of PLL.

Lentiviral downregulation of TG2

TG2 was downregulated in primary rat astrocytes by lentiviral transduction with TG2 specific shRNA. Therefore, astrocytes were plated on a laminin (2 $\mu\text{g}/\text{cm}^2$; Sigma-Aldrich) coated 12-well plate (0.2×10^6 cells/well, Nunc). The following day, 0.2×10^6 infectious units of virus (IFU) of lentiviral particles (Santa Cruz, sc-270266-V for rat TG2 shRNA, or sc-108080 for control scrambled shRNA) were added to the cells. The next day, the same amount of lentiviral particles was added to the cells. The medium was changed the following day, and the cells were left for 5 days before treatment with TNF- α +IL-1 β .

Sample collection

Whole cell lysates

Whole cell lysates were obtained by homogenizing the cells in ice-cold lysis buffer containing 20 mM tris-HCl pH 7.5, 137 mM NaCl, 1 mM ethylenediaminetetraacetic acid, 1% NP40, 1% sodium deoxycholate, 0.1% sodium dodecyl sulfate (SDS), 50 mM dithiothreitol (DTT), 100 μM phenylmethanesulfonyl fluoride (PMSF), 7.5 μM pepstatin-A, 10 μM leupeptin, and 0.75 μM aprotinin (all from Sigma-Aldrich).

Transglutaminase activity lysates

For transglutaminase (TG) activity measurements, cells were collected in tris-buffered saline (TBS), pH 7.5, containing 100 μM PMSF, 7.5 μM pepstatin-A, 10 μM leupeptin, and 0.75 μM aprotinin (all from Sigma-Aldrich).

ECM lysates

For the collection of ECM samples, cells were removed after washing with TBS at room temperature (RT), by incubation (5 min) with 50 mM NH_4OH + 0.05% Triton-100 (Sigma-Aldrich). This was followed by incubation (5 min) with 50 mM NH_4OH . Any remaining cell fractions were washed away by repeated wash steps with TBS. The ECM was incubated for 1 h at 37 °C with DNase (10 U/ml,

Promega) in a buffer (pH 7.9) containing 4.84 g/L tris, 0.58 g/L NaCl, 0.57 g/L MgCl_2 , and 1.95 g/L CaCl_2 . The ECM was again washed with TBS and collected in ice-cold lysis buffer as described for the whole cell lysates.

Sample preparation

After sample collection, homogenates of whole cell lysate samples and TG activity samples were sonicated (Branson "Sonifier 250"; output 1, duty 30%, 8 pulses) and cleared by centrifugation (20,000 g for 10 min at 4 °C), and protein concentrations of supernatants were determined by the BCA method (Pierce Biotechnology, Perbio Science, Etten-Leur, The Netherlands). Samples were heated for 10 min at 95 °C. ECM samples were not subjected to sonication or heating to maintain the integrity of the ECM aggregates.

Western blotting

Equal amounts of protein (20 μg) were subjected to 8% SDS-polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane (Li-Cor Biosciences, Lincoln, NE, USA). Membranes were blocked at RT with Odyssey blocking buffer (Li-Cor Biosciences) diluted 1:1 with TBS. Membranes were incubated overnight at 4 °C with mouse anti-TG2 (ab3, Neomarkers 1:2000), rabbit anti-fibronectin (Neomarkers, 1:1000), or mouse anti- β -actin (Abcam, 1:5,000) antibodies. After several washes with TBS containing 0.1% triton (TBS-T), blots were incubated for 1 h at RT with corresponding goat anti-rabbit or donkey anti-mouse IRDye 800CW IgG's (1:10,000, Li-Cor Biosciences) for subsequent antigen detection. All antibody solutions were prepared in the Odyssey blocking buffer (Li-Cor Biosciences) diluted 1:1 with TBS-T. After washing with TBS-T and TBS, membranes were scanned for fluorescence emission at 800 nm, using an Odyssey infrared imaging system (Li-Cor Biosciences). Bands were visualized, and their signal intensities were measured using the Odyssey Sa Infrared scanning software (version 1.1, Li-Cor Biosciences).

Transglutaminase activity assay

TG activity was measured using the commercially available TG Covtest Transglutaminase Colorimetric Microassay (Covalab, Villeurbanne, France), which uses immobilized CBZ-Gln-Gly as the first substrate and biotinylated-cadaverine (biotin-Cd) as second substrate of the enzyme. To determine the amount of TG2 activity among the TG activity measured, samples were pre-incubated with a selective inhibitor of TG2 (Z-DON, 1 μM , Zedira) diluted in 0.001% DMSO or only with DMSO for 30 min at RT. The assay was performed following the manufacturer's instructions [64]. In brief, 10 μg /well of each sample was incubated with 50 μl /well of biotin-Cd solution containing CaCl_2 for 1 h at 37°C on the CBZ-Gln-Gly coated plates. At the end of the incubation period, plates were washed three times with TBS (pH 7.5) containing 0.1% Tween 20.

Then, 100 μ l/well of streptavidin-labeled horseradish peroxidase (HRP) diluted to 1:2000 was added to the wells and incubated for 1 h at RT. After washing, peroxidase activity was revealed using 100 μ l/well of 0.01% H_2O_2 as HRP substrate and (0.1 mg/ml) tetramethylbenzidine as electron acceptor (chromogen). The reaction was stopped by the addition of 2.5 N H_2SO_4 , and TG activity was detected by absorbance measurement of streptavidin-labeled peroxidase activity in each well on a microplate reader (SpectraMax 250, Molecular Devices, Sunnyvale, CA, USA) at 450 nm. Guinea pig TG2 (T5398, Sigma-Aldrich) was used as a standard. One unit of guinea pig TG2 will catalyze the formation of 1.0 μ mole of hydroxamate per min from Na-Z-Gln-Gly and hydroxylamine at pH 6.0 at 37 °C.

Immunocytochemistry

Astrocytes (4.0×10^4 /well) were plated on laminin (2 μ g/ cm^2 , Sigma-Aldrich) or PLL (2 μ g/ cm^2 , Sigma-Aldrich) coated 8-well Lab-Tek Permanox chamber slides (Nunc) and treated with TGF- β 1 (for ECM stainings only) or a combination of TNF- α + IL-1 β cytokines (20 ng/ml each) for 48 h. Alternatively, exogenous recombinant guinea pig TG2 (0.13 and 1.3 μ M, Sigma-Aldrich) was added for 48 h to untreated astrocytes. Astrocytes were fixed in ice-cold methanol for 10 mins and washed in TBS. Non-specific binding was blocked with 3% bovine serum albumin in TBS containing 0.05% triton (TBS-T, pH 7.6) for 30 min at RT. Subsequently, astrocytes were double-labeled with mouse anti-TG2 (ab1, Neomarkers, 1:300) and the astrocytic marker rabbit anti-GFAP (DAKO, 1:4000) or with rabbit anti-fibronectin (Millipore, 1:200) to examine TG2 localization in astrocytes and any colocalization with fibronectin, respectively. Triton was alternatively omitted from all the incubation steps during double stainings to detect TG2 and fibronectin on the cell surface. Prior to ECM stainings, astrocytes were removed as described before. Triton was not omitted for the staining of the ECM. Secondary antibodies used were the appropriate Alexa Fluor[®] (488 or 594)-conjugated IgG's (Molecular Probes, 1:400). After washing in TBS, astrocytes were cover-slipped in polyvinyl alcohol mounting medium with DABCO (Sigma-Aldrich) and examined using a confocal microscope (Leica TSC-SP2-AOBS; Leica Microsystems, Wetzlar, Germany).

Filter trap assay

ECM lysates were treated for 30 min at RT with 150 mM DTT to break down disulfide bonds and to obtain stable aggregates. Equal amounts of ECM samples (12 μ g) were applied to a cellulose acetate membrane (GE water and process technologies) using a filter trap assay apparatus (FTA, Bio-Dot SF microfiltration apparatus, Bio-Rad). This apparatus was coupled to a vacuum-pump in order to wash through the monomers and dimers of ECM

aggregates and trap aggregates larger than the pore-size of 5 μ m of the membrane. Approximately 5 min after sample application, the membrane was washed in PBS and incubated overnight at 4 °C with rabbit anti-fibronectin (Neomarkers, 1:1000) antibodies. For subsequent antigen detection, blots were incubated for 2 h with corresponding goat anti-rabbit IRDye 800CW IgG's (1:10,000, Li-Cor Biosciences). After washing with TBS-T and TBS, membranes were scanned for fluorescence emission at 800 nm, using the same Odyssey infrared imaging system (Li-Cor Biosciences) as used for western blotting.

Statistical analysis

Normal distribution of the data was tested using the Shapiro-Wilk procedure. One-way analysis of variance was performed followed by Dunnett's post hoc test for multiple comparisons by using IBM SPSS software, version 20.0 (IBM Corp., NY, USA). $P < 0.05$ was considered to represent statistically significant differences, and a P value between 0.05 and 0.1 was regarded as a statistical trend.

Results

Astrocytes had higher TG2 and fibronectin protein levels than microglia

We first examined whether primary rat astrocytes and microglia produced TG2 and fibronectin. Under control conditions and after treatment with LPS, astrocytes expressed more TG2 and fibronectin than microglia (Additional file 1). Upon subsequent treatment of astrocytes and microglia with a variety of inflammatory mediators, TG2 protein levels were significantly increased after LPS, TNF- α +IL-1 β , or IL-4 treatment ($p < 0.001$ for all, Fig. 1a, b). Little protein was found in untreated microglia. Incubation of microglia with the various mediators resulted in significant changes in TG2 protein levels (Fig. 1d, e). LPS or TNF- α +IL-1 β treatment resulted in a significant increase in TG2 protein level ($p < 0.01$ for LPS and $p = 0.001$ for TNF- α +IL-1 β , Fig. 1d, e). Moreover, IL-4 treatment showed a trend ($p = 0.080$) toward an increase in TG2 protein level (Fig. 1e). Other conditions studied, i.e., TGF- β 1, TGF- β 2, IL-1 β , TNF- α , IL-6, and IL-10, did not show significant effects on TG2 protein level in astrocytes or microglia (Fig. 1a–e). Double bands at approximately 78 kDa were visible and quantified for TG2 in most conditions studied. This has been shown by others as well [65, 66] and could represent the full-length TG2 protein and the short TG2 protein that is truncated at the 3' end [67]. Fibronectin levels were significantly increased in astrocytes after treatment with TGF- β 1 ($P < 0.01$, Fig. 1a, c), and TGF- β 2 showed a trend ($P = 0.062$, Fig. 1c) toward an increased fibronectin level. Other conditions studied, i.e., LPS, IL-1 β , TNF- α , TNF- α +IL-1 β , IL-4, IL-6, and IL-10, did not result in significant effects on fibronectin protein levels in astrocytes (Fig. 1a, c). Moreover, none of the treatments affected the level of




Fig. 1 TG2 and fibronectin protein levels in astrocytes and microglia. Primary rat astrocytes and microglia were treated for 48 h with a range of inflammatory mediators. TG2 and fibronectin (FN) protein levels in astrocytes were detected by western blotting (**a**), which were subsequently semi-quantified (**b**, **c**). TG2 and fibronectin (FN) protein levels in microglia were detected by western blotting (**d**), which were subsequently semi-quantified (**e**, **f**). Representative blots of three independent experiments are shown. Each bar represents the mean + standard error of the mean of signal intensities from blots of three separate experiments that were calculated as average percentage compared to control (** $p < 0.01$ and *** $p < 0.001$)

fibronectin in microglia (Fig. 1d, f). Considering the minimal amount of fibronectin detected in microglia (Fig. 1d), we decided to continue our study with a focus on astrocytes.

TG2 expression and activity in astrocytes were increased by inflammatory conditions

We subsequently visualized TG2 by immunohistochemistry in astrocytes. Under control conditions, little TG2 immunoreactivity was observed (Fig. 2a), but a clear increase was apparent in TG2 immunoreactivity in specific GFAP positive

astrocytes after 48 h of TNF- α +IL-1 β treatment (indicated by arrows in Fig. 2b). Inflammatory conditions not only affected expression of TG2 in astrocytes, but also increased its activity. In particular, LPS ($P < 0.01$) and TNF- α +IL-1 β ($P < 0.001$) increased TG activity by astrocytes (Fig. 2c) which was reduced when adding the TG2 specific inhibitor Z-DON to the activity assay, indicating that the measured TG activity is mainly due to TG2. Of interest is that although IL-4 increased TG2 protein level, it showed a trend but did not significantly affect TG activity (Fig. 2c).





Fig. 2 TG2 expression and activity in astrocytes. Primary rat astrocytes were treated for 48 h with medium alone as a control condition (CTRL) (a) or with TNF- α +IL-1 β (b). TG2 immunoreactivity (green) was detected in astrocytes (as indicated by arrows in b) which were visualized by the astrocytic marker glial fibrillary acidic protein (GFAP, red). Representative images of three independent experiments are shown. Scale bar represents 50 μ m. TG activity was examined in protein samples of astrocytes that were treated for 48 h with a range of inflammatory mediators. Subsequent derived astrocyte protein lysates were pre-incubated with a selective inhibitor of TG2 (Z-DON, 1 μ M) diluted in 0.001% DMSO or only with DMSO for 30 min at room temperature (c). Each bar represents the mean + standard error of the mean of measurements from three separate experiments that were calculated as average percentage compared to control (** P < 0.01 and *** P < 0.001)

TG2 and fibronectin were deposited in the ECM by astrocytes

Astrocytes were subsequently plated onto the ECM protein laminin to mimic the situation of an MS lesion, where laminin expression is increased [22]. Moreover, laminin is of interest because it is expressed by cultured astrocytes from healthy subjects and MS patients [29], and laminin is a myelination permissive ECM molecule [68–70]. Under those conditions, we again examined the effect of inflammatory conditions on TG2 and fibronectin levels. LPS or TNF- α +IL-1 β treatment were as effective in enhancing TG2 and fibronectin protein levels (P < 0.05 for both treatments, Fig. 3a–c) as they were in astrocytes cultured

on PLL (Fig. 1). However, IL-4 did not affect TG2 protein levels in astrocytes cultured on laminin (Fig. 3a, b) in contrast to astrocytes cultured on PLL (Fig. 1a, b), while fibronectin levels were not affected in astrocytes cultured on both ECM proteins (Fig. 1a, c and Fig. 3a, c). TGF- β 1 treatment did not affect TG2 protein levels in astrocytes cultured on either PLL or laminin (Fig. 1a, b and Fig. 3a, b), whereas fibronectin protein levels in astrocytes cultured on laminin were slightly enhanced (Fig. 3a, c) but more significantly when astrocytes were cultured on PLL (Fig. 1a, c). We next examined the effect of different inflammatory conditions on the extracellular deposition of endogenously produced TG2 and fibronectin by astrocytes cultured on




laminin. Astrocytes were therefore first cultured on laminin coated chamber slides and treated with TGF-β1 or a combination of TNF-α+IL-1β cytokines for 48 h. The astrocytes were then removed and the ECM, produced by these astrocytes during the 48 h incubation period, was immunohistochemically stained for TG2 and fibronectin. Without treatment, little extracellular TG2 and fibronectin immunoreactivity in the ECM could be detected (Fig. 3d). Similarly, after TGF-β1 treatment of the cells, TG2 in the ECM was almost undetectable (Fig. 3e), but fibronectin deposition was clearly visible (Fig. 3e) and showed fibril-like structures. Upon treatment with TNF-α+IL-1β, extracellular TG2 immunoreactivity was present in the ECM (Fig. 3f). This treatment did not visibly affect the amount of fibronectin deposited in the ECM, though co-localization between TG2 and fibronectin was apparent under this condition (indicated by arrows in Fig. 3f).

Exogenous TG2 affected astrocyte-derived fibronectin deposition and cross-linking

To follow-up on the observation that TG2 can be extracellularly deposited by astrocytes, we determined if extracellular TG2 can have an effect on fibronectin deposition. As proof of principle, we applied exogenous

TG2 to untreated astrocytes plated onto laminin and studied fibronectin deposition and aggregation. We observed that within a pathophysiological range of TG2 concentrations, the concentration of 0.13 μM exogenously added TG2 was visible in either the ECM or interacting with astrocytes, but no visible effect on fibronectin expression or deposition was observed (Fig. 4b). When the TG2 concentration of 1.3 μM was added to the cells, it resulted in a more general interaction with astrocytes and was possibly incorporated into the astrocytes (Fig. 4c). In addition, this concentration of TG2 resulted in morphologically altered fibronectin deposition which appeared with a more fibrillary structure (as indicated by arrows in Fig. 4c). Moreover, adding 0.64 μM TG2 to the deposited ECM, after removal of astrocytes, resulted in a significant increase in astrocyte-derived fibronectin aggregates in the ECM as detected by FTA (*P* < 0.001, Fig. 4d). Pre-incubation of the ECM with the specific TG2 inhibitor Z-DON partially, but significantly, reduced this increase in fibronectin aggregates (*P* < 0.05, Fig. 4d), indicating that TG2 activity contributes to an altered morphology of the fibronectin deposited and to fibronectin aggregation.



Endogenous TG2 affected the morphological appearance of fibronectin

We next determined whether endogenously produced TG2, enhanced by MS relevant inflammatory stimuli, is involved in fibronectin deposition and aggregation. As already found that $\text{TNF-}\alpha + \text{IL-1}\beta$ did not affect fibronectin protein levels in astrocytes (Fig. 3c), we now observed that it neither significantly affected aggregation of extracellular fibronectin (Fig. 5a, b). Interestingly, the morphological appearance of extracellular fibril-like fibronectin deposits (Fig. 5c, see arrows) was clearly altered, i.e., shorter and thinner fibrils, upon co-incubation with the bona fide TG2 inhibitor ERW1041E (48 h) (Fig. 5d). Also, after lentiviral downregulation of TG2 (37% TG2 protein left of control scrambled shRNA treatment, data not shown), the morphological fibril-like fibronectin deposition, as found in astrocytes lentivirally transduced with control scrambled shRNA (Fig. 5e, see arrows), was altered (Fig. 5f).

Discussion

In the present study, we demonstrated that certain inflammatory mediators enhance TG2 protein levels in astrocytes and microglial cells. In addition, fibronectin protein levels were elevated in astrocytes, but not microglial cells. Although exogenous TG2 contributed to aggregation of fibronectin produced by astrocytes, endogenously produced TG2 contributed to the appearance of morphological fibril-like fibronectin, but was not involved in fibronectin aggregation under inflammatory conditions in vitro.

At first, the present study extends our previous observations in astroglia to microglia and additional inflammatory treatment effects were studied. We now observed that astrocytes produced more TG2 than microglia. To determine which inflammatory mediators affect TG2 protein levels in rat astrocytes and microglial cells, the effect of a wide range of inflammatory mediators was studied. Of these LPS, $\text{TNF-}\alpha + \text{IL-1}\beta$ and IL-4 most prominently




Fig. 5 (See legend on next page.)

(See figure on previous page.)

Fig. 5 Endogenous TG2 affects the morphological appearance of fibronectin. Fibronectin (FN) aggregates on laminin (LAM) coated plastic were studied after removal of astrocytes that were allowed to produce extracellular matrix during 48 h of TNF- α +IL-1 β treatment (**a, b**). Representative blots of four independent experiments are shown. Each bar represents the mean + standard error of the mean of signal intensities from blots of four separate experiments that were calculated as average percentage compared to control. Primary rat astrocytes were treated with TNF- α +IL-1 β for 48 h plated on laminin coated plastic. Immunocytochemical double stainings showed immunoreactivity of fibronectin (FN, red) and TG2 (green) in astrocytes (**c, d, e, and f**). Astrocytes were co-incubated with TNF- α +IL-1 β and the TG2-specific inhibitor ERW1041E (10 μ M for 48 h) or DMSO (**d**) or treated with TNF- α +IL-1 β after lentiviral downregulation of TG2 (TG2 knockdown (KD)) compared to scrambled knockdown (scrambled KD) (**f**). Fibril-like fibronectin deposition (indicated by arrows in **c** and **e**) was morphological altered after treatment with ERW1041E and after lentiviral downregulation of TG2. Scale bar represents 20 μ m

enhanced TG2 protein levels in both cell types, with a higher fold increase in astrocytes. It has been shown before by us and others that astrocytes can produce TG2 [50, 71, 72]. A recent study indicated that mouse microglial cells and not astrocytes are the primary source of TG2 mRNA [73]. Their contrasting observation may be due to differences in species, rats in our study and mice in theirs, and developmental stage of the cells that can affect mRNA expression [74–76]. The responsiveness of glial cells to various inflammatory mediators can be explained by the presence of inflammatory factor-related response elements in the promoter region of the TG2 gene [47, 77, 78], including NF- κ B (nuclear factor-kappa B), a transcription factor involved in the regulation of expression of many inflammatory mediators [79].

Astroglial scar formation is considered to contribute to the scar formation in MS lesions. As part of that process, astro- and microglial cells produce ECM proteins to stabilize the matrix which then becomes a non-permissive environment for regeneration, by depositing ECM proteins that are often inhibitory to regeneration, e.g., chondroitin sulfate proteoglycans [24, 51, 80], hyaluronan [19, 81, 82], and fibronectin [29, 30, 69]. In the present study, we observed that rat astrocytes had higher fibronectin protein levels than microglial cells. This is in line with other observations showing that ECM proteins are predominantly produced by astrocytes and to a lesser extent by microglia and other glial cells [8, 29, 52, 53].

We thus subsequently focused on the regulation of TG2 and fibronectin production in astrocytes during inflammatory conditions. Besides intracellular expression of TG2, we observed TG2 protein on the surface of astrocytes which was also regulated by inflammatory mediators (data not shown). This is in agreement with our previous finding that TG2 is upregulated in the presence of IL-1 β and TNF- α at the cell surface of rat astrocytes, where we showed that this coincided with an increased adhesion to and interaction of astrocytes with the ECM protein fibronectin, creating an environment for astroglial scarring [50]. In addition to TG2 protein production, its activity is essential to mediate possible cross-linking of fibronectin. We now showed that TG2 activity was present and increased by LPS or TNF- α +IL-1 β , but was not significantly affected by IL-4 treatment.

This last observation was unexpected, as the amount of TG2 protein was enhanced, and in human macrophages, IL-4 does increase TG2 activity [83]. This may suggest that in astrocytes, IL-4-generated TG2 has a function different from protein cross-linking. In agreement with our finding that astrocytes respond to inflammatory conditions by increasing their TG2 expression and activity, others have shown, in different cell types and tissues, such as lung and liver cells and cartilage tissue, that inflammatory cytokines, including TNF- α and IL-1 β , can indeed upregulate TG2 expression and activity [47, 77, 84–86].

It is of interest to note that the effects some cytokines had on astrocytes that were cultured on PLL, such as an increase in TG2 protein levels after IL-4 treatment and an increase in fibronectin protein levels after TGF- β 1 treatment, were no longer present when astrocytes were cultured on laminin. This is in line with previous work that showed that ECM composition determines astrocyte responses to inflammatory stimuli [87] and extends on our previous observations [50]. As in brain lesions induced by damage, i.e., MS, more ECM proteins such as laminin are present [22, 27], we considered it relevant to pursue our studies with astrocytes exposed to laminin in the culture dish. The subsequent effect of inflammatory mediators in our study on either TG2 or fibronectin was not confined to protein expression inside the cells, but extracellular deposition of both TG2 and fibronectin was also affected by TNF- α +IL-1 β and TGF- β 1. Under inflammatory conditions, including MS, it is well known that astrocytes contribute to ECM production and deposition [8, 13, 21, 88–90]. In our present study, we showed that the addition of exogenous TG2 to astrocytes resulted in a fibril-like morphology of the fibronectin deposited and increased fibronectin aggregation. This protein aggregation was partially reduced by the TG2-specific inhibitor Z-DON, which suggests that TG2 activity can, at least partly, mediate aggregation of astrocyte-derived fibronectin, shown here as fibronectin fibrils. Alternatively, a non-enzymatic, non-covalent interaction of TG2 with fibronectin [91] might contribute to the altered morphology of the fibronectin. Such TG2-fibronectin interaction stabilizes the ECM by enhancing fibronectin matrix formation and interaction with other proteins, such as integrins, syndecan-4,

growth factor receptors, and other cell surface or ECM proteins [32, 92–94].

Under inflammatory conditions, as present in MS lesions, when endogenous TG2 protein levels were enhanced by TNF- α +IL-1 β treatment of astrocytes, fibronectin deposits showed morphologically a fibril-like structure. This was similar as shown by addition of exogenous TG2, while now fibronectin aggregation was absent. These observations are suggestive for a role of TG2 in the assembly of fibronectin into a fibril-like structure without being aggregated. Indeed, treatment of astrocytes with the TG2 inhibitor ERW1041E or lentiviral downregulation of TG2 with TG2-specific shRNA in astrocytes altered the morphology of TNF- α +IL-1 β -induced fibronectin deposits, by reducing fibril size and thickness, indicating that TG2 is involved in this process *in vitro*. Though, no clear fibronectin aggregates were observed, which were evident when TG2 was exogenously added to the astrocytes. This is in line with the observations that TG2 can lead to a switch from intramolecular cross-linking to intermolecular protein cross-linking giving rise to protein aggregate formation [63]. Thus, we cannot exclude that the level of TG2 protein present in astrocytes under inflammatory conditions *in vitro* is able to alter the conformation of fibronectin to a fibril-like structure, but it is not able to form cross-linked protein aggregates. In support of this, we also did not observe fibronectin aggregation in the cuprizone model of demyelination in which astrocytes become hypertrophic and express TG2 and fibronectin [49]. However, fibronectin deposition is found in MS lesions as a complex network of fibrils of high molecular weight aggregates, which impairs remyelination [29]. Fibronectin aggregation, as defined by deoxycholate-insolubility, is likely the result of strong, non-covalent protein-protein interactions [31, 95, 96]. As a consequence, we might hypothesize that during inflammatory lesion formation in MS and concomitant deposition of ECM proteins, astrocyte-derived TG2, only in the presence of infiltrating leukocytes or their derived factors, can increase the formation of fibronectin fibril-like deposits which aggregate and may contribute to the process of scarring and subsequent impaired remyelination.

Conclusions

The results presented here demonstrate that under inflammatory conditions *in vitro*, TG2 protein levels are enhanced in astrocytes and microglia. In addition, in particular, astrocytes produce ECM proteins, i.e., fibronectin, that can be cross-linked and aggregated by exogenously added TG2. Endogenously produced TG2, enhanced by inflammatory stimuli, is involved in the appearance of morphological fibril-like fibronectin deposits, but does not lead to cross-linked fibronectin

aggregates. Thus, our *in vitro* observations suggest that during MS lesion formation, when inflammatory mediators are produced, astrocyte-derived TG2 may contribute to ECM rearrangement and subsequent astroglial scarring. The possibility to intervene with the formation of fibronectin fibril-like deposits may open avenues to reduce the astroglial scarring process, not only in MS but also in other CNS disorders, when scarring prohibits regeneration [97].

Additional file

Additional file 1: TG2 and fibronectin protein expression in rat astrocytes and microglia. Untreated (Ctrl) and 48 h LPS-treated primary rat astrocytes (A) expressed more fibronectin (FN) and more TG2 compared to untreated (Ctrl) and LPS-treated primary rat microglia (M). Representative blot of three independent experiments is shown. (TIFF 195 kb)

Abbreviations

biotin-Cd: Biotinylated-cadaverine; CNS: Central nervous system; CTRL: Control; DTT: Dithiothreitol; ECM: Extracellular matrix; FN: Fibronectin; FTA: Filter trap assay; GFAP: Glial fibrillary acidic protein; h: Hours; HRP: Horseradish peroxidase; IL-10: Interleukin-10; IL-1 β : Interleukin-1 beta; IL-4: Interleukin-4; IL-6: Interleukin-6; LAM: Laminin; LPS: Lipopolysaccharide; Min: Minutes; MS: Multiple sclerosis; NF- κ B: Nuclear factor-kappa B; PLL: Poly-L-lysine; PMSF: Phenylmethanesulfonyl fluoride; RT: Room temperature; SDS: Sodium dodecyl sulfate; TBS: Tris-buffered saline; TBS-T: Tris-buffered saline 0.1%/0.05% Triton; TG: Transglutaminase; TG2: Tissue transglutaminase; TGF- β 1: Transforming growth factor-beta 1; TGF- β 2: Transforming growth factor-beta 2; TNF- α : Tumor necrosis factor-alpha

Acknowledgements

Not applicable

Funding

This project was financially supported by the Dutch MS Research Foundation (project number 09-693MS).

Availability of data and materials

The datasets acquired during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Authors' contributions

A-MvD and WB designed and supervised the project with input from BD, NEP, and JJPB, and JGJMB performed the described experiments. NEP performed statistical analysis. NEP, A-MvD, and WB interpreted the data and prepared the manuscript. All authors read and approved the final manuscript.

Ethics approval

Animal studies were conducted with an approval of the Animal Experiment Committee of the VU University Medical Center (ID FGA 11-03).

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department Anatomy and Neurosciences, VU University Medical Center, Amsterdam Neuroscience, De Boelelaan 1108, 1081, HZ, Amsterdam, The

Netherlands. ²Department of Cell Biology, University Medical Center Groningen, Groningen, The Netherlands.

Received: 27 July 2017 Accepted: 11 December 2017

Published online: 28 December 2017

References

- Trapp BD, Peterson J, Ransohoff RM, Rudick R, Mörk S, Bö L. Axonal transection in the lesions of multiple sclerosis. *N Engl J Med*. 1998;338:278–85.
- Bö L, Mörk S, Kong PA, Nyland H, Pardo CA, Trapp BD. Detection of MHC class II-antigens on macrophages and microglia, but not on astrocytes and endothelia in active multiple sclerosis lesions. *J Neuroimmunol*. 1994;51:135–46.
- Lassmann H, Brück W, Lucchinetti CF. The immunopathology of multiple sclerosis: an overview. *Brain Pathol*. 2007;17:210–8.
- Koning N, Bö L, Hoek RM, Huitinga I. Downregulation of macrophage inhibitory molecules in multiple sclerosis lesions. *Ann Neurol*. 2007;62:504–14.
- Brück W, Porada P, Poser S, Rieckmann P, Hanefeld F, Kretschmarch HA, et al. Monocyte/macrophage differentiation in early multiple sclerosis lesions. *Ann Neurol*. 1995;38:788–96.
- Frohman EM, Racke MK, Raine CS. Multiple sclerosis—the plaque and its pathogenesis. *N Engl J Med*. 2006;354:942–55.
- Bramow S, Frischer JM, Lassmann H, Koch-Henriksen N, Lucchinetti CF, Sørensen PS, et al. Demyelination versus remyelination in progressive multiple sclerosis. *Brain*. 2010;133:2983–98.
- Wiese S, Karus M, Faissner A. Astrocytes as a source for extracellular matrix molecules and cytokines. *Front Pharmacol*. 2012;3:120.
- Meeuwssen S, Persoon-Deen C, Bsibsi M, Ravid R, van Noort JM. Cytokine, chemokine and growth factor gene profiling of cultured human astrocytes after exposure to proinflammatory stimuli. *Glia*. 2003;43:243–53.
- Smith GM, Strunz C. Growth factor and cytokine regulation of chondroitin sulfate proteoglycans by astrocytes. *Glia*. 2005;52:209–18.
- Ransohoff RM, Brown MA. Innate immunity in the central nervous system. *J Clin Invest*. 2012;122:1164–71.
- Williams A, Piaton G, Lubetzki C. Astrocytes—friends or foes in multiple sclerosis? *Glia*. 2007;55:1300–12.
- Fawcett JW, Asher RA. The glial scar and central nervous system repair. *Brain Res Bull*. 1999;49:377–91.
- Kotter MR, Stadelmann C, Hartung H-P. Enhancing remyelination in disease—can we wrap it up? *Brain*. 2011;134:1882–900.
- Motavaf M, Sadeghizadeh M, Javan M. Attempts to overcome remyelination failure: toward opening new therapeutic avenues for multiple sclerosis. *Cell Mol Neurobiol*. 2017;37:1335–48.
- Stankoff B, Joachim J, Hartung H, Ku P. Repair strategies for multiple sclerosis: challenges, achievements and perspectives. *Curr Opin Neurol*. 2016;29:286–92.
- Lublin FD, Reingold SC, Cohen JA, Cutter GR, Sørensen PS, Thompson AJ, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. *Neurology*. 2014;83:278–86.
- Gutowski NJ, Newcombe J, Cuzner ML. Tenascin-R and C in multiple sclerosis lesions: relevance to extracellular matrix remodelling. *Neuropathol Appl Neurobiol*. 1999;25:207–14.
- Back SA, Tuohy TMF, Chen H, Wallingford N, Craig A, Struve J, et al. Hyaluronan accumulates in demyelinated lesions and inhibits oligodendrocyte progenitor maturation. *Nat Med*. 2005;11:966–72.
- Van Horsen J, Bö L, Vos CMP, Virtanen I, de Vries HE. Basement membrane proteins in multiple sclerosis-associated inflammatory cuffs: potential role in influx and transport of leukocytes. *J Neuropathol Exp Neurol*. 2005;64:722–9.
- Silver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci*. 2004;5:146–56.
- Van Horsen J, Dijkstra CD, de Vries HE. The extracellular matrix in multiple sclerosis pathology. *J Neurochem*. 2007;103:1293–301.
- Lau LW, Cua R, Keough MB, Haylock-Jacobs S, Yong VW. Pathophysiology of the brain extracellular matrix: a new target for remyelination. *Nat Rev Neurosci*. 2013;14:722–9.
- Lau LW, Keough MB, Haylock-Jacobs S, Cua R, Döring A, Sloka S, et al. Chondroitin sulfate proteoglycans in demyelinated lesions impair remyelination. *Ann Neurol*. 2012;72:419–32.
- Van der Laan LJW, De Groot CJA, Elices MJ, Dijkstra CD. Extracellular matrix proteins expressed by human adult astrocytes in vivo and in vitro: an astrocyte surface protein containing the CS1 domain contributes to binding of lymphoblasts. *J Neurosci Res*. 1997;50:539–48.
- Price J, Hynes RO. Astrocytes in culture synthesize and secrete a variant form of fibronectin. *J Neurosci*. 1985;5:2205–11.
- Liesi P, Kaakkola S, Dahl D, Vaheri A. Laminin is induced in astrocytes of adult brain by injury. *EMBO J*. 1984;3:683–6.
- Liesi P, Kirkwood T, Vaheri A. Fibronectin is expressed by astrocytes cultured from embryonic and early postnatal rat brain. *Exp Cell Res*. 1986;163:175–85.
- Stoffels JMJ, de Jonge JC, Stancic M, Nomden A, van Strien ME, Ma D, et al. Fibronectin aggregation in multiple sclerosis lesions impairs remyelination. *Brain*. 2013;136:116–31.
- Sisková Z, Yong VW, Nomden A, van Strien M, Hoekstra D, Baron W. Fibronectin attenuates process outgrowth in oligodendrocytes by mislocalizing MMP-9 activity. *Mol Cell Neurosci*. 2009;42:234–42.
- Stoffels JMJ, Zhao C, Baron W. Fibronectin in tissue regeneration: timely disassembly of the scaffold is necessary to complete the build. *Cell Mol Life Sci*. 2013;70:4243–53.
- Aeschlimann D, Thomazy V. Protein crosslinking in assembly and remodelling of extracellular matrices: the role of transglutaminases. *Connect Tissue Res*. 2000;41:1–27.
- Fesus L, Piacentini M. Transglutaminase 2: an enigmatic enzyme with diverse functions. *Trends Biochem Sci*. 2002;27:534–9.
- Griffin M, Casadio R, Bergamini CM. Transglutaminases: nature's biological glues. *Biochem J*. 2002;396:377–96.
- Iismaa SE, Mearns BM, Lorand L, Graham RM. Transglutaminases and disease: lessons from genetically engineered mouse models and inherited disorders. *Physiol Rev*. 2009;99:1–1023.
- Mehta K, Kumar A, Kim HL. Transglutaminase 2: a multi-tasking protein in the complex circuitry of inflammation and cancer. *Biochem Pharmacol*. 2010;80:1921–9.
- Numinskaya MV, Belkin AM. Cellular functions of tissue transglutaminase. *Int Rev Cell Mol Biol*. 2012;294:1–97.
- Park D, Choi SS, Ha K-S. Transglutaminase 2: a multi-functional protein in multiple subcellular compartments. *Amino Acids*. 2010;39:619–31.
- Thomazy V, Fesus L. Differential expression of tissue transglutaminase in human cells. *Cell Tissue Res*. 1989;255:215–24.
- Balklava Z, Verderio E, Collighan R, Gross S, Adams J, Griffin M. Analysis of tissue transglutaminase function in the migration of Swiss 3T3 fibroblasts: the active-state conformation of the enzyme does not affect cell motility but is important for its secretion. *J Biol Chem*. 2002;277:16567–75.
- Lorand L, Graham RM. Transglutaminases: crosslinking enzymes with pleiotropic functions. *Nat Rev Mol Cell Biol*. 2003;4:140–56.
- Chou C-Y, Streets AJ, Watson PF, Huang L, Verderio EAM, Johnson TS. A crucial sequence for transglutaminase type 2 extracellular trafficking in renal tubular epithelial cells lies in its N-terminal beta-sandwich domain. *J Biol Chem*. 2011;286:27825–35.
- Wang Z, Collighan RJ, Gross SR, Danen E, Orend G, Telci D, et al. RGD-independent cell adhesion via a tissue transglutaminase-fibronectin matrix promotes fibronectin fibril deposition and requires syndecan-4/2 $\alpha 5 \beta 1$ integrin co-signaling. *J Biol Chem*. 2010;285:40212–29.
- Akimov SS, Belkin AM. Cell surface tissue transglutaminase is involved in adhesion and migration of monocytic cells on fibronectin. *Blood*. 2001;98:1567–76.
- Telci D, Wang Z, Li X, Verderio EAM, Humphries MJ, Baccarini M, et al. Fibronectin-tissue transglutaminase matrix rescues RGD-impaired cell adhesion through syndecan-4 and beta1 integrin co-signaling. *J Biol Chem*. 2008;283:20937–47.
- Collighan RJ, Griffin M. Transglutaminase 2 cross-linking of matrix proteins: biological significance and medical applications. *Amino Acids*. 2009;36:659–70.
- Gundemir S, Colak G, Tucholski J, Johnson GWW. Transglutaminase 2: a molecular Swiss army knife. *Biochim Biophys Acta*. 2012;1823:406–19.
- Van Strien ME, Drukarch B, Bol JG, van der Valk P, van Horsen J, Gerritsen WH, et al. Appearance of tissue transglutaminase in astrocytes in multiple sclerosis lesions: a role in cell adhesion and migration? *Brain Pathol*. 2011;21:44–54.
- Espitia Pinzón N, Sanz-Morello B, Brevé JJP, Bol JGJM, Drukarch B, Bauer J, et al. Astrocyte-derived tissue transglutaminase affects fibronectin deposition, but not aggregation, during cuprizone-induced demyelination. *Sci Rep*. 2017;7:40995.
- Van Strien ME, Brevé JJP, Fratantoni S, Schreurs MWJ, Bol JGJM, Jongenelen CAM, et al. Astrocyte-derived tissue transglutaminase interacts with fibronectin: a role in astrocyte adhesion and migration? *PLoS One*. 2011;6:e25037.

51. Harlow DE, Macklin WB. Inhibitors of myelination: ECM changes, CSPGs and PTPs. *Exp Neurol*. 2014;251:39–46.
52. Fitch MT, Silver J. CNS injury, glial scars, and inflammation: inhibitory extracellular matrices and regeneration failure. *Exp Neurol*. 2008;209:294–301.
53. Sherman LS, Back S a. A “GAG” reflex prevents repair of the damaged CNS. *Trends Neurosci*. 2008;31:44–52.
54. Galtrey CM, Kwok JCF, Carulli D, Rhodes KE, Fawcett JW. Distribution and synthesis of extracellular matrix proteoglycans, hyaluronan, link proteins and tenascin-R in the rat spinal cord. *Eur J Neurosci*. 2008;27:1373–90.
55. Morgenstern DA, Asher RA, Fawcett JW. Chondroitin sulphate proteoglycans in the CNS injury response. *Prog Brain Res*. 2002;137:313–32.
56. Rhodes KE, Fawcett JW. Chondroitin sulphate proteoglycans: preventing plasticity or protecting the CNS? *J Anat*. 2004;204:33–48.
57. Ledebauer A, Breve JJP, Wierinckx A, van der Jagt S, Bristow AF, Leysen JE, et al. Expression and regulation of interleukin-10 and interleukin-10 receptor in rat astroglial and microglial cells. *Eur J Neurosci*. 2002;16:1175–85.
58. Prins M, Dutta R, Baselmans B, Brevé JJP, Bol JGJM, Deckard SA, et al. Discrepancy in CCL2 and CCR2 expression in white versus grey matter hippocampal lesions of multiple sclerosis patients. *Acta Neuropathol Commun*. 2014;2:98.
59. Dafik L, Albertelli M, Stammaes J, Sollid LM, Khosla C. Activation and inhibition of transglutaminase 2 in mice. *PLoS One*. 2012;7:e30642.
60. Dafik L, Khosla C. Dihydroisoxazole analogs for labeling and visualization of catalytically active transglutaminase 2. *Chem Biol*. 2011;18:58–66.
61. Watts RE, Siegel M, Khosla C. Structure-activity relationship analysis of the selective inhibition of transglutaminase 2 by dihydroisoxazoles. *J Med Chem*. 2006;49:7493–501.
62. Konno T, Morii T, Hirata A, Sato SI, Oiki S, Ikura K. Covalent blocking of fibril formation and aggregation of intracellular amyloidogenic proteins by transglutaminase-catalyzed intramolecular cross-linking. *Biochemistry*. 2005;44:2072–9.
63. Segers-Nolten IMJ, Wilhelmus MMM, Veldhuis G, Van Rooijen BD, Drukarch B, Subramaniam V. Tissue transglutaminase modulates α -synuclein oligomerization. *Protein Sci*. 2008;17:1395–402.
64. Sakly W, Thomas V, Quash G, El Alaoui S. A role for tissue transglutaminase in alpha-gliadin peptide cytotoxicity. *Clin Exp Immunol*. 2006;146:550–8.
65. Kumar S, Mehta K. Tissue transglutaminase constitutively activates HIF-1 α promoter and nuclear factor- κ B via a non-canonical pathway. *PLoS One*. 2012;7:e49321.
66. Boroughs LK, Antonyak MA, Cerione RA. A novel mechanism by which tissue transglutaminase activates signaling events that promote cell survival. *J Biol Chem*. 2014;289:10115–25.
67. Antonyak MA, Jansen JM, Miller AM, Ly TK, Endo M, Cerione RA. Two isoforms of tissue transglutaminase mediate opposing cellular fates. *Proc Natl Acad Sci U S A*. 2006;103:18609–14.
68. Colognato H, Ffrench-Constant C, Feltri ML. Human diseases reveal novel roles for neural laminins. *Trends Neurosci*. 2005;28:480–6.
69. Sisková Z, Baron W, de Vries H, Hoekstra D. Fibronectin impedes “myelin” sheet-directed flow in oligodendrocytes: a role for a beta 1 integrin-mediated PKC signaling pathway in vesicular trafficking. *Mol Cell Neurosci*. 2006;33:150–9.
70. BATTERY PC, Ffrench-Constant C. Laminin-2/integrin interactions enhance myelin membrane formation by oligodendrocytes. *Mol Cell Neurosci*. 1999;14:199–212.
71. Monteagudo A, Ji C, Akbar A, Keillor JW, Johnson GWW. Inhibition or ablation of transglutaminase 2 impairs astrocyte migration. *Biochem Biophys Res Commun*. 2017;482:942–7.
72. Campisi A, Caccamo D, Raciti G, Cannavò G, Macaione V, Currò M, et al. Glutamate-induced increases in transglutaminase activity in primary cultures of astroglial cells. *Brain Res*. 2003;978:24–30.
73. Bennett ML, Bennett C, Liddelov SA, Ajami B, Zamanian JL, Fernhoff NB, et al. New tools for studying microglia in the mouse and human CNS. *Proc Natl Acad Sci U S A*. 2016;113:E1738–46.
74. Hoffmann R. Gene expression patterns in human and mouse B cell development. *Curr Top Microbiol Immunol*. 2005;294:19–29.
75. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Meth*. 2008;5:621–8.
76. Wilhelm BT, Marguerat S, Watt S, Schubert F, Wood V, Goodhead I, et al. Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution. *Nature*. 2008;453:1239–43.
77. Kuncio GS, Tsyganskaya M, Zhu J, Liu S, Nagy L, Thomazy V, et al. TNF- α modulates expression of the tissue transglutaminase gene in liver cells. *Am J Phys*. 1998;274:240–5.
78. Ikura K, Shinagawa R, Suto N, Sasaki R. Increase caused by interleukin-6 in promoter activity of guinea pig liver transglutaminase gene. *Biosci Biotechnol Biochem*. 1994;58:1540–1.
79. Ientile R, Currò M, Caccamo D. Transglutaminase 2 and neuroinflammation. *Amino Acids*. 2015;47:19–26.
80. Fitch MT, Silver J. Glial cell extracellular matrix: boundaries for axon growth in development and regeneration. *Cell Tissue Res*. 1997;290:379–84.
81. Sloane JA, Batt C, Ma Y, Harris ZM, Trapp B, Vartanian T. Hyaluronan blocks oligodendrocyte progenitor maturation and remyelination through TLR2. *Proc Natl Acad Sci*. 2010;107:11555–60.
82. Chang A, Staugaitis SM, Dutta R, Batt CE, Easley KE, Chomyk AM, et al. Cortical remyelination: a new target for repair therapies in multiple sclerosis. *Ann Neurol*. 2012;72:918–26.
83. Martinez FO, Helming L, Milde R, Varin A, Melgert BN, Drajer C, et al. Genetic programs expressed in resting and IL-4 alternatively activated mouse and human macrophages: similarities and differences. *Blood*. 2013;121:57–70.
84. Johnson K, Hashimoto S, Lotz M, Pritzker K, Terkeltaub R. Interleukin-1 induces pro-mineralizing activity of cartilage tissue transglutaminase and factor XIIIa. *Am J Pathol*. 2001;159:149–63.
85. Chen R, Gao B, Huang C, Olsen B, Rotundo RF, Blumenstock F, et al. Transglutaminase-mediated fibronectin multimerization in lung endothelial matrix in response to TNF- α . *Am J Phys*. 2000;279:161–74.
86. Codarri L, Fontana A, Becher B. Cytokine networks in multiple sclerosis: lost in translation. *Curr Opin Neurol*. 2010;23:205–11.
87. Johnson KM, Milder R, Crocker SJ. Extracellular matrix composition determines astrocyte responses to mechanical and inflammatory stimuli. *Neurosci Lett*. 2015;1848:3047–54.
88. Klausmeyer A, Conrad R, Faissner A, Wiese S. Influence of glial-derived matrix molecules, especially chondroitin sulfates, on neurite growth and survival of cultured mouse embryonic motoneurons. *J Neurosci Res*. 2011;89:127–41.
89. Voskuhl RR, Peterson RS, Song B, Ao Y, Morales LBJ, Tiwari-Woodruff S, et al. Reactive astrocytes form scar-like perivascular barriers to leukocytes during adaptive immune inflammation of the CNS. *J Neurosci*. 2009;29:11511–22.
90. Sofroniew MV, Vinters HV. Astrocytes: biology and pathology. *Acta Neuropathol*. 2010;119:7–35.
91. Kanchan K, Fuxreiter M, Fésüs L. Physiological, pathological, and structural implications of non-enzymatic protein–protein interactions of the multifunctional human transglutaminase 2. *Cell Mol Life Sci*. 2015;72:3009–35.
92. Forsprecher J, Wang Z, Nelea V, Kaartinen MT. Enhanced osteoblast adhesion on transglutaminase 2-crosslinked fibronectin. *Amino Acids*. 2009;36:747–53.
93. Belkin AM. Extracellular TG2: emerging functions and regulation. *FEBS J*. 2011;278:4704–16.
94. Akimov SS, Krylov D, Fleischman LF, Belkin AM. Tissue transglutaminase is an integrin-binding adhesion coreceptor for fibronectin. *J Cell Biol*. 2000;148:825–38.
95. Chen H, Mosher DF. Formation of sodium dodecyl sulfate-stable fibronectin multimers. *J Biol Chem*. 1996;271:9084–9.
96. Ohashi T, Erickson HP. Revisiting the mystery of fibronectin multimers: the fibronectin matrix is composed of fibronectin dimers cross-linked by non-covalent bonds. *Matrix Biol*. 2009;28:170–5.
97. Sun D, Jakobs TC. Structural remodeling of astrocytes to the injured CNS. *Neuroscientist*. 2012;18:567–88.