



Published in final edited form as:

*J Neurosci Res.* 2018 January ; 96(1): 31–44. doi:10.1002/jnr.24088.

## Connexins & Pannexins: at the Junction of Neuro-Glial Homeostasis & Disease

Andrew S. Lapato<sup>1,3</sup> and Seema K. Tiwari-Woodruff<sup>1,2,3</sup>

<sup>1</sup>Division of Biomedical Sciences, School of Medicine, University of California Riverside, Riverside, CA 92521

<sup>2</sup>Neuroscience Graduate Program, University of California Riverside, Riverside, CA 92521

<sup>3</sup>Center for Glial-Neuronal Interactions, University of California Riverside, Riverside, CA 92521

### Abstract

In the central nervous system (CNS), connexin and pannexin gap junctions and hemichannels are an integral component of homeostatic neuronal excitability and synaptic plasticity. Neuronal connexin (Cx) gap junctions form electrical synapses across biochemically similar GABAergic networks, allowing rapid and extensive inhibition in response to principle neuron excitation. Glial Cx gap junctions link astrocytes and oligodendrocytes in the pan-glial network that is responsible for removing excitotoxic ions and metabolites. In addition, Cx gap junctions between glia help constrain excessive excitatory activity in neurons and facilitate astrocyte Ca<sup>2+</sup> slow wave propagation. Pannexins (Panxs) do not form gap junctions in vivo, but Panx hemichannels participate in autocrine and paracrine gliotransmission alongside connexin hemichannels. ATP and other gliotransmitters released by Cx and Panx hemichannels maintain physiologic glutamatergic tone by strengthening synapses and mitigating aberrant high frequency bursting. Under pathological depolarizing and inflammatory conditions, gap junctions and hemichannels become dysregulated, resulting in aberrant neuronal firing and seizure. In this review, we present known contributions of Cxs and Panxs to physiologic neuronal excitation and explore how disruption of gap junctions and hemichannels lead to aberrant glutamatergic transmission, purinergic signaling, and seizures.

### Keywords

connexin; pannexin; synaptic plasticity; epilepsy; seizure; hemichannel; gap junction; gliotransmission; purinergic signaling; electrical synapse; pan-glial network; neuronal excitability

---

Corresponding Author: \*Seema K. Tiwari-Woodruff, Ph.D. Division of Biomedical Sciences School of Medicine, University of California Riverside Rm 205, 311 School of Medicine Research Building 900 University Ave, Riverside, CA 92521, USA Work #: (951) 827-7819, seema.tiwari-woodruff@medsch.ucr.edu; seema-tiwari-woodruff@axonremyelination.org.

**Conflict of interest statement:** The authors have no conflicts of interest, including any financial, personal, or other relationships with people or organizations that could influence the present article.

**Author contributions:** Lapato, AS and Tiwari-Woodruff, SK conducted the literature review, prepared the manuscript, and composed figures.

## 1. Introduction

Accruing evidence indicates that connexin (Cx) and pannexin (Panx) transmembrane channels are crucial to the coordination and maintenance of physiologic CNS activity. In neurons, Cxs electrochemically couple neurons by electrical synapses (Galarreta and Hestrin 1999), while glial Cxs mediate numerous functions ranging from K<sup>+</sup> buffering to direct modulation of glutamatergic activity (Battfeld et al. 2016; Chever et al. 2014a; Kamasawa et al. 2005; Kofuji and Newman 2004). Alone, Cxs and Panxs represent a mechanism for robust autocrine and paracrine signaling through release of gliotransmitters, which are essential to synaptic strength and plasticity (Prochnow et al. 2012; Thompson et al. 2008). Dysregulation of Cx and Panx activity is implicated in neurodegenerative disease (Markoullis et al. 2012a) and may be etiologic in some human epilepsy (Bedner et al. 2015). In addition, Cx and Panx sensitivity to inflammatory mediators suggests that alteration in neuronal excitability may be present in a range of disease states. In the following sections, we will describe the structure, function, regulation, and distribution of CNS Cx and Panx molecules. We will also summarize evidence for their functions related to neuronal excitability under homeostatic conditions and examine their role as effectors of pathological glutamatergic transmission.

## 2. Structure and function of connexins and pannexins

Structurally, the Cx and Panx family of proteins comprise a group of transmembrane pores that are permeable to ions, metabolites, second messengers, and purine signaling mediators up to 1.5 kDa (Loewenstein 1981) with divergent peptide sequences but homologous topology. Each channel forming complex is composed of six monomers containing four membrane-spanning domains linked by two extracellular loops that mediate docking with complimentary Cx hexamers (Figure 1). Post-translational modification of Cx monomers largely takes place at the site of the intracellular carboxyl tail (May et al. 2013) and is thought to regulate non-channel functions of Cxs such as adhesion, migration (Giepmans et al. 2001; Pannasch et al. 2014), and proliferation (Cheng et al. 2004; Santiago et al. 2010). Adhesion and migration mediated by GJs are of particular importance in the developing CNS, where they facilitate neocortical neuron migration by providing points of contact with radial glia (Elias et al. 2007) and movement of subventricular zone-derived cells along the rostral migratory stream (Marins et al. 2009). Phosphorylation of the Cx intracellular tail plays a role in gating pore permeability, thereby allowing dynamic opening and closing under a range of conditions (Zador et al. 2008). However, Cx and Panx biology may extend beyond these functions into a variety of intracellular regulatory processes (reviewed in (Esseltine and Laird 2016)).

When uncoupled to a complimentary hexamer, these pores are termed hemichannels (HCs) and allow exchange of cellular contents with the extracellular space (Simard and Nedergaard 2004). Opening of Cx and Panx HCs has traditionally been thought of as deleterious to CNS homeostasis (Paul et al. 1991), occurring under pathological conditions leading to excitotoxic cell death such as ischemia (John et al. 1999; Kondo et al. 2000) and excessive depolarization (Ebihara and Steiner 1993; Valiunas and Weingart 2000). More recent evidence, however, has shed light on Cx and Panx HC physiologic activities including their

function as a platform for robust autocrine and paracrine signaling between and amongst glia and neurons (Klaassen et al. 2011; Prochnow et al. 2012).

Gliotransmission continues to draw attention to Cx and Panx HCs, whereby paracrine signaling by purinergic mediators and a host of other gliotransmitters released by these channels have proven essential to synaptic strength and plasticity (Cherian et al. 2005; Montero and Orellana 2015; Stout et al. 2002). When HCs are coupled with complementary HCs on adjacent cells, they form gap junctions (GJs), which act in pairs to facilitate direct intercellular communication of cytoplasmic molecules. Of note, Panx1 HCs have not been shown to form GJs in vivo without substantial manipulation (MacVicar and Thompson 2010; Penuela et al. 2007; Sosinsky et al. 2011), which is thought to be due to post-translational N-linked glycosylation of the second extracellular loop of each Panx subunit (Figure 1). In neurons, GJs are the substrate of electrical synapses formation (Pereda et al. 2013), while GJs among glia couple cells in the panglial network, which participates in buffering excitotoxic metabolites produced by depolarizing activity.

Regulation of GJ and HC open probability is affected by a variety of circumstances. Under homeostatic conditions, neuronal activity leading to decreased extracellular  $\text{Ca}^{2+}$  (Thompson et al. 2008; Torres et al. 2012) and increased  $\text{K}^{+}$  (Santiago et al. 2011) encourages opening of HCs and autocrine purinergic signaling via adenosine triphosphate (ATP) release (Kawamura et al. 2010). However, pathological environments also open HCs and GJs aberrantly. Neurodegenerative diseases such as experimental autoimmune encephalomyelitis (Markoullis et al. 2012a), multiple sclerosis (Masaki 2015), and epilepsy (Bedner et al. 2015) cause uncoupling of pan-glial network GJs, with the resulting dysregulated gliotransmitter release contributing to abnormal neuronal activity (Thompson et al. 2008). Inflammatory mediators, including the cytokines interleukin (IL)- $1\beta$  and tumor necrosis factor (TNF) $\alpha$  drive HC open probability, as application of either cytokine or the toll-like receptor (TLR) 4 ligand lipopolysaccharide (LPS) induces increased hemichannel—mediated uptake of ethidium bromide in astrocytes both in vitro and slice (Froger et al. 2010; Retamal et al. 2007).

Expression of Cxs and Panxs is distributed throughout the CNS (Table 1) where they participate in a variety of functions. In addition to their channel activity which promotes metabolic coupling amongst macroglia (Niu et al. 2016), Cx GJs are critical to glial survival and stabilization of associated Cx HCs (Magnotti et al. 2011; May et al. 2013). Mutations in oligodendrocyte-expressed Cx genes result in phenotypes indistinguishable from inherited hypomyelinating leukodystrophies, which are characterized by impairment of myelin sheath formation, inflammation, and sensorimotor neurological deficits (Magnotti et al. 2011; May et al. 2013; Schiza et al. 2015). For example, a mutation in the Cx47 promoter results in Pelizaeus–Merzbacher-like disease (Gotoh et al. 2014), while altered Cx32 expression leads to symptomology closely resembling Charcot-Marie-Tooth disease (Sargiannidou et al. 2015).

Further emphasizing the importance of glial coupling, astrocyte-oligodendrocyte coupling is reduced in Cx47—KO mice and is associated with myelin vacuolation in the optic nerve (Odermatt et al. 2003). Mice lacking Cx32 and Cx47 exhibit more pronounced myelin

pathology, which is accompanied by loss of oligodendrocytes and action tremors progressing into tonic-clonic seizures and mortality by the sixth postnatal week (Menichella et al. 2003; Odermatt et al. 2003). Loss and activation of astrocytes, oligodendrocyte loss, myelin vacuolation, inflammation, and invasion of phagocytic cells are also common to disorders of GJ coupling (Magnotti et al. 2011; May et al. 2013; Tress et al. 2012), suggesting a role for intact pan-glial network GJ connections in supporting its members' survival and homeostatic functions. However, while non-channel properties of Cxs have been explored in the context of neuronal differentiation (Cheng et al. 2004; Kunze et al. 2009) and glutamatergic transmission (Chever et al. 2014b), the mechanism whereby Cxs support glial survival remains poorly understood.

### 3. Connexins and pannexins in synaptic plasticity

#### 3.1 Electrical synapses, excitability, and learning

At the forefront of Cx and Panx neurobiology research is their role in neuronal excitability and their ability to modify synaptic activity. Neuronal GJs, which couple cells by electrical synapses, are numerous in the embryonic CNS, but become increasingly limited during development (Belluardo et al. 2000; Bruzzone et al. 2003; Kreuzberg et al. 2008; Rash et al. 2000; Rozental et al. 1998; Schutte et al. 1998; Swayne and Bennett 2016; Vis et al. 1998; Weickert et al. 2005). In the adult mammalian brain, GJs are restricted primarily to inhibitory GABAergic networks and associated principle neurons (Apostolides and Trussell 2013; Galarreta and Hestrin 1999), with Cx36 becoming the predominant isoform expressed (Belluardo et al. 2000; Condorelli et al. 2000). Cx36 GJs allow rapid electrochemical transmission and influence synchronization of interconnected neurons (Deans et al. 2001; Landisman et al. 2002). Coupled neurons of the reticular thalamic nucleus and suprachiasmatic nucleus of the thalamus (SCN) display inherent Cx36-dependent desynchronizing properties (Sevetson and Haas 2015; Wang et al. 2014). While synchronization of SCN neurons is required for light-dark cycle detection and circadian rhythm, these examples illustrate the overriding strength of inhibitory networks coupled by GJs and hint at their physiologic role in resisting development of epileptiform synchronization.

Traditionally, GJ-mediated electrical synapses have been considered static structures. However, recent research casts doubt on this long-standing assumption. Like glutamatergic chemical synapses, GJ-mediated electrical synapses in the rat inferior olive also display activity dependent strengthening (Turecek et al. 2014). In this study, the authors observed that the N-Methyl-D-aspartate receptor (NMDAR) NR1 subunit co-localized with neuropilar Cx36 GJs. Agonism of NMDARs resulted in increased coupled potential upon stimulation that was accompanied by elevated uptake of a tracer dye (Turecek et al. 2014), indicating that GJ—NMDAR complexes are sensitive to synaptic activity and become upregulated or opened in response. As electrical synapses facilitate rapid and widespread inhibition, these data suggest that depolarizing inputs modify GJ strength to set a basal inhibitory tone on excitatory networks.

Unsurprisingly, in addition to their function in limiting excitability, neuronal GJs also play a major role in learning and memory. Knockout of Cx36 resulted in diminished novel object

recognition that was exaggerated by environments containing complex stimuli (Frisch et al. 2005). In addition, Cx36—KO mice exhibited impaired short-term and long-term memory in Y-maze testing, and reduced motor learning by rotorod (Frisch et al. 2005). Interestingly, field excitatory post-synaptic potentials (fEPSPs) recorded from cornu ammonis 1 (CA1) hippocampal pyramidal neurons after Schaffer collateral stimulation were also decreased in Cx36—KO mice (Wang and Belousov 2011), suggesting coordinated GJ-coupled interneuron activity is an essential component of LTP inducing events. This is supported by reports finding that the power of gamma and theta frequencies, which are thought to be mediated by GABAergic interneurons (Lasztozci and Klausberger 2014), are increased in the hippocampus following tetanic stimulation of Schaffer collaterals (Bikbaev and Manahan-Vaughan 2008).

Similarly, knockout of Cx31.1, which is expressed in dopaminergic neurons of the substantia nigra pars compacta (Vandecasteele et al. 2006) and striatal output neurons (Venance et al. 2004) that play a role in sensory-motor control and novelty-induced exploration (Leussis and Bolivar 2006), resulted in elevated exploration of novel environments and impaired performance in novel object recognition tasks (Dere et al. 2008). Together, these studies point to a central role for neuronal coupling via GJs in memory formation and consolidation. This may be attributable to autoinhibition of GABAergic networks by remaining inhibitory chemical synapses and reduced long term potentiation (LTP), which is thought to represent the cellular substrate of learning (Morris et al. 2003).

### 3.2 The pan-glial network: beyond buffering

The most abundant source of GJs and HCs in the brain come from the highly interconnected lattice of glial cells interspersed throughout the CNS (Figure 2). Chief among the tasks of this syncytium of astrocytes, oligodendrocytes, and endothelial cells is the modification of neuronal excitation through the spatial buffering of ions and metabolites by GJ connected cells (Battefeld et al. 2016; Bedner et al. 2015; Kamasawa et al. 2005; Kofuji and Newman 2004; Magnotti et al. 2011; Nagy et al. 2001; Wasseff and Scherer 2011). In this model, inward  $K^+$  currents are generated by active uptake by astrocytes and oligodendrocytes near depolarizing cells and sustained by conduction of buffered  $K^+$  to blood vessels for elimination (Figure 2). Failure to remove sufficient  $K^+$  results in adequate change in resting membrane potential to allow spontaneous action potentials. However, while this activity is critical to physiologic excitatory activity, the pan-glial network exercises additional GJ and HC mediated effects on neuronal excitability both directly and indirectly.

Astrocyte  $Ca^{2+}$  waves are associated with neuronal activity and are conducted through astrocyte GJs and HCs (Torres et al. 2012). Glutamatergic activity induces decline in extracellular  $Ca^{2+}$  accompanied by rapid elevation of intracellular  $Ca^{2+}$  in both astrocytes and neurons. After a delay, astrocytes then exhibit a secondary “slow” intracellular  $Ca^{2+}$  wave initiated by ATP release by Panx1 HCs acting in an autocrine fashion (Torres et al. 2012). These waves are not attributable to glutamate signaling and require Cx30 and Cx43 GJs to travel between neighboring astrocytes (Torres et al. 2012). Importantly, astrocyte slow waves are associated with activation of the metabotropic purine receptor P2Y1 on nearby inhibitory interneurons, leading to increased inhibitory post synaptic currents (IPSCs)

(Torres et al. 2012). This suggests that  $\text{Ca}^{2+}$  slow waves may be involved in augmenting feedback inhibition through purinergic signaling, expanding the ways that they modify glutamatergic transmission.

Additional modulation of excitatory activity is exerted by astrocytes through Cx dependent regulation of synaptic glutamate release. Interestingly, astrocyte expressed Cx isoforms appear to affect glutamatergic transmission by distinct mechanisms. Basal quantal release of glutamate is set by Cx43 at excitatory synapses without altering the threshold for depolarization or the activity of excitatory amino acid transporters (Chever et al. 2014b). Instead, postsynaptic glutamate release is augmented by purinergic signaling mediated by astrocyte HCs (Chever et al. 2014a).

In contrast, Cx30 buffers synaptic glutamate through non-GJ mediated activities (Pannasch et al. 2014). Organotypic slices from Cx30—KO mice exhibit decreased glutamatergic activity independent of presynaptic quantal release or postsynaptic glutamate sensitivity. Instead, Cx30 regulates astrocyte process migration and glutamate transporter 1 (GLT-1) efficacy. In a 2014 study, Pannasch et al. observed that Cx30—KO astrocyte processes enriched in GLT-1 invade excitatory synapses more deeply than wild type astrocytes, thereby reducing glutamate within the synaptic cleft (Pannasch et al. 2014). This activity is independent of Cx30's GJ function, possibly relying on regulation by proteins complexed to Cx30 GJs.

Oligodendrocytes are commonly overlooked in the composition of the pan-glia network, instead thought of as merely myelinating projection axons. However, emerging evidence indicates that oligodendrocytes are also essential to maintaining physiologic generation of action potentials independent of their myelinating function (Dermietzel et al. 1989; Domercq et al. 2010; Nagy et al. 2003). Oligodendrocytes comprise a varying fraction of pan-glia network participants that form biocytin-permeable homotypic GJs with other oligodendrocytes and  $\text{NG2}^+/\text{Olig2}^+\text{CNPase}^+$  oligodendrocyte precursors in white matter (Maglione et al. 2010) as well as heterotypic GJs with astrocytes throughout the CNS (Griemsmann et al. 2015).

In myelinated fiber tracts, oligodendrocytes participate in spatial  $\text{K}^+$  buffering at the level of periaxonal myelin, which conducts ions through reflexive Cx32 GJs within myelin layers closest to the axolemma and strings of Cx32 GJs connecting adjacent paranodal loops (Kamasawa et al. 2005). Nearby astrocyte processes then siphon  $\text{K}^+$  from myelin and oligodendrocyte somata for elimination through heterotypic Cx47—43 and Cx32—30 GJs (Kamasawa et al. 2005). In genetic ablation studies, mice lacking both Cx32 and Cx47 exhibit periaxonal myelin vacuolation in the optic nerve that is worsened by retinal ganglion activity (Menichella et al. 2006). Interestingly, the authors of this study also found that mice lacking the inward rectifier  $\text{K}^+$  channel Kir4.1 deficient mice exhibited myelin vacuolation in spinal cord grey matter (Menichella et al. 2006), possibly due to osmotic damage due to failure to disperse rising  $\text{K}^+$  concentration within the myelin sheath.

In a recent study from Bettefeld et al., satellite oligodendrocytes (sOLs) associated with the axon initial segment of somatosensory cortex layer V pyramidal neurons were predicted to

dampen neuronal bursting by computational modelling (Battfeld et al. 2016). Their model was validated experimentally and found to be mediated by activity dependent inward rectifying  $K^+$  (Kir) currents in sOLs. Furthermore, GJ blockade using the non-specific Cx/Panx inhibitor carbenoxolone (CBX) abolished Kir currents and reduced voltage coupling between sOLs and adjacent astrocytes (Battfeld et al. 2016). These findings support the importance of oligodendrocytes in maintaining homeostatic extracellular ion concentrations, with implications for disease states involving white matter injury, such as MS.

### 3.3 Hemichannels and purinergic signaling in synaptic plasticity, learning, and memory

Purinergic signaling molecules released by Cx and Panx HCs acting on neuronal and glial P2X and P2Y receptors comprise a fundamental component of synaptic plasticity (Kim and Kang 2011; Prochnow et al. 2012; Thompson et al. 2008; Zoidl et al. 2007). Measurement of CA1 fEPSPs after CA3 Schaffer collateral tetanic stimulation shows that LTP is abnormally increased in mice lacking Panx1 HCs (Panx1—KO) (Ardiles et al. 2014; Prochnow et al. 2012). Interestingly, in addition to elevated fEPSPs, stimulation of Panx1—KO Schaffer collateral projections results in absent long term depression (LTD) in CA1 pyramidal neurons (Ardiles et al. 2014; Prochnow et al. 2012). Behaviorally, Panx1—KO mice also perform worse at novel object recognition and spatial recall tasks (Prochnow et al. 2012). Thus, purines released by Panx1 HCs represent a negative feedback mechanism dampening exaggerated LTP, which has been shown to impair spatial learning (Moser et al. 1998). Remarkably, physiologic LTP is restored in Panx1—KO mice when Schaffer collateral stimulation is accompanied by a mix of gliotransmitters including adenosine and ATP, possibly reflecting hyperpolarization through ATP-dependent  $K^+$  channels (Kawamura et al. 2004) or adenosine mediated inhibition through  $A_1$  and  $A_2$  receptors (Boison and Aronica 2015; Diogenes et al. 2014).

Astrocyte HCs also participate in purinergic signaling and learning. In a 2014 study from Chever et al., blockade of Cx43 HCs with the Cx43 HC-specific inhibitor gap26 resulted in reduced fEPSPs recorded in CA1 neurons after Schaffer collateral stimulation (Chever et al. 2014a). These results were replicated when Cx43 HCs were left intact, but P2 receptors were blocked using the P2 antagonists RB2 and PPADS (Chever et al. 2014a), suggesting purinergic signaling molecules released by Cx43 HCs were responsible for modifying glutamatergic activity after stimulation.

A similar in vivo study examining the role of Cx43 HCs in fear conditioning supports these findings. In a 2012 study by Stehberg et al., microinjection of the Cx43 HC specific blocking peptide TAT-Cx43L2 or gap27 into the basolateral amygdala resulted in amnesia to fear conditioning training (Stehberg et al. 2012). Interestingly, co-administration of these blocking peptides with a gliotransmitter cocktail including glutamate, glutamine, lactate, D-serine, glycine, and ATP restored fear conditioning memory. This effect was not observed when either peptide was injected several hours after training, indicating that Cx43 HCs participated in short term fear memory consolidation (Stehberg et al. 2012). While this study did not implicate a specific gliotransmitter in mediating the observed effects, it bolsters slice recording findings that implicate HCs in learning and memory.

## 4. A little too excited: connexin & pannexin dysregulation and seizures

### 4.1 Electrical synapses in seizures

Seizures are characterized by aberrant hypersynchronous excitatory activity that may be localized to a specific population of neurons or generalized throughout several brain regions. Classically, seizure development is thought to occur when inhibitory GABAergic transmission is compromised or overwhelmed by glutamatergic activity. Under homeostatic conditions, a range of mechanisms exist that confer resistance to seizures, including principle cell inhibition and desynchronizing activity of GABAergic networks (Apostolides and Trussell 2013; Deans et al. 2001; Landisman et al. 2002). Damage to or dysfunction of these networks may be involved in seizure initiation and propagation (Schwaller et al. 2004; Toyoda et al. 2015).

As the primary Cx isoform expressed in the adult CNS, Cx36 GJ-coupled GABAergic populations are critical for setting basal inhibitory tone in several brain regions (Belluardo et al. 2000; Deans et al. 2001; Turecek et al. 2014; Vandecasteele et al. 2006; Venance et al. 2004), and may be indispensable to maintaining physiologic resistance to epileptiform activity. However, the contribution of Cx36—coupling to seizures remains uncertain due to disparate—and occasionally contrary—reporting. Studies utilizing global ablation of Cx36 GJs have yielded data indicating an anti-epileptogenic (Jacobson et al. 2010), pro-epileptogenic (Maier et al. 2002), and bystander (Beaumont and Maccaferri 2011; Voss et al. 2010b) role for these molecules.

In support of an anti-epileptogenic function, Cx36—KO mice exhibit increased susceptibility to seizure induction in the pentylenetetrazol (PTZ) model of epilepsy (Jacobson et al. 2010). Consistent with these findings, pharmacological GJ blockade by CBX or quinine, which preferentially inhibits Cx36 and Cx50 (Srinivas et al. 2001), resulted in increased cortical epileptiform activity in organotypic slice culture using the low-Mg<sup>2+</sup> model of seizure induction (Voss et al. 2009). Jacobson et al. hypothesize that this phenomenon may be the result of autoinhibition of GABAergic Cx36-coupled neurons (Jacobson et al. 2010). In this study, the authors suggest that although the electrical synapses responsible for excitation of linked inhibitory networks would be absent in Cx36—KO animals, GABAergic chemical synapses on nearby inhibitory interneurons would remain. The resulting inhibition of inhibitory cells could feasibly give rise to aberrant excitatory activity by failing to suppress collateral excitation of surrounding pyramidal neurons or providing feedback inhibition (Jacobson et al. 2010). This hypothesis is supported by lines of evidence that indicate that in Cx36—KO mice, hippocampal epileptiform activity induced by kainate exposure may be partially attributable to reduced GABA<sub>A</sub> receptor activity (Pais et al. 2003).

In contrast, Maier et al. found that Cx36—KO hippocampi were less susceptible to seizures induced by the K<sup>+</sup> channel inhibitor, 4-aminopyridine (4-AP) (Maier et al. 2002). Their recordings from Cx36—KO slices showed less frequent spontaneous sharp wave events than wild type hippocampi with fewer and slower ripples. These data suggest that Cx36 GJs help coordinate the excitatory hypersynchronization thought to generate sharp wave-ripple complexes (Schlingloff et al. 2014), with their activity becoming pathological after 4-AP



exposure. Surprisingly, while ongoing seizurelike events were reduced in Cx36—KO mice, a portion of the slices recorded from failed to exhibit any activity whatsoever in response to 100 $\mu$ M 4-AP, indicating that Cx36 blockade could represent an effective method of managing seizures. Corroborating this, relatively selective Cx36 GJ inhibition by quinine or CBX application after 4-AP-induced seizures suppressed the amplitude of epileptiform activity and reduced involvement of the contralateral cortex in vivo (Gajda et al. 2005). However, to qualify these results, although their summed ictal activity was lower than control animals, quinine treated rats displayed more frequent seizures of shorter duration.

Additional complication in understanding Cx36-mediated activity in seizures comes from studies that find it dispensable to seizure initiation and resistance. Mefloquine, a quinine derivative inhibitor of Cx36 and Cx50 GJs at low doses (Cruikshank et al. 2004), failed to elicit any change in seizure-like event amplitude, frequency, or duration in low Mg<sup>2+</sup> and aconitine treated neocortical slice preparations from wild type mice compared control treated and Cx36—KOs (Voss et al. 2010b). Similarly, in a 2011 study, Beaumont and Maccaferri found that 4-AP treated CA1 interneurons of Cx36—KO mice exhibit no difference in epileptiform GABAergic currents relative to wild type despite uncoupling of interneurons across hippocampal strata (Beaumont and Maccaferri 2011). Furthermore, the authors demonstrated that CBX, but not mefloquine, inhibits GABA<sub>A</sub> receptors independently of the allosteric regulatory site of benzodiazepines, calling into question the use of CBX to study Cx contributions to seizures.

The use of Cx36—KO mice also presents an important limitation in the study of epilepsy. In the low Mg<sup>2+</sup> model of seizure, slices from these mice exhibit a GJ-independent increase in picrotoxin-sensitive GABAergic augmentation (Voss et al. 2010a). Wild type slices treated with mefloquine prior to seizure induction did not exhibit picrotoxin or etomidate-sensitive augmentation, indicating these effects were the result of compensatory changes in Cx36—KO animals. While differences in model may underlie the variegated findings in these reports, they also highlight the need for selective Cx36 inhibitors in the study of electrical synapse dysregulation and its contribution to neuronal hyperexcitability.

#### 4.2 Inflammation, seizure, and pan-glial network gap junction dysregulation

Pan-glial network maintenance of a sufficiently high seizure threshold in the homeostatic CNS relies on various GJ and non-GJ functions (Chever et al. 2014b; Pannasch et al. 2014). Spatial buffering of K<sup>+</sup> and other metabolites released during periods of increased neuronal depolarization is critical to constraining hypersynchronous excitatory activity and reducing generalization of epileptiform bursting when initiated (Battfeld et al. 2016; Bedner et al. 2015; Kamasawa et al. 2005; Kofuji and Newman 2004; Magnotti et al. 2011; Wasseff and Scherer 2011). Astrocytes are central to the pan-glial network's ability to carry out this function and recent evidence illustrates that dysregulation of astrocyte Cx43 GJs may drive the pathogenesis of some human temporal lobe epilepsy.

In a 2015 study from Bedner et al., hippocampal astrocytes from surgical specimens of sclerotic hippocampi from epileptic patient anomalously expressed ionotropic glutamate receptors, but not glutamate transporters (Bedner et al. 2015). In addition, dye loading experiments in these astrocytes revealed severely restricted diffusion of biocytin into

neighboring cells, suggesting they had become uncoupled from the pan-gial network (Bedner et al. 2015). These results were replicated in vivo using the 3 month post-intrahippocampal kainate (KA) injection mouse model of epileptogenesis (Bedner et al. 2015).

In the same study, Cx43-eYFP labeled astrocyte uncoupling was induced by a single injection of KA and persisted for at least six months and preceded neuronal apoptosis (Bedner et al. 2015). Astrocyte uncoupling was recapitulated in organotypic slice and in vivo by application of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), and interleukin (IL)-1 $\beta$  (Bedner et al. 2015), which have been identified in serum and cerebrospinal fluid of epileptic patients (Vezzani et al. 2011). Interestingly, another study using 7 days post-intrahippocampal KA injection model showed an increase in dye coupling of hippocampal reactive astrocytes and an increase in GFAP and Cx43 protein (Takahashi et al. 2010), indicating time-dependent changes in coupling of astrocyte GJs after seizure.

Seizure activity could also be due to loss of oligodendrocyte Cxs and reduction of Cx43 expression as observed in immune-mediated mouse models of MS (Brand-Schieber et al. 2005; Markoullis et al. 2012a). Furthermore, demyelinating MS lesions show evidence of GJ dysregulation, where inflamed white matter tracts and adjacent normal appearing tissue exhibit decreased Cx43 expression and oligodendrocyte uncoupling from reactive astrocytes (Markoullis et al. 2012b; Markoullis et al. 2014). In addition, MS patients are three to six times more likely to develop epilepsy than the overall population (Poser and Brinar 2003). While the mechanism that predisposes a subset of these patients to seizures remains unclear, a recent study identified altered expression of the astrocytic water channel implicated in epileptogenesis, aquaporin (AQP)4 (Binder et al. 2012), and extensive infiltration of microglia/macrophages into the CA1 of mice that experienced seizures following chronic cuprizone (CPZ)-induced demyelination (Lapato et al. 2017). However, although Cx47 redistribution has been noted in CPZ demyelination (Parenti et al. 2010), whether GJ dysfunction is etiologic in seizure development in this model is still unknown.

Transcriptome analysis of Cx and Panx expression following seizure induction using a cobalt (Co<sup>2+</sup>) model of seizure genesis demonstrated significant upregulation of Panx1, Panx2, and Cx43 mRNAs and post-translational phosphorylation of Cx43 (Mylvaganam et al. 2010), which is associated with increased GJ open probability and seizure (Zador et al. 2008). Protein and mRNA changes were independent of Co<sup>2+</sup> administration and required epileptiform discharges, since Co<sup>2+</sup> application in the presence of tetrodotoxin, a voltage gated Na<sup>+</sup> channel blocker, ablated transcriptome changes (Mylvaganam et al. 2010).

Intracellular Ca<sup>2+</sup> slow wave propagation and HC mediated purinergic signaling amongst GJ coupled astrocytes also influences seizure development (Kekesi et al. 2015; Torres et al. 2012). In the low Mg<sup>2+</sup> model of seizure, slow Ca<sup>2+</sup> waves become synchronized across nearby astrocytes, which become paired to synchronized neurons during seizure-like events (Kekesi et al. 2015). This aberrant neuron-gial synchrony is GJ dependent, as CBX and anti-Cx43 antibodies reduce epileptiform activity-induced Ca<sup>2+</sup> slow wave coordination, increased interictal periods, and completely ablated seizures in a number of trials (Kekesi et al. 2015). However, this study does not distinguish between Cx43 GJs and HCs, so whether

the observed anti-epileptic effects of these molecules is due to inhibited  $\text{Ca}^{2+}$  second messenger exchange across coupled astrocytes or impaired gliotransmitter release remains to be demonstrated.

### 4.3 Microglia, hemichannels, and inflammation

Microglia express Cx32 and Cx36 under homeostatic conditions, but because microglia do not form heterotypic or homotypic GJs (Wasseff and Scherer 2014), their physiologic role remains unclear (Dobrenis et al. 2005; Maezawa and Jin 2010). Research investigating microglial Cx and Panx HC function often occurs in the context of disease, where they have been implicated in excitotoxic glutamatergic signaling (Abudara et al. 2015; Eugenin et al. 2001; Mandolesi et al. 2013; Takaki et al. 2012; Takeuchi et al. 2006). Using organotypic slices, Adubara et al. demonstrated that LPS application augments astrocyte Cx43 HC opening, resulting in enhanced synaptic glutamate (Abudara et al. 2015). The authors identified TLR4 activation by LPS, which triggers secretion of inflammatory mediators such as  $\text{TNF}\alpha$ , inducible nitric oxide synthase, and  $\text{IL-1}\beta$  (Akira and Takeda 2004), was required for HC dependent glutamate release. Interestingly, fEPSP amplitude was suppressed in LPS treated slices despite increased synaptic glutamate, but was restored by blockade of Cx43 by gap26 or  $\text{TNF}\alpha/\text{IL-1}\beta$  inhibitors  $\text{IL-1RA}$  and s $\text{TNF-}\alpha\text{R1}$  (Abudara et al. 2015). While decreased glutamatergic transmission and its recovery following  $\text{IL-1RA}$  administration may partially be explained by  $\text{IL-1}\beta$ 's dampening effect on post-synaptic glutamate sensitivity (Mandolesi et al. 2013), this study illustrates how microglia modify neuronal activity in pathology.

Microglial Panx1 and Cx HCs also release purinergic signaling molecules under inflammatory conditions. In a 2015 report by Orellana et al., restraint stress increased hippocampal microglial, astrocyte, and neuronal ethidium bromide uptake, which was abrogated by application of the Panx1 HC blocking peptide  $^{10}\text{panx1}$  but not Cx43 HC blockers (Orellana et al. 2015). These changes in dye uptake were abolished by application of ionotropic purinergic receptor  $\text{P2X}_7$  and NMDAR antagonists, but not metabotropic purinergic  $\text{P2Y}_1$  receptor blockade, and were accompanied by increased extracellular glutamate and ATP (Orellana et al. 2015). This suggests that ionotropic purinergic signaling downstream of Panx1-mediated gliotransmitter release may be a vehicle of microglial HC dysfunction during inflammation, such as those generated by chronic stress (Walker et al. 2013).

In addition to TLR4 ligands, microglial HCs open in response to inflammatory cytokines. Application of LPS or  $\text{TNF}\alpha$  to primary microglia in vitro results in upregulation of glutaminase and glutamate release (Takeuchi et al. 2006). Furthermore, supernatant from glutamate—releasing microglial cultures induces downregulation of the astrocyte glutamate aspartate transporter, GLAST (Takaki et al. 2012). Together, these results indicate that microglial HCs alter extracellular glutamate concentration and astrocyte-dependent glutamate buffering in pathology. However, because in vitro models often lack the complex regulatory environment of the CNS, in vivo studies are required to validate these findings.

#### 4.4 Purinergic signaling in epilepsy

Like Cx and Panx HCs, the P2X<sub>7</sub> ionotropic purinergic receptor is a relatively large, ATP-gated transmembrane channel permeable to molecules 800 kDa or smaller, including low to medium weight cations such as K<sup>+</sup> and Ca<sup>2+</sup> (Sperlagh and Illes 2014) and cytokines (i.e. IL-1 $\beta$ ) (Monif et al. 2016). Activation of NMDARs results in a burst of ATP release by synaptic and astrocyte-derived Panx1 and Cx43 HCs, respectively, that amplify depolarization (Kim and Kang 2011; Prochnow et al. 2012; Thompson et al. 2008). Activity dependent ATP signaling is facilitated by Panx1 coupling to P2X<sub>7</sub> in membrane complexes (Bravo et al. 2015; Pan et al. 2015), thereby ensuring that NMDAR activation is augmented by the ensuing purinergic drive.

Dysregulation of this interaction has been implicated in the pathogenesis of seizure, but evidence is highly model dependent. Increased Panx1 protein expression was detected in lobectomy specimens from epileptic patients, while Panx2 was decreased (Jiang et al. 2013), indicating that purinergic signaling dysregulation may occur in these patients. Organotypic slice studies support this finding, showing that in the low Mg<sup>2+</sup> model of seizure, blockade of Panx1 by CBX and NMDAR antagonists attenuated epileptiform burst frequency and amplitude (Thompson et al. 2008). This suggests that feed-forward augmentation of NMDAR currents can initiate runaway seizure like activity. However, while the putative mechanism of epileptiform bursting is NMDAR activation in this model, synchronization of principle neurons GABAergic interneuron populations may have contributed to neuronal hyper synchronization through low Mg<sup>2+</sup> sensitive Cx36 GJs (Palacios-Prado et al. 2013). Thus, selective inhibition of Panx1 HCs may be required to untangle specific contribution of Panx1 to seizures in this model.

Alternatively, P2X<sub>7</sub>—KO mice display augmented seizure susceptibility when challenged with the muscarinic agonist pilocarpine that is not attributable to changes in glutamatergic or GABAergic transmission (Kim and Kang 2011). P2X<sub>7</sub> or Panx1 blockade in wild type mice reproduced the decreased seizure threshold observed in knockout animals (Kim and Kang 2011). Physiologic seizure resistance was restored in P2X<sub>7</sub>—KO mice by inhibition of intracellular Ca<sup>2+</sup> release by the ryanodine receptor antagonist dantrolene (Kim and Kang 2011). In contrast, P2X<sub>7</sub> antagonism by JNJ-42253432 resulted in a less severe seizure profile in the kainate model of epilepsy, with decreased seizure severity, but not frequency in Sprague-Dawley rats (Amhaoul et al. 2016). These studies suggest that purinergic signaling in epilepsy may be complicated by the choice of model and potentially species examined. Further research is required to fully illuminate how Panx1 function in seizures relates to the complex neuropathology observed in the epileptic CNS.

#### 5. Concluding remarks

Evidence increasingly indicates that Cx and Panx GJs and HCs are critical to maintaining physiologic neuronal excitability, resistance to seizure, and may be central to hippocampus and amygdala based learning (Apostolides and Trussell 2013; Battefeld et al. 2016; Prochnow et al. 2012; Stehberg et al. 2012). In GABAergic interneurons, Cx36 GJs allow for rapid and expansive inhibition via electrically coupled inhibitory syncytia that are calibrated to depolarizing activity (Apostolides and Trussell 2013; Deans et al. 2001;

Turecek et al. 2014). In glia, GJs link participants in the pan-glial network, which supports physiologic resting membrane potential through spatial buffering of K<sup>+</sup> and constrains epileptiform bursting (Battfeld et al. 2016; Bedner et al. 2015; Kamasawa et al. 2005; Kofuji and Newman 2004; Magnotti et al. 2011; Wasseff and Scherer 2011). Purinergic and glutamatergic gliotransmission, which continues to accrue attention for its role in modifying neuronal activity, relies heavily on autocrine and paracrine signaling by Cx and Panx HCs (Kim and Kang 2011; Prochnow et al. 2012; Thompson et al. 2008).

Inflammation-induced dysregulation of Cx GJs not only impacts glutamatergic neurotransmission (Abudara et al. 2015; Mandolesi et al. 2013; Schwaller et al. 2004; Takaki et al. 2012; Takeuchi et al. 2006; Toyoda et al. 2015), but also underlies disorders of excitation, such as epilepsy (Bedner et al. 2015). However, the role of Cx and Panx HCs in purinergic signaling and excitotoxicity remains conflicted, exhibiting a model-dependent effect on the result of ionotropic purinergic receptor activation (Amhaoul et al. 2016; Kim and Kang 2011; Thompson et al. 2008). Furthermore, the contribution of microglial Cxs and Panxs to homeostatic or disease processes remains largely unknown. Apart from a handful of studies (Abudara et al. 2015; Orellana et al. 2015), existing evidence regarding microglial HC and GJ function in pathology is restricted to in vitro work and focuses on TLR4 signaling (Takaki et al. 2012; Takeuchi et al. 2006), which may produce artifacts not seen in vivo (Wasseff and Scherer 2014).

An important limitation to many studies of individual Cx and Panx function is the lack of selective and specific GJ and HC inhibitors. Mimetic peptides and antibodies that inhibit Panx1 (<sup>10</sup>panx1) and Cx43 (gap26, gap27, and TAT-Cx43L2) HCs with relatively high selectivity are now available (Table 2). However, many of these peptides also inhibit GJs, making blockade of HCs or GJs alone difficult in situ (Giaume et al. 2013). Additionally, peptides targeting other Cx and Panx isoforms have not been developed, making their study more arduous and reliant on knockout models. However, as the tools available for their study become more varied and available, the study of Cxs and Panxs is likely to more easily uncover the homeostatic and pathological properties of HCs and GJs.

## Acknowledgments

This work was generously supported by the National Multiple Sclerosis Society Grants NMSS-RG-4853A3/2 and NIH R01-7R01NS081141 Grants to S.K.T.-W.

## References

- Abudara V, Roux L, Dallerac G, Matias I, Dulong J, Mothet JP, Rouach N, Giaume C. Activated microglia impairs neuroglial interaction by opening Cx43 hemichannels in hippocampal astrocytes. *Glia*. 2015; 63(5):795–811. [PubMed: 25643695]
- Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol*. 2004; 4(7):499–511. [PubMed: 15229469]
- Amhaoul H, Ali I, Mola M, Van Eetveldt A, Szewczyk K, Missault S, Bielen K, Kumar-Singh S, Rech J, Lord B, Ceusters M, Bhattacharya A, Dedeurwaerdere S. P2X7 receptor antagonism reduces the severity of spontaneous seizures in a chronic model of temporal lobe epilepsy. *Neuropharmacology*. 2016; 105:175–185. [PubMed: 26775823]
- Apostolides PF, Trussell LO. Regulation of interneuron excitability by gap junction coupling with principal cells. *Nat Neurosci*. 2013; 16(12):1764–1772. [PubMed: 24185427]

- Ardiles AO, Flores-Munoz C, Toro-Ayala G, Cardenas AM, Palacios AG, Munoz P, Fuenzalida M, Saez JC, Martinez AD. Pannexin 1 regulates bidirectional hippocampal synaptic plasticity in adult mice. *Front Cell Neurosci.* 2014; 8:326. [PubMed: 25360084]
- Battefeld A, Klooster J, Kole MH. Myelinating satellite oligodendrocytes are integrated in a glial syncytium constraining neuronal high-frequency activity. *Nat Commun.* 2016; 7:11298. [PubMed: 27161034]
- Beaumont M, Maccaferri G. Is connexin36 critical for GABAergic hypersynchronization in the hippocampus? *J Physiol.* 2011; 589(Pt 7):1663–1680. [PubMed: 21300748]
- Bedner P, Dupper A, Huttman K, Muller J, Herde MK, Dublin P, Deshpande T, Schramm J, Haussler U, Haas CA, Henneberger C, Theis M, Steinhäuser C. Astrocyte uncoupling as a cause of human temporal lobe epilepsy. *Brain.* 2015; 138(Pt 5):1208–1222. [PubMed: 25765328]
- Belluardo N, Mudo G, Trovato-Salinaro A, Le Gurun S, Charollais A, Serre-Beinier V, Amato G, Haefliger JA, Meda P, Condorelli DF. Expression of connexin36 in the adult and developing rat brain. *Brain Res.* 2000; 865(1):121–138. [PubMed: 10814742]
- Bikbaev A, Manahan-Vaughan D. Relationship of hippocampal theta and gamma oscillations to potentiation of synaptic transmission. *Front Neurosci.* 2008; 2(1):56–63. [PubMed: 18982107]
- Binder DK, Nagelhus EA, Ottersen OP. Aquaporin-4 and epilepsy. *Glia.* 2012; 60(8):1203–1214. [PubMed: 22378467]
- Boison D, Aronica E. Comorbidities in Neurology: Is adenosine the common link? *Neuropharmacology.* 2015; 97:18–34. [PubMed: 25979489]
- Brand-Schieber E, Werner P, Iacobas DA, Iacobas S, Beelitz M, Lowery SL, Spray DC, Scemes E. Connexin43, the major gap junction protein of astrocytes, is down-regulated in inflamed white matter in an animal model of multiple sclerosis. *J Neurosci Res.* 2005; 80(6):798–808. [PubMed: 15898103]
- Bravo D, Maturana CJ, Pelissier T, Hernandez A, Constandil L. Interactions of pannexin 1 with NMDA and P2X7 receptors in central nervous system pathologies: Possible role on chronic pain. *Pharmacol Res.* 2015; 101:86–93. [PubMed: 26211949]
- Bruzzone R, Hormuzdi SG, Barbe MT, Herb A, Monyer H. Pannexins, a family of gap junction proteins expressed in brain. *Proc Natl Acad Sci USA.* 2003; 100(23):13644–13649. [PubMed: 14597722]
- Cheng AW, Tang HY, Cai JL, Zhu M, Zhang XY, Rao M, Mattson MP. Gap junctional communication is required to maintain mouse cortical neural progenitor cells in a proliferative state. *Developmental Biology.* 2004; 272(1):203–216. [PubMed: 15242801]
- Cherian PP, Siller-Jackson AJ, Gu S, Wang X, Bonewald LF, Sprague E, Jiang JX. Mechanical strain opens connexin 43 hemichannels in osteocytes: a novel mechanism for the release of prostaglandin. *Mol Biol Cell.* 2005; 16(7):3100–3106. [PubMed: 15843434]
- Chever O, Lee CY, Rouach N. Astroglial connexin43 hemichannels tune basal excitatory synaptic transmission. *J Neurosci.* 2014a; 34(34):11228–11232. [PubMed: 25143604]
- Chever O, Pannasch U, Ezan P, Rouach N. Astroglial connexin 43 sustains glutamatergic synaptic efficacy. *Philos Trans R Soc Lond B Biol Sci.* 2014b; 369(1654):20130596. [PubMed: 25225090]
- Condorelli DF, Belluardo N, Trovato-Salinaro A, Mudo G. Expression of Cx36 in mammalian neurons. *Brain Res Brain Res Rev.* 2000; 32(1):72–85. [PubMed: 10751658]
- Cruikshank SJ, Hopperstad M, Younger M, Connors BW, Spray DC, Srinivas M. Potent block of Cx36 and Cx50 gap junction channels by mefloquine. *Proc Natl Acad Sci USA.* 2004; 101(33):12364–12369. [PubMed: 15297615]
- Deans MR, Gibson JR, Sellitto C, Connors BW, Paul DL. Synchronous activity of inhibitory networks in neocortex requires electrical synapses containing connexin36. *Neuron.* 2001; 31(3):477–485. [PubMed: 11516403]
- Dere E, Zheng-Fischhofer Q, Viggiano D, Gironi Carnevale UA, Ruocco LA, Zlomuzica A, Schnichels M, Willecke K, Huston JP, Sadile AG. Connexin31.1 deficiency in the mouse impairs object memory and modulates open-field exploration, acetylcholine esterase levels in the striatum, and cAMP response element-binding protein levels in the striatum and piriform cortex. *Neuroscience.* 2008; 153(2):396–405. [PubMed: 18384970]

- Dermietzel R, Traub O, Hwang TK, Beyer E, Bennett MV, Spray DC, Willecke K. Differential expression of three gap junction proteins in developing and mature brain tissues. *Proc Natl Acad Sci USA*. 1989; 86(24):10148–10152. [PubMed: 2557621]
- Diogenes MJ, Neves-Tome R, Fucile S, Martinello K, Scianni M, Theofilas P, Lopatar J, Ribeiro JA, Maggi L, Frenguelli BG, Limatola C, Boison D, Sebastiao AM. Homeostatic control of synaptic activity by endogenous adenosine is mediated by adenosine kinase. *Cereb Cortex*. 2014; 24(1):67–80. [PubMed: 22997174]
- Dobrenis K, Chang HY, Pina-Benabou MH, Woodroffe A, Lee SC, Rozental R, Spray DC, Scemes E. Human and mouse microglia express connexin36, and functional gap junctions are formed between rodent microglia and neurons. *J Neurosci Res*. 2005; 82(3):306–315. [PubMed: 16211561]
- Domercq M, Perez-Samartin A, Aparicio D, Alberdi E, Pampliega O, Matute C. P2X7 receptors mediate ischemic damage to oligodendrocytes. *Glia*. 2010; 58(6):730–740. [PubMed: 20029962]
- Ebihara L, Steiner E. Properties of a nonjunctional current expressed from a rat connexin46 cDNA in *Xenopus* oocytes. *J Gen Physiol*. 1993; 102(1):59–74. [PubMed: 7690837]
- Elias LA, Wang DD, Kriegstein AR. Gap junction adhesion is necessary for radial migration in the neocortex. *Nature*. 2007; 448(7156):901–907. [PubMed: 17713529]
- Esseltine JL, Laird DW. Next-Generation Connexin and Pannexin Cell Biology. *Trends in Cell Biology*. 2016
- Eugenin EA, Eckardt D, Theis M, Willecke K, Bennett MV, Saez JC. Microglia at brain stab wounds express connexin 43 and in vitro form functional gap junctions after treatment with interferon-gamma and tumor necrosis factor-alpha. *Proc Natl Acad Sci USA*. 2001; 98(7):4190–4195. [PubMed: 11259646]
- Frisch C, De Souza-Silva MA, Sohl G, Guldenagel M, Willecke K, Huston JP, Dere E. Stimulus complexity dependent memory impairment and changes in motor performance after deletion of the neuronal gap junction protein connexin36 in mice. *Behav Brain Res*. 2005; 157(1):177–185. [PubMed: 15617784]
- Froger N, Orellana JA, Calvo CF, Amigou E, Kozoriz MG, Naus CC, Saez JC, Giaume C. Inhibition of cytokine-induced connexin43 hemichannel activity in astrocytes is neuroprotective. *Mol Cell Neurosci*. 2010; 45(1):37–46. [PubMed: 20684043]
- Gajda Z, Szupera Z, Blazso G, Szente M. Quinine, a blocker of neuronal cx36 channels, suppresses seizure activity in rat neocortex in vivo. *Epilepsia*. 2005; 46(10):1581–1591. [PubMed: 16190928]
- Galarreta M, Hestrin S. A network of fast-spiking cells in the neocortex connected by electrical synapses. *Nature*. 1999; 402(6757):72–75. [PubMed: 10573418]
- Giaume C, Leybaert L, Naus CC, Saez JC. Connexin and pannexin hemichannels in brain glial cells: properties, pharmacology, and roles. *Front Pharmacol*. 2013; 4:88. [PubMed: 23882216]
- Giepmans BN, Verlaan I, Hengeveld T, Janssen H, Calafat J, Falk MM, Moolenaar WH. Gap junction protein connexin-43 interacts directly with microtubules. *Curr Biol*. 2001; 11(17):1364–1368. [PubMed: 11553331]
- Gotoh L, Inoue K, Helman G, Mora S, Maski K, Soul JS, Bloom M, Evans SH, Goto Y, Caldovic L, Hobson GM, Vanderver A. GJC2 promoter mutations causing Pelizaeus-Merzbacher-like disease. *Mol Genet Metab*. 2014; 111(3):393–398. [PubMed: 24374284]
- Griemsmann S, Hoft SP, Bedner P, Zhang J, von Staden E, Beinhauer A, Degen J, Dublin P, Cope DW, Richter N, Crunelli V, Jabs R, Willecke K, Theis M, Seifert G, Kettenmann H, Steinhauser C. Characterization of Panglial Gap Junction Networks in the Thalamus, Neocortex, and Hippocampus Reveals a Unique Population of Glial Cells. *Cereb Cortex*. 2015; 25(10):3420–3433. [PubMed: 25037920]
- Jacobson GM, Voss LJ, Melin SM, Mason JP, Cursons RT, Steyn-Ross DA, Steyn-Ross ML, Sleigh JW. Connexin36 knockout mice display increased sensitivity to pentylentetrazol-induced seizure-like behaviors. *Brain Res*. 2010; 1360:198–204. [PubMed: 20833151]
- Jiang T, Long H, Ma Y, Long L, Li Y, Li F, Zhou P, Yuan C, Xiao B. Altered expression of pannexin proteins in patients with temporal lobe epilepsy. *Mol Med Rep*. 2013; 8(6):1801–1806. [PubMed: 24146091]

- John SA, Kondo R, Wang SY, Goldhaber JI, Weiss JN. Connexin-43 hemichannels opened by metabolic inhibition. *J Biol Chem.* 1999; 274(1):236–240. [PubMed: 9867835]
- Kamasawa N, Sik A, Morita M, Yasumura T, Davidson KGV, Nagy JI, Rash JE. Connexin-47 and connexin-32 in gap junctions of oligodendrocyte somata, myelin sheaths, paranodal loops and Schmidt-Lanterman incisures: Implications for ionic homeostasis and potassium siphoning. *Neuroscience.* 2005; 136(1):65–86. [PubMed: 16203097]
- Kawamura M, Gachet C, Inoue K, Kato F. Direct excitation of inhibitory interneurons by extracellular ATP mediated by P2Y1 receptors in the hippocampal slice. *J Neurosci.* 2004; 24(48):10835–10845. [PubMed: 15574734]
- Kawamura M Jr, Ruskin DN, Masino SA. Metabolic autocrine regulation of neurons involves cooperation among pannexin hemichannels, adenosine receptors, and KATP channels. *J Neurosci.* 2010; 30(11):3886–3895. [PubMed: 20237259]
- Kekesi O, Ioja E, Szabo Z, Kardos J, Heja L. Recurrent seizure-like events are associated with coupled astroglial synchronization. *Front Cell Neurosci.* 2015; 9:215. [PubMed: 26150770]
- Kim JE, Kang TC. The P2X7 receptor-pannexin-1 complex decreases muscarinic acetylcholine receptor-mediated seizure susceptibility in mice. *J Clin Invest.* 2011; 121(5):2037–2047. [PubMed: 21505260]
- Klaassen LJ, Sun Z, Steijaert MN, Bolte P, Fahrenfort I, Sjoerdsma T, Klooster J, Claassen Y, Shields CR, Ten Eikelder HM, Janssen-Bienhold U, Zoidl G, McMahon DG, Kamermans M. Synaptic transmission from horizontal cells to cones is impaired by loss of connexin hemichannels. *PLoS Biol.* 2011; 9(7):e1001107. [PubMed: 21811399]
- Kofuji P, Newman EA. Potassium buffering in the central nervous system. *Neuroscience.* 2004; 129(4):1045–1056. [PubMed: 15561419]
- Kondo RP, Wang SY, John SA, Weiss JN, Goldhaber JI. Metabolic inhibition activates a non-selective current through connexin hemichannels in isolated ventricular myocytes. *J Mol Cell Cardiol.* 2000; 32(10):1859–1872. [PubMed: 11013130]
- Kreuzberg MM, Deuchars J, Weiss E, Schober A, Sonntag S, Wellershaus K, Draguhn A, Willecke K. Expression of connexin30.2 in interneurons of the central nervous system in the mouse. *Mol Cell Neurosci.* 2008; 37(1):119–134. [PubMed: 17942321]
- Kunze A, Congreso MR, Hartmann C, Wallraff-Beck A, Huttmann K, Bedner P, Requardt R, Seifert G, Redecker C, Willecke K, Hofmann A, Pfeifer A, Theis M, Steinhauser C. Connexin expression by radial glia-like cells is required for neurogenesis in the adult dentate gyrus. *Proc Natl Acad Sci U S A.* 2009; 106(27):11336–11341. [PubMed: 19549869]
- Landisman CE, Long MA, Beierlein M, Deans MR, Paul DL, Connors BW. Electrical synapses in the thalamic reticular nucleus. *J Neurosci.* 2002; 22(3):1002–1009. [PubMed: 11826128]
- Lapato AS, Szu J, Hasselmann JPC, Khalaj AJ, Binder DK, Tiwari-Woodruff SK. Chronic demyelination-induced seizures. *Neuroscience.* 2017
- Lasztozci B, Klausberger T. Layer-specific GABAergic control of distinct gamma oscillations in the CA1 hippocampus. *Neuron.* 2014; 81(5):1126–1139. [PubMed: 24607232]
- Leussis MP, Bolivar VJ. Habituation in rodents: a review of behavior, neurobiology, and genetics. *Neurosci Biobehav Rev.* 2006; 30(7):1045–1064. [PubMed: 16774787]
- Loewenstein WR. Junctional intercellular communication: the cell-to-cell membrane channel. *Physiol Rev.* 1981; 61(4):829–913. [PubMed: 6270711]
- MacVicar BA, Thompson RJ. Non-junction functions of pannexin-1 channels. *Trends Neurosci.* 2010; 33(2):93–102. [PubMed: 20022389]
- Maezawa I, Jin LW. Rett syndrome microglia damage dendrites and synapses by the elevated release of glutamate. *J Neurosci.* 2010; 30(15):5346–5356. [PubMed: 20392956]
- Maglione M, Tress O, Haas B, Karram K, Trotter J, Willecke K, Kettenmann H. Oligodendrocytes in mouse corpus callosum are coupled via gap junction channels formed by connexin47 and connexin32. *Glia.* 2010; 58(9):1104–1117. [PubMed: 20468052]
- Magnotti LM, Goodenough DA, Paul DL. Deletion of oligodendrocyte Cx32 and astrocyte Cx43 causes white matter vacuolation, astrocyte loss and early mortality. *Glia.* 2011; 59(7):1064–1074. [PubMed: 21538560]



- Maier N, Güldenagel M, Söhl G, Siegmund H, Willecke K, Draguhn A. Reduction of high-frequency network oscillations (ripples) and pathological network discharges in hippocampal slices from connexin 36-deficient mice. *J Physiol.* 2002; 541(2):521–528. [PubMed: 12042356]
- Mandolesi G, Musella A, Gentile A, Grasselli G, Haji N, Sepman H, Fresegna D, Bullitta S, De Vito F, Musumeci G, Di Sanza C, Strata P, Centonze D. Interleukin-1beta alters glutamate transmission at purkinje cell synapses in a mouse model of multiple sclerosis. *J Neurosci.* 2013; 33(29):12105–12121. [PubMed: 23864696]
- Marins M, Xavier AL, Viana NB, Fortes FS, Froes MM, Menezes JR. Gap junctions are involved in cell migration in the early postnatal subventricular zone. *Dev Neurobiol.* 2009; 69(11):715–730. [PubMed: 19565626]
- Markoullis K, Sargiannidou I, Gardner C, Hadjisavvas A, Reynolds R, Kleopa KA. Disruption of oligodendrocyte gap junctions in experimental autoimmune encephalomyelitis. *Glia.* 2012a; 60(7):1053–1066. [PubMed: 22461072]
- Markoullis K, Sargiannidou I, Schiza N, Hadjisavvas A, Roncaroli F, Reynolds R, Kleopa KA. Gap junction pathology in multiple sclerosis lesions and normal-appearing white matter. *Acta Neuropathol.* 2012b; 123(6):873–886. [PubMed: 22484441]
- Markoullis K, Sargiannidou I, Schiza N, Roncaroli F, Reynolds R, Kleopa KA. Oligodendrocyte gap junction loss and disconnection from reactive astrocytes in multiple sclerosis gray matter. *J Neuropathol Exp Neurol.* 2014; 73(9):865–879. [PubMed: 25101702]
- Masaki K. Early disruption of glial communication via connexin gap junction in multiple sclerosis, Balo's disease and neuromyelitis optica. *Neuropathology.* 2015; 35(5):469–480. [PubMed: 26016402]
- May D, Tress O, Seifert G, Willecke K. Connexin47 protein phosphorylation and stability in oligodendrocytes depend on expression of Connexin43 protein in astrocytes. *J Neurosci.* 2013; 33(18):7985–7996. [PubMed: 23637189]
- Menichella DM, Goodenough DA, Sirkowski E, Scherer SS, Paul DL. Connexins are critical for normal myelination in the CNS. *J Neurosci.* 2003; 23(13):5963–5973. [PubMed: 12843301]
- Menichella DM, Majdan M, Awatramani R, Goodenough DA, Sirkowski E, Scherer SS, Paul DL. Genetic and physiological evidence that oligodendrocyte gap junctions contribute to spatial buffering of potassium released during neuronal activity. *J Neurosci.* 2006; 26(43):10984–10991. [PubMed: 17065440]
- Monif M, Reid CA, Powell KL, Drummond KJ, O'Brien TJ, Williams DA. Interleukin-1beta has trophic effects in microglia and its release is mediated by P2X7R pore. *J Neuroinflammation.* 2016; 13(1):173. [PubMed: 27364756]
- Montero TD, Orellana JA. Hemichannels: new pathways for gliotransmitter release. *Neuroscience.* 2015; 286:45–59. [PubMed: 25475761]
- Morris RG, Moser EI, Riedel G, Martin SJ, Sandin J, Day M, O'Carroll C. Elements of a neurobiological theory of the hippocampus: the role of activity-dependent synaptic plasticity in memory. *Philos Trans R Soc Lond B Biol Sci.* 2003; 358(1432):773–786. [PubMed: 12744273]
- Moser EI, Krobot KA, Moser MB, Morris RG. Impaired spatial learning after saturation of long-term potentiation. *Science.* 1998; 281(5385):2038–2042. [PubMed: 9748165]
- Mylvaganam S, Zhang L, Wu C, Zhang ZJ, Samoilova M, Eubanks J, Carlen PL, Poulter MO. Hippocampal seizures alter the expression of the pannexin and connexin transcriptome. *J Neurochem.* 2010; 112(1):92–102. [PubMed: 19840216]
- Nagy JI, Ionescu AV, Lynn BD, Rash JE. Coupling of astrocyte connexins Cx26, Cx30, Cx43 to oligodendrocyte Cx29, Cx32, Cx47: Implications from normal and connexin32 knockout mice. *Glia.* 2003; 44(3):205–218. [PubMed: 14603462]
- Nagy JI, Li X, Rempel J, Stelmack G, Patel D, Staines WA, Yasumura T, Rash JE. Connexin26 in adult rodent central nervous system: demonstration at astrocytic gap junctions and colocalization with connexin30 and connexin43. *J Comp Neurol.* 2001; 441(4):302–323. [PubMed: 11745652]
- Niu J, Li T, Yi C, Huang N, Koulakoff A, Weng C, Li C, Zhao CJ, Giaume C, Xiao L. Connexin-based channels contribute to metabolic pathways in the oligodendroglial lineage. *Journal of Cell Science.* 2016; 129(9):1902–1914. [PubMed: 27006115]


- Odermatt B, Wellershaus K, Wallraff A, Seifert G, Degen J, Euwens C, Fuss B, Bussow H, Schilling K, Steinhauser C, Willecke K. Connexin 47 (Cx47)-deficient mice with enhanced green fluorescent protein reporter gene reveal predominant oligodendrocytic expression of Cx47 and display vacuolized myelin in the CNS. *J Neurosci.* 2003; 23(11):4549–4559. [PubMed: 12805295]
- Orellana JA, Moraga-Amaro R, Diaz-Galarce R, Rojas S, Maturana CJ, Stehberg J, Saez JC. Restraint stress increases hemichannel activity in hippocampal glial cells and neurons. *Front Cell Neurosci.* 2015; 9:102. [PubMed: 25883550]
- Pais I, Hormuzdi SG, Monyer H, Traub RD, Wood IC, Buhl EH, Whittington MA, LeBeau FE. Sharp wave-like activity in the hippocampus in vitro in mice lacking the gap junction protein connexin 36. *J Neurophysiol.* 2003; 89(4):2046–2054. [PubMed: 12686578]
- Palacios-Prado N, Hoge G, Marandykina A, Rimkute L, Chapuis S, Paulauskas N, Skeberdis VA, O'Brien J, Pereda AE, Bennett MV, Bukauskas FF. Intracellular magnesium-dependent modulation of gap junction channels formed by neuronal connexin36. *J Neurosci.* 2013; 33(11):4741–4753. [PubMed: 23486946]
- Pan HC, Chou YC, Sun SH. P2X7 R-mediated Ca(2+) -independent d-serine release via pannexin-1 of the P2X7 R-pannexin-1 complex in astrocytes. *Glia.* 2015; 63(5):877–893. [PubMed: 25630251]
- Pannasch U, Freche D, Dallerac G, Ghezali G, Escartin C, Ezan P, Cohen-Salmon M, Benchenane K, Abudara V, Dufour A, Lubke JH, Deglon N, Knott G, Holcman D, Rouach N. Connexin 30 sets synaptic strength by controlling astroglial synapse invasion. *Nat Neurosci.* 2014; 17(4):549–558. [PubMed: 24584052]
- Parenti R, Cicirata F, Zappala A, Catania A, La Delia F, Cicirata V, Tress O, Willecke K. Dynamic expression of Cx47 in mouse brain development and in the cuprizone model of myelin plasticity. *Glia.* 2010; 58(13):1594–1609. [PubMed: 20578039]
- Paul DL, Ebihara L, Takemoto LJ, Swenson KI, Goodenough DA. Connexin46, a novel lens gap junction protein, induces voltage-gated currents in nonjunctional plasma membrane of *Xenopus* oocytes. *J Cell Biol.* 1991; 115(4):1077–1089. [PubMed: 1659572]
- Penuela S, Bhalla R, Gong XQ, Cowan KN, Celetti SJ, Cowan BJ, Bai D, Shao Q, Laird DW. Pannexin 1 and pannexin 3 are glycoproteins that exhibit many distinct characteristics from the connexin family of gap junction proteins. *J Cell Sci.* 2007; 120(Pt 21):3772–3783. [PubMed: 17925379]
- Pereda AE, Curti S, Hoge G, Cachepe R, Flores CE, Rash JE. Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity. *Biochim Biophys Acta.* 2013; 1828(1):134–146. [PubMed: 22659675]
- Poser CM, Brinar VV. Epilepsy and multiple sclerosis. *Epilepsy & Behavior.* 2003; 4(1):6–12. [PubMed: 12609222]
- Prochnow N, Abdulazim A, Kurtenbach S, Wildforster V, Dvorianchikova G, Hanske J, Petrasch-Parwez E, Shestopalov VI, Dermietzel R, Manahan-Vaughan D, Zoidl G. Pannexin1 stabilizes synaptic plasticity and is needed for learning. *PLoS One.* 2012; 7(12):e51767. [PubMed: 23284764]
- Rash JE, Staines WA, Yasumura T, Patel D, Furman CS, Stelmack GL, Nagy JI. Immunogold evidence that neuronal gap junctions in adult rat brain and spinal cord contain connexin-36 but not connexin-32 or connexin-43. *P Natl Acad Sci USA.* 2000; 97(13):7573–7578.
- Retamal MA, Froger N, Palacios-Prado N, Ezan P, Saez PJ, Saez JC, Giaume C. Cx43 hemichannels and gap junction channels in astrocytes are regulated oppositely by proinflammatory cytokines released from activated microglia. *J Neurosci.* 2007; 27(50):13781–13792. [PubMed: 18077690]
- Rozenal R, Morales M, Mehler MF, Urban M, Kremer M, Dermietzel R, Kessler JA, Spray DC. Changes in the properties of gap junctions during neuronal differentiation of hippocampal progenitor cells. *J Neurosci.* 1998; 18(5):1753–1762. [PubMed: 9465000]
- Samoilova M, Wentlandt K, Adamchik Y, Velumian AA, Carlen PL. Connexin 43 mimetic peptides inhibit spontaneous epileptiform activity in organotypic hippocampal slice cultures. *Exp Neurol.* 2008; 210(2):762–775. [PubMed: 18284929]
- Santiago MF, Alcamí P, Striedinger KM, Spray DC, Scemes E. The Carboxyl-terminal Domain of Connexin43 Is a Negative Modulator of Neuronal Differentiation. *Journal of Biological Chemistry.* 2010; 285(16):11836–11845. [PubMed: 20164188]

- Santiago MF, Veliskova J, Patel NK, Lutz SE, Caille D, Charollais A, Meda P, Scemes E. Targeting pannexin1 improves seizure outcome. *PLoS One*. 2011; 6(9):e25178. [PubMed: 21949881]
- Sargiannidou I, Kim GH, Kyriakoudi S, Eun BL, Kleopa KA. A start codon CMT1X mutation associated with transient encephalomyelitis causes complete loss of Cx32. *Neurogenetics*. 2015; 16(3):193–200. [PubMed: 25771809]
- Schiza N, Sargiannidou I, Kagiava A, Karaiskos C, Nearchou M, Kleopa KA. Transgenic replacement of Cx32 in gap junction-deficient oligodendrocytes rescues the phenotype of a hypomyelinating leukodystrophy model. *Hum Mol Genet*. 2015; 24(7):2049–2064. [PubMed: 25524707]
- Schlingloff D, Kali S, Freund TF, Hajos N, Gulyas AI. Mechanisms of sharp wave initiation and ripple generation. *J Neurosci*. 2014; 34(34):11385–11398. [PubMed: 25143618]
- Schutte M, Chen SH, Buku A, Wolosin JM. Connexin50, a gap junction protein of macroglia in the mammalian retina and visual pathway. *Experimental Eye Research*. 1998; 66(5):605–613. [PubMed: 9628808]
- Schwaller B, Tetko IV, Tandon P, Silveira DC, Vreugdenhil M, Henzi T, Potier MC, Celio MR, Villa AE. Parvalbumin deficiency affects network properties resulting in increased susceptibility to epileptic seizures. *Mol Cell Neurosci*. 2004; 25(4):650–663. [PubMed: 15080894]
- Sevetson J, Haas JS. Asymmetry and modulation of spike timing in electrically coupled neurons. *J Neurophysiol*. 2015; 113(6):1743–1751. [PubMed: 25540226]
- Simard M, Nedergaard M. The neurobiology of glia in the context of water and ion homeostasis. *Neuroscience*. 2004; 129(4):877–896. [PubMed: 15561405]
- Sosinsky GE, Boassa D, Dermietzel R, Duffy HS, Laird DW, MacVicar B, Naus CC, Penuela S, Scemes E, Spray DC. Pannexin channels are not gap junction hemichannels. *Channels (Austin)*. 2011:5.
- Sperlagh B, Illes P. P2X7 receptor: an emerging target in central nervous system diseases. *Trends Pharmacol Sci*. 2014; 35(10):537–547. [PubMed: 25223574]
- Srinivas M, Hopperstad MG, Spray DC. Quinine blocks specific gap junction channel subtypes. *Proc Natl Acad Sci USA*. 2001; 98(19):10942–10947. [PubMed: 11535816]
- Stehberg J, Moraga-Amaro R, Salazar C, Becerra A, Echeverria C, Orellana JA, Bultynck G, Ponsaerts R, Leybaert L, Simon F, Saez JC, Retamal MA. Release of gliotransmitters through astroglial connexin 43 hemichannels is necessary for fear memory consolidation in the basolateral amygdala. *FASEB J*. 2012; 26(9):3649–3657. [PubMed: 22665389]
- Stout CE, Costantin JL, Naus CC, Charles AC. Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *J Biol Chem*. 2002; 277(12):10482–10488. [PubMed: 11790776]
- Swayne LA, Bennett SAL. Connexins and pannexins in neuronal development and adult neurogenesis. *BMC Cell Biology*. 2016; 17(1):S10.
- Takahashi D, Vargas JR, Wilcox K. Increased coupling and altered glutamate transport currents in astrocytes following kainic-acid-induced status epilepticus. *Neurobiol Dis*. 2010; 40(3):573–585. [PubMed: 20691786]
- Takaki J, Fujimori K, Miura M, Suzuki T, Sekino Y, Sato K. L-glutamate released from activated microglia downregulates astrocytic L-glutamate transporter expression in neuroinflammation: the ‘collusion’ hypothesis for increased extracellular L-glutamate concentration in neuroinflammation. *J Neuroinflammation*. 2012; 9:275. [PubMed: 23259598]
- Takeuchi H, Jin S, Wang J, Zhang G, Kawanokuchi J, Kuno R, Sonobe Y, Mizuno T, Suzumura A. Tumor necrosis factor- $\alpha$  induces neurotoxicity via glutamate release from hemichannels of activated microglia in an autocrine manner. *J Biol Chem*. 2006; 281(30):21362–21368. [PubMed: 16720574]
- Thompson RJ, Jackson MF, Olah ME, Rungta RL, Hines DJ, Beazely MA, MacDonald JF, MacVicar BA. Activation of pannexin-1 hemichannels augments aberrant bursting in the hippocampus. *Science*. 2008; 322(5907):1555–1559. [PubMed: 19056988]
- Torres A, Wang FS, Xu QW, Fujita T, Dobrowolski R, Willecke K, Takano T, Nedergaard M. Extracellular Ca<sup>2+</sup> Acts as a Mediator of Communication from Neurons to Glia. *Science Signaling*. 2012; 5(208)

- Toyoda I, Fujita S, Thamattoor AK, Buckmaster PS. Unit Activity of Hippocampal Interneurons before Spontaneous Seizures in an Animal Model of Temporal Lobe Epilepsy. *J Neurosci*. 2015; 35(16): 6600–6618. [PubMed: 25904809]
- Tress O, Maglione M, May D, Pivneva T, Richter N, Seyfarth J, Binder S, Zlomuzica A, Seifert G, Theis M, Dere E, Kettenmann H, Willecke K. Pannal gap junctional communication is essential for maintenance of myelin in the CNS. *J Neurosci*. 2012; 32(22):7499–7518. [PubMed: 22649229]
- Turecek J, Yuen GS, Han VZ, Zeng XH, Bayer KU, Welsh JP. NMDA receptor activation strengthens weak electrical coupling in mammalian brain. *Neuron*. 2014; 81(6):1375–1388. [PubMed: 24656255]
- Valiunas V, Weingart R. Electrical properties of gap junction hemichannels identified in transfected HeLa cells. *Pflugers Arch*. 2000; 440(3):366–379. [PubMed: 10954323]
- Vandecasteele M, Glowinski J, Venance L. Connexin mRNA expression in single dopaminergic neurons of substantia nigra pars compacta. *Neurosci Res*. 2006; 56(4):419–426. [PubMed: 17014920]
- Venance L, Glowinski J, Giaume C. Electrical and chemical transmission between striatal GABAergic output neurones in rat brain slices. *J Physiol*. 2004; 559(Pt 1):215–230. [PubMed: 15235091]
- Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in epilepsy. *Nat Rev Neurol*. 2011; 7(1):31–40. [PubMed: 21135885]
- Vis JC, Nicholson LF, Faull RL, Evans WH, Severs NJ, Green CR. Connexin expression in Huntington's diseased human brain. *Cell Biol Int*. 1998; 22(11-12):837–847. [PubMed: 10873295]
- Voss LJ, Jacobson G, Sleight JW, Steyn-Ross A, Steyn-Ross M. Excitatory effects of gap junction blockers on cerebral cortex seizure-like activity in rats and mice. *Epilepsia*. 2009; 50(8):1971–1978. [PubMed: 19486358]
- Voss LJ, Melin S, Jacobson G, Sleight JW. GABAergic compensation in connexin36 knockout mice evident during low-magnesium seizure-like event activity. *Brain Res*. 2010a; 1360:49–55. [PubMed: 20831861]
- Voss LJ, Mutsaerts N, Sleight JW. Connexin36 gap junction blockade is ineffective at reducing seizure-like event activity in neocortical mouse slices. *Epilepsy Res Treat*. 2010b; 2010:310753. [PubMed: 22937225]
- Walker FR, Nilsson M, Jones K. Acute and chronic stress-induced disturbances of microglial plasticity, phenotype and function. *Curr Drug Targets*. 2013; 14(11):1262–1276. [PubMed: 24020974]
- Wang MH, Chen N, Wang JH. The coupling features of electrical synapses modulate neuronal synchrony in hypothalamic superchiasmatic nucleus. *Brain Res*. 2014; 1550:9–17. [PubMed: 24440632]
- Wang Y, Belousov AB. Deletion of neuronal gap junction protein connexin 36 impairs hippocampal LTP. *Neurosci Lett*. 2011; 502(1):30–32. [PubMed: 21798314]
- Wasseff SK, Scherer SS. Cx32 and Cx47 mediate oligodendrocyte:astrocyte and oligodendrocyte:oligodendrocyte gap junction coupling. *Neurobiology of Disease*. 2011; 42(3): 506–513. [PubMed: 21396451]
- Wasseff SK, Scherer SS. Activated microglia do not form functional gap junctions in vivo. *J Neuroimmunol*. 2014; 269(1-2):90–93. [PubMed: 24582000]
- Weickert S, Ray A, Zoidl G, Dermietzel R. Expression of neural connexins and pannexin1 in the hippocampus and inferior olive: a quantitative approach. *Brain Res Mol Brain Res*. 2005; 133(1): 102–109. [PubMed: 15661370]
- Zador Z, Weiczner R, Mihaly A. Long-lasting dephosphorylation of connexin 43 in acute seizures is regulated by NMDA receptors in the rat cerebral cortex. *Mol Med Rep*. 2008; 1(5):721–727. [PubMed: 21479476]
- Zoidl G, Petrasch-Parwez E, Ray A, Meier C, Bunse S, Habbes HW, Dahl G, Dermietzel R. Localization of the pannexin1 protein at postsynaptic sites in the cerebral cortex and hippocampus. *Neuroscience*. 2007; 146(1):9–16. [PubMed: 17379420]


### Significance

Connexins and pannexins contribute to normal excitatory activity in the central nervous system (CNS). Cells linked by pairs of connexin hemichannels, called gap junctions, regulate levels of potentially toxic metabolites produced during neuronal activity. Unpaired connexin and pannexin hemichannels facilitate learning and memory by providing feedback to neurons through the release of signaling molecules. In this review, we describe how CNS cells utilize connexins and pannexins to perform these tasks. In addition, we discuss how disruption of connexin and pannexin activity leads to abnormal excitation and diseases such as epilepsy.



**Figure 1. Connexin and pannexin structure and organization**

**A** Connexin and pannexin hemichannels are hexamers composed of six isoform subunits. Connexin hemichannels may be paired with homotypic or heterotypic hemichannels on adjacent cells to allow exchange of cytoplasmic contents up to 1.5 kDa as gap junctions. Pannexin hemichannels are not thought to form gap junctions due to N-linked glycosylation patterns. **B, C** Connexins and pannexins are structurally and functionally homologous, but have distinct amino acid sequences. Each subunit possesses four transmembrane domains linked by one intracellular and two extracellular loops. The carboxyl and amine terminals extend into the cytoplasm. The carboxyl tail is the site of regulatory modification and phosphorylation. **C** Pannexin monomers are structurally related to connexins but glycosylation of the extracellular loop closest to the carboxyl terminal prohibits pannexin hemichannel assembly into functional gap junctions.



**Figure 2. The pan-glia network participates in regulating excitatory neuronal transmission through gap junction and hemichannel-mediated functions**

Connexin gap junctions control glutamatergic activity indirectly through generation of inward  $K^+$  currents & spatial  $K^+$  buffering, activity-dependent astrocyte  $Ca^{2+}$  slow wave propagation, and regulation of synaptic invasion by GLT-1 enriched astrocyte process. Model depicts coupling partners for homotypic and heterotypic glial GJs and their cellular expression. Cx43 and Panx1 HCs contribute to synaptic plasticity, inhibitory feedback, and glutamatergic tone by autocrine and paracrine release of synaptic gliotransmitters, including glutamate and ATP.

**Table 1**

Cellular distribution of Cx and Panx isoforms expressed by glia and neurons in the adult mammalian CNS. Astrocyte expressed Cx isoforms participate in HC-mediated gliotransmission and astrocyte-oligodendrocyte GJs within the pan-glial network. Cx36 HCs are expressed by microglia under homeostatic conditions, while Cx32 and Cx43 are upregulated in response to inflammation. Oligodendrocyte Cx isoforms contribute to GJ coupling with pan-glial network members and between myelin layers. Neuronal Cx heterogeneity reflects CNS regional specialization. Panx1 HCs are found in all CNS populations listed, but Panx2 is only identified in neurons.

Cell Type	Connexins	Pannexins	References
Astrocytes	Cx26, Cx30, Cx43,	Panx1	Dermietzel et al. 1989, Nagy et al. 2001, Zoidl et al. 2007
Microglia	Cx32 <sup>a</sup> , Cx36, Cx43 <sup>a</sup>	Panx1	Orellana et al. 2015, Dobrenis et al. 2005, Eugenin et al. 2001, Takeuchi et al. 2006
Oligodendrocytes	Cx29, Cx32, Cx47	Panx1	Dermietzel et al. 1989, Domercq et al. 2010, Nagy et al. 2003
Neurons	Cx30.2, Cx31.1, Cx32, Cx36, Cx40, Cx45, Cx50	Panx1, Panx2	Bruzzone et al. 2003, Dere et al. 2008, Kreuzberg et al. 2008, Rash et al. 2000, Rozental et al. 1998, Schutte et al. 1998, Vis et al. 1998, Weickert et al. 2005

<sup>a</sup> expressed by activated microglia



**Table 2**

Summary of genetic and pharmacologic manipulations used to identify the function of specific Cx and Panx isoforms. Key results are indicated beside manipulation. Note that selective inhibitors are only available for HCs, but many will also block GJs over time (Samoilova et al. 2008).

Ma	nipulation	Mechanism	Outcomes	References
Cx30	KO	genetic ablation	↓ CA1 fEPSPs after Schaffer collateral stimulation, ↓ LTP; ↓ astrocyte-astrocyte dye coupling	Abudara et al., 2015, Pannasch et al., 2014
Cx31.1	KO	genetic ablation	↓ novel object recognition, ↑ novel environment exploration	Dere et al., 2008
Cx36	KO	genetic ablation	absent interneuron voltage coupling, ↓ IPSP amplitude; ↓ novel object recognition, ↓ short-term memory by Y-maze, ↓ motor learning by Rotarod; ↑ seizures w/ PTZ induction; ↓ CA1 fEPSPs after Schaffer collateral stimulation	Apostolides, 2013, Deans et al., 2001 Frisch et al., 2005, Jacobson et al., 2010, Wang and Belousov, 2011
Cx43	Astrocyte KO	GFAP:Cre Cx43 <sup>fl/fl</sup>	↓ dye coupling, hypertrophic astrocytes, ↓ vesicular glutamate release	Chever et al., 2014b
	gap26/27	EC loop mimetic peptide; HC inhibitor	↓ Ethidium bromide uptake, ↓ excitatory post synaptic current amplitude, ↓ ATP release; ↑ fEPSP amplitude after LPS	Abudara et al., 2015, Chever et al., 2014a
	TAT-Cx43L2	tail mimetic peptide; HC inhibitor	↓ fear conditioning memory with intra-amygdala injection	Stehberg et al., 2012
	anti-Cx43 antibodies	HC blockade	↓ Ca <sup>2+</sup> slow wave synchronization, ↓ seizure number & frequency	Kekesi et al., 2015
Cx30 + Cx43	Cx30 + astrocyte Cx43 DKO	Cx30—KO + GFAP:Cre Cx43 <sup>fl/fl</sup>	↑ astrocyte intracellular Ca <sup>2+</sup> after LPS, ↓ dye uptake after LPS; ↓ Ca <sup>2+</sup> slow waves	Abudara et al., 2015, Torres et al., 2012
Panx1	KO	genetic ablation	absent LTD; ↑ CA1 fEPSPs after Schaffer collateral stimulation, ↑ LTP, ↓ novel object recognition ↓ spatial memory recall	Ardiles et al., 2014, Prochnow et al., 2012
	<sup>10</sup> panx1	EC loop mimetic peptide; Panx1 HC inhibitor	After chronic restraint stress: ↓ ethidium bromide uptake, ↓ glutamate release, ↓ ATP release	Orellana et al., 2015