



Published in final edited form as:

Gynecol Oncol. 2018 January ; 148(1): 235. doi:10.1016/j.ygyno.2017.11.030.

Corrigendum to “Indications for and complications of transfusion and the management of gynecologic malignancies” [Gynecol. Oncol. 146 (2017) 416–426]

Paulina Cybulska^a, Cheryl Goss^b, William P. Tew^{c,d}, Rekha Parameswaran^{e,f}, and Yukio Sonoda^{a,g,*}

^aGynecology Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY, USA

^bHematology and Coagulation Laboratory Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA

^cGynecologic Medical Oncology Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

^dDepartment of Medicine, Weill Cornell Medical College, New York, NY, USA

^eTransfusion Medicine Service, Department of Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

^fHematology Service, Department of Medicine, Division of Hematologic Oncology, Memorial Sloan Kettering Cancer Center, New York, NY, USA

^gDepartment of Obstetrics and Gynecology, Weill Cornell Medical College, New York, NY, USA

The authors regret that there was an error in Section 3.3, paragraph 2 of the above paper. The following sentence is incorrect:

Time to recurrence was 17 months in patients who had undergone a transfusion compared with 11 months for those who had not ($p = 0.03$).

The correct sentence is:

Time to recurrence was 11 months in patients who had undergone a transfusion compared with 17 months for those who had not ($p = 0.03$).

The online version has been updated.

The authors would like to apologise for any inconvenience caused.

*Corresponding author at: Gynecology Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY, USA. sonoday@mskcc.org (Y. Sonoda).