


Immune checkpoint inhibitors: new strategies to checkmate cancer

R. A. M. Wilson,* T. R. J. Evans,^{†‡}
A. R. Fraser ^{*§} and R. J. B. Nibbs*

**Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK, [†]Cancer Research UK Beatson Institute, Glasgow, UK, [‡]Institute of Cancer Sciences, University of Glasgow, Glasgow, UK, and [§]Advanced Therapeutics, Scottish National Blood Transfusion Service, Edinburgh, UK*

Accepted for publication 13 November 2017
Correspondence: R. J. B. Nibbs, University of Glasgow, Sir Graeme Davies Building, 120 University Place, Glasgow G12 8TA, UK.
E-mail: robert.nibbs@glasgow.ac.uk

Introduction

There are extensive interactions between tumour cells and the components of the immune system. The process of immune surveillance ensures that aberrant cells with tumorigenic potential can undergo immune destruction before they develop into cancers, and from the very earliest stages of tumour development the tumour microenvironment (TME) contains often-substantial populations of leucocytes, including various subsets of T cells [1,2]. Effector T cell responses against tumour antigens are induced early during tumour development. Local dendritic cells (DCs) can acquire and process tumour proteins from lysed tumour cells, including mutated versions of normal proteins, and present peptides from these proteins to naive T cells in secondary lymphoid organs. This has the potential of generating potent tumour-specific effector T cells that could home to the tumour and facilitate selective tumour cell killing. However, tumours evolve diverse mechanisms of immune evasion and immunosuppression to prevent or restrain anti-tumour T cell responses [3,4]. This includes changing their antigen profile (to make them unrecognizable to effector T cells [5,6]), blocking T cell recruitment [7,8] and exploiting immunosuppressive leucocytes, such

Summary

Immune checkpoint inhibitors (ICIs) targeting cytotoxic T lymphocyte-associated protein-4 (CTLA-4) or programmed cell death protein 1 (PD-1) receptors have demonstrated remarkable efficacy in subsets of patients with malignant disease. This emerging treatment modality holds great promise for future cancer treatment and has engaged pharmaceutical research interests in tumour immunology. While ICIs can induce rapid and durable responses in some patients, identifying predictive factors for effective clinical responses has proved challenging. This review summarizes the mechanisms of action of ICIs and outlines important preclinical work that contributed to their development. We explore clinical data that has led to disease-specific drug licensing, and highlight key clinical trials that have revealed ICI efficacy across a range of malignancies. We describe how ICIs have been used as part of combination therapies, and explore their future prospects in this area. We conclude by discussing the incorporation of these new immunotherapeutics into precision approaches to cancer therapy.

Keywords: antibodies, cancer, T cell, tumour immunology

as regulatory T cells (T_{regs}) and tumour-associated macrophages [9–11]. In addition, cancer cells and/or the TME can produce molecules that actively inhibit any tumour-specific T cells that manage to enter the tumour [12–14]. All these processes are potential targets for therapeutic interventions that aim to induce, reinstate or enhance anti-tumour T cell responses. There have been several exciting recent successes with cancer immunotherapies: principal among these are antibodies targeting the ‘immune checkpoint’ proteins cytotoxic T lymphocyte-associated protein-4 (CTLA-4) or programmed cell death protein 1 (PD-1).

Enhancing anti-tumour T cell responses with anti-CTLA-4 antibodies

Full activation of naive T cells results in clonal expansion and the development of effector functions. This requires positive signals from several membrane receptors. ‘Signal one’ comes from the T cell receptor (TCR) after it engages cognate antigen displayed on target cell major histocompatibility complex (MHC), but secondary positive signals (co-stimulation) are also essential, while further signals through cytokine receptors shape effector T cell phenotype [15–18]. Co-stimulatory signals are delivered principally

through CD28 receptors after they bind CD80 and CD86 (also known as B7-1 and B7-2, respectively), which are expressed abundantly on mature DCs. CTLA-4 also binds to CD80/86 with higher affinity than CD28, but it regulates co-stimulation negatively [19,20]. CTLA-4 is induced transiently on activated T cells, peaking 2–3 days after initial activation [19], and is expressed strongly by T_{reg} : naturally occurring T_{reg} (nT_{reg}) are the major cell type constitutively expressing this molecule [9]. The receptor CTLA-4 is reported to transduce negative intracellular signals [21,22] and enhance T cell motility [23,24], but perhaps more significant is its ability to outcompete CD28 molecules for CD80/86 binding [19,24,25]; displace CD28 to distal regions of the immunological synapse [26,27]; and strip CD80/86 from the surface of dendritic cells [28,29]. These immunosuppressive activities prevent or restrain T cell activation, and ensure that autoreactive T cells, or those that bind weakly to antigen, fail to become activated and instead enter a state of unresponsiveness (anergy).

The immunosuppressive properties of CTLA-4 are evident in *Ctla4*-deficient mice, which develop fatal lymphoproliferative disease and multi-organ failure due to unopposed T cell activation and loss of immunological tolerance [30,31]. These phenotypes are rescued by blocking or deleting CD80/86 or CD28. Moreover, germline heterozygous mutations in *CTLA4* in humans are linked to severe immune dysregulation [32,33], and a *CTLA4* variant is associated with early-onset Crohn's disease and autoimmunity [34]. Individuals with *CTLA4* mutations have defective T_{reg} function, hyperactivated effector T cells and reduced numbers of circulating B cells [32,33].

It is well established that anti-CTLA-4 monoclonal antibodies can block the immunosuppressive properties of CTLA-4 and enhance anti-tumour T cell responses in animal models [35,36]. Blockade induces a reduction in T_{regs} and an increase in effector T cells within the TME, and simultaneous blockade of CTLA-4 on these cells can synergize to enhance anti-tumour responses in animal models [37–43]. Anti-CTLA-4 antibodies can also bind Fc γ receptors (Fc γ R) to mediate antibody-dependent cell-mediated cytotoxicity against CTLA-4⁺ T_{regs} , leading to their selective depletion [40–43]. This is largely dependent upon the presence of atypical, Fc γ R-expressing macrophages in the tumour [40–43] and this mechanism of intratumoural T_{reg} depletion has been identified in humans treated with ipilimumab [44].

The functional effects of anti-CTLA-4 antibodies on anti-tumour responses in a patient depend upon the characteristics of the patient's tumour, the underlying TME and the nature of any previous or concurrent therapies. Predicting responses is challenging and, given the profound immunological consequences of *Ctla4* deletion or genetic variation in *CTLA4*, the potential side effects of administering anti-CTLA-4 antibodies may be significant. However, the preclinical research and clinical trial data from CTLA4

blockade has provided a proof-of-principle and laid the groundwork for further targeted disruption of other immune checkpoints, such as the programmed cell death-1/programmed cell death ligand-1 (PD-1/PL-L1) axis.

Dismantling tumour defence by inhibiting PD-1 activity

PD-1 is a surface receptor for the cognate ligands PD-L1 and PD-L2. It is expressed predominantly by activated T cells, but is also found on other leucocyte subsets, including activated B cells, DCs, monocytes and natural killer (NK) cells [45–47]. PD-L1 can be found on activated T and B cells, DCs, macrophages and many tissue cells [47–50], while PD-L2 appears limited to DCs and macrophages and some stromal cells [47]. On effector T cells, PD-1 ligation causes dephosphorylation of key signalling molecules that lie downstream of the TCR, thereby dampening TCR-mediated T cell activation [51,52]. It has also been identified recently that PD-1/PD-L1 interactions trigger dephosphorylation of CD28 preferentially over the TCR, and is the primary mechanism of T cell suppression [53]. PD-1 also enhances T cell motility to limit T cell/DC interactions [54]. PD-L1 has also been reported to bind to CD80 and might, like CTLA-4, compete with CD28 [55]. Deletion of *Pd-1* leads to autoimmune disease [56,57], but this develops later in life than the more severe, early-onset disease in *Ctla4*^{-/-} mice [30,31]. Expression of PD-L2 in lymphoid organs is thought to help maintain tolerance [51], while PD-L1, which is up-regulated in response to interferons, particularly IFN- γ , protects tissues from excessive immune cell activity [48,49]. This occurs during chronic viral infection when persistent viral antigen exposure causes T cell exhaustion, a form of anergy [58,59]. In this context, blockade of PD-L1 by administration of anti-PD-L1 antibodies causes a resurgence of effector T cell activity [60], predominantly through reactivation of the CD28 signalling pathway [61]. Thus, the physiological functions of the PD-1 pathway are to maintain T cell tolerance and suppress effector T cell responses in peripheral tissues.

Cancers can exploit the PD-1 pathway, up-regulating PD-1 ligands to suppress T cell-mediated cytotoxicity. Up-regulated PD-1 expression by tumour-infiltrating lymphocytes is associated with poor outcomes in many human cancers, and correlates with an exhausted T cell phenotype that impedes tumour immunity [62–67]. Early *in-vivo* experiments demonstrated that myeloma cells were targeted more effectively when *Pd-1* was absent, and that anti-PD-L1 antibodies could suppress myeloma in wild-type mice [68]. In other mouse models, transplantation of *Pd-1*^{-/-} T cells [69] or administration of anti-PD-L1 antibodies [68,69] induced regression of established tumours. PD-L1 expression by infiltrating myeloid cells has also been shown to impair anti-tumour T cell immunity [70,71] but, interestingly, this may be due in part to the induction and




Fig. 1. History of checkpoint inhibitors: key milestones. Timeline showing when checkpoint inhibitors were approved for the treatment of specific cancers in the United States, Europe and Japan. Tumour type is indicated by the colour of each immune checkpoint inhibitor (ICI) depicted in the figure, according to the key in the top left. Abbreviations: 1L = first-line; 2L = second-line; NSCLC = non-small-cell lung cancer; NSQ = non-squamous; PD-L1 = programmed death ligand 1; RCC = renal cell carcinoma; R/M = recurrent/metastatic; SCCHN = squamous cell carcinoma of the head and neck; SQ = squamous. ¹ US Food and Drug Administration. <http://www.fda.gov>, accessed 11 November 2016. ² European Medicines Agency. <http://www.ema.europa.eu>, accessed 11 November 2016. ³ONO Pharmaceutical Co. Ltd [press release], 4 July 2014. ⁴ONO Pharmaceutical Co. Ltd [press release], March 23, 2015. ⁵ ONO Pharmaceutical Co, Ltd. [press release], 17 December 17, 2015. ⁶Merck [press release], 27 June 2016, accessed 8 August 2016. ⁷Bristol-Myers Squibb Company [press release], 22 November 2016. ⁸Merck [press release], 10 December 2016.

regulation of peripherally induced T_{reg} subsets (iT_{reg}) by the PD-L1-expressing myeloid cells [72]. Disruption of this process may represent an additional mechanism underpinning the enhanced anti-tumour immunity that can be induced by PD-1 blockade.

Translating efficacy from bench to bedside

CTLA-4 and PD-1 provide critical but discrete mechanisms of physiological immunoregulation. A wealth of experimental data shows that antibodies blocking these checkpoints can enhance anti-tumour immune responses in animal models. Despite some concerns over potential toxicities, this drove the development of humanized monoclonal antibodies targeting these molecules. These novel immunotherapies are referred to as ‘immune checkpoint inhibitors’ (ICIs). These have now been licensed in a number of different tumour types (Fig. 1), and are discussed briefly here.

Anti-CTLA-4: licensing ipilimumab for metastatic melanoma

Ipilimumab (Bristol Myers Squibb, New York, NY, USA), an anti-CTLA-4 antibody, was licensed for malignant melanoma in 2011 [71]. This followed a landmark Phase III clinical trial involving approximately 600 patients with previously treated malignant melanoma [73] (Table 1). There were three treatment arms: ipilimumab plus gp100 vaccine,

ipilimumab alone or gp100 vaccine alone. The gp100 vaccine targets a protein expressed abundantly by melanomas [74,75]. The treatments containing ipilimumab conferred greatest median overall survival (OS): 10.0 months (ipilimumab/gp100) and 10.1 months (ipilimumab monotherapy) *versus* 6.4 months (gp100 vaccine alone). Severe immune-related adverse effects (IAEs) were reported in 10–15% of patients receiving ipilimumab, and were of longer duration in the ipilimumab/gp100 cohort [73]. Established protocols for managing immune-mediated toxicities were adopted from previous trials [76–78]. Subsequent analysis of data from 1861 metastatic melanoma patients in 12 individual clinical trials of ipilimumab showed that OS at 3 years was 20% for previously treated patients and 26% for treatment-naïve patients [79]. The survival curve plateaued at approximately 3 years, with evidence in some patients of long-term progression-free survival of up to 10 years.

PD-1 blockade moves into the clinic

The safety and efficacy of pembrolizumab (Merck), a monoclonal antibody targeting PD-1, was demonstrated in patients with ipilimumab-refractory malignant melanoma. In an expansion cohort of a Phase I trial, 173 patients were treated every 3 weeks with pembrolizumab [92]. Equivalent efficacy, as determined by overall response rates (ORR) [93], was observed at both dosing regimens (ORR = 26%). Pembrolizumab was well tolerated: only 12% of patients

Table 1. Key clinical trials of immune checkpoint inhibitor (ICIs) in patients with malignant melanoma and their associated survival data

Trial name and/or Phase [Ref]	Trial drugs (dose)	Patient no.	1-year OS rate (%)	18-month OS rate (%)	2-year OS rate (%)	Median OS (months)
Phase III [73]	Ipi (3 mg/kg) versus gp100 vaccine	676	45.6 versus 25.3	33.2 versus 16.3	23.5 versus 13.7	10.1 versus 6.4
Phase II [76]	Ipi (10 mg/kg) + budesonide versus Ipi (10 mg/kg)	115	55.9 versus 62.4	n.r.	40.5 versus 41.7	17.7 versus 19.3
Phase II [77]	Ipi (10 versus 3 versus 0.3 mg/kg)	217	48.6 versus 39.3 versus 39.6	34.5 versus 30.2 versus 23.0	29.8 versus 24.2 versus 18.4	11.4 versus 8.7 versus 8.6
Phase II [78]	Ipi (10mg/kg)	226	47.2	39.4	32.8	10.2
KEYNOTE-006	Pembro (10mg/kg every 2 weeks)	834	74.1 versus 68.4	n.r.	55.1 versus 55.3	n.a. versus n.a.
Phase III [80,81]	versus Pembro (10mg/kg every 3 weeks) versus Ipi (3mg/kg every 3 weeks)		versus 58.2		versus 43.0	versus 16.0
Phase I [82,83]	Pembro (1–10 mg/kg)	104	62	n.r.	43	16.8
Phase III [84]	Niv (3 mg/kg) versus dacarbazine (1000 mg/m ²)	418	72.9 versus 42.1	n.a.	n.a.	n.a. versus 10.8
Phase I [85,86]	Cohort 1: Niv (0.3 mg/kg) + Ipi (3 mg/kg)	14	56	n.a.	n.a.	14.8
	Cohort 2: Niv (1 mg/kg) + Ipi (3 mg/kg)	17	94	n.a.	n.a.	N.R.
	Cohort 2a: Niv (3 mg/kg) + Ipi (1 mg/kg)	16	89	n.a.	n.a.	N.R.
	Cohort 3: Niv (3 mg/kg) + Ipi (mg/kg)	6	100	n.a.	n.a.	N.R.
CheckMate-069	Niv (1 mg/kg) + Ipi (3 mg/kg) versus Ipi (3 mg/kg)	142	73.4 versus 64.8	n.r.	63.8 versus 53.6	N.R.
Phase II [87,88]	Niv (3 mg/kg)	316	n.a.	n.a.	59	37.6
CheckMate-067	Niv (1 mg/kg) + Ipi (3 mg/kg), then Niv (3 mg/kg) maintenance	314	n.a.	n.a.	64	n.a.
Phase III [89,90]	Ipi (3 mg/kg)	315	n.a.	n.a.	45	19.9
CheckMate-064	Niv (3 mg/kg), then Ipi (3 mg/kg)	68	76	n.a.	n.a.	n.a.
Phase II [91]	Ipi (3 mg/kg), then Niv (3 mg/kg)	70	54	n.a.	n.a.	16.9

Ipi = ipilimumab; pembro = pembrolizumab; niv = nivolumab; OS = overall survival; n.a. = not available; n.r. = not reported.

developed drug-related grade 3/4 toxicities. Given the promising clinical responses, good safety profiles and the absence of other effective therapies, pembrolizumab was licensed for ipilimumab-resistant advanced melanoma in 2014 [94].

In 2015, Robert and colleagues demonstrated significant improvements in progression-free survival, OS, response rates and treatment-related adverse events of pembrolizumab monotherapy compared with ipilimumab in advanced melanoma [80] (Table 1). A total of 834 patients were randomized to either pembrolizumab every 2 weeks, pembrolizumab every 3 weeks or four cycles of ipilimumab every 3 weeks. The 12- and 24-month OS was significantly greater in the pembrolizumab groups, and progression-free survival at 6 months was 47.3, 46.4 and 26.5%, respectively, with more durable responses seen in the pembrolizumab groups at 7.9 months [81]. Lower rates of significant immune-mediated toxicities were also seen in the pembrolizumab groups.

A Phase I clinical trial of another anti-PD-1 monoclonal antibody, nivolumab (Bristol Myers Squibb), was conducted in 296 patients with melanoma, non-small-cell lung cancer (NSCLC), renal cell carcinoma (RCC) or other selected treatment-refractory malignancies [82]. Melanoma patients were treated every 2 weeks with increasing doses of 0.1–10 mg/kg and showed overall objective response rates of 28%, with the highest response rate (41%) at a dose of 3 mg/kg. Responses were durable in many responders. Grade 3/4 treatment-related adverse events occurred in 14% of patients, with IAEs in 6%. Longer-term follow-up of melanoma patients showed 1- and 2-year survival rates of 62 and 43%, respectively, with an OS of 16.8 months [83]. Long-term safety evaluation was comparable to the original analysis, with 22% experiencing grade 3/4 treatment-related adverse events and 5% having grade 3/4 IAEs. Toxicities were not cumulative and occurred almost exclusively in the first 6 months of therapy.

In an open-label, randomized, Phase III study involving patients with ipilimumab-refractory melanoma, nivolumab was associated with a higher ORR than chemotherapy (32 *versus* 11%) [95]. Subsequently, nivolumab was compared with dacarbazine in patients with previously untreated metastatic melanoma without proto-oncogene B-Raf (BRAF) mutation [84]. Nivolumab gave superior 1-year survival (72 *versus* 42%), progression-free survival (5.1 *versus* 2.2 months), and ORRs (40 *versus* 13.9%). The survival benefit with nivolumab was observed across prespecified subgroups, including those defined by PD-L1 status. Grade 3/4 drug-related adverse events occurred in 11.7% of nivolumab-treated patients and 17.6% of those receiving dacarbazine.

Combination regimens of ICIs in patients with advanced melanoma

The distinct mechanisms of actions of anti-PD-1 and anti-CTLA-4 infer that their co-administration could have

enhanced efficacy (Fig. 2). The first Phase I study combining two ICIs, nivolumab and ipilimumab, was undertaken in patients with malignant melanoma and used an initial dose-escalation trial design to identify safe doses of these drugs when used in combination regimens [85]. One mg/kg nivolumab and 3 mg/kg ipilimumab were the maximum doses associated with acceptable levels of adverse events. This regimen yielded objective responses in 21 of 52 patients (40%). Durable responses, ranging from 60.1 to 72.1 weeks, were ongoing in 19 patients at the time of publication. Impressively, 16 patients had tumour reduction $\geq 80\%$ at 12 weeks, including five with a complete response, although treatment-related grade 3/4 adverse events were reported in 53% of patients. Updated OS data were 82 and 75% for 1- and 2-year survival, respectively (Table 1) [86]. Thus, this preliminary trial data indicated that simultaneous targeting of PD-1 and CTLA-4 could be tolerated and result in durable responses.

The Checkmate 069 trial compared nivolumab plus ipilimumab, with ipilimumab monotherapy in patients with untreated malignant melanoma (Table 1). Greater objective responses were seen with combination therapy (61 *versus* 11% with ipilimumab alone), with complete responses in 22% of patients on combination therapy but none with ipilimumab alone [87]. One- and 2-year OS was 73.4 and 63.8%, respectively, in the combination cohort, compared with 64.8% and 53.6% with ipilimumab alone [88]. Importantly, 2-year survival was not affected significantly by BRAF mutation or tumour PD-L1 expression status at diagnosis. A large subsequent Phase III trial (Checkmate 067) compared nivolumab alone, ipilimumab alone and nivolumab plus ipilimumab: median progression-free survival was 6.9, 2.9 and 11.5 months, respectively [89]. Notably, in patients with PD-L1-negative tumours, progression-free survival was greater with combination therapy than either monotherapy. Furthermore, with a minimum follow-up of 36 months, the median overall survival has not been reached in the combination regimen arm and was 37.6 months in the nivolumab arm of the study compared with 19.9 months with ipilimumab monotherapy [hazard ratio (HR) = 0.55, $P < 0.001$ for combination *versus* ipilimumab; HR = 0.65, $P < 0.001$ for nivolumab *versus* ipilimumab]. The overall survival at 3 years was 58, 52 and 34% for the combination regimen, nivolumab and ipilimumab monotherapy, respectively. However, treatment-related grade 3/4 toxicities occurred in 59, 21 and 28% of patients treated with the combination regimen, nivolumab and ipilimumab, respectively [90].

A Phase II trial (Checkmate 064) has also revealed intriguing results of switching between nivolumab and ipilimumab [91]. Melanoma patients underwent induction with either nivolumab (3 mg/kg every 2 weeks; six doses) and then ipilimumab (3 mg/kg every 3 weeks; four doses), or the reverse sequence. This was followed by maintenance therapy with nivolumab (3 mg/kg every 2 weeks) in both

cohorts. Greater efficacy was observed in the nivolumab followed by ipilimumab cohort, with significantly greater 1-year OS than the reverse sequence (Table 1). As expected from previous combination trials, the frequency of IAEs was relatively high (43–50%), although this was not affected by the order of drug administration.

Collectively, these trials have allowed ICIs, either alone or in combination, to herald in a new era of immunotherapy for the treatment of malignant melanoma. Impressive

durable responses have led to unprecedented improvements for some patients with metastatic disease. Administering ICIs in combination with other therapeutics will no doubt expand their use further in cancer patients.

New therapeutic approaches for non-small-cell lung cancer

Anti-CTLA-4 therapy has been trialled in patients with NSCLC, but promising outcomes akin to those seen in patients with melanoma have not been forthcoming. One initial trial used tremelimumab, an anti-CTLA-4 antibody, but it showed minimal efficacy and a significant side-effect burden [96] and, when used in combination with chemotherapy, anti-CTLA-4 antibodies have consistently demonstrated only modest improvements in responses in NSCLC patients [97,98]. However, antibodies targeting the PD-1/PD-L1 axis have shown greater efficacy and marked reductions in side effects, so have largely replaced anti-CTLA-4 antibodies as the ICI of choice in NSCLC.

In 2015, nivolumab received Food and Drug Administration (FDA) approval for use in patients with advanced squamous-cell NSCLC who progressed during or after platinum chemotherapy (Table 2) [94]. This followed a Phase III trial (CheckMate-017) of nivolumab compared with docetaxel chemotherapy in patients with progressive disease following first-line chemotherapy [99]. Median OS was 9.2 months in the nivolumab cohort, compared with 6.0 months in those receiving docetaxel. At 12 months, OS was 42% with nivolumab and 24% with docetaxel.




Fig. 2. Immune checkpoint inhibitor (ICI) blockade reverses tumour-mediated immune suppression. (a) Established tumours block immune attack through a variety of mechanisms, including inhibition of tumour-specific cytotoxic T lymphocyte (CTL) and CD4 T cell activation and function (1). This is driven by tumour over-expression of programmed cell death protein 1 (PD-L1), interacting with tumour-specific T cell PD-1 receptor (2) and T cell anergy induced by tumour-mediated T cell expression of cytotoxic T lymphocyte-associated protein 4 (CTLA-4) inhibitory receptor (4). In addition, tolerogenic dendritic cells (DC) drive regulatory T cell (T_{reg}) induction and expansion via CTLA-4 (3) and accumulation of T_{regs} then contributes to the immunosuppressive milieu of the tumour microenvironment (TME). (b) After ICI therapy, there is reactivation and proliferation of tumour-specific CTLs via blockade of the PD-1 axis (1), and return of functional cytotoxicity, resulting in perforin release and tumour cell killing (2). As tumour damage increases, the TME is disrupted allowing macrophages to deplete T_{regs} via fragment crystalline receptor (FcR) binding of anti-CTLA-4 antibody (3). Tumour antigen release is driven by immune lysis of tumour cells which are processed by conventional DC and presented to naive T cells in context of checkpoint inhibition of CTLA-4, enhancing CTL proliferation and function (4). Tumour damage and antigen release is also supplemented by concomitant use of conventional chemo/radiotherapy, which can reveal new tumour-associated antigens and contribute to anti-tumour immune responses (5).

Pretreatment tumour PD-L1 expression was not predictive of efficacy. A similar Phase III trial (CheckMate-057) demonstrated considerable improvements in survival for patients with advanced non-squamous NSCLC treated with nivolumab compared to those receiving docetaxel [100]. OS at 12 and 18 months was higher for patients on nivolumab compared with those on docetaxel. An additional Phase II trial (CheckMate-063) supports the survival benefit of nivolumab, reporting a 1-year OS of 40.8% [101]. Similarly, compared with docetaxel, pembrolizumab showed greater efficacy than docetaxel in advanced NSCLC in a Phase II/III trial (KEYNOTE-010) [102,103]. Patients with $\geq 1\%$ PD-L1 expression on the tumour received 2 mg/kg pembrolizumab, 10 mg/kg pembrolizumab or docetaxel, resulting in median OS of 10.4, 12.7 or 8.5 months, respectively. Greater efficacy was seen in patients with high tumour PD-L1 expression. These results echo findings from a similar Phase I study (KEYNOTE-001) investigating pembrolizumab in NSCLC [104], while a Phase III trial (KEYNOTE-024) demonstrated survival benefits, durable responses, increased response rates and reduced treatment-related adverse events with pembrolizumab compared with platinum-based chemotherapy in previously untreated patients, with $\geq 50\%$ tumoral PD-L1 expression [105]. This has led to pembrolizumab being licensed for treatment of naive metastatic NSCLC with $\geq 50\%$ PD-L1 expression [92]. In contrast, nivolumab did not improve progression-free or overall survival compared with chemotherapy in patients ($n = 423$) with previously untreated stage IV or recurrent NSCLC and with a PD-1 expression of 5% or more [106]. This suggests that robust predictive markers are required to select patients optimally for anti-PD-1 therapy, and that the sequence of administration with other standard therapies may also be relevant for optimal treatment strategies.

In the Phase II for patients with atezolizumab *versus* docetaxel for previously treated non-small-cell lung cancer (POPLAR) trial, atezolizumab (Genentech, South San Francisco, CA, USA), an antibody targeting the PD-L1 rather than PD-1, showed improved OS for NSCLC patients who had progressed following platinum chemotherapy [107]. This study included prospective evaluation of tumour PD-L1 expression, and used immune gene expression to define and quantify effector T cell activity within the TME. It was clear that efficacy was related closely to both PD-L1 expression and effector T cell abundance in the tumour. Future studies should consider incorporating this pretreatment analysis of the presence of active tumour-infiltrating effector T cells that may be a predictive marker of ICI efficacy. The clinical outcomes of POPLAR were reiterated in atezolizumab *versus* docetaxel for patients with previously treated non-small-cell lung cancer (OAK), a Phase III clinical trial demonstrating improved median OS with atezolizumab *versus* docetaxel (13.8 *versus* 9.6 months) in NSCLC patients progressing on platinum

chemotherapy, although improvement in OS was seen regardless of tumour PD-L1 expression [108,109]. As a result of these trials, in 2016 the FDA approved atezolizumab for the treatment of metastatic NSCLC that has progressed on platinum chemotherapy [94].

Significantly, PD-L1/PD-1 inhibitors not only improve OS but they are also associated with fewer treatment-related adverse events (\geq grade 3) than chemotherapy [99,100,102,105–108]. Moreover, they may have broad usage: although strongest responses were seen in patients with high tumour PD-L1 expression, they also showed similar, if not greater, efficacy than chemotherapy in patients with very low tumour PD-L1 expression [99–104,107,108]. It has not yet been possible to stratify patient responses accurately based on PD-L1 expression, although this typically involves the analysis of a single pretreatment tumour biopsy, sometimes taken long before starting checkpoint blockade therapy. In the Checkmate-063 trial, for example, there was a median of 1.3 years between biopsy and commencing nivolumab [101]. Thus, more detailed studies are required to explore the prognostic value of tumour PD-L1 expression, and ideally analysis of effector T cell abundance and phenotype should also be included.

ICIs offer new hope for treating NSCLC which, historically, has poor survival outcomes in patients with metastatic disease, and they have become standard second-line therapies in NSCLC patients. They are also being used as first-line treatments, particularly for patients with PD-L1-expressing tumours, although optimal ICI treatment strategies, with or without chemotherapy, still need to be defined more precisely in this context [111]. Trials using a combination of anti-CTLA4 and anti-PD-L1 antibodies have begun in NSCLC patients, with two Phase I studies demonstrating safety profiles comparable to those seen in melanoma patients (Table 2) [110,112]. The results of ongoing Phase II/III trials are eagerly anticipated [113–115]. Other studies have focused upon including ICIs into treatment strategies either in combination or sequentially after other existing therapies in NSCLC. For example, patients ($n = 713$) with locally advanced unresectable NSCLC were assigned randomly (2 : 1) to receive consolidation therapy with either the anti-PDL-1 antibody durvalumab or placebo every 2 weeks for up to 12 months after definitive chemoradiotherapy (chemotherapy with concurrent radiotherapy). Patients treated with durvalumab had a significantly improved progression-free survival (16.8 *versus* 5.6 months; HR = 0.52, $P < 0.001$) and objective response rates 28.4 *versus* 16%; $P < 0.001$) [116].

ICIs show efficacy in other cancers

ICIs have been, or are being, investigated in clinical trials across a broad range of other cancers in the hope that responses seen in patients with melanoma or NSCLC can be reproduced in patients with other tumour types.

Table 2. Key trials of immune checkpoint inhibitor (ICIs) targeting the programmed cell death-1 (PD-1) pathway in patients with non-small-cell lung cancer (NSCLC)

Trial name, Phase [Ref]	Disease	Trial drugs (dose)	Patient no.	1-year OS rate (%)	Median OS (months)	Objective responses rate (%)	Median PFS (months)	Correlation between tumour PD-L1 expression and response?	Treatment-related adverse events (grade \geq 3) (%)
CheckMate-063 Phase III [101]	Sq NSCLC	Niv (3 mg/kg)	117	40.8	8.2	14.5	1.9	Yes	17
CheckMate-017 Phase III [99]	Sq NSCLC	Niv (3 mg/kg) <i>versus</i> docetaxel	272	42 <i>versus</i> 24	9.2 <i>versus</i> 6.0	20 <i>versus</i> 9	3.5 <i>versus</i> 2.8	n.r.	7 <i>versus</i> 55
CheckMate-057 Phase III [100]	Non-Sq NSCLC	Niv (3 mg/kg) <i>versus</i> docetaxel	582	51 <i>versus</i> 39	12.2 <i>versus</i> 9.4	19 <i>versus</i> 12	2.3 <i>versus</i> 4.2	Yes	10 <i>versus</i> 54
KEYNOTE-001 Phase I [104]	NSCLC	Pembro (2–10 mg/kg)	495	n.r.	12.0	19.4	3.7	Yes	9.5
KEYNOTE-010 Phase II/III [102,103]	>1% PD-L1 + NSCLC	Pembro (2 mg/kg) <i>versus</i> Pembro (10 mg/kg) <i>versus</i> Docetaxel	1034	43.2 <i>versus</i> 52.3 <i>versus</i> 34.6	10.4 <i>versus</i> 12.7 <i>versus</i> 8.5	n.r.	3.9 <i>versus</i> 4.0 <i>versus</i> 4.9	Yes	13 <i>versus</i> 16 <i>versus</i> 35
KEYNOTE-024 Phase II [105]	>50% PD-L1 + NSCLC	Pembro (200 mg) <i>versus</i> docetaxel	305	n.r.	n.r.	44.8 <i>versus</i> 27.8	10.3 <i>versus</i> 6.0	High PD-L1 a prerequisite for inclusion in trial	26.6 <i>versus</i> 53.3
POPLAR Phase II [106]	NSCLC	Atezo (1200 mg) <i>versus</i> docetaxel	277	47.9 <i>versus</i> 37.8	12.6 <i>versus</i> 9.7	15 <i>versus</i> 15	2.7 <i>versus</i> 3.0	Yes	40 <i>versus</i> 53
OAK Phase III [107,108]	NSCLC	Atezo <i>versus</i> docetaxel	1225	55 <i>versus</i> 41	13.8 <i>versus</i> 9.6	14 <i>versus</i> 13	2.8 <i>versus</i> 4.0	Yes	37 <i>versus</i> 54
Phase 1b [109]	NSCLC	Durvalumab + tremelimumab	102	n.r.	n.r.	0–23	n.r.	n.r.	36
CheckMate-012 Phase I [110]	NSCLC	Niv (3 mg/kg) + Ipi <i>versus</i> Niv (3 mg/kg) + Ipi	77	Not calculated <i>versus</i> 69	n.r.	47 <i>versus</i> 38	8.1 <i>versus</i> 3.9	Yes	37 <i>versus</i> 33

Sq = squamous; non-sq = non-squamous; niv = nivolumab; pembro = pembrolizumab; atezo = atezolizumab; ipi = ipilimumab; OS = overall survival; PFS = progression-free survival; n.r. = not reported.

PD-L1 expression has long been known to be associated with tumour aggressiveness and poorer outcomes in patients with renal cell carcinoma (RCC) [62,117]. Nivolumab demonstrated efficacy in a Phase II trial of 168 patients with metastatic RCC [118]. They received 0.2, 2 or 10 mg/kg of nivolumab: median OS was 18.2, 25.5 or 24.7 months, respectively. In a subsequent Phase III study, patients that had previously received anti-angiogenic therapy had a median OS of 25.0 months on nivolumab compared with 19.6 months on everolimus [a mechanistic target of rapamycin (mTOR) inhibitor recommended after failed anti-angiogenic therapy] [119]. Nivolumab was subsequently approved for use in patients with advanced RCC after failing anti-angiogenic therapy [94]. Interestingly, high tumour PD-L1 expression was not associated with improved responses to therapy in this study: median OS in patients with PD-L1 expression $\geq 1\%$ was 21.6 and 18.8 months (nivolumab and everolimus, respectively), but 27.4 *versus* 21.2 months in patients with $\leq 1\%$ tumour PD-L1 expression. This emphasizes the need to consider carefully the suitability of using PD-L1 expression alone to predict responses.

Following two clinical trials [120,121], atezolizumab, and more recently nivolumab, have been granted FDA approval for bladder cancer patients with disease progression within 12 months of treatment with platinum-containing chemotherapy [94]. The Phase II trial using atezolizumab improved objective response rates compared with a historical control [120]. Approximately 15% of patients responded and in this case, PD-L1 expression on tumour-associated immune cells correlated with objective responses. Ongoing responses in 84% of responders demonstrate excellent treatment durability, and for a small fraction (5%) a complete response was observed. This trial attempted to define features of the TME associated with responses. In addition to PD-L1 expression, it determined CD8⁺ T cell infiltration, mutational load and molecular subtype. Several 'immune cell' genes were associated with complete and partial responses to atezolizumab, including the IFN- γ -inducible genes *CXCL9* and *CXCL10*, which encode chemokines that direct leucocyte and CD8⁺ T cell homing. Mutational load was also significantly higher in responders than non-responders (12.4 versus 6.4 mutations per Mb). High mutational load has also been correlated with responses to anti-CTLA-4 in malignant melanoma [122], and may predict better survival in melanoma patients treated with anti-PD1 therapy [123]. This is most probably because increased mutation will increase the generation of neoantigens recognizable by T cells. Thus, in addition to characterizing PD-L1 expression and the immune cell infiltrate, efforts to predict ICI responses should take into account the mutational landscape of the tumour.

Tumours with mismatch-repair (MMR) defects contain a large number of somatic mutations, so may be

particularly sensitive to ICIs. Indeed, in a recent Phase II trial pembrolizumab was administered every 2 weeks to patients with MMR-deficient colorectal cancers ($n = 11$), MMR-proficient colorectal cancers ($n = 21$) or MMR-deficient cancers that were not colorectal ($n = 9$) [124]. The objective response rate and progression-free survival rate at 20 weeks were 40 and 78% for MMR-deficient colorectal cancers; 71 and 67% for non-colorectal MMR-deficient cancers; and 0 and 11% for MMR-proficient colorectal cancers. This apparent sensitivity of MMR-deficient tumours supports the rationale of combining DNA repair inhibitors with ICIs in clinical trials, and a Phase I/II study combining the poly ADP ribose polymerase (PARP) inhibitor, olaparib, with durvalumab in patients with selected advanced malignancies is ongoing [125].

PD-1 blockade has shown positive outcomes in classical Hodgkin's lymphoma. Nivolumab and pembrolizumab are now FDA-approved for treatment of patients who have relapsed after other therapies. Two clinical trials have reported ORRs of 65 and 87%, with complete response rates in 16 and 17% of patients [126,127]. Nivolumab and pembrolizumab are also approved for use in patients with recurrent or metastatic squamous cell carcinoma of the head and neck (SCCHN) whose disease has progressed after administration of platinum-based therapeutics [94]. Pembrolizumab treatment gave a clinically significant ORR and durable responses were evident in some patients [128], while nivolumab improved overall survival compared to standard single-agent therapy (methotrexate, docetaxel or cetuximab) [129]. However, in a Phase III study of pembrolizumab *versus* standard of care in patients with recurrent or metastatic SCCHN, pembrolizumab yielded a 19% reduction in the risk of death compared with standard of care but this did not meet the prespecified efficacy boundary, although it is possible that subsequent immunotherapy use in the standard of care arm may have confounded the overall survival analysis [130]. Nonetheless, greater differences in overall survival, progression-free survival and objective response rates were seen in patients with PD-1-expressing tumours [130].

Based on a Phase II clinical trial involving 88 patients, avelumab (Merck–Pfizer, New York, NY, USA), a PD-L1 blocking antibody, received FDA approval for use in patients with metastatic Merkel cell carcinoma [94,131]. This is the first ever drug approved for this rare skin cancer, in which Merkel cell polyomavirus infection is a major aetiological factor [132]. Nearly a third of patients showed objective responses, with eight complete responses reported: five grade 3 treatment-related adverse events were noted [131]. Pembrolizumab has also been trialed on 26 patients with this disease, with an ORR of 56% being reported, including four complete responses [133].

In a large Phase III trial involving 1132 patients with extensive late-stage small-cell lung cancer (SCLC),

ipilimumab with chemotherapy *versus* placebo with chemotherapy demonstrated no significant difference in OS [134]. A previous Phase I/II trial had also given disappointing results in SCLC, with no survival advantage for patients receiving ipilimumab [135]. However, nivolumab, either alone or in combination with ipilimumab, showed anti-tumour activity with durable responses and manageable safety profiles in previously treated patients with SCLC [136]. Patients were administered nivolumab (3 mg/kg); nivolumab (1 mg/kg) plus ipilimumab (3 mg/kg); or nivolumab (3 mg/kg) plus ipilimumab (1 mg/kg). Objective responses were seen in 10, 23 and 19% patients, respectively. Pembrolizumab is currently under investigation in patients with PD-L1+ ($\geq 1\%$ expression) SCLC, who have progressed during or after platinum-based chemotherapy [137]. Initial results indicate good tolerability and, while the data are preliminary, the initial report shows that 25% of SCLC patients developed a partial response.

A Phase III trial compared ipilimumab against placebo, following radiation, in patients with metastatic castration-resistant prostate cancer that progressed on docetaxel chemotherapy [138]. They reported no significant improvement in OS: median OS was 11.2 months for ipilimumab and 10.0 months for placebo. However, subgroup analysis revealed improved OS outcomes for patients with better prognostic factors. This is perhaps unsurprising, as it is now established that immunotherapies often take longer to show measurable efficacy [139]. Final results from a completed Phase III trial comparing placebo with ipilimumab in minimally symptomatic, previously untreated patients with metastatic castration-resistant prostate cancer should reveal whether there is a role for ICIs in this malignancy [140]. Finally, promising responses have also been observed with anti-PD-1 antibody treatment in patients with refractory gastric cancer [141,142] or hepatocellular carcinoma [143]: results of randomized trials are eagerly awaited.

Thus, in addition to melanoma and NSCLC, ICI monotherapies, particularly those targeting the PD-1 pathway, are yielding positive outcomes in a range of metastatic cancers. Toxicities are evident and need to be managed, and the long-term health implications of ICI monotherapy have yet to be determined. This might be significant, given the improvements in patient survival seen with ICIs, although the issue of resistance to ICIs raises the challenge of how to treat cancers which fail multiple therapies.

Future prospects for ICIs in combination therapies

Clinical trial data have justifiably generated considerable excitement concerning the effectiveness of ICIs. In many cases responses are durable, perhaps even curative, possibly because they release systemic personalized memory T cell responses against multiple patient-specific tumour antigens. However, only subsets of patients respond to ICI monotherapy. Combining ICIs can improve outcomes, and

further ICI combination trials, with careful side-effect monitoring, are certainly merited. There is also a large number of ongoing clinical trials exploring ICIs in combination with other treatment modalities, such as radiotherapy, chemotherapy or molecular targeted therapies [144]. The administration of ICIs after radiotherapy attempts to augment the 'abscopal effect', where localized radiation-induced tumour cell death can deliver anti-tumour immune responses against distant tumour deposits (Fig. 1) [145–150]. ICIs even show activity against tumours viewed classically as poorly immunogenic, such as pancreatic adenocarcinoma and glioblastoma, with efforts aimed at administering ICIs while either modifying the TME, up-regulating neoantigen abundance [151] or, in glioblastoma, potentiating immune responses using adenovirus [152]. Modulating the chemokine composition of the TME [153], or blocking chemokine receptors [154], can also enhance ICI efficacy in animal models, and these approaches may translate to humans. Preliminary data in animal tumour models have shown that depletion of T_{regs} via modified anti-CD25 antibody plus anti-PD-1 antibody can induce complete regression of established tumours [155]. In short, combination therapies have the potential to further improve the activity and scope of ICIs, leading to the induction of durable effector T cell responses in more patients, and throughout a greater range of malignancies.

Integrating ICIs into precision oncology

The future of oncology will undoubtedly involve a precision approach to patient care, with treatment tailored to an individual's disease to maximize efficacy and ameliorate side effects as far as possible. Numerous factors determine a patient's response to ICIs. These include, but are not limited to, the distribution of the target immune checkpoint protein; the immune cell context within the tumour; the availability, immunogenicity and frequency of tumour antigens; the pharmacodynamics and pharmacokinetics of the ICI; and the amenability of the TME for effective drug delivery [156]. The analysis of these factors must be incorporated into the design of future clinical trials so that combinations of biomarkers that predict response to ICI immunotherapy can be identified and exploited. It will also be important to clarify further the reciprocal relationship between tumour cells and the TME in the context of ICI therapies, particularly if they are to be implemented across a range of different malignancies. Clinical trials have evolved from when tumours were classified according to histological observations, and comprehensive molecular, cellular and genetic phenotyping is now possible. Genomic and transcriptomic analysis of pretreatment melanoma has identified a transcriptional signature associated with innate resistance to anti-PD1 therapy, and demonstrated that mutations in the DNA repair gene *BRCA2* are enriched in patients that do respond [123]. The analysis of serial

tumour biopsies taken before, during and after ICI therapy will further enhance understanding of how anti-tumour responses are induced by ICIs, and how tumours adapt to evade or limit these responses. Interestingly, one study has identified mutations in melanoma cells in patients whose disease progressed after initially responding to PD-1 blockade [157], and at least one trial is under way that aims to understand the selection pressures and evolutionary changes that ICIs impose upon tumour cells [158].

Concluding remarks

The development and application of ICIs has been a major advance in the treatment of cancer, dramatically rekindling academic, clinical and commercial interest in anti-cancer immunotherapies. As monotherapies, they show considerable promise and remarkable responses have been seen in some patients. However, it is clear that there is improved efficacy when these ICIs are used as part of a concerted therapeutic approach for management of malignant disease. The key issue to be resolved is generation of sufficient evaluable data from combinatorial trials to define effective parameters for treatment. Results from current trials where ICIs are combined with chemotherapy, radiotherapy or with other immunotherapies are starting to demonstrate which malignancies are particularly susceptible to immune reactivation. These studies are also providing clear evidence that, although ICIs have produced dramatic improvements in some patients, they are not necessarily sufficient to mediate long-term remission or cure in most. Identification and validation of suitable prognostic markers in patient cohorts will also help to stratify treatment modalities and target therapy to those patients most likely to respond. The corollary to this is that suitable prognostic criteria could also minimize the IAEs associated with ICI treatment and immune reactivation [159]. The management of side effects and the evolution of resistance are ongoing concerns, but further advances are anticipated as these new medicines are incorporated into combination therapies and integrated into precision oncology. Progress is likely to be swift, given the current explosion in clinical trials of ICIs and the rapidly expanding number of preclinical studies involving these drugs will no doubt stimulate further clinical applications. The range of new ICIs becoming available, and the novel checkpoint targets being identified and characterized in preclinical studies, means that the resistance or relapse seen in some patients could be reversed by using alternative checkpoint inhibitors. There is also a significant new approach to tumour immunotherapy with the recent FDA approval for autologous gene-modified chimeric antigen receptor (CAR) T cell therapy. This cell therapy (Kymriah; Novartis, Basel, Switzerland) targets CD19 in B cell acute lymphocytic leukaemia and has shown significant efficacy [160]. There are several other CAR-T cell therapies in development and combination

therapy with ICIs is already planned, offering the potential for even greater benefits. ICIs are important new weapons in the oncologist's arsenal, and exciting times lie ahead in the field of cancer immunotherapy.

Acknowledgements

The authors would like to thank Alison Schroeer of Schroeer Scientific Illustration for assistance with the preparation of Fig. 2.

Disclosure

The authors confirm that they have no competing financial interests or activities.

References

- Gajewski TF, Schreiber H, Fu YX. Innate and adaptive immune cells in the tumor microenvironment. *Nat Immunol* 2013; **14**: 1014–22.
- Speiser DE, Ho PC, Verdeil G. Regulatory circuits of T cell function in cancer. *Nat Rev Immunol* 2016; **16**:599–611.
- Dunn GP, Old LJ, Schreiber RD. The immunobiology of cancer immunosurveillance and immunoediting. *Immunity* 2004; **21**: 137–48.
- Prendergast GC. Immune escape as a fundamental trait of cancer: focus on IDO. *Oncogene* 2008; **27**:3889–900.
- Igney FH, Krammer PH. Immune escape of tumors: apoptosis resistance and tumor counterattack. *J Leukoc Biol* 2002; **71**: 907–20.
- Swann JB, Smyth MJ. Immune surveillance of tumors. *J Clin Invest* 2007; **117**:1137–46.
- Bouzin C, Brouet A, De Vriese J, Dewever J, Feron O. Effects of vascular endothelial growth factor on the lymphocyte–endothelium interactions: identification of caveolin-1 and nitric oxide as control points of endothelial cell anergy. *J Immunol* 2007; **178**:1505–11.
- Buckanovich RJ, Facciabene A, Kim S *et al.* Endothelin B receptor mediates the endothelial barrier to T cell homing to tumors and disables immune therapy. *Nat Med* 2008; **14**: 28–36.
- Oleinika K, Nibbs RJ, Graham GJ, Fraser AR. Suppression, subversion and escape: the role of regulatory T cells in cancer progression. *Clin Exp Immunol* 2013; **171**:36–45.
- Facciabene A, Motz GT, Coukos GT. Regulatory cells: key players in tumor immune escape and angiogenesis. *Cancer Res* 2012; **72**:2162–71.
- Grivnikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell* 2010; **140**:883–99.
- Dong H, Strome SE, Salomao DR *et al.* Tumor-associated B7-H1 promotes T cell apoptosis: a potential mechanism of immune evasion. *Nat Med* 2002; **8**:793–800.
- Zou W, Chen L. Inhibitory B7-family molecules in the tumour microenvironment. *Nat Rev Immunol* 2008; **8**:467–77.
- Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012; **12**:252–64.
- Harding FA, McArthur JG, Gross JA, Raulet DH, Allison JP. CD28-mediated signalling co-stimulates murine T cells and

- prevents induction of anergy in T cell clones. *Nature* 1992; **356**:607–9.
- 16 McAdam AJ, Schweitzer AN, Sharpe AH. The role of B7 co-stimulation in activation and differentiation of CD4⁺ and CD8⁺ T cells. *Immunol Rev* 1998; **165**:231–47.
 - 17 Acuto O, Michel F. CD28-mediated co-stimulation: a quantitative support for TCR signalling. *Nat Rev Immunol* 2003; **3**: 939–51.
 - 18 Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat Rev Immunol* 2013; **13**: 227–42.
 - 19 Krummel MF, Allison JP. CD28 and CTLA-4 have opposing effects on the response of T cells to Stimulation. *J Exp Med* 1995; **182**:459–65.
 - 20 Walunas TL, Lenschow DJ, Bakker CY *et al.* CTLA-4 can function as a negative regulator of T cell activation. *Immunity* 1994; **1**:405–13.
 - 21 Alegre ML, Frauwirth KA, Thompson CB. T cell regulation by CD28 and CTLA-4. *Nat Rev Immunol* 2001; **1**:220–8.
 - 22 Lee KM, Chuang E, Griffin M *et al.* Molecular basis of T cell inactivation by CTLA-4. *Science* 1998; **282**:2263–6.
 - 23 Schneider H, Downey J, Smith A *et al.* Reversal of the TCR stop signal by CTLA-4. *Science* 2006; **313**:1972–5.
 - 24 Walker LS, Sansom DM. The emerging role of CTLA4 as a cell-extrinsic regulator of T cell responses. *Nat Rev Immunol* 2011; **11**:852–63.
 - 25 Greene JL, Leytze GM, Emswiler J *et al.* Covalent dimerization of CD28/CTLA-4 and oligomerization of CD80/CD86 regulate T cell costimulatory interactions. *J Biol Chem* 1996; **271**: 26762–71.
 - 26 Egen JG, Allison JP. Cytotoxic T lymphocyte antigen-4 accumulation in the immunological synapse is regulated by TCR signal strength. *Immunity* 2002; **16**:23–35.
 - 27 Yokosuka T, Kobayashi W, Takamatsu M *et al.* Spatiotemporal basis of CTLA-4 costimulatory molecule-mediated negative regulation of T cell activation. *Immunity* 2010; **33**:326–39.
 - 28 Qureshi OS, Zheng Y, Nakamura K *et al.* Trans-endocytosis of CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science* 2011; **332**:600–3.
 - 29 Walker LS, Sansom DM. Confusing signals: recent progress in CTLA-4 biology. *Trends Immunol* 2015; **36**:63–70.
 - 30 Tivol EA, Borriello F, Schweitzer AN, Lynch WP, Bluestone JA, Sharpe AH. Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity* 1995; **3**:541–7.
 - 31 Waterhouse P, Penninger JM, Timms E *et al.* Lymphoproliferative disorders with early lethality in mice deficient in CTLA-4. *Science* 1995; **270**:985–8.
 - 32 Kuehn HS, Ouyang W, Lo B *et al.* Immune dysregulation in human subjects with heterozygous germline mutations in CTLA4. *Science* 2014; **345**:1623–7.
 - 33 Schubert D, Bode C, Kenefeck R *et al.* Autosomal dominant immune dysregulation syndrome in humans with CTLA4 mutations. *Nat Med* 2014; **20**:1410–6.
 - 34 Zeissig S, Petersen B-S, Tomczak M *et al.* Early-onset Crohn's disease and autoimmunity associated with a variant in CTLA-4. *Gut* 2015; **64**:1889–97.
 - 35 Leach DR, Krummel MF, Allison JP. Enhancement of antitumor immunity by CTLA-4 blockade. *Science* 1996; **271**: 1734–6.
 - 36 Hurwitz AA, Foster BA, Kwon ED *et al.* Combination immunotherapy of primary prostate cancer in a transgenic mouse model using CTLA-4 blockade. *Cancer Res* 2000; **60**:2444–8.
 - 37 Selby MJ, Engelhardt JJ, Quigley M *et al.* Anti-CTLA-4 antibodies of IgG2a isotype enhance antitumor activity through reduction of intratumoral regulatory T cells. *Cancer Immunol Res* 2013; **1**:32–42.
 - 38 Quezada SA, Peggs KS, Curran MA, Allison JP. CTLA4 blockade and GM-CSF combination immunotherapy alters the intratumor balance of effector and regulatory T cells. *J Clin Invest* 2006; **116**:1935–45.
 - 39 Peggs KS, Quezada SA, Chambers CA, Korman AJ, Allison JP. Blockade of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of anti-CTLA-4 antibodies. *J Exp Med* 2009; **206**:1717–25.
 - 40 Simpson TR, Li F, Montalvo-Ortiz W *et al.* Fc-dependent depletion of tumor-infiltrating regulatory T cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. *J Exp Med* 2013; **210**:1695–710.
 - 41 Bulliard Y, Jolicoeur R, Windman M *et al.* Activating Fc γ receptors contribute to the antitumor activities of immunoregulatory receptor-targeting antibodies. *J Exp Med* 2013; **210**: 1685–93.
 - 42 Bulliard Y, Jolicoeur R, Zhang J, Dranoff G, Wilson NS, Brogdon JL. OX40 engagement depletes intratumoral Tregs via activating Fc γ Rs, leading to antitumor efficacy. *Immunol Cell Biol* 2014; **92**:475–80.
 - 43 Śledzińska A, Menger L, Bergerhoff K, Peggs KS, Quezada SA. Negative immune checkpoints on T lymphocytes and their relevance to cancer immunotherapy. *Mol Oncol* 2015; **9**:1936–65.
 - 44 Romano E, Kusio-Kobialka M, Foukas PG *et al.* Ipilimumab-dependent cell-mediated cytotoxicity of regulatory T cells *ex vivo* by nonclassical monocytes in melanoma patients. *Proc Natl Acad Sci USA* 2015; **112**:6140–5.
 - 45 Ishida Y, Agata Y, Shibahara K, Honjo T. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* 1992; **11**:3887–95.
 - 46 Agata Y, Kawasaki A, Nishimura H *et al.* Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. *Int Immunol* 1996; **8**:765–72.
 - 47 Keir ME, Butte MJ, Freeman GJ, Sharpe AH. PD-1 and its ligands in tolerance and immunity. *Annu Rev Immunol* 2008; **26**:677–704.
 - 48 Yamazaki T, Akiba H, Iwai H *et al.* Expression of programmed death 1 ligands by murine T cells and APC. *J Immunol* 2002; **169**:5538–45.
 - 49 Eppihimer MJ, Gunn J, Freeman GJ *et al.* Expression and regulation of the PD-L1 immunoinhibitory molecule on microvascular endothelial cells. *Microcirculation* 2002; **9**:133–45.
 - 50 McDermott DF, Atkins MB. PD-1 as a potential target in cancer therapy. *Cancer Med* 2013; **2**:662–73.
 - 51 Latchman Y, Wood CR, Chernova T *et al.* PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat Immunol* 2001; **2**:261–8.
 - 52 Freeman GJ, Long AJ, Iwai Y *et al.* Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* 2000; **192**:1027–34.

- 53 Hui E, Cheung J, Zhu J *et al.* T cell costimulatory receptor CD28 is a primary target for PD-1-mediated inhibition. *Science* 2017; **355**:1428–33.
- 54 Fife BT, Pauken KE, Eagar TN *et al.* Interactions between PD-1 and PD-L1 promote tolerance by blocking the TCR-induced stop signal. *Nat Immunol* 2009; **10**:1185–92.
- 55 Butte MJ, Keir ME, Phamduy TB, Sharpe AH, Freeman GJ. Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses. *Immunity* 2007; **27**:111–22.
- 56 Nishimura H, Nose M, Hiai H, Minato N, Honjo T. Development of lupus-like autoimmune diseases by disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999; **11**:141–51.
- 57 Nishimura H, Okazaki T, Tanaka Y *et al.* Autoimmune dilated cardiomyopathy in PD-1 receptor deficient mice. *Science* 2001; **291**:319–22.
- 58 Zajac AJ, Blattman JN, Murali-Krishna K *et al.* Viral immune evasion due to persistence of activated T cells without effector function. *J Exp Med* 1998; **188**:2205–13.
- 59 Gallimore A, Glithero A, Godkin A *et al.* Induction and exhaustion of lymphocytic choriomeningitis virus-specific cytotoxic T lymphocytes visualized using soluble tetrameric major histocompatibility complex class I-peptide complexes. *J Exp Med* 1998; **187**:1383–93.
- 60 Barber DL, Wherry EJ, Masopust D *et al.* Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature* 2006; **439**:682–7.
- 61 Kamphorst AO, Wieland A, Nasti T *et al.* Rescue of exhausted CD8 T cells by PD-1-targeted therapies is CD28-dependent. *Science* 2017; **355**:1423–7.
- 62 Thompson RH, Gillett MD, Cheville JC *et al.* Costimulatory B7-H1 in renal cell carcinoma patients: indicator of tumor aggressiveness and potential therapeutic target. *Proc Natl Acad Sci USA* 2004; **101**:17174–9.
- 63 Thompson RH, Dong H, Lohse CM *et al.* PD-1 is expressed by tumor-infiltrating immune cells and is associated with poor outcome for patients with renal cell carcinoma. *Clin Cancer Res* 2007; **13**:1757–61.
- 64 Ahmadzadeh M, Johnson LA, Heemskerk B *et al.* Tumor antigen-specific CD8 T cells infiltrating the tumor express high levels of PD-1 and are functionally impaired. *Blood* 2009; **114**:1537–44.
- 65 Zhang Y, Huang S, Gong D, Qin Y, Shen Q. Programmed death-1 upregulation is correlated with dysfunction of tumor-infiltrating CD8+ T lymphocytes in human non-small cell lung cancer. *Cell Mol Immunol* 2010; **7**:289–95.
- 66 Hino R, Kabashima K, Kato Y *et al.* Tumor cell expression of programmed cell death-1 ligand 1 is a prognostic factor for malignant melanoma. *Cancer* 2010; **116**:1757–66.
- 67 Shi F, Shi M, Zeng Z *et al.* PD-1 and PD-L1 upregulation promotes CD8(+) T cell apoptosis and postoperative recurrence in hepatocellular carcinoma patients. *Int J Cancer* 2011; **128**:887–96.
- 68 Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, Minato N. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci USA* 2002; **99**:12293–7.
- 69 Blank C, Brown I, Peterson AC *et al.* PD-L1/B7H-1 inhibits the effector phase of tumor rejection by T cell receptor (TCR) transgenic CD8+ T cells. *Cancer Res* 2004; **64**:1140–5.
- 70 Kuang DM, Zhao Q, Peng C *et al.* Activated monocytes in peritumoral stroma of hepatocellular carcinoma foster immune privilege and disease progression through PD-L1. *J Exp Med* 2009; **206**:1327–37.
- 71 Curiel TJ, Wei S, Dong H *et al.* Blockade of B7-H1 improves myeloid dendritic cell-mediated antitumor immunity. *Nat Med* 2003; **9**:562–7.
- 72 Francisco LM, Salinas VH, Brown KE *et al.* PD-L1 regulates the development, maintenance, and function of induced regulatory T cells. *J Exp Med* 2009; **206**:3015–29.
- 73 Hodi FS, O'Day SJ, McDermott DF *et al.* Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010; **363**:711–23.
- 74 Rosenberg SA, Yang JC, Schwartzentruber DJ *et al.* Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nat Med* 1998; **4**:321–7.
- 75 Smith FO, Downey SG, Klapper JA *et al.* Treatment of metastatic melanoma using interleukin-2 alone or in conjunction with vaccines. *Clin Cancer Res* 2008; **14**:5610–8.
- 76 Weber J, Thompson JA, Hamid O *et al.* A randomized, double-blind, placebo-controlled, phase II study comparing the tolerability and efficacy of ipilimumab administered with or without prophylactic budesonide in patients with unresectable stage III or IV melanoma. *Clin Cancer Res* 2009; **15**:5591–8.
- 77 Wolchok JD, Neyns B, Linette G *et al.* Ipilimumab monotherapy in patients with pretreated advanced melanoma: a randomized, double-blind, multicentre, phase 2, dose-ranging study. *Lancet Oncol* 2010; **11**:155–64.
- 78 O'Day SJ, Maio M, Chiarion-Sileni V *et al.* Efficacy and safety of ipilimumab monotherapy in patients with pretreated advanced melanoma: a multicenter single-arm phase II study. *Ann Oncol* 2010; **21**:1712–7.
- 79 Schadendorf D, Hodi FS, Robert C *et al.* Pooled analysis of long-term survival data from phase II and phase III trials of ipilimumab in unresectable or metastatic melanoma. *J Clin Oncol* 2015; **33**:1889–94.
- 80 Robert C, Schachter J, Long GV *et al.* Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med* 2015; **372**:2521–32.
- 81 Schachter J, Ribas A, Long GV *et al.* Pembrolizumab versus ipilimumab for advanced melanoma: final overall survival analysis of KEYNOTE-006. *J Clin Oncol* 2016; **34**:
- 82 Topalian SL, Hodi FS, Brahmer JR *et al.* Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 2012; **366**:2443–54.
- 83 Topalian SL, Sznol M, McDermott DF *et al.* Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab. *J Clin Oncol* 2014; **32**:1020–30.
- 84 Robert C, Long GV, Brady B *et al.* Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med* 2015; **372**:320–30.
- 85 Wolchok JD, Kluger H, Callahan MK *et al.* Nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med* 2013; **369**:122–33.
- 86 Sznol M, Kluger HM, Callahan MK *et al.* Survival, response duration, and activity by BRAF mutation (MT) status of nivolumab (NIVO, anti-PD-1, BMS-936558, ONO-4538) and

- ipilimumab (IPI) concurrent therapy in advanced melanoma (MEL). *J Clin Oncol* 2014; **32**:9003.
- 87 Postow MA, Chesney J, Pavlick AC *et al.* Nivolumab and ipilimumab versus ipilimumab in untreated melanoma. *N Engl J Med* 2015; **372**:2006–17.
- 88 Hodi FS, Chesney J, Pavlick AC *et al.* Combined nivolumab and ipilimumab versus ipilimumab alone in patients with advanced melanoma: 2-year overall survival outcomes in a multicentre, randomised, controlled, phase 2 trial. *Lancet Oncol* 2016; **17**:1558–68.
- 89 Larkin J, Chiarion-Sileni V, Gonzalez R *et al.* Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* 2015; **373**:23–43.
- 90 Wolchok JD, Chiarion-Sileni V, Gonzalez R *et al.* Overall survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2017; **377**:1345–56.
- 91 Weber JS, Gibney G, Sullivan RJ *et al.* Sequential administration of nivolumab and ipilimumab with a planned switch in patients with advanced melanoma (CheckMate 064): an open-label, randomised, phase 2 trial. *Lancet Oncol* 2016; **17**:943–55.
- 92 Robert C, Ribas A, Wolchok JD *et al.* Anti-programmed-death-receptor-1 treatment with pembrolizumab in ipilimumab-refractory advanced melanoma: a randomised dose-comparison cohort of a phase 1 trial. *Lancet* 2014; **384**:1109–17.
- 93 Eisenhauer EA, Therasse P, Bogaerts J *et al.* New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009; **45**:228–47.
- 94 FDA.gov. Hematology/oncology (cancer) approvals and safety notifications. Available at: <http://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm279174.htm> (accessed 13 October 2017).
- 95 Weber JS, D'Angelo SP, Minor D *et al.* Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2015; **16**:375–84. doi: 10.1016/S1470-2045(15)70076-8.
- 96 Zatlouk P, Heo DS, Park K *et al.* Randomized phase II clinical trial comparing tremelimumab (CP-675,206) with best supportive care (BSC) following first-line platinum-based therapy in patients (pts) with advanced non-small cell lung cancer (NSCLC). *J Clin Oncol* 2009; **27**:abstr8071.
- 97 Lynch TJ, Bondarenko I, Luft A *et al.* Ipilimumab in combination with paclitaxel and carboplatin as first-line treatment in stage IIIB/IV non-small-cell lung cancer: results from a randomized, double-blind, multicenter phase II study. *J Clin Oncol* 2012; **30**:2046–2054.
- 98 Govindan R, Szczesna A, Ahn M-J *et al.* Phase III trial of ipilimumab combined with paclitaxel and carboplatin in advanced squamous non-small-cell lung cancer. *J Clin Oncol* 2017; **35**:3449–57.
- 99 Brahmer J, Reckamp KL, Baas P *et al.* Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 2015; **373**:123–35.
- 100 Borghaei H, Paz-Ares L, Horn L *et al.* Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015; **373**:1627–39.
- 101 Rizvi NA, Mazières J, Planchard D *et al.* Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol* 2015; **16**:257–65.
- 102 Herbst RS, Baas P, Kim D-W *et al.* Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 2016; **387**:1540–50.
- 103 Herbst RS, Baas P, Kim D-W *et al.* Pembrolizumab (pembro) vs docetaxel (doce) for previously treated, PD-L1-expressing NSCLC: updated outcomes of KEYNOTE-010. *Ann Oncol* 2016; **387**: 1540–50.
- 104 Garon EB, Rizvi NA, Hui R *et al.* Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 2015; **372**:2018–28.
- 105 Reck M, Rodríguez-Abreu D, Robinson AG *et al.* Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* 2016; **375**:1823–33.
- 106 Carbone DP, Reck M, Paz-Ares L *et al.* First-line nivolumab in stage IV or recurrent non-small-cell lung cancer. *New Engl J Med* 2017; **376**:2415–2426.
- 107 Fehrenbacher L, Spira A, Ballinger M *et al.* Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet* 2016; **387**:1837–46.
- 108 Rittmeyer A, Barlesi F, Waterkamp D *et al.* Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet* 2017; **389**:255–65.
- 109 Barles F, Park K, Ciardiello F *et al.* Primary analysis from OAK, a randomized phase III study comparing atezolizumab with docetaxel in 2L/3L NSCLC. *Ann Oncol* 2016; **27**:vi552–87.
- 110 Hellmann MD, Rizvi NA, Goldman JW *et al.* Nivolumab plus ipilimumab as first-line treatment for advanced non-small-cell lung cancer (CheckMate 012): results of an open-label, phase 1, multicohort study. *Lancet Oncol* 2017; **18**:31–41.
- 111 Remon J, Pardo N, Martinez-Martí A *et al.* Immune-checkpoint inhibition in first-line treatment of advanced non-small cell lung cancer patients: current status and future approaches. *Lung Cancer* 2017; **106**:70–75.
- 112 Antonia S, Goldberg SB, Balmanoukian A *et al.* Safety and antitumour activity of durvalumab plus tremelimumab in non-small cell lung cancer: a multicentre, phase 1b study. *Lancet Oncol* 2016; **17**:299–308.
- 113 A global study to assess the effects of MEDI4736, given as monotherapy or in combination with tremelimumab determined by PD-L1 expression versus standard of care in patients with locally advanced or metastatic non small cell lung cancer (ARCTIC). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/show/NCT02352948> (accessed 13 October 2017).
- 114 Phase III open label first line therapy study of MEDI 4736 (Durvalumab) with or without tremelimumab versus soc in non small-cell lung cancer (NSCLC). (MYSTIC). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/show/NCT02453282> (accessed 13 October 2017).
- 115 Study of 1st line therapy study of durvalumab with tremelimumab versus SoC in non small-cell lung cancer (NSCLC) (NEPTUNE). (NEPTUNE). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/show/NCT02542293> (accessed 13 October 2017).

- 116 Antonia SJ, Villegas A, Daniel D *et al.* Durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. *New Engl J Med* 2017; **77**:1919–29.
- 117 Thompson RH, Kuntz SM, Leibovich BC *et al.* Tumor B7-H1 is associated with poor prognosis in renal cell carcinoma patients with long-term follow-up. *Cancer Res* 2006; **66**:3381–5.
- 118 Motzer RJ, Rini BI, McDermott DF *et al.* Nivolumab for metastatic renal cell carcinoma: results of a randomized phase II trial. *J Clin Oncol* 2015; **33**:1430–7.
- 119 Motzer RJ, Escudier B, McDermott DF *et al.* Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Engl J Med* 2015; **373**:1803–13.
- 120 Rosenberg JE, Hoffman-Censits J, Powles T *et al.* Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet* 2016; **387**:1909–20.
- 121 Sharma P, Callahan MK, Bono P *et al.* Nivolumab monotherapy in recurrent metastatic urothelial carcinoma (CheckMate 032): a multicentre, open-label, two-stage, multi-arm, phase 1/2 trial. *Lancet Oncol* 2016; **17**:1590–98.
- 122 Snyder A, Makarov V, Merghoub T *et al.* Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med* 2014; **371**:2189–99.
- 123 Hugo W, Zartsky JM, Sun L *et al.* Genomic and transcriptomic features of response to anti-PD-1 therapy in metastatic melanoma. *Cell* 2016; **165**:35–44.
- 124 Le DT, Uram JN, Wang H *et al.* PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015; **372**:2509–20.
- 125 Domchek S, Bang Y-J, Coukos K *et al.* MEDIOLA: a phase I/II, open-label trial of olaparib in combination with durvalumab (MEDI4736) in patients with advanced solid tumours. *Ann Oncol* 2016; **27**:1103TiP.
- 126 Ansell SM, Lesokhin AM, Borrello I *et al.* PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med* 2015; **374**:311–19.
- 127 Armand P, Shipp MA, Ribrag V *et al.* Programmed death-1 blockade with pembrolizumab in patients with classical Hodgkin lymphoma after brentuximab vedotin failure. *J Clin Oncol* 2016; **34**:373–39.
- 128 Chow LQ, Haddad R, Gupta S *et al.* Antitumor activity of pembrolizumab in biomarker-unselected patients with recurrent and/or metastatic head and neck squamous cell carcinoma: results from the phase Ib KEYNOTE-012 expansion cohort. *J Clin Oncol* 2016; **34**:3838–45.
- 129 Ferris RL, Blumenschein G, Fayette J *et al.* Nivolumab for recurrent squamous-cell carcinoma of the head and neck. *N Engl J Med* 2016; **375**:1856–67.
- 130 Cohen EE, Harrington KJ, Le Tourneau C *et al.* Pembrolizumab versus standard of care for recurrent or metastatic head and neck squamous cell carcinoma: phase 3 KEYNOTE-040 trial. *Ann Oncol* 2017; **28**: 628LBA45.
- 131 Kaufman HL, Russell J, Hamid O *et al.* Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. *Lancet Oncol* 2016; **17**:1374–85.
- 132 Liu W, MacDonald M, You J. Merkel cell polyomavirus infection and Merkel cell carcinoma. *Curr Opin Virol* 2016; **20**: 20–27.
- 133 Nghiem PT, Bhatia S, Lipson EJ *et al.* PD-1 blockade with pembrolizumab in advanced Merkel-cell carcinoma. *N Engl J Med* 2016; **374**:2542–52.
- 134 Reck M, Luft A, Szczesna A *et al.* Phase III randomized trial of ipilimumab plus etoposide and platinum versus placebo plus etoposide and platinum in extensive-stage small-cell lung cancer. *J Clin Oncol* 2016; **34**:3740–48.
- 135 Reck M, Bondarenko I, Luft A *et al.* Ipilimumab in combination with paclitaxel and carboplatin as first-line therapy in extensive-disease-small-cell lung cancer: results from a randomized, double-blind, multicenter phase 2 trial. *Ann Oncol* 2013; **24**:75–83.
- 136 Antonia SJ, López-Martin JA, Bendell J *et al.* Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol* 2016; **17**:88395.
- 137 Ott PA, Fernandez ME, Hirt S *et al.* Pembrolizumab (MK-3475) in patients (pts) with extensive-stage small cell lung cancer (SCLC): preliminary safety and efficacy results from KEYNOTE-028. *J Clin Oncol* 2015; **33**: abstr7502.
- 138 Kwon ED, Drake CG, Scher HI *et al.* Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol* 2014; **15**:700–12.
- 139 Wolchok JD, Hoos A, O'Day S *et al.* Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin Cancer Res* 2009; **15**:7412–20.
- 140 Phase 3 study of immunotherapy to treat advanced prostate cancer. *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/ct2/show/NCT01057810> (accessed 14 April 2017).
- 141 Muro K, Chung HC, Shankaran V *et al.* Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial. *Lancet Oncol* 2016; **17**:717–26.
- 142 Le DT, Bendell JC, Calvo E *et al.* Safety and activity of nivolumab monotherapy in advanced and metastatic (A/M) gastric or gastroesophageal junction cancer (GC/GEC): results from the CheckMate-032 study. *J Clin Oncol* 2016; **34**:6.
- 143 El-Khoueiry AB, Melero I, Crocenzi TS *et al.* Phase I/II safety and antitumor activity of nivolumab in patients with advanced hepatocellular carcinoma (HCC): CA209-040. *J Clin Oncol* 2015; **33**:LBA101.
- 144 *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/ct2/results?term=checkpoint+inhibitors&Search=Search> (accessed 13 October 2017).
- 145 Twyman-Saint Victor C, Rech AJ, Maity A *et al.* Radiation and dual checkpoint blockade activate non-redundant immune mechanisms in cancer. *Nature* 2015; **520**:373–7.
- 146 Postow MA, Callahan MK, Barker CA *et al.* Immunologic correlates of the abscopal effect in a patient with melanoma. *N Engl J Med* 2012; **366**:925–31.
- 147 Schoenfeld JD, Mahadevan A, Floyd SR *et al.* Ipilimumab and cranial radiation in metastatic melanoma patients: a case series and review. *J Immunother Cancer* 2015; **3**:50.
- 148 Simone CB, Burri SH, Heinzerling JH. Novel radiotherapy approaches for lung cancer: combining radiation therapy with targeted and immunotherapies. *Transl Lung Cancer Res* 2015; **4**:545–52.

- 149 Palma DA, Louie AV, Rodrigues GB. New strategies in stereotactic radiotherapy for oligometastases. *Clin Cancer Res* 2015; **21**:5198–204.
- 150 Okwan-Duodu D, Pollack BP, Lawson D, Khan MK. Role of radiation therapy as immune activator in the era of modern immunotherapy for metastatic malignant melanoma. *Am J Clin Oncol* 2015; **38**:119–25.
- 151 Immune checkpoint inhibition in combination with radiation therapy in pancreatic cancer patients (CheckPAC). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/ct2/show/NCT02866383?term=pancreas+AND+ipilimumab&rank=5> (accessed 13 October 2017).
- 152 Combination Adenovirus + Pembrolizumab to Trigger Immune Virus Effects (CAPTIVE). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/ct2/show/NCT02798406?term=checkpoint+inhibitors&rank=5> (accessed 20 August 2016).
- 153 Peng D, Kryczek I, Nagarsheth N *et al.* Epigenetic silencing of TH1-type chemokines shapes tumour immunity and immunotherapy. *Nature* 2015; **527**:249–53.
- 154 Steele CW, Karim SA, Leach JDG *et al.* CXCR2 inhibition profoundly suppresses metastases and augments immunotherapy in pancreatic ductal adenocarcinoma. *Cancer Cell* 2016; **29**:832–45.
- 155 Arce Vargas F, Furness AJS, Solomon I *et al.* Fc-optimized anti-CD25 depletes tumor-infiltrating regulatory T cells and synergizes with PD-1 blockade to eradicate established tumors. *Immunity* 2017; **46**:577–586.
- 156 Sharma P, Allison JP. The future of immune checkpoint therapy. *Science* 2015; **348**: 56–61.
- 157 Zaretsky JM, Garcia-Diaz A, Shin DS *et al.* Mutations Associated with Acquired Resistance to PD-1 Blockade in Melanoma. *N Engl J Med* 2016; **375**:819–29.
- 158 Selection Pressure and Evolution Induced by Immune Checkpoint Inhibitors and Other Immunologic Therapies (SPECIAL). *ClinicalTrials.gov*. Available at: <https://clinicaltrials.gov/ct2/show/NCT02724488?term=checkpoint+inhibitors&rank=1> (accessed 20 August 2016).
- 159 Ma W, Gilligan BM, Yuan J, Li T. Current status and perspectives in translational biomarker research for PD-1/PD-L1 immune checkpoint blockade therapy. *J Hematol Oncol* 2016; **9**:47.
- 160 Khalil D, Smith EL, Brentjens RJ, Wolchok JD. The future of cancer treatment: immunomodulation, CARs and combination immunotherapy. *Nature Rev Clin Oncol* 2016; **13**: 273–90.