



HHS Public Access

Author manuscript

Int J Cancer Manag. Author manuscript; available in PMC 2018 January 17.

Published in final edited form as:

Int J Cancer Manag. 2017 ; 10(9): . doi:10.5812/ijcm.9430.

Halting *Schistosoma haematobium* - associated bladder cancer

Monica C. Botelho^{1,2,*}, Helena Alves², and Joachim Richter³

¹INSA, National Institute of Health Dr. Ricardo Jorge, Porto, Portugal

²I3S, Instituto de Investigação e Inovação da Universidade do Porto, Portugal

³Institute of Tropical Medicine and International Health, Charité – Universitätsmedizin Berlin, Germany

Abstract

Background—At present schistosomiasis is endemic in 78 countries affecting more than 260 million people. Schistosomiasis haematobia alone affects more than 112 millions.

Material and Methods—We performed a computerized search of Pubmed database with keywords: bladder cancer cost and schistosomiasis mass treatment.

Results—Bladder cancer is an important sequelae of this infection. In low-resource countries, where this disease is endemic, individuals inflicted with bladder cancer have very limited access to treatment and death is most probably certain.

Conclusion—Mass treatment with praziquantel is an easy, safe and inexpensive treatment that could save the lives of thousands and reduce the morbidity of millions.

1. Background

Schistosomiasis is an ancient scourge of mankind, depicted in papyri from Pharaonic Egypt and known from human remains over 2000 years old from China. Human schistosomiasis is caused by six species: *Schistosoma (S.) haematobium*, *S. mansoni*, *S. japonicum*, *S. intercalatum*, *S. guineensis*, *S. mekongi*. A further species *S. malayensis* has been identified to infect humans in a limited focus in Malaysia (1).

Schistosomiasis was originally called bilharziasis in homage to Theodor Bilharz, a young German pathologist who first described the disease and its association with *Schistosoma* worms, working at Cairo in Egypt. Blood-dwelling *Trematoda* (phylum Platyhelminthes) of the genus *Schistosoma* cause this chronic and debilitating disease (2). Schistosomes present a heteroxenic life cycle, requiring as secondary host an invertebrate freshwater snail. Geographic distribution and maintenance of human infection depend on the presence of a suitable snail host. Adult worms are sexually distinct, a characteristic that separates them from other flukes that are hermaphrodites. Males present a ventral infolding from the ventral sucker to the posterior end forming the gynecophoric canal, where females are kept during

*Corresponding author. Mónica C. Botelho, PhD, INSA - National Institute of Health Dr. Ricardo Jorge, Rua Alexandre Herculano, 321, 4000-055, Porto, Portugal, Tel.: +351 223 401 114; fax: +351 223 401 109, monicabotelho@hotmail.com; monica.botelho@insa.min-saude.pt.

copulation (copulatory groove). *Schistosoma* life cycle differs from that of other flatworms also by the infecting process of the definitive host: the penetration of cercariae through the skin. Of the *Schistosoma* species pathogenic to humans, only *Schistosoma haematobium* causes urinary schistosomiasis (3). Snails of the genus *Bulinus* are the principle intermediate hosts for *S. haematobium*. Animals may play a role in *S. haematobium* infections as they may hybridize with closely related zoonotic *Schistosoma spp.* such as *S. bovis* (4).

As of 1989 schistosomiasis was endemic in 76 countries (5). Recent WHO report declared schistosomiasis endemic in 78 countries (6,7). The estimated total number of people requiring treatment for schistosomiasis for 2013 was 261 008 019, of whom 121 170 936 (46.4%) were school-age children (5–14 years of age) (7).

S. haematobium is endemic in 53 countries in the Middle East and most of the African continent, including the islands of Madagascar and Mauritius. Due to successful eradication programs the infection is no more of significant public health significance in Egypt, Lebanon, Oman, Syria, Tunisia and Turkey because transmission is low or nonexistent. A disputed and ill-defined focus exists in India and requires further confirmation (2). After more than 50 years in which no more autochthonous cases of schistosomiasis were recorded in Europe *S. haematobium* infection has recently emerged in Corsica (8).

In this review we discuss the basis for making beneficial therapeutic purposes in prevention, treatment and reducing the effects of schistosomiasis, such as bladder cancer mortality.

2. Material and Methods

In this article, the cost of bladder cancer treatment and the success of mass treatment for eradicating schistosomiasis are reviewed. For the literature review, we have used standard search strategies involving the querying of PubMed online database by using terms including “treatment cost bladder cancer” and “schistosomiasis mass treatment”. No specific key words were required as inclusion criteria. The reference lists of each article have been reviewed in details to find additional articles. Articles found were reviewed independently in full text.

3. Results

3.1. Schistosomiasis associated bladder cancer

Squamous cell carcinoma of the bladder (SCC) is a malignant, poorly differentiated neoplasm. SCC is the common form of bladder cancer in rural Africa where *S. haematobium* is prevalent (9, 10). By contrast, the majority of bladder cancer in developing countries and regions not endemic for urogenital schistosomiasis is transitional cell carcinoma (TCC), which arises from the transitional epithelium lining of the bladder. The parasite eggs trapped in the bladder wall release antigens and other metabolites (presumably evolved to expedite egress to the urine, and hence to the external environment). The phenomenon leads to hematuria and to chronic inflammation, in turn increasing the risk of SCC of the bladder. The epidemiological association between SCC of the bladder with schistosomiasis haematobia is based both on case control studies and on the correlation of bladder cancer incidence with prevalence of *S. haematobium* infection within diverse geographic areas. The

incidence of urogenital schistosomiasis-associated SCC is estimated in 3–4 cases per 100 000, yearly (11). Schistosomiasis haematobia is a chronic infection. The adult, egg-producing schistosomes live for many years, re-infections frequently occur, and schistosomiasis associated bladder SCC appears relatively early, often by the mid-decades of life (TCC usually presents in the later decades of life). In its recent monograph, IARC (International Agency for Research on Cancer) confirmed that chronic infection with *S. haematobium* causes cancer of the urinary bladder (12).

Our group has been working on the identification of parasite derived compounds that might be implicated in the carcinogenesis of *S. haematobium*. The majority of these compounds are catechol estrogens. The genotoxic effects of these estrogen metabolites might be attributed to oxidation of catechol estrogens to quinones, followed by redox cycling and formation of reactive oxygen species that in turn react with DNA (13). Given the context of the unarguable link between *S. haematobium* infection and bladder cancer, the presence of putative carcinogenic molecules in *S. haematobium* eggs hopefully may have practical consequences for new approaches to disease control (14,15). Metabolism of estrogens and the production of depurinating estrogen–DNA adducts can be implicated in a pathway underlying *S. haematobium*-promoted host cell DNA damage, leading eventually to cell transformation. The carcinogenic effect of this estrogen–DNA adduct mediated pathway could explain the link between chronic schistosomiasis haematobia and SCC of the bladder. We anticipate that these findings will contribute to understand how schistosomiasis haematobia leads to SCC of the bladder (13).

3.2. Treatment cost of bladder cancer

Although costs of medical care for bladder cancer have been investigated extensively, patient time costs associated with cancer care have rarely been estimated systematically. Yabroff and colleagues [16] identified 763,527 US patients with cancer from 1995 to 2001. Net time cost estimates for the initial phase of care were applied to national estimates of numbers of new cancers in 2005 to obtain national time costs for the initial phase of care. According to these authors in 2005, patient time costs for the initial phase of care were 2.3 billion dollars (16).

In another example, the diagnosis and treatment of bladder cancer represents a significant financial burden to the population in the United States. Therapeutic advances in bladder cancer care have come at a high cost to payers, providers, and patients. In this study Noyes et al (17) describe the principals of cost-effectiveness evaluation in healthcare and provide recommendations for a more economical use of resources in bladder cancer care. Although several studies have demonstrated that bladder cancer is a common disease associated with substantial economic burden for patients and society, the evidence supporting the cost-effectiveness of many interventions in bladder cancer care is limited and of insufficient quality. In addition, very little is known about quality of life, the preferred outcome measure for economic evaluations, associated with bladder cancer states and treatments. Moreover, current clinical guidelines for bladder cancer care do not incorporate economic factors when evaluating clinical pathways (17).

In Canada, bladder cancer is the most costly malignancy to treat per patient. Treatment options for bladder cancer are immediate cystectomy and conservative therapy with intravesical Bacillus Calmette-Guerin (BCG). The corresponding mean per-patient discounted lifetime costs (in 2005 Canadian dollars) were \$37,600 and \$42,400, respectively (18). In 2006 the annual cost of care for all patients with muscle-invasive BC (MIBC) was \$35.72M, 70% more than the \$21.03M for patients with non-MIBC. The major cost drivers, regardless of disease stage, were diagnostic/surveillance and complications, accounting for up to 43% and 37% of BC care costs, respectively. Co-morbidity-adjusted incremental annual resource costs per patient with MIBC were more than four times greater than those for patients with non-MIBC, similar to those of OC controls ($P = 0.490-0.913$), except for inpatient ($P = 0.002$) and hospice ($P < 0.001$) costs, which were both statistically significantly lower. Annual adjusted incremental Medicare reimbursements totalled \$36.3M for non-MIBC and \$96.1 million for MIBC (19)

In 2007 Konety and Allareddy (20) identified 6,577 US patients undergoing radical cystectomy for bladder cancer from the Nationwide Inpatient Sample of the Healthcare Cost and Utilization Project (1998 to 2002). Of commonly performed urological cancer procedures radical cystectomy is associated with the highest morbidity and mortality. According to these authors the impact of each individual type of complication or a combination of them on various outcome measures, such as mortality, charges and length of stay, is unclear. They attempted to quantify the impact of specific post-cystectomy complications and combinations thereof in terms of mortality, charges and length of stay. Although most patients undergoing cystectomy are older and have multiple comorbidities, the postoperative complications with the most significant impact were those directly related to surgery (primary complications). Secondary complications (cardiac, respiratory, vascular, etc) appear to have less of an impact on most common outcome measures. Median total charges was 41,905 dollars and median length of stay was 9 days (20).

Calculations of the treatment costs are limited by the fact that SCC may cause different costs than TCC and that costs depend on the current exchange rate of a country's currency with the USD. However, figures allow an approximate estimation of costs SCC would produce in industrialized countries if properly treated according to current standards.

3.3. Mass treatment of schistosomiasis and preventive chemotherapy

It has been shown previously that patients treated with praziquantel, pathological lesions of the urinary tract as shown ultrasonographically disappear (21). In these patients, the resolution of bladder lesions induced by urinary schistosomiasis resulted in reversibility of urinary tract obstructions one year after treatment with praziquantel (22,23,24). Other chronic sequelae like Symmers' fibrosis has been shown to be reversed by praziquantel (25).

We anticipate that the incidence of SCC in endemic areas will decrease to that of non-endemic regions if patients with urinary schistosomiasis are regularly treated from childhood on in spite of constant re-exposure, as it has been demonstrated (6,7).

The literature provides us with several reports in which mass treatment with praziquantel with the purpose of preventive chemotherapy of schistosomiasis is effective. An intervention

study was conducted in Khamir, north of Sana'a Yemen, for control of urinary schistosomiasis using chemotherapy. Prevalence of *S. haematobium* infection 14 months post-intervention fell from 58.9% to 5.8% and frequency of heavy infection from 40.0% to 18.9% (26). Schistosomiasis and soil-transmitted helminthiasis (STH) are among the neglected tropical diseases in Africa. A national control program for these diseases was initiated in Uganda during March 2003. Annual treatment with praziquantel and albendazole was given to schoolchildren in endemic areas and to adults in selected communities where local prevalence of *S. mansoni* in school children was high. Two rounds of treatment significantly reduced the prevalence of *S. mansoni* infection in school children across three regions in the country from 33.4–49.3% to 9.7–29.6%, and intensity of infection from 105.7–386.8 eggs per gram of faeces (epg) to 11.6–84.1 epg. The proportion of children with heavy *S. mansoni* infection was significantly reduced from 15% (95% CI 13.4–16.8%) to 2.3% (95% CI 1.6–3.0%). In this work the authors show that annual anthelmintic treatment delivered to school children and to adults at high risk in Uganda can significantly reduce the prevalence and intensity of infection for schistosomiasis and STH, and potentially also significantly reduce levels of environmental transmission of infection (27).

Also in the case of genital infection with *S. haematobium* it was shown that preventive chemotherapy could be useful. Genital infection with *S. haematobium* is prevalent in sub-Saharan Africa. Epidemiological studies have observed that genital schistosomiasis is associated with an increased odd of HIV infection among women. Ndeffo Mbah and collaborators (28) estimated that, in *S. haematobium* high-risk communities, targeted annual treatment of school-age children could reduce HIV prevalence by 20% (95% CI: 12–31%) in Angola, 16% (95% CI: 10–32%) in Kenya, and 6% (95% CI: 3–18%) in Zambia after the first 20 years of intervention (28).

The latest success story in the literature was designed to assess the impact of a decade of biennial mass administration of praziquantel on schistosomiasis in school-age children in Burkina Faso. In 2013, in a national assessment based on 22 sentinel sites, 3514 schoolchildren aged 7–11 years were checked for *S. haematobium* and *S. mansoni* infection by the examination of urine and stool samples, respectively. Ouedraogo et al (29) analysed the observed prevalence and intensity of infections and compared these with the relevant results of earlier surveys in Burkina Faso. Less than 1% of the children in six regions had heavy *S. haematobium* infections *S. mansoni* was only detected in two regions. By mass use of preventive chemotherapy, Burkina Faso may have eliminated schistosomiasis as a public health problem in eight regions and controlled schistosome-related morbidity in another three regions (29).

On the other hand, the number of people treated for schistosomiasis worldwide in 2013 was 39 485 376. This represents only 12.7% of the population requiring preventive chemotherapy for schistosomiasis globally. Schistosomiasis treatments in 2013 were reported by 9 fewer countries than in 2012. In 2013, 160 million praziquantel tablets were delivered to countries in the African Region for implementation of treatment. This would have been enough to treat about 60 million people. The fact that little more than 26 million people were treated in 2013 suggests that availability of praziquantel was not the only limiting factor for schistosomiasis control in the region (7). Considering that there are at

least 112 million people infected with schistosomiasis haematobia (13,30,31,32), and given that 3–4 in 100 000 individuals yearly develop bladder cancer (see above), the incidence of SCC attributable to schistosomiasis expected would approximately amount to 5000 patients yearly with bladder cancer caused by *S. haematobium* infection worldwide. The bladder cancer treatment of these patients would cost at least 20 million USD per year (see above).

In contrast, given that the individual cost per treatment including logistic costs amounts to 0.32 USD and that a biennial treatment is necessary to prevent serious morbidity due to schistosomiasis, the annual cost per schistosomiasis patient amounts to 0.16 USD (33). Mass therapy which prevents SCC in schistosomiasis patients would than cost 17.92 million USD per year and a chronic often incurable disease like SCC would be not only “treated” but safely prevented (34). If only school age children ((46.4%; 5–14 years of age) were targeted the cost would drop to 9.61 million USD. Affected populations would benefit from other effects of praziquantel on other complications and sequelae of schistosomiasis, including renal impairment, increased transmission of HIV, HBV, HTLV and other STDs sterility, complicated pregnancy, bleeding from esophageal varices, to mention just a few (35,36,37).

4. Conclusions

Schistosomiasis is a neglected tropical disease which in early stages is a disease easy to treat and thus to eliminate schistosomiasis induced SCC. The problem to solve is political rather than medical. More commitment is urgently needed to tackle this disease. Leading policy makers should be acquainted with these data. Are there simpler and more cost-effective ways of combating cancer than that to eliminate schistosomiasis induced SCC by preventive praziquantel mass therapy? Preventing schistosomiasis is the best treatment and least expensive way of fighting bladder cancer in endemic areas.

Acknowledgments

The authors would like to thank do the millions of people infected with schistosomiasis who inspired us to study this disease.

References

1. Adamson PB. Schistosomiasis in antiquity. *Med Hist.* 1976; 20:176–88. [PubMed: 781425]
2. Steinmann P, Keiser J, Bos R, Tanner M, Utzinger J. Schistosomiasis and water resources development: systematic review, meta-analysis and estimates of people at risk. *Lancet Infect Dis.* 2006; 6:411–425. DOI: 10.1016/S1473-3099(06)70521-7 [PubMed: 16790382]
3. Botelho MC, Machado JC, Brindley, Correia da Costa JM. Targeting molecular signaling pathways of *Schistosoma haematobium* infection in bladder cancer. *Virulence.* 2011; 2:267–279. [PubMed: 21788729]
4. Moné H, Holtfreter MC, Allienne JF, Mintsá-Nguéma R, Ibikounlé M, Boissier J, Berry A, Mitta G, Richter J, Mouahid G. Introgressive hybridizations of *Schistosoma haematobium* by *Schistosoma bovis* at the origin of the first case report of schistosomiasis in Corsica (France, Europe). *Parasitol Res.* 2015; 114:4127–4133. DOI: 10.1007/s00436-015-4643-4 [PubMed: 26268566]
5. Mott KE. Contrasts in the control of schistosomiasis. *Mem Inst Oswaldo Cruz.* 1989; 84:3–19. [PubMed: 2517866]
6. World Health Organization. Weekly epidemiological record, 90th year. 2015; 90:25–32. [30 January 2015] 5, 2015, <http://www.who.int/wer>.

7. WHO. Schistosomiasis: progress Report 2001–2011, strategic plan 2012–2020. Geneva: World Health Organization; 2013. <http://www.who.int> [accessed Jan 29, 2016]
8. Holtfreter MC, Moné H, Müller-Stöver I, Mouahid G, Richter J. Schistosoma haematobium infections acquired in Corsica, France, August 2013. Euro Surveill. 2014 Jun 5.19(22):20821. [PubMed: 24925456]
9. Mostafa MH, Sheweita SA, O'Connor PJ. Relationship between schistosomiasis and bladder cancer. Clin. Microbiol. Rev. 1999; 12:97–111. [PubMed: 9880476]
10. Zhong X, Isharwal S, Naples JM, Shiff C, Veltri RW, Shao C, Bosompem KM, Sidransky D, Hoque MO. Hypermethylation of genes detected in urine from Ghanaian adults with bladder pathology associated with Schistosoma haematobium infection. PLoS One. 2013; 8(3):e59089.doi: 10.1371/journal.pone.0059089 [PubMed: 23527093]
11. Shiff C, Veltri R, Naples J, Quartey J, Otchere J, Anyan W, Marlow C, Wiredu E, Adjei A, Brakohiapa E, Bosompem K. Ultrasound verification of bladder damage is associated with known biomarkers of bladder cancer in adults chronically infected with Schistosoma haematobium in Ghana. Trans R Soc Trop Med Hyg. 2006; 100(9):847–854. DOI: 10.1016/j.trstmh.2005.10.010 [PubMed: 16443246]
12. IARC Biological agents. IARC monographs on the evaluation of carcinogenic risks to humans/WHO. Vol. 100. IARC; 2012. Volume 100 B. A review of human carcinogens; p. 1-441.
13. Botelho MC, Alves H, Barros A, Rinaldi G, Brindley PJ, Sousa M. The role of estrogens and estrogen receptor signaling pathways in cancer and infertility: the case of schistosomes. Trends Parasitol. 2015; 31:246–250. DOI: 10.1016/j.pt.2015.03.005 [PubMed: 25837311]
14. Botelho MC, Vale N, Gouveia MJ, Rinaldi G, Santos J, Santos LL, Gomes P, Brindley PJ, Correia da Costa JM. Tumour-like phenotypes in urothelial cells after exposure to antigens from eggs of Schistosoma haematobium: an oestrogen-DNA adducts mediated pathway? Int J Parasitol. 2013; 43:17–26. DOI: 10.1016/j.ijpara.2012.10.023 [PubMed: 23260770]
15. Botelho MC, Soares R, Vale N, Ribeiro R, Camilo V, Almeida R, Medeiros R, Gomes P, Machado JC, Correia da Costa JM. Schistosoma haematobium: identification of new estrogenic molecules with estradiol antagonistic activity and ability to inactivate estrogen receptor in mammalian cells. Exp Parasitol. 2010; 126:526–535. DOI: 10.1016/j.exppara.2010.06.012 [PubMed: 20547157]
16. Yabroff KR, Davis WW, Lamont EB, Fahey A, Topor M, Brown ML, Warren JL. Patient time costs associated with cancer care. J Natl Cancer Inst. 2007; 99:14–23. DOI: 10.1093/jnci/djk001 [PubMed: 17202109]
17. Noyes K, Singer EA, Messing EM. Healthcare economics of bladder cancer: cost-enhancing and cost-reducing factors. Curr Opin Urol. 2008; 18:533–539. DOI: 10.1097/MOU.0b013e32830b8910 [PubMed: 18670280]
18. Kulkarni GS, Alibhai SM, Finelli A, Fleshner NE, Jewett MA, Lopushinsky SR, Bayoumi AM. Cost-effectiveness analysis of immediate radical cystectomy versus intravesical Bacillus Calmette-Guerin therapy for high-risk, high-grade (T1G3) bladder cancer. Cancer. 2009; 115:5450–5459. DOI: 10.1002/cncr.24634 [PubMed: 19685529]
19. Cooksley CD, Avritscher EB, Grossman HB, Sabichi AL, Dinney CP, Pettaway C, Elting LS. Clinical model of cost of bladder cancer in the elderly. Urology. 2008; 71:519–525. DOI: 10.1016/j.urology.2007.10.056 [PubMed: 18342201]
20. Konety BR, Allareddy V. Influence of post-cystectomy complications on cost and subsequent outcome. J Urol. 2007; 177:280–287. DOI: 10.1016/j.juro.2006.08.074 [PubMed: 17162064]
21. Doehring E, Ehrich JH. Effectiveness of praziquantel as an antihelmintic agent in the treatment of bilharziasis. Monatsschr Kinderheilkd. 1986; 134:282–284. [PubMed: 3088435]
22. Doehring E, Ehrich JH, Bremer HJ. Reversibility of urinary tract abnormalities due to Schistosoma haematobium infection. Kidney Int. 1986; 30:582–585. [PubMed: 3097374]
23. Richter J. Evolution of schistosomiasis-induced pathology after therapy and interruption of exposure to schistosomes: a review of ultrasonographic studies. Acta Trop. 2000; 77:111–131. [PubMed: 10996127]
24. Richter J. The impact of chemotherapy on morbidity due to schistosomiasis. Acta Trop. 2003; 86:161–183. [PubMed: 12745135]

25. Homeida MA, el Tom I, Nash T, Bennett JL. Association of the therapeutic activity of praziquantel with the reversal of Symmers' fibrosis induced by *Schistosoma mansoni*. *Am J Trop Med Hyg.* 1991; 45:360–65. [PubMed: 1928571]
26. Nagi MA. Evaluation of a programme for control of schistosoma haematobium infection in Yemen. *East Mediterr Health J.* 2005; 11:977–987. [PubMed: 16761668]
27. Koukounari A, Kabatereine N, Fleming F, Kazibwe F, Tukahebwa E, Stothard JR, Webster JP, Fenwick A. Parasitological impact of 2-year preventive chemotherapy on schistosomiasis and soil-transmitted helminthiasis in Uganda. *BMC Med.* 2007; 5:27.doi: 10.1186/1741-7015-5-27 [PubMed: 17767713]
28. Ndeffo Mbah ML, Gilbert JA, Galvani AP. Evaluating the potential impact of mass praziquantel administration for HIV prevention in *Schistosoma haematobium* high-risk communities. *Epidemics.* 2014; 7:22–7. DOI: 10.1016/j.epidem.2014.04.002 [PubMed: 24928666]
29. Ouedraogo H, Drabon F, Zongo D, Bagayan M, Bamba I, Pima T, Yago-Wienne F, Toubalie E, Zhang Y. Schistosomiasis in school-age children in Burkina Faso after a decade of preventive chemotherapy. Publication: Bulletin of the World Health Organization. 2015; BLT.15.161885. doi: 10.2471/BLT.15.161885
30. van der Werf MJ, et al. Quantification of clinical morbidity associated with schistosome infection in sub-Saharan Africa. *Acta Trop.* 2003; 86:125–139. [PubMed: 12745133]
31. Hotez PJ, Fenwick A, Kjetland EF. Africa's 32 cents solution for HIV/AIDS. *PLoS Negl Trop Dis.* 2009; 3:e430.doi: 10.1371/journal.pntd.0000430 [PubMed: 19479041]
32. King CH. Parasites and poverty: the case of schistosomiasis. *Acta Trop.* 2010; 113:95–104. DOI: 10.1016/j.actatropica.2009.11.012 [PubMed: 19962954]
33. Gabrielli AF1, Touré S, Sellin B, Sellin E, Ky C, Ouedraogo H, Yaogho M, Wilson MD, Thompson H, Sanou S, Fenwick A. A combined school- and community-based campaign targeting all school-age children of Burkina Faso against schistosomiasis and soil-transmitted helminthiasis: performance, financial costs and implications for sustainability. *Acta Trop.* 2006; 99:234–242. DOI: 10.1016/j.actatropica.2006.08.008 [PubMed: 16997268]
34. Fenwick A. Waterborne infectious diseases—could they be consigned to history? *Science.* 2006; 313:1077–81. DOI: 10.1126/science.1127184 [PubMed: 16931751]
35. Poggensee G, Sikelu M, Saría M, Richter J, Krantz I, Feldmeier H. Schistosomiasis of the lower reproductive tract without egg excretion in urine. *Am J Trop Med Hyg.* 1998; 59(9):782–783. [PubMed: 9840597]
36. Richter J, Bode JG, Blondin D, Kircheis G, Kubitz R, Holtfreter MC, Müller-Stöver I, Breuer M, Hüttig F, Antoch G, Häussinger D. Severe liver fibrosis caused by *Schistosoma mansoni*: management and treatment with a transjugular intrahepatic portosystemic shunt. *Lancet Infect Dis.* 2015; 15:731–737. DOI: 10.1016/S1473-3099(15)70009-5 [PubMed: 25769268]
37. Koukounari A, Donnelly CA, Sacko M, et al. The impact of single versus mixed schistosome species infections on liver, spleen and bladder morbidity within Malian children pre- and postpraziquantel treatment. *BMC Infect Dis.* 2010; 10:227.doi: 10.1186/1471-2334-10-227 [PubMed: 20670408]