



HHS Public Access

Author manuscript

Occup Environ Med. Author manuscript; available in PMC 2019 February 01.

Published in final edited form as:

Occup Environ Med. 2018 February ; 75(2): 79–89. doi:10.1136/oemed-2017-104431.

Occupational pesticide exposure and subclinical hypothyroidism among male pesticide applicators

Catherine C. Lerro¹, Laura E. Beane Freeman¹, Curt T. DellaValle^{1,2}, Muhammad G. Kibriya³, Briseis Aschebrook-Kilfoy³, Farzana Jasmine³, Stella Koutros¹, Christine G. Parks⁴, Dale P. Sandler⁴, Michael C.R. Alavanja^{1,5}, Jonathan N. Hofmann^{1,*}, and Mary H. Ward^{1,*}

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, USA

²Environmental Working Group, Washington, DC, USA

³Department of Public Health Sciences, The University of Chicago, Chicago, IL, USA

⁴National Institute of Environmental Health Sciences, Research Triangle Park, NC, USA

⁵Department of Biology, Hood College, Frederick, MD, USA

Abstract

Objectives—Animal studies suggest that exposure to pesticides may alter thyroid function; however, few epidemiologic studies have examined this association. We evaluated the relationship between individual pesticides and thyroid function in 679 men enrolled in a sub-study of the Agricultural Health Study, a cohort of licensed pesticide applicators.

Methods—Self-reported lifetime pesticide use was obtained at cohort enrollment (1993–1997). Intensity-weighted lifetime days were computed for 33 pesticides, which adjusts cumulative days of pesticide use for factors that modify exposure (e.g., use of personal protective equipment). Thyroid-stimulating hormone (TSH), thyroxine (T4), triiodothyronine (T3), and anti-thyroid peroxidase autoantibodies (anti-TPO) were measured in serum collected 2010–2013. We used multivariate logistic regression to estimate odds ratios (ORs) and 95% confidence intervals (CIs) for subclinical hypothyroidism (TSH>4.5mIU/L) compared to normal TSH (0.4– 4.5mIU/L) and for anti-TPO positivity. We also examined pesticide associations with TSH, T4, and T3 in multivariate linear regression models.

Results—Higher exposure to the insecticide aldrin (third and fourth quartile of intensity-weighted days vs. no exposure) was positively associated with subclinical hypothyroidism (OR_{Q3}=4.15, 95% CI=1.56–11.01, OR_{Q4}=4.76, 95% CI=1.53–14.82, $p_{\text{trend}}<0.01$), higher TSH ($p_{\text{trend}}=0.01$), and lower T4 ($p_{\text{trend}}=0.04$). Higher exposure to the herbicide pendimethalin was

Address for correspondence and reprints: Catherine Lerro, PhD, MPH, Occupational and Environmental Epidemiology Branch, National Cancer Institute, 9609 Medical Center Drive, 6E116, MSC 7991, Bethesda, MD 20892-7991, (t) 240-276-7813, (f) 240-276-7835, lerrocc@mail.nih.gov.

*These authors contributed equally to this work

CONTRIBUTORSHIP

All contributors meet the criteria for authorship.

COMPETING INTERESTS

None declared.

associated with subclinical hypothyroidism (fourth quartile vs. no exposure: $OR_{Q4}=2.78$, 95% $CI=1.30-5.95$, $p_{trend}=0.02$), higher TSH ($p_{trend}=0.04$), and anti-TPO positivity ($p_{trend}=0.01$). The fumigant methyl bromide was inversely associated with TSH ($p_{trend}=0.02$) and positively associated with T4 ($p_{trend}=0.01$).

Conclusions—Our results suggest that long-term exposure to aldrin, pendimethalin, and methyl bromide may alter thyroid function among male pesticide applicators.

Keywords

Thyroid disease; Hypothyroidism; Thyroid Stimulating Hormone; Agriculture; Pesticides

BACKGROUND

Thyroid dysfunction is relatively common in the United States (US) adult population. Approximately 5%, or 10.4 million adults, self-reported thyroid disease in a survey conducted in 1988–1994,¹ though prevalence estimates vary depending on population characteristics and diagnostic criteria.²³ Thyroid disease includes both hypothyroidism (i.e. thyroid gland fails to secrete adequate thyroid hormone) and hyperthyroidism (i.e. elevated blood levels of thyroid hormones);⁴ diagnosis of hypothyroidism is more common.²⁵ Established risk factors for thyroid disease include female sex, increasing age, prior benign thyroid nodules or goiter, autoimmunity, iodine deficiency, and ionizing radiation exposure.⁶ If untreated, thyroid disease may result in increased risk for atrial fibrillation, bone loss, and elevated cholesterol.³

Thyroid function is assessed by measurement of thyroid-stimulating hormone (TSH) and the thyroid hormones triiodothyronine (T3) and thyroxine (T4). TSH regulates thyroid hormone secretion and promotes thyroid gland growth. TSH is also regulated by T4 and T3 through negative feedback mechanisms. Anti-thyroid peroxidase (anti-TPO) autoantibodies are the most common autoantibodies present in the sera of individuals with autoimmune thyroid disease.⁷

Subclinical hypothyroidism is a subset of hypothyroid disease where laboratory evidence of thyroid dysfunction is present without symptoms characteristic of overt disease.⁸ It is characterized by TSH concentrations above the upper limit of the reference range (0.4–4.5 mIU/L) with normal free T4 levels. Occurring in approximately 3.9% of US adults who do not report being diagnosed with thyroid disease, subclinical hypothyroidism represents early, mild thyroid failure that can progress to more clinically significant disease.¹⁶

There is growing evidence that environmental exposures, including certain pesticides, have biological properties that may influence thyroid function. For example, dichlorodiphenyltrichloroethane (DDT) is structurally similar to thyroid hormones⁹ and can competitively bind to thyroid hormone receptors.¹⁰ Also, fungicides belonging to the ethylenebis(dithiocarbamate) (EBDC) class are metabolized to ethylene thiourea, a chemical known to inhibit thyroid peroxidase. Additional posited mechanisms of pesticide-induced thyroid disruption include inhibition of iodine uptake, binding to transport proteins, interference with iodothyronine deiodinases, increased clearance of thyroid hormones,

interference with cellular uptake of thyroid hormones, and interference with thyroid gene expression.¹¹¹²

Few epidemiologic studies have evaluated thyroid disease and long-term occupational exposure to pesticides. A large study conducted among male pesticide applicators in the Agricultural Health Study (AHS) found higher occupational use of the pesticides aldrin, chlordane, DDT, lindane, parathion, alachlor, and 2,4-dichlorophenoxyacetic acid (2,4-D) was associated with self-reported hypothyroidism.¹³ Among female spouses of AHS applicators, use of chlordane, paraquat, benomyl, and maneb/mancozeb was associated with self-reported hypothyroidism; maneb/mancozeb use was also associated with hyperthyroidism.¹⁴ Several epidemiologic studies have evaluated occupational pesticide exposure and subclinical thyroid function. EBDC fungicide-exposed pesticide applicators in Mexico demonstrated higher TSH compared to unexposed controls.¹⁵ Several epidemiologic studies have found changes in TSH, T3, and/or T4 to be associated with self-reported pesticide use.^{16–18} However, a common limitation of these studies was their inability to examine associations with individual pesticides, instead examining overall pesticide use or broad chemical groupings (e.g. herbicides, fungicides).

These previous studies of occupational pesticide exposures and thyroid function have been limited by low power, lack of information on specific pesticides, or reliance on self-reported thyroid disease. In the present study, we examine whether subclinical hypothyroidism, TSH, T3, and T4 levels, and thyroid autoantibodies are associated with occupational exposures to pesticides among male pesticide applicators without self-reported thyroid disease in Iowa and North Carolina.

METHODS

Study Design

We evaluated thyroid dysfunction in a subset of AHS participants who enrolled in the Biomarkers of Exposure and Effect in Agriculture (BEEA) study between June 2010 and September 2013; the design and enrollment methods of the BEEA study have been described.¹⁹ Briefly, BEEA participants were selected from among male pesticide applicators with a private license in the AHS cohort.²⁰ Those who continuously resided in Iowa or North Carolina, were over 50 years of age at the time of BEEA enrollment, had never been diagnosed with any type of cancer other than non-melanoma skin cancer, and had completed the enrollment AHS questionnaire (1993–1997), as well as two follow-up interviews (1999–2003 and 2005–2010) were eligible for BEEA. All participants provided written informed consent at BEEA enrollment. A trained phlebotomist conducted a computer-assisted personal interview and collected a non-fasting blood sample. Samples were shipped in temperature-controlled containers overnight to a central processing facility, where serum was isolated, aliquoted, and stored at -80°C . We sampled 684 of 955 eligible BEEA participants who had provided a blood sample, did not have a history of clinical thyroid disease (as reported on previous AHS questionnaires), and were not taking thyroid medications at the time of phlebotomy. Among the selected participants, we excluded five with laboratory results indicative of subclinical hyperthyroidism ($\text{TSH} < 0.4 \text{ mIU/L}$), leaving 679 participants for analysis.

TSH, thyroid hormones and anti-thyroid peroxidase autoantibodies

TSH was measured using a multiplexing kit from Millipore (The Human Pituitary panel, Cat # STTHMAG-21K). Total T4 and T3 were measured using another multiplexing kit from the same manufacturer (Steroid/Thyroid Hormone Magnetic Bead Panel, Cat # STTHMAG-21K). Coefficients of variation for blinded duplicate quality control (QC) samples (N=30) were 9.5%, 11.7%, and 15.8% for TSH, T4, and T3, respectively. Anti-TPO autoantibodies were measured using a Luminex-based kit from Applied Biosystems, Inc. (Cat # PP1000). We categorized results as negative or positive based on the raw Luminex median fluorescence intensity (MFI), with MFI 10 to 55 times higher than the median indicating anti-TPO positivity. In QC analyses of blind duplicate samples (N=30), we found 100% agreement.

For a subset of the QC samples with adequate serum (N=23) we compared the multiplex assay results with measurements from standard clinical laboratory immunoassays for TSH (Roche Diagnostics, Cat # 11731459122) and T4 (Roche Diagnostics, Cat # 20739006322). The assay results were highly correlated (TSH Spearman $\rho=0.76$, T4 Spearman $\rho=0.71$).

We defined normal thyroid function as having TSH levels of 0.4–4.5 mIU/L and subclinical hypothyroidism as TSH>4.5 mIU/L.²¹ We evaluated pesticide associations comparing those with subclinical hypothyroidism to those with normal TSH levels. We also evaluated TSH, T4, and T3 as continuous outcomes, and anti-TPO as a binary outcome (negative or positive).

Exposure Assessment

Information on use of individual pesticides came from two AHS cohort study questionnaires that were completed at enrollment (<http://www.aghealth.nih.gov/collaboration/questionnaires.html>). All applicators completed the first enrollment questionnaire, and provided information about ever/never use of 50 pesticides, as well as duration (years) and frequency (average days/year) of use for a subset of 22 pesticides. In addition, 63% of the applicators in this analysis returned the second (take-home) enrollment questionnaire, and provided information about duration and frequency of use for the remaining 28 pesticides. On average, there were 17 years of follow-up from enrollment to specimen collection.

For each pesticide, the number of cumulative days of use was calculated by multiplying the number of days the pesticide was used per year by total years of use. Intensity-weighted lifetime exposure days were computed by multiplying an intensity-weighting factor by lifetime exposure days. The intensity-weighting factor incorporates information that may influence exposure, i.e. the repair and cleaning of equipment, application method, whether the applicator mixed pesticides, and the use personal protective equipment.²² For herbicides and insecticides, we examined agents with ≥ 20 exposed cases of subclinical hypothyroidism. Because use of fumigants and fungicides was less prevalent, we examined agents with >5 exposed cases of subclinical hypothyroidism. Herbicides and insecticides were classified as unexposed (no reported use) and quartiles of intensity-weighted exposure days among all applicators., Fungicides and fumigants were classified as unexposed (no reported use), low (< median), or high exposure based on the intensity-weighted exposure days among all

applicators. Similarly, for analyses of anti-TPO positivity, we classified pesticide intensity-weighted days as unexposed, low, or high exposed based on the median among all applicators reporting use. Applicators with missing information for a particular pesticide were excluded from that analysis. We evaluated a total of 33 pesticides (16 herbicides, 13 insecticides, two fungicides, two fumigants).

Statistical Analysis

We compared sociodemographic, farming, and other health and behavioral characteristics for persons with normal TSH and those with subclinical hypothyroidism. We utilized logistic regression models to estimate odds ratios (OR) and 95% confidence intervals (95% CI) for the association of each pesticide with subclinical hypothyroidism and anti-TPO positivity. To evaluate the relationship between each pesticide active ingredient and TSH, T4, and T3 levels modeled continuously, we estimated β coefficients and 95% CIs using multivariate linear regression models. TSH, T4, and T3 levels were natural log (ln) transformed to satisfy the assumptions of our statistical models. We exponentiated the resulting β coefficients, which reflect the ratio of the geometric mean for individuals in each exposure category relative to the geometric mean for unexposed individuals. Tests for trend used the median of each exposure category treated as a continuous variable. All models were adjusted for age at sample collection (continuous), state of residence (Iowa or North Carolina), body mass index at sample collection (BMI, continuous), and smoking status (current, former, never). Models were also adjusted for use of correlated pesticides (no exposure, quartiles of exposure, missing) at AHS study enrollment (Spearman $\rho > 0.40$ calculated using continuous intensity-weighted lifetime days; Table S1). Other variables considered as potential confounders included month and time of day of blood draw (modeled as cubic terms), alcohol consumption, educational attainment, and race. These variables were ultimately not included in the final models because they did not impact the observed associations ($< 20\%$ change in main effect). We also examined pesticide use reported during follow-up interviews conducted after AHS enrollment up to time of BEEA sample collection. For the pesticides queried at enrollment, pesticide use had either stopped completely (i.e. ingredient registration was cancelled) or was markedly lower, the one exception being the commonly used herbicide glyphosate. However, including additional recent exposure to these pesticides (i.e. exposure at follow-up interviews) did not substantively alter our findings ($< 20\%$ change in main effect). To evaluate pesticide associations with subclinical hypothyroidism, TSH, and thyroid hormones among those without evidence of an autoimmune etiology, we performed analyses excluding anti-TPO positive participants.

Analyses were conducted using SAS version 9.3 (SAS Institute, Inc., Cary, NC). All tests were two-sided with $\alpha = 0.05$.

RESULTS

Among the 679 pesticide applicators in our analysis, 127 (19%) met our definition for subclinical hypothyroidism (Table 1). The prevalence increased with age: 17%, 16%, 21%, and 32% of men ages 50–59, 60–69, 70–79, and 80+ had subclinical hypothyroidism,

respectively. Men with subclinical hypothyroidism were more likely to be positive for anti-TPO ($p < 0.01$). Age, race, state, BMI, smoking status, alcohol consumption, and education did not vary substantially by subclinical hypothyroidism status. Men with subclinical hypothyroidism were less likely to have raised animals or to have applied pesticides in the last 12 months, although these differences were not statistically significant.

Among the insecticides (Table 2), aldrin use was positively associated with subclinical hypothyroidism (fourth quartile of intensity-weighted lifetime days of use vs. no use: $OR_{Q4} = 4.76$, 95% CI=1.53–14.82, $p_{trend} < 0.01$) and TSH ($p_{trend} = 0.01$). Increasing use of aldrin was also associated with lower T4 ($p_{trend} = 0.04$) and T3 ($p_{trend} = 0.08$). Diazinon use was associated with elevated odds of subclinical hypothyroidism (third quartile vs. no use: $OR_{Q3} = 2.64$, 95% CI=1.00–6.98, $p_{trend} = 0.10$) and higher TSH levels ($p_{trend} = 0.07$), with no significant monotonic trend. Terbufos use in all quartiles was associated with elevated T3, though we observed no significant exposure-response trend ($p_{trend} = 0.10$), and no association with TSH or T4.

Applicators reporting use of pendimethalin had increased odds of subclinical hypothyroidism (fourth quartile of intensity-weighted lifetime days of use vs. no use: $OR_{Q4} = 2.78$, 95% CI=1.30–5.95, $p_{trend} = 0.02$) and significantly increased TSH with increasing use ($p_{trend} = 0.04$) (Table 3). Higher use of S-Ethyl dipropylthiocarbamate (EPTC) was statistically significantly associated with increased TSH (fourth quartile vs. no use: $\exp(\beta)_{Q4} = 1.24$, 95% CI=1.00–1.54, $p_{trend} = 0.04$), and was non-significantly associated with subclinical hypothyroidism (fourth quartile vs. no use: $OR_{Q4} = 2.05$, 95% CI=0.91–4.63, $p_{trend} = 0.05$). Increasing 2,4,5-trichlorophenoxyacetic acid (2,4,5-T) intensity-weighted lifetime days of use was associated with significantly decreased T4 ($p_{trend} = 0.03$). Cyanazine was associated with elevated odds of subclinical hypothyroidism (third quartile vs. no use: $OR_{Q3} = 2.53$, 95% CI=1.34–4.78, $p_{trend} = 0.35$) and increased TSH (third quartile vs. no use: $\exp(\beta)_{Q3} = 1.19$, 95% CI=1.02–1.39, $p_{trend} = 0.11$) without evidence for a monotonic trend. Increasing use of petroleum distillates was associated with significantly decreased T3 ($p_{trend} = 0.045$), and non-significantly elevated TSH ($p_{trend} = 0.09$).

No evaluated fumigant or fungicide was significantly associated with subclinical hypothyroidism (Table 4). However, methyl bromide use was associated with significantly lower TSH (>median vs. unexposed: $\exp(\beta)_{high} = 0.75$, 95% CI=0.56–1.00, $p_{trend} = 0.02$), higher T4 ($p_{trend} = 0.01$), and higher T3 ($p_{trend} = 0.06$). In contrast, captan intensity-weighted lifetime days of use above the median was associated with borderline significantly higher TSH levels ($\exp(\beta)_{high} = 1.23$, 95% CI=1.00–1.52, $p_{trend} = 0.05$) compared to unexposed.

Results for pesticide use and anti-TPO positivity are shown in Table S2. Compared to men with no exposure, intensity-weighted lifetime days of pendimethalin use above the median was significantly associated with anti-TPO positivity ($OR_{high} = 2.69$, 95% CI=1.29–5.60, $p_{trend} = 0.01$) compared to unexposed applicators. A few additional pesticides had non-significantly elevated associations at exposure greater than the median intensity-weighted days, including aldrin ($OR_{high} = 2.31$, 95% CI=0.79–6.73, $p_{trend} = 0.12$), DDT ($OR_{high} = 2.28$, 95% CI=0.73–7.15, $p_{trend} = 0.09$), and atrazine ($OR_{high} = 2.15$, 95% CI 0.75–4.88, $p_{trend} = 0.29$).

Excluding individuals with anti-TPO positivity (N=65), we generally observed weaker associations for subclinical hypothyroidism, TSH, T4, and T3 with pesticide exposure that were in the same direction as those among all subjects (Table S3). The associations between aldrin and cyanazine with subclinical hypothyroidism, methyl bromide with TSH, and terbufos with T3 remained statistically significant; however, other reported associations were substantially attenuated including pendimethalin with hypothyroidism (fourth quartile of intensity-weighted days vs. no exposure: $OR_{Q4}=1.86$, 95% CI=0.73–4.73, $p_{trend}=0.25$) and EPTC with TSH (fourth quartile vs. no exposure: $exp(\beta)_{Q4}=1.07$, 95% CI=0.86–1.32, $p_{trend}=0.45$).

DISCUSSION

Our study is the first to comprehensively examine occupational use of a wide range of pesticide active ingredients and the prevalence of subclinical hypothyroidism, TSH, thyroid hormones, and thyroid autoantibodies. The insecticide aldrin and the herbicide pendimethalin were associated with subclinical hypothyroidism and higher TSH levels. Pendimethalin was also significantly associated with anti-TPO positivity, suggesting that it may influence TSH through an autoimmune pathway. The prevalence of subclinical hypothyroidism appeared to be higher among BEEA participants compared to similarly-aged thyroid disease-free adults in the general US population in NHANES.¹ However, a direct comparison of our results with these published data from NHANES should be interpreted cautiously, as the NHANES prevalence estimates are based on measurements from 1988–1994 in a racially diverse population including men and women, utilizing a different laboratory assay.

Insecticides

In our study, high use of the insecticide aldrin was positively associated with subclinical hypothyroidism and with levels of TSH, and inversely associated with T4. Aldrin is an organochlorine (OC) insecticide that was first produced in the US in the 1950s and was primarily used on corn.²³ Most uses for aldrin were banned in the US in 1974, and its use has similarly been banned or restricted in many other countries.²⁴ However, OCs have long half-lives and persist in the environment. Due to their lipophilic nature they biomagnify in the food chain and are detectable in human tissue and serum many years after exposure.²⁵ Our findings are consistent with a previous investigation among male applicators in the AHS in which higher prior use of aldrin was associated with self-reported hypothyroidism.¹³ In contrast to our findings, a study of agricultural workers in Brazil reported no association between serum levels of aldrin or dieldrin, the major metabolite of aldrin, with TSH, T3, or T4 in men or women.²⁶ Another Brazilian study of agricultural workers found no association between serum levels of aldrin and dieldrin and TSH; however, dieldrin levels were associated with significantly lower free T4.¹⁸ Evidence of a negative correlation between dieldrin and free T4 was also reported in a small study of women in India.²⁷ Rats exposed at low levels to a mixture of persistent pesticides and other contaminants, including aldrin and several other OCs, demonstrated alterations in thyroid gland morphology, serum thyroid hormone levels, and hepatic thyroid hormone levels.²⁸ Additionally, studies suggest

that aldrin is associated with follicular thyroid tumors in male and female rats, though these associations were only observed at low doses.²⁹

Herbicides

We observed associations with subclinical hypothyroidism and TSH for pendimethalin, a selective dinitroaniline herbicide used for control of broadleaf and grassy weeds that is registered for both crop and residential uses.³⁰ Since 2003, pendimethalin has been among the top ten most commonly applied pesticides in the US for agricultural use, and among the top five for residential use.³¹ In our study, higher pendimethalin use was also associated with anti-TPO positivity, suggesting that exposure may influence TSH through an autoimmune pathway. Pendimethalin has been shown *in vivo* to enhance the metabolism and excretion of T3 and T4, increase TSH, increase thyroid cell growth, and cause thyroid follicular cell adenomas.^{30,32} In previous analyses in the AHS cohort, pendimethalin was not associated with self-reported hypothyroidism in men or women.^{13,14} Though the associations with subclinical hypothyroidism, TSH, and anti-TPO at the highest levels of exposure are interesting, caution should be taken in interpreting these results. These findings are based on small numbers of exposed individuals and the exposure-response trends do not appear to be monotonic.

We noted significantly increased TSH levels among applicators with the highest EPTC use, with a significant exposure-response trend. Goldner et al. found no significant associations with EPTC and self-reported hypothyroidism among AHS men or women;^{13,14} however, there was evidence of non-significantly elevated risk among men with the highest exposure (above median intensity-weighted days of use). Recent studies evaluating EPTC and endocrine disruption found only modest effects on the hypothalamic-pituitary-thyroidal axis, noting decreases in serum T4 in rats at high doses and decreased thyroid weight in dogs.³³

When individuals with anti-TPO positivity were removed from the analyses, the associations for pendimethalin and EPTC with subclinical hypothyroidism and TSH were attenuated, suggesting that the effect of these herbicides on TSH may be stronger among participants with anti-TPO positivity. It has been hypothesized that individuals with thyroid autoantibodies are more susceptible to chemical exposures known to interfere with thyroid production or metabolism due to inability to properly compensate by increasing hormone production.³⁴ Direct studies are needed to investigate this mechanism.

2,4,5-T use was associated with significantly decreased T4 and a suggestive increase in subclinical hypothyroidism with no monotonic trend. First developed in the 1940s, US registrations of 2,4,5-T have been cancelled since 1985 due to serious health effects (e.g. birth defects, cancer) caused by contamination with the dioxin 2,3,7,8,-tetrachlorodibenzo-*p*-dioxin (TCDD) during the manufacturing process.^{35,36} Goldner et al. previously noted a significant positive association with self-reported hypothyroidism and ever use of 2,4,5-T among male pesticide applicators.¹³ Additionally, US Air Force veterans who sprayed Agent Orange, an herbicide containing 2,4-D and TCDD-contaminated 2,4,5-T, had significantly increased TSH even 20+ years after exposure compared to unexposed veterans. Based on the exposure information in our study, it is not possible to disentangle the separate effects of TCDD and 2,4,5-T on TSH and thyroid hormones. However, there is extensive animal

literature indicating that chronic exposure to TCDD is associated with lower T4 and higher TSH.⁹³⁷ It is possible that the associations observed for 2,4,5-T are due to confounding by TCDD.

Fumigants and fungicides

Higher intensity-weighted days of methyl bromide use were associated with lower TSH, significantly higher T4, and non-significantly elevated T3. Methyl bromide was first registered for use in the US as a soil fumigant in 1961, but has been banned since 2005 under the Clean Air Act and the Montreal Protocol on Substances that Deplete the Ozone Layer.³⁸ Methyl bromide has demonstrated neurotoxic effects,³⁸ though its potential for thyroid and other endocrine disruption has not been well-studied. In the AHS, methyl bromide use was associated with non-significantly elevated odds of self-reported hypothyroidism among women,¹⁴ hyperthyroidism among women and thyroid disease among men were not evaluated.¹³¹⁴ Despite relatively few individuals having been exposed to methyl bromide in our analysis, we noted consistent associations across TSH, T4, and T3.

We found that high use of the fungicide captan in our study was associated with higher TSH. These findings are consistent with those from an earlier AHS study,¹³ where ever use of captan was associated with non-significantly elevated risk of self-reported hypothyroidism among male pesticide applicators. Fungicide exposure has been associated with increased TSH levels among pesticide applicators in Minnesota.¹⁶³⁹ Thyroid cancer mortality was also significantly higher in the same agricultural areas of Minnesota with high fungicide use compared to non-agricultural areas.⁴⁰ While these studies did not investigate individual fungicides, captan was reportedly among the most commonly used fungicides during this time period in Minnesota.⁴⁰ Due to the small number of exposed applicators with subclinical hypothyroidism (N=1), we were not able to evaluate the fungicides maneb and mancozeb. These fungicides are applied in orchards and other settings where captan may be used, and were shown to be associated with both hypo- and hyperthyroidism among AHS spouses.¹⁴ However, in our study and in the cohort overall, use of captan and maneb/mancozeb are not correlated (Spearman ρ 0.02 and 0.17, respectively); it is unlikely that maneb/mancozeb use is confounding the relationship between captan and TSH in our study.

Strengths and Limitations

The characterization of altered thyroid function based on measured TSH and thyroid hormone levels was a strength of our investigation, as it allowed us to examine past pesticide exposure and subclinical changes in thyroid function. However, lack of laboratory measurement of free T4 was an important limitation and impacted our classification of subclinical hypothyroidism, which is often defined as TSH concentrations above the upper limit of the reference range (0.4–4.5 mIU/L) with normal free T4 levels. Individuals with TSH and free T4 outside the reference range may be considered to have overt hypothyroidism requiring medical attention;⁸ we were not able to make this distinction in our study. We excluded participants with self-reported thyroid disease or thyroid medication. These individuals may have normal TSH and thyroid hormone levels due to treatment, potentially biasing results toward the null. Use of a single blood sample to characterize TSH and thyroid hormones is a potential limitation; however, TSH has been shown to be

relatively stable over time (e.g., up to three years).⁴¹ To rule out confounding due to seasonality or diurnal variation we controlled for time of day and month of blood draw in our statistical models; neither was an important confounder. Measurement of multiple markers of thyroid function was a strength of our study, and we focused on results where we saw concordance between markers (e.g. higher TSH and lower T4). However, for some analyses we had limited statistical power to detect associations, particularly for subclinical hypothyroidism and pesticides with low prevalence of use.

Another strength is the detailed pesticide exposure assessment for many pesticides reported at AHS study enrollment.²⁰ Though exposure misclassification is a potential concern with self-reported pesticide use, the AHS intensity-weighting algorithm has been shown to correlate well with urinary pesticide measures.^{22,42} Additionally, due to the prospective cohort design and objective nature of the outcome measures, any exposure misclassification would likely be non-differential with respect to health outcomes. We were also able to adjust for use of other pesticides, minimizing potential confounding by use of correlated chemicals. For this study, we limited our analyses to self-reported lifetime days of use as reported at study enrollment. The evidence is unclear as to whether recent or chronic exposures have the greatest impact on thyroid function. Our strongest association was with cumulative exposure to aldrin where recent exposures would not be possible, as it has been banned in the US since 1974. For chemicals with ongoing use (e.g. pendimethalin), adjustment for use reported in follow-up questionnaires did not alter the reported main effects. For most pesticides, exposures reported at enrollment accounted for the majority of applicators' cumulative lifetime use. Future studies should examine windows of pesticide and other chemical exposures in relation to thyroid dysfunction and thyroid disease incidence.

CONCLUSIONS

Lifetime use of several pesticides was associated with increased TSH levels and subclinical hypothyroidism in our study population of male pesticide applicators. Our findings, taken together with limited epidemiologic studies and experimental evidence from animal models, suggest that certain pesticides may act as thyroid hormone disruptors. Furthermore, the agreement of our findings with previous results in the AHS for aldrin and self-reported hypothyroidism¹³ suggest that subclinical hormone disruption may potentially lead to overt hypothyroid disease among those exposed to this chemical. Thyroid disease and thyroid hormone disruption can impact human health, and may increase risk of other diseases including thyroid cancer,⁴³⁻⁴⁶ breast cancer,^{47,48} non-Hodgkin lymphoma,⁴⁹ and cardiovascular disease.⁵⁰ Most studies of pesticides and endocrine disruption have been conducted only in animals; more epidemiologic studies are necessary to understand how chronic occupational exposures impact thyroid function in men and women. Additionally, future work should attempt to assess whether short-term exposure to pesticides impacts thyroid hormone function in a similar manner as long-term use.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge Amy Miller, Kate Torres, Sandor Balogh, Shane Neese, Linda Gowen, Himanshi Singh, and Marsha Dunn (Westat, Inc., Rockville, Maryland) for study coordination, data management, and field research efforts for the BEEA study. We also thank Charles Lynch, Debra Lande, Debra Podaril, and Jennifer Hamilton from the field research team at the University of Iowa for their efforts on this study. AHS data in this analysis are based on releases *P1REL201506.00*, *P2REL201209* and *P3REL201209*.

FUNDING

This work was supported by the intramural research program of the National Institutes of Health, the National Cancer Institute (Z01-CP010119), and the National Institute of Environmental Health Sciences (Z01-ES049030).

References

1. Hollowell JG, Staehling NW, Flanders WD, et al. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab.* 2002; 87:489–99. [PubMed: 11836274]
2. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med.* 2000; 160:526–34. [PubMed: 10695693]
3. Surks MI, Ortiz E, Daniels GH, et al. Subclinical thyroid disease: scientific review and guidelines for diagnosis and management. *JAMA.* 2004; 291:228–38. [PubMed: 14722150]
4. Singer PA, Cooper DS, Levy EG, et al. Treatment guidelines for patients with hyperthyroidism and hypothyroidism. Standards of Care Committee, American Thyroid Association. *Jama.* 1995; 273:808–12. [PubMed: 7532241]
5. Bjoro T, Holmen J, Kruger O, et al. Prevalence of thyroid disease, thyroid dysfunction and thyroid peroxidase antibodies in a large, unselected population. The Health Study of Nord-Trondelag (HUNT). *Eur J Endocrinol.* 2000; 143:639–47. [PubMed: 11078988]
6. Vanderpump, MP., Tunbridge, WM. *The epidemiology of thyroid diseases.* Philadelphia: Lippincott Williams and Wilkins; 2000.
7. Charades T, Chapal N, Bresson D, et al. The human anti-thyroid peroxidase autoantibody repertoire in Graves' and Hashimoto's autoimmune thyroid diseases. *Immunogenetics.* 2002; 54:141–57. [PubMed: 12073143]
8. Cooper DS, Biondi B. Subclinical thyroid disease. *Lancet.* 2012; 379:1142–54. [PubMed: 22273398]
9. Brucker-Davis F. Effects of environmental synthetic chemicals on thyroid function. *Thyroid.* 1998; 8:827–56. [PubMed: 9777756]
10. Kuiper GG, Lemmen JG, Carlsson B, et al. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology.* 1998; 139:4252–63. [PubMed: 9751507]
11. Boas M, Feldt-Rasmussen U, Skakkebaek NE, Main KM. Environmental chemicals and thyroid function. *European Journal of Endocrinology.* 2006; 154:599–611. [PubMed: 16645005]
12. Zoeller RT. Environmental chemicals impacting the thyroid: Targets and consequences. *Thyroid.* 2007; 17:811–817. [PubMed: 17956155]
13. Goldner WS, Sandler DP, Yu F, et al. Hypothyroidism and pesticide use among male private pesticide applicators in the agricultural health study. *J Occup Environ Med.* 2013; 55:1171–8. [PubMed: 24064777]
14. Goldner WS, Sandler DP, Yu F, Hoppin JA, Kamel F, Levan TD. Pesticide use and thyroid disease among women in the Agricultural Health Study. *Am J Epidemiol.* 2010; 171:455–64. [PubMed: 20061368]
15. Steenland K, Cedillo L, Tucker J, et al. Thyroid hormones and cytogenetic outcomes in backpack sprayers using ethylenebis(dithiocarbamate) (EBDC) fungicides in Mexico. *Environmental Health Perspectives.* 1997; 105:1126–1130. [PubMed: 9349837]
16. Garry VF, Holland SE, Erickson LL, Burroughs BL. Male reproductive hormones and thyroid function in pesticide applicators in the Red River Valley of Minnesota. *J Toxicol Environ Health A.* 2003; 66:965–86. [PubMed: 12775511]

17. Toft G, Flyvbjerg A, Bonde JP. Thyroid function in Danish greenhouse workers. *Environ Health*. 2006; 5:32. [PubMed: 17147831]
18. Piccoli C, Cremonese C, Koifman RJ, Koifman S, Freire C. Pesticide exposure and thyroid function in an agricultural population in Brazil. *Environ Res*. 2016; 151:389–398. [PubMed: 27540871]
19. Hofmann JN, Beane Freeman LE, Lynch CF, et al. The Biomarkers of Exposure and Effect in Agriculture (BEEA) Study: Rationale, Design, Methods, and Participant Characteristics. *J Toxicol Environ Health A*. 2015; 78:1338–47. [PubMed: 26555155]
20. Alavanja MC, Sandler DP, McMaster SB, et al. The Agricultural Health Study. *Environ Health Perspect*. 1996; 104:362–9. [PubMed: 8732939]
21. LeFevre ML. Screening for thyroid dysfunction: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2015; 162:641–50. [PubMed: 25798805]
22. Coble J, Thomas KW, Hines CJ, et al. An Updated Algorithm for Estimation of Pesticide Exposure Intensity in the Agricultural Health Study. *International Journal of Environmental Research and Public Health*. 2011; 8:4608–4622. [PubMed: 22408592]
23. Abbot, DC., Kimbrough, RD., Kroes, R., et al. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man: Some organochlorine pesticides. Lyon, France: International Agency for Research on Cancer; 1974.
24. Carcinogenicity of Aldrin and Dieldrin (EPA/600/6–87/006). Washington, DC: Carcinogen Assessment Group, Office of Health and Environmental Assessment, Office of Research and Development, United States Environmental Protection Agency; 1987.
25. Kutz, FW., Wood, PH., Bottimore, DP. Organochlorine Pesticides and Polychlorinated Biphenyls in Human Adipose Tissue*. In: Ware, GW., editor. *Reviews of Environmental Contamination and Toxicology*. New York, NY: Springer New York; 1991. p. 1-82.
26. Freire C, Koifman RJ, Sarcinelli PN, Simoes Rosa AC, Clapauch R, Koifman S. Long-term exposure to organochlorine pesticides and thyroid status in adults in a heavily contaminated area in Brazil. *Environ Res*. 2013; 127:7–15. [PubMed: 24183346]
27. Rathore M, Bhatnagar P, Mathur D, Saxena GN. Burden of organochlorine pesticides in blood and its effect on thyroid hormones in women. *Sci Total Environ*. 2002; 295:207–15. [PubMed: 12186288]
28. Wade MG, Parent S, Finnson KW, et al. Thyroid toxicity due to subchronic exposure to a complex mixture of 16 organochlorines, lead, and cadmium. *Toxicol Sci*. 2002; 67:207–18. [PubMed: 12011480]
29. Bioassays of aldrin and dieldrin for possible carcinogenicity. *Natl Cancer Inst Carcinog Tech Rep Ser*. 1978; 21:1–184. [PubMed: 12844187]
30. Reregistration Eligibility Decision Facts: Pendimethalin. Washington, DC: US Environmental Protection Agency; 1997.
31. Grube, A., Donaldson, D., Kiely, T., Wu, L. Pesticide Industry Sales and Usage: 2006 and 2007 Market Estimates. Washington, DC: Office of Pesticide Programs, US Environmental Protection Agency; 2011.
32. Hurley PM. Mode of carcinogenic action of pesticides inducing thyroid follicular cell tumors in rodents. *Environ Health Perspect*. 1998; 106:437–45. [PubMed: 9681970]
33. Chemical: EPTC. Washington, DC: Office of Pesticide Programs, Office of Science Coordination and Policy, U.S. Environmental Protection Agency; 2015. Endocrine Disruptor Screening Program: Weight of Evidence Analysis of Potential Interaction with the Estrogen, Androgen, or Thyroid Pathways.
34. Brent GA. Environmental exposures and autoimmune thyroid disease. *Thyroid*. 2010; 20:755–61. [PubMed: 20578899]
35. EPA Fact Sheet: Regulatory Status of 2,4,5-T. San Francisco, CA: US Environmental Protection Agency; 1978.
36. 2,4-D. US Environmental Protection Agency; 2016. <https://www.epa.gov/ingredients-used-pesticide-products/24-d> accessed

37. Pohl, H., Lladós, F., Ingerman, L., et al. Sciences DoTaHH. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 1998. Toxicological Profile for Chlorinated Dibenzo-p-dioxins (CDDs).
38. Prevention P, and Toxic Substances. Washington, DC: US Environmental Protection Agency; 2009. Amended Reregistration Eligibility Decision for Methyl Bromide (soil and non-food structural uses).
39. Garry VF. Biomarkers of thyroid function, genotoxicity and agricultural fungicide use. *Journal of Biochemical and Molecular Toxicology*. 2005; 19:175–175. [PubMed: 15977190]
40. Schreinemachers DM, Creason JP, Garry VF. Cancer mortality in agricultural regions of Minnesota. *Environmental Health Perspectives*. 1999; 107:205–211. [PubMed: 10064550]
41. Arslan AA, Gu Y, Zeleniuch-Jacquotte A, et al. Reproducibility of serum pituitary hormones in women. *Cancer Epidemiol Biomarkers Prev*. 2008; 17:1880–3. [PubMed: 18708375]
42. Thomas KW, Dosemeci M, Coble JB, et al. Assessment of a pesticide exposure intensity algorithm in the agricultural health study. *J Expo Sci Environ Epidemiol*. 2010; 20:559–69. [PubMed: 19888312]
43. Boelaert K, Horacek J, Holder RL, Watkinson JC, Sheppard MC, Franklyn JA. Serum thyrotropin concentration as a novel predictor of malignancy in thyroid nodules investigated by fine-needle aspiration. *Journal of Clinical Endocrinology & Metabolism*. 2006; 91:4295–4301. [PubMed: 16868053]
44. Kim SS, Lee BJ, Lee JC, et al. Preoperative serum thyroid stimulating hormone levels in well-differentiated thyroid carcinoma is a predictive factor for lateral lymph node metastasis as well as extrathyroidal extension in Korean patients: a single-center experience. *Endocrine*. 2011; 39:259–265. [PubMed: 21161440]
45. Meinhold CL, Ron E, Schonfeld SJ, et al. Nonradiation risk factors for thyroid cancer in the US Radiologic Technologists Study. *Am J Epidemiol*. 2010; 171:242–52. [PubMed: 19951937]
46. Franceschi S, Preston-Martin S, Dal Maso L, et al. A pooled analysis of case-control studies of thyroid cancer. IV. Benign thyroid diseases. *Cancer Causes Control*. 1999; 10:583–95. [PubMed: 10616827]
47. Hardefeldt PJ, Eslick GD, Edirimanne S. Benign thyroid disease is associated with breast cancer: a meta-analysis. *Breast Cancer Res Treat*. 2012; 133:1169–77. [PubMed: 22434524]
48. Sogaard M, Farkas DK, Ehrenstein V, Jorgensen JO, Dekkers OM, Sorensen HT. Hypothyroidism and hyperthyroidism and breast cancer risk: a nationwide cohort study. *Eur J Endocrinol*. 2016; 174:409–14. [PubMed: 26863886]
49. Fallah M, Liu X, Ji J, Forsti A, Sundquist K, Hemminki K. Autoimmune diseases associated with non-Hodgkin lymphoma: a nationwide cohort study. *Ann Oncol*. 2014; 25:2025–30. [PubMed: 25081899]
50. Osman F, Gammage MD, Franklyn JA. Thyroid disease and its treatment: short-term and long-term cardiovascular consequences. *Curr Opin Pharmacol*. 2001; 1:626–31. [PubMed: 11757819]

SUMMARY

1. What is already known about this subject?

Studies in laboratory animals have suggested that occupational exposure to individual pesticides may alter thyroid function. Posited mechanisms include competitive binding to thyroid hormone receptors and interference with thyroid enzymes. However, few epidemiologic studies have examined this association.

2. What are the new findings?

In a large study of U.S. male pesticide applicators, we observed associations between long-term occupational exposure to certain pesticides with thyroid hormone levels and subclinical hypothyroidism.

3. How might it impact on clinical practice in the foreseeable future?

For many of the pesticides examined, this is the first report in a human population; thus, there are no immediate implications for clinical practice. Future studies are needed to confirm these findings.

Table 1

Selected characteristics of pesticide applicators at BEEA sample collection (unless noted) stratified by subclinical hypothyroidism (TSH 0.4–4.5 mIU/L or >4.5 mIU/L)

	Total N=679 N (%)	Subclinical hypothyroidism		p-value ^a
		No N=552 N (%)	Yes N=127 N (%)	
Age				
50–59	254 (37.41)	210 (38.04)	44 (34.65)	0.12
60–69	231 (34.02)	193 (34.96)	38 (29.92)	
70–79	156 (22.97)	123 (22.28)	33 (25.98)	
80+	38 (5.60)	26 (4.71)	12 (9.45)	
Race ^b				
Non-Hispanic white	664 (97.80)	540 (97.80)	124 (97.60)	0.90
Other	15 (2.20)	12 (2.20)	3 (2.40)	
State ^b				
Iowa	582 (85.71)	474 (85.87)	108 (85.04)	0.81
North Carolina	97 (14.29)	78 (14.13)	19 (14.96)	
Body mass index				
Normal (<25)	118 (17.38)	97 (17.57)	21 (16.54)	0.86
Overweight (25–29.9)	301 (44.33)	242 (43.84)	59 (46.46)	
Obese (30+)	260 (38.29)	213 (38.59)	47 (37.01)	
Tobacco smoking ^c				
Never	407 (59.94)	325 (58.88)	82 (64.57)	0.46
Former	230 (33.87)	191 (34.6)	39 (30.71)	
Current	42 (6.19)	36 (6.52)	6 (4.72)	
Alcohol consumption (last 7 days)				
None	322 (47.42)	257 (46.56)	65 (51.18)	0.63
<5 drinks	206 (30.34)	171 (30.98)	35 (27.56)	
5+ drinks	151 (22.24)	124 (22.46)	27 (21.26)	
Education ^b				
Less than high school	26 (3.83)	18 (3.26)	8 (6.3)	0.29
High school graduate	321 (47.28)	260 (47.1)	61 (48.03)	
Vocational school/some college	157 (23.12)	133 (24.09)	24 (18.9)	
College grad	162 (23.86)	132 (23.91)	30 (23.62)	
Missing/don't know	13 (1.91)	9 (1.63)	4 (3.15)	
Raised Farm Animals in the last 12 months				
Did not farm	104 (15.32)	79 (14.31)	25 (19.69)	0.06
No	274 (40.35)	217 (39.31)	57 (44.88)	
Yes	301 (44.33)	256 (46.38)	45 (35.43)	
Applied pesticides occupationally in the last 12 months				
No	192 (28.28)	148 (26.81)	44 (34.65)	0.08

	Total N=679 N (%)	Subclinical hypothyroidism		p-value ^a
		No N=552 N (%)	Yes N=127 N (%)	
Yes	487 (71.72)	404 (73.19)	83 (65.35)	
Off-farm job in the last 12 months				
No	446 (65.68)	365 (66.12)	81 (63.78)	0.62
Yes	233 (34.32)	187 (33.88)	46 (36.22)	
Anti-TPO antibodies				
Negative	614 (90.43)	516 (93.48)	98 (77.17)	<0.01
Positive	65 (9.57)	36 (6.52)	29 (22.83)	

BEEA: Biomarkers of Exposure and Effect in Agriculture; TSH: Thyroid stimulating hormone

^aChi-square test for homogeneity

^bCollected at AHS enrollment

^cCollected at the second AHS follow-up

Table 2
Multivariate^a regression models examining association between intensity-weighted days of specific insecticides and subclinical hypothyroidism (TSH>4.5 mIU/L) and natural log of TSH, T4, and T3

	TSH and Thyroid Hormones										
	Subclinical hypothyroidism					TSH		T4		T3	
	Intensity-weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	
Organochlorine											
Aldrin ^d	0	85	1.00 (ref)	533	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	26-210	5	0.99 (0.31, 3.18)	25	1.10 (0.84, 1.45)	1.07 (0.96, 1.20)	1.07 (0.96, 1.20)	1.09 (0.97, 1.23)	1.09 (0.97, 1.23)	1.09 (0.97, 1.23)	
	>210-392	5	1.53 (0.44, 5.28)	19	0.89 (0.65, 1.22)	1.01 (0.89, 1.15)	1.01 (0.89, 1.15)	0.97 (0.84, 1.11)	0.97 (0.84, 1.11)	0.97 (0.84, 1.11)	
	>392-1117	9	4.15 (1.56, 11.0) [†]	20	1.53 (1.16, 2.03) [†]	1.03 (0.92, 1.16)	1.03 (0.92, 1.16)	1.00 (0.89, 1.13)	1.00 (0.89, 1.13)	1.00 (0.89, 1.13)	
	>1117-10339	9	4.76 (1.53, 14.8) [†]	21	1.33 (0.97, 1.81)	0.87 (0.76, 0.98) [†]	0.87 (0.76, 0.98) [†]	0.88 (0.77, 1.01)	0.88 (0.77, 1.01)	0.88 (0.77, 1.01)	
	P _{trend}		<0.01 [†]		0.01 [†]	0.04 [†]	0.04 [†]	0.08	0.08	0.08	
Chlordane^d											
	0	92	1.00 (ref)	521	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	24-210	5	1.13 (0.39, 3.27)	23	0.94 (0.72, 1.23)	1.00 (0.89, 1.11)	1.00 (0.89, 1.11)	0.95 (0.85, 1.06)	0.95 (0.85, 1.06)	0.95 (0.85, 1.06)	
	>210-315	4	1.45 (0.43, 4.91)	16	0.80 (0.58, 1.10)	1.01 (0.89, 1.15)	1.01 (0.89, 1.15)	0.99 (0.87, 1.14)	0.99 (0.87, 1.14)	0.99 (0.87, 1.14)	
	>315-980	5	1.35 (0.45, 4.11)	20	1.10 (0.82, 1.47)	1.02 (0.91, 1.15)	1.02 (0.91, 1.15)	1.03 (0.91, 1.16)	1.03 (0.91, 1.16)	1.03 (0.91, 1.16)	
	>980-13020	6	1.80 (0.57, 5.73)	19	1.08 (0.78, 1.48)	0.96 (0.84, 1.09)	0.96 (0.84, 1.09)	1.07 (0.93, 1.22)	1.07 (0.93, 1.22)	1.07 (0.93, 1.22)	
	P _{trend}		0.30		0.68	0.56	0.56	0.35	0.35	0.35	
DDT^d											
	0	92	1.00 (ref)	531	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	18-240	5	0.88 (0.27, 2.83)	22	0.95 (0.71, 1.28)	1.02 (0.90, 1.15)	1.02 (0.90, 1.15)	1.06 (0.94, 1.20)	1.06 (0.94, 1.20)	1.06 (0.94, 1.20)	
	>240-662	5	0.87 (0.28, 2.76)	21	1.17 (0.87, 1.56)	1.02 (0.91, 1.15)	1.02 (0.91, 1.15)	1.02 (0.90, 1.16)	1.02 (0.90, 1.16)	1.02 (0.90, 1.16)	
	>662-2625	5	0.69 (0.22, 2.21)	22	0.91 (0.68, 1.22)	1.10 (0.98, 1.24)	1.10 (0.98, 1.24)	1.05 (0.93, 1.19)	1.05 (0.93, 1.19)	1.05 (0.93, 1.19)	
	>2625-172800	6	0.85 (0.26, 2.81)	22	1.03 (0.75, 1.41)	1.11 (0.98, 1.26)	1.11 (0.98, 1.26)	1.02 (0.89, 1.16)	1.02 (0.89, 1.16)	1.02 (0.89, 1.16)	
	P _{trend}		0.87		0.88	0.14	0.14	0.90	0.90	0.90	
Heptachlor^d											
	0	93	1.00 (ref)	550	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	26-182	5	1.65 (0.50, 5.39)	19	1.33 (0.98, 1.82)	0.86 (0.76, 0.98)	0.86 (0.76, 0.98)	0.89 (0.78, 1.02)	0.89 (0.78, 1.02)	0.89 (0.78, 1.02)	
	>182-474	6	2.15 (0.67, 6.89)	18	1.21 (0.89, 1.64)	1.06 (0.93, 1.20)	1.06 (0.93, 1.20)	0.93 (0.82, 1.06)	0.93 (0.82, 1.06)	0.93 (0.82, 1.06)	
	>474-980	6	1.27 (0.36, 4.42)	18	1.14 (0.82, 1.59)	0.93 (0.81, 1.07)	0.93 (0.81, 1.07)	0.99 (0.86, 1.14)	0.99 (0.86, 1.14)	0.99 (0.86, 1.14)	

TSH and Thyroid Hormones									
Subclinical hypothyroidism					TSH		T4		T3
Intensity- weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)
>980-30680	4	0.63 (0.16, 2.46)	18	1.00 (0.73, 1.38)	1.06 (0.93, 1.20)	1.06 (0.92, 1.22)			
P _{trend}		0.75		0.68	0.73	0.57			
Organophosphate									
Chlorpyrifos	0		60	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	25-407		15	0.99 (0.52, 1.87)	0.94 (0.81, 1.09)	1.01 (0.95, 1.07)	1.02 (0.96, 1.09)	1.02 (0.96, 1.09)	1.02 (0.96, 1.09)
	>407-1005		18	1.22 (0.67, 2.22)	0.99 (0.85, 1.15)	1.03 (0.97, 1.09)	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)
	>1005-2807		20	1.35 (0.75, 2.41)	1.03 (0.89, 1.19)	1.07 (1.01, 1.14) [†]	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)
	>2807-53708		12	0.71 (0.36, 1.41)	1.00 (0.86, 1.16)	1.02 (0.96, 1.09)	1.01 (0.95, 1.07)	1.01 (0.95, 1.07)	1.01 (0.95, 1.07)
P _{trend}		0.39		0.88	0.36	0.91			
Diazinon ^d									
	0		86	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	28-236		4	1.32 (0.42, 4.11)	1.16 (0.87, 1.54)	0.95 (0.84, 1.06)	1.00 (0.88, 1.13)	1.00 (0.88, 1.13)	1.00 (0.88, 1.13)
	>236-420		7	2.44 (0.95, 6.30)	1.04 (0.79, 1.36)	1.04 (0.93, 1.16)	1.04 (0.93, 1.17)	1.04 (0.93, 1.17)	1.04 (0.93, 1.17)
	>420-882		7	2.64 (1.00, 6.98) [‡]	1.40 (1.06, 1.86) [‡]	0.98 (0.87, 1.10)	0.98 (0.87, 1.11)	0.98 (0.87, 1.11)	0.98 (0.87, 1.11)
	>882-13423		6	1.88 (0.68, 5.17)	1.22 (0.92, 1.62)	1.10 (0.98, 1.23)	1.03 (0.92, 1.17)	1.03 (0.92, 1.17)	1.03 (0.92, 1.17)
P _{trend}		0.10		0.07	0.15	0.62			
Fonofos									
	0		78	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	28-368		11	1.23 (0.60, 2.52)	0.95 (0.80, 1.14)	0.98 (0.91, 1.05)	0.98 (0.91, 1.06)	0.98 (0.91, 1.06)	0.98 (0.91, 1.06)
	>368-982		11	1.16 (0.56, 2.38)	0.94 (0.79, 1.13)	1.06 (0.98, 1.13)	1.03 (0.96, 1.11)	1.03 (0.96, 1.11)	1.03 (0.96, 1.11)
	>982-2646		13	1.35 (0.68, 2.69)	0.98 (0.82, 1.17)	0.99 (0.92, 1.06)	1.05 (0.97, 1.13)	1.05 (0.97, 1.13)	1.05 (0.97, 1.13)
	>2646-21948		10	1.09 (0.52, 2.31)	1.16 (0.97, 1.39)	1.03 (0.95, 1.10)	0.96 (0.89, 1.04)	0.96 (0.89, 1.04)	0.96 (0.89, 1.04)
P _{trend}		0.73		0.11	0.53	0.51			
Malathion ^d									
	0		29	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	20-276		12	1.04 (0.47, 2.28)	0.90 (0.76, 1.07)	1.00 (0.93, 1.08)	1.02 (0.95, 1.10)	1.02 (0.95, 1.10)	1.02 (0.95, 1.10)
	>276-780		15	1.20 (0.55, 2.60)	1.02 (0.86, 1.22)	1.00 (0.93, 1.08)	1.03 (0.95, 1.11)	1.03 (0.95, 1.11)	1.03 (0.95, 1.11)
	>780-2250		14	0.95 (0.42, 2.13)	0.90 (0.75, 1.07)	1.03 (0.96, 1.12)	1.02 (0.94, 1.10)	1.02 (0.94, 1.10)	1.02 (0.94, 1.10)
	>2250-117600		18	1.50 (0.69, 3.27)	1.10 (0.91, 1.33)	0.99 (0.91, 1.07)	1.00 (0.92, 1.09)	1.00 (0.92, 1.09)	1.00 (0.92, 1.09)
P _{trend}		0.30		0.16	0.65	0.81			

TSH and Thyroid Hormones									
	Intensity- weighted days	Subclinical hypothyroidism			TSH		T4		T3
		N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	
Phorate ^d	0	79	1.00 (ref)	429	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	28-236	6	0.81 (0.32, 2.05)	36	0.95 (0.77, 1.18)	1.01 (0.93, 1.10)	1.01 (0.93, 1.10)	1.00 (0.91, 1.09)	
	>236-588	6	0.76 (0.30, 1.92)	38	1.03 (0.84, 1.27)	1.02 (0.94, 1.11)	1.02 (0.94, 1.11)	1.04 (0.96, 1.14)	
	>588-1302	7	0.98 (0.41, 2.35)	38	1.16 (0.94, 1.43)	1.01 (0.93, 1.10)	1.01 (0.93, 1.10)	0.94 (0.87, 1.03)	
	>1302-37758	8	1.10 (0.47, 2.58)	37	1.03 (0.83, 1.27)	1.03 (0.94, 1.12)	1.03 (0.94, 1.12)	0.98 (0.90, 1.07)	
P _{trend}			0.80		0.54	0.55	0.43		
Terbufos	0	62	1.00 (ref)	309	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	34-463	18	1.17 (0.63, 2.16)	86	0.93 (0.80, 1.08)	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)	1.08 (1.01, 1.15) [†]	
	>463-1259	15	0.87 (0.46, 1.64)	86	0.97 (0.83, 1.12)	1.01 (0.95, 1.07)	1.01 (0.95, 1.07)	1.03 (0.96, 1.09)	
	>1259-3368	12	0.65 (0.33, 1.29)	86	0.91 (0.78, 1.06)	1.00 (0.94, 1.07)	1.00 (0.94, 1.07)	1.07 (1.00, 1.14) [†]	
	>3368-17748	16	0.95 (0.51, 1.78)	86	0.99 (0.85, 1.15)	1.02 (0.96, 1.08)	1.02 (0.96, 1.08)	1.07 (1.00, 1.13) [†]	
P _{trend}			0.72		0.97	0.69	0.10		
Carbamate	0	71	1.00 (ref)	397	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	28-200	7	1.03 (0.42, 2.55)	39	0.85 (0.69, 1.05)	1.02 (0.93, 1.11)	1.02 (0.93, 1.11)	1.08 (0.99, 1.19)	
	>200-662	10	1.79 (0.78, 4.11)	38	1.06 (0.86, 1.32)	1.01 (0.92, 1.10)	1.01 (0.92, 1.10)	1.03 (0.94, 1.13)	
	>662-2315	9	1.42 (0.59, 3.45)	40	1.18 (0.95, 1.47)	1.06 (0.97, 1.16)	1.06 (0.97, 1.16)	1.03 (0.94, 1.14)	
	>2315-227850	7	0.77 (0.27, 2.23)	38	1.06 (0.83, 1.36)	0.96 (0.87, 1.07)	0.96 (0.87, 1.07)	0.92 (0.83, 1.03)	
P _{trend}			0.52		0.58	0.45	0.08		
Carbofuran	0	67	1.00 (ref)	413	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	28-294	14	1.67 (0.86, 3.24)	58	0.95 (0.81, 1.13)	0.96 (0.90, 1.03)	0.96 (0.90, 1.03)	1.00 (0.94, 1.08)	
	>294-858	13	1.59 (0.80, 3.13)	56	1.07 (0.90, 1.28)	1.04 (0.97, 1.12)	1.04 (0.97, 1.12)	1.00 (0.93, 1.08)	
	>858-2016	12	1.32 (0.66, 2.65)	56	1.05 (0.89, 1.25)	0.97 (0.90, 1.04)	0.97 (0.90, 1.04)	1.01 (0.94, 1.08)	
	>2016-33837	14	1.58 (0.81, 3.07)	59	1.01 (0.85, 1.19)	1.03 (0.96, 1.10)	1.03 (0.96, 1.10)	1.03 (0.96, 1.11)	
P _{trend}			0.21		0.79	0.55	0.40		
Pyrethroid	0	89	1.00 (ref)	460	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	
	Permethrin								

	TSH and Thyroid Hormones					
	Subclinical hypothyroidism			TSH		T3
	Intensity- weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)
24–230	9	0.93 (0.43, 2.01)	50	0.91 (0.76, 1.09)	0.96 (0.90, 1.04)	0.98 (0.90, 1.05)
>230–599	5	0.49 (0.19, 1.28)	50	0.81 (0.68, 0.98) [‡]	1.03 (0.96, 1.11)	0.98 (0.91, 1.06)
>599–2021	10	1.09 (0.52, 2.29)	50	1.05 (0.88, 1.26)	1.03 (0.96, 1.11)	1.03 (0.96, 1.12)
>2021–612000	10	1.14 (0.54, 2.39)	50	1.13 (0.94, 1.35)	1.02 (0.95, 1.10)	1.03 (0.95, 1.11)
P _{trend}		0.66		0.13	0.50	0.37

TSH: Thyroid stimulating hormone; T4: thyroxine; T3: triiodothyronine

^a Adjusted for age, state, body mass index, smoking, correlated pesticides

^b Exposed cases

^c Total exposed

^d Detailed information for these chemicals was collected on the take-home questionnaire at enrollment

[‡] p<0.05 compared with never users

Table 3 Multivariate^a regression models examining association between intensity-weighted days of specific herbicides and subclinical hypothyroidism (TSH>4.5 mIU/L) and natural log of TSH, T4, and T3

	TSH and Thyroid Hormones									
	Subclinical Hypothyroidism					TSH		T4		T3
	Intensity-weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	
Dinitroaniline										
Pendimethalin ^d	0	68	1.00 (ref)	412	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	28-208	10	1.77 (0.81, 3.87)	36	1.01 (0.82, 1.24)	0.98 (0.90, 1.07)	0.98 (0.90, 1.07)	1.01 (0.92, 1.11)	1.01 (0.92, 1.11)	1.01 (0.92, 1.11)
	>208-375	6	0.96 (0.38, 2.45)	35	0.89 (0.72, 1.11)	1.02 (0.94, 1.12)	1.02 (0.94, 1.12)	1.02 (0.93, 1.12)	1.02 (0.93, 1.12)	1.02 (0.93, 1.12)
	>375-1920	5	0.78 (0.29, 2.09)	36	1.05 (0.85, 1.30)	1.04 (0.96, 1.14)	1.04 (0.96, 1.14)	1.03 (0.94, 1.12)	1.03 (0.94, 1.12)	1.03 (0.94, 1.12)
	>1920-14105	12	2.78 (1.30, 5.95) [†]	35	1.25 (1.01, 1.55) [†]	0.98 (0.90, 1.07)	0.98 (0.90, 1.07)	0.96 (0.87, 1.05)	0.96 (0.87, 1.05)	0.96 (0.87, 1.05)
	P _{trend}		0.02 [†]		0.04 [†]	0.81	0.81	0.46	0.46	0.46
Trifluralin										
	0	55	1.00 (ref)	308	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	25-632	15	0.94 (0.47, 1.87)	87	0.99 (0.84, 1.15)	0.99 (0.92, 1.05)	0.99 (0.92, 1.05)	0.99 (0.93, 1.06)	0.99 (0.93, 1.06)	0.99 (0.93, 1.06)
	>632-1411	13	0.74 (0.35, 1.53)	87	0.90 (0.76, 1.05)	1.05 (0.98, 1.12)	1.05 (0.98, 1.12)	1.01 (0.94, 1.08)	1.01 (0.94, 1.08)	1.01 (0.94, 1.08)
	>1411-3906	23	1.42 (0.74, 2.74)	87	1.05 (0.90, 1.24)	0.98 (0.92, 1.05)	0.98 (0.92, 1.05)	0.98 (0.91, 1.05)	0.98 (0.91, 1.05)	0.98 (0.91, 1.05)
	>3906-113400	18	1.09 (0.55, 2.17)	87	1.14 (0.97, 1.34)	1.05 (0.99, 1.13)	1.05 (0.99, 1.13)	1.04 (0.97, 1.11)	1.04 (0.97, 1.11)	1.04 (0.97, 1.11)
	P _{trend}		0.65		0.05	0.14	0.14	0.28	0.28	0.28
Thiocarbamate										
Butylate^d										
	0	66	1.00 (ref)	390	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	43-319	7	1.19 (0.45, 3.16)	40	0.94 (0.75, 1.16)	1.00 (0.91, 1.09)	1.00 (0.91, 1.09)	0.98 (0.89, 1.08)	0.98 (0.89, 1.08)	0.98 (0.89, 1.08)
	>319-882	8	1.30 (0.50, 3.40)	39	1.10 (0.88, 1.38)	0.99 (0.91, 1.09)	0.99 (0.91, 1.09)	1.00 (0.91, 1.10)	1.00 (0.91, 1.10)	1.00 (0.91, 1.10)
	>882-2184	9	1.23 (0.50, 3.02)	39	1.20 (0.96, 1.49)	0.96 (0.88, 1.05)	0.96 (0.88, 1.05)	0.99 (0.90, 1.09)	0.99 (0.90, 1.09)	0.99 (0.90, 1.09)
	>2184-22925	12	1.78 (0.74, 4.28)	42	1.11 (0.89, 1.39)	0.99 (0.90, 1.08)	0.99 (0.90, 1.08)	0.97 (0.88, 1.07)	0.97 (0.88, 1.07)	0.97 (0.88, 1.07)
	P _{trend}		0.21		0.24	0.72	0.72	0.53	0.53	0.53
EPTC										
	0	87	1.00 (ref)	510	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	33-225	8	1.65 (0.71, 3.81)	34	1.01 (0.82, 1.25)	1.00 (0.92, 1.09)	1.00 (0.92, 1.09)	1.02 (0.93, 1.12)	1.02 (0.93, 1.12)	1.02 (0.93, 1.12)
	>225-588	6	1.12 (0.44, 2.82)	32	0.93 (0.75, 1.16)	0.99 (0.91, 1.09)	0.99 (0.91, 1.09)	1.00 (0.91, 1.1)	1.00 (0.91, 1.1)	1.00 (0.91, 1.1)

		TSH and Thyroid Hormones						
		Subclinical Hypothyroidism			TSH		T3	
		N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)
Intensity-weighted days								
>588-1792		10	2.10 (0.96, 4.58)	34	1.09 (0.88, 1.35)	0.88 (0.81, 0.96) [†]	0.88 (0.80, 0.96) [†]	
>1792-16461		9	2.05 (0.91, 4.63)	33	1.24 (1.00, 1.54) [†]	1.03 (0.94, 1.13)	1.03 (0.94, 1.13)	
P _{trend}			0.05		0.04 [†]	0.97	0.93	
Phenoxy								
2,4-D		25	1.00 (ref)	118	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
68-878		22	0.75 (0.39, 1.44)	136	0.94 (0.81, 1.10)	0.99 (0.93, 1.05)	0.98 (0.92, 1.05)	
>878-2604		26	0.81 (0.43, 1.55)	138	1.01 (0.86, 1.18)	0.99 (0.92, 1.05)	0.99 (0.93, 1.06)	
>2604-7229		24	0.70 (0.36, 1.36)	134	1.04 (0.88, 1.22)	0.97 (0.91, 1.04)	0.97 (0.91, 1.04)	
>7229-192780		29	0.80 (0.40, 1.58)	138	1.09 (0.92, 1.29)	0.98 (0.91, 1.05)	0.95 (0.88, 1.02)	
P _{trend}			0.85		0.14	0.64	0.14	
2,4,5 T ^d								
0		88	1.00 (ref)	518	1.00 (ref)	1.00 (ref)	1.00 (ref)	
25-263		7	2.58 (0.94, 7.06)	21	1.04 (0.78, 1.38)	0.93 (0.83, 1.04)	1.04 (0.93, 1.17)	
>263-458		5	1.23 (0.41, 3.70)	22	0.99 (0.74, 1.31)	0.95 (0.85, 1.06)	0.98 (0.87, 1.10)	
>458-1392		6	1.72 (0.63, 4.65)	22	1.28 (0.98, 1.68)	1.03 (0.93, 1.15)	1.00 (0.89, 1.12)	
>1392-29869		7	2.10 (0.78, 5.71)	21	1.03 (0.78, 1.37)	0.88 (0.78, 0.98) [†]	0.90 (0.80, 1.02)	
P _{trend}			0.16		0.68	0.03 [†]	0.09	
Triazine								
Atrazine		19	1.00 (ref)	101	1.00 (ref)	1.00 (ref)	1.00 (ref)	
25-980		23	0.76 (0.35, 1.63)	142	0.86 (0.73, 1.02)	1.02 (0.95, 1.10)	1.06 (0.99, 1.14)	
>980-2604		20	0.55 (0.25, 1.23)	144	0.79 (0.66, 0.94) [†]	1.03 (0.96, 1.11)	1.01 (0.94, 1.09)	
>2604-6962		33	1.12 (0.50, 2.48)	140	0.84 (0.70, 1.02)	0.97 (0.90, 1.05)	0.98 (0.91, 1.06)	
>6962-113400		32	1.16 (0.49, 2.78)	145	0.88 (0.72, 1.07)	0.99 (0.91, 1.07)	1.00 (0.92, 1.09)	
P _{trend}			0.28		0.97	0.45	0.50	
Cyanazine								
0		48	1.00 (ref)	294	1.00 (ref)	1.00 (ref)	1.00 (ref)	
25-455		16	0.89 (0.44, 1.80)	90	0.97 (0.83, 1.14)	0.98 (0.92, 1.05)	0.97 (0.91, 1.04)	
>455-1313		8	0.43 (0.18, 1.01)	89	0.97 (0.83, 1.13)	1.00 (0.94, 1.06)	1.04 (0.98, 1.11)	
>1313-3689		28	2.53 (1.34, 4.78) [†]	90	1.19 (1.02, 1.39) [†]	1.00 (0.94, 1.07)	1.00 (0.94, 1.07)	

		TSH and Thyroid Hormones						
		Subclinical Hypothyroidism			TSH		T3	
	Intensity-weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)
	>3689–113400	20	1.26 (0.61, 2.61)	89	1.13 (0.95, 1.34)	1.04 (0.97, 1.12)	1.04 (0.97, 1.12)	1.04 (0.97, 1.12)
	P _{trend}		0.25		0.11	0.23	0.30	
Metribuzin ^d	0	61	1.00 (ref)	353	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	33–221	5	0.48 (0.17, 1.35)	48	0.89 (0.73, 1.08)	1.06 (0.98, 1.15)	1.06 (0.97, 1.15)	1.06 (0.97, 1.15)
	>221–385	9	0.94 (0.39, 2.30)	50	0.88 (0.72, 1.08)	0.97 (0.90, 1.06)	0.93 (0.85, 1.02)	0.93 (0.85, 1.02)
	>385–1000	12	1.56 (0.67, 3.67)	43	1.25 (1.01, 1.55) [†]	0.94 (0.86, 1.03)	1.01 (0.92, 1.11)	1.01 (0.92, 1.11)
	>1000–14850	9	0.73 (0.27, 1.95)	49	0.89 (0.71, 1.11)	0.96 (0.88, 1.06)	0.98 (0.88, 1.08)	0.98 (0.88, 1.08)
	P _{trend}		0.75		0.62	0.28	0.64	
Chloracetamide								
Alachlor	0	36	1.00 (ref)	218	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	25–502	29	2.34 (1.25, 4.37) [†]	107	1.03 (0.89, 1.20)	1.00 (0.94, 1.06)	0.98 (0.92, 1.05)	0.98 (0.92, 1.05)
	>502–1358	12	0.69 (0.33, 1.44)	106	0.91 (0.78, 1.05)	1.02 (0.96, 1.09)	0.98 (0.92, 1.04)	0.98 (0.92, 1.04)
	>1358–4463	21	0.98 (0.50, 1.92)	107	1.06 (0.91, 1.24)	0.97 (0.91, 1.03)	0.98 (0.92, 1.05)	0.98 (0.92, 1.05)
	>4463–68162	21	0.99 (0.48, 2.01)	106	0.99 (0.84, 1.17)	1.04 (0.97, 1.11)	0.99 (0.93, 1.07)	0.99 (0.93, 1.07)
	P _{trend}		0.66		0.98	0.28	0.95	
Metolachlor	0	54	1.00 (ref)	317	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	25–455	19	1.47 (0.79, 2.71)	84	1.02 (0.88, 1.18)	0.98 (0.92, 1.05)	0.97 (0.91, 1.03)	0.97 (0.91, 1.03)
	>455–1240	16	1.23 (0.64, 2.37)	81	1.08 (0.93, 1.26)	1.04 (0.98, 1.11)	0.97 (0.91, 1.03)	0.97 (0.91, 1.03)
	>1240–4232	10	0.65 (0.30, 1.39)	83	0.92 (0.79, 1.08)	0.98 (0.92, 1.04)	0.98 (0.92, 1.05)	0.98 (0.92, 1.05)
	>4232–68162	21	1.44 (0.72, 2.88)	83	1.18 (0.99, 1.40)	0.97 (0.91, 1.05)	0.98 (0.91, 1.05)	0.98 (0.91, 1.05)
	P _{trend}		0.41		0.07	0.43	0.74	
Other Herbicides								
Chlorimuron-ethyl ^d	0	74	1.00 (ref)	435	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
	30–118	6	1.13 (0.44, 2.89)	32	0.88 (0.70, 1.10)	0.99 (0.90, 1.08)	0.96 (0.87, 1.06)	0.96 (0.87, 1.06)
	>118–315	8	1.62 (0.69, 3.84)	30	0.94 (0.75, 1.18)	1.01 (0.92, 1.11)	1.03 (0.93, 1.13)	1.03 (0.93, 1.13)
	>315–588	7	1.35 (0.56, 3.26)	33	1.03 (0.83, 1.28)	1.02 (0.93, 1.12)	0.99 (0.90, 1.09)	0.99 (0.90, 1.09)
	>588–7560	7	1.45 (0.60, 3.50)	32	1.05 (0.84, 1.31)	1.04 (0.95, 1.14)	0.99 (0.90, 1.09)	0.99 (0.90, 1.09)

	TSH and Thyroid Hormones													
	Subclinical Hypothyroidism					TSH					T4		T3	
	Intensity-weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	
	P _{trend}		0.3		0.66		0.35		0.81					
Dicamba	0	42	1.00 (ref)	220	1.00 (ref)		1.00 (ref)		1.00 (ref)					
	36-350	20	0.93 (0.50, 1.73)	109	0.86 (0.75, 1.00) [‡]		0.98 (0.92, 1.04)		0.95 (0.90, 1.02)					
	>350-1046	15	0.68 (0.35, 1.33)	103	0.94 (0.81, 1.09)		0.96 (0.91, 1.02)		0.99 (0.92, 1.05)					
	>1046-2699	19	0.91 (0.48, 1.72)	106	0.97 (0.84, 1.13)		1.00 (0.94, 1.06)		1.01 (0.95, 1.08)					
	>2699-107823	21	1.06 (0.57, 1.98)	106	0.98 (0.84, 1.14)		0.98 (0.92, 1.04)		0.97 (0.91, 1.03)					
	P _{trend}		0.62		0.59		0.86		0.59					
Glyphosate	0	32	1.00 (ref)	179	1.00 (ref)		1.00 (ref)		1.00 (ref)					
	20-315	25	1.28 (0.71, 2.32)	116	1.02 (0.88, 1.17)		1.01 (0.95, 1.07)		0.98 (0.92, 1.04)					
	>315-907	24	1.07 (0.60, 1.93)	130	1.00 (0.87, 1.15)		1.01 (0.95, 1.07)		1.01 (0.95, 1.07)					
	>907-2622	21	0.95 (0.51, 1.77)	123	1.01 (0.88, 1.17)		1.01 (0.96, 1.07)		0.99 (0.93, 1.05)					
	>2622-113400	25	1.21 (0.66, 2.24)	123	1.14 (0.99, 1.33)		0.99 (0.94, 1.05)		0.97 (0.91, 1.03)					
	P _{trend}		0.70		0.05		0.68		0.41					
Imazethapyr	0	62		360										
	33-263	12	1.08 (0.53, 2.17)	70	0.90 (0.77, 1.06)		0.97 (0.91, 1.04)		0.97 (0.91, 1.04)					
	>263-600	14	1.22 (0.63, 2.37)	73	1.12 (0.96, 1.31)		0.96 (0.90, 1.02)		0.95 (0.89, 1.02)					
	>600-1176	14	1.20 (0.61, 2.33)	73	0.97 (0.83, 1.14)		0.94 (0.88, 1.00) [‡]		0.96 (0.90, 1.03)					
	>1176-22500	16	1.54 (0.81, 2.95)	71	1.15 (0.98, 1.35)		0.97 (0.91, 1.03)		0.98 (0.92, 1.05)					
	P _{trend}		0.19		0.09		0.24		0.56					
Petroleum distillates ^d	0	74	1.00 (ref)	413	1.00 (ref)		1.00 (ref)		1.00 (ref)					
	25-372	6	1.27 (0.49, 3.30)	27	0.93 (0.73, 1.18)		0.98 (0.88, 1.08)		0.99 (0.90, 1.10)					
	>372-1110	3	0.56 (0.16, 1.92)	27	1.16 (0.91, 1.47)		0.99 (0.89, 1.09)		1.00 (0.90, 1.11)					
	>1110-3973	7	1.54 (0.62, 3.84)	27	1.07 (0.84, 1.36)		1.01 (0.92, 1.12)		1.03 (0.93, 1.14)					
	>3973-253333	5	0.97 (0.35, 2.69)	27	1.22 (0.96, 1.56)		1.00 (0.91, 1.11)		0.89 (0.80, 0.99)					
	P _{trend}		0.95		0.09		0.93		0.045 [‡]					

TSH: Thyroid stimulating hormone; T4: thyroxine; T3: triiodothyronine

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

β Adjusted for age, state, body mass index, smoking, correlated pesticides

q Exposed cases

c Total exposed

d Detailed information for these chemicals was collected on the take-home questionnaire at enrollment

\hat{d} $p < 0.05$ compared with never users

Table 4
Multivariate^a regression models examining association between intensity-weighted days of specific fumigants/fungicides and subclinical hypothyroidism (TSH>4.5 mIU/L) and natural log of TSH, T4, and T3

	TSH and Thyroid Hormones									
	Subclinical hypothyroidism			TSH			T4			T3
	Intensity-weighted days	N ^b	OR (95% CI)	N ^c	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)	Expβ (95% CI)
Fumigant										
Carbon tetrachloride/	0	113	1.00 (ref)	617	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)	1.00 (ref)
Carbon disulfide ^d	6-170	4	1.70 (0.51, 5.61)	14	1.06 (0.76, 1.47)	0.97 (0.85, 1.10)	0.95 (0.82, 1.09)			
	>170-18563	3	1.08 (0.29, 4.08)	14	1.01 (0.73, 1.41)	0.94 (0.82, 1.07)	0.89 (0.77, 1.02)			
	P _{trend}		0.87		0.93	0.33	0.09			
Methyl bromide										
	0	117	1.00 (ref)	622	1.00 (ref)		1.00 (ref)			
	12-1139	6	1.32 (0.44, 4.02)	24	1.31 (0.98, 1.74)	0.89 (0.80, 1.00)	0.94 (0.83, 1.06)			
	>1139-102000	3	0.45 (0.11, 1.81)	26	0.75 (0.56, 1.00) [†]	1.14 (1.01, 1.28) [†]	1.11 (0.98, 1.26)			
	P _{trend}		0.23		0.02 [†]	0.01 [†]	0.06			
Fungicide										
Captan										
	0	103	1.00 (ref)	566	1.00 (ref)	1.00 (ref)	1.00 (ref)			
	2-12	3	0.40 (0.12, 1.35)	36	0.87 (0.71, 1.07)	1.07 (0.98, 1.17)	1.07 (0.98, 1.16)			
	>12-46500	10	1.89 (0.87, 4.11)	34	1.23 (1.00, 1.52) [†]	0.99 (0.91, 1.09)	0.99 (0.90, 1.08)			
	P _{trend}		0.09		0.05 [†]	0.87	0.73			
Metaxyl^d										
	0	105	1.00 (ref)	560	1.00 (ref)	1.00 (ref)	1.00 (ref)			
	3-141	6	1.06 (0.41, 2.72)	28	1.06 (0.84, 1.34)	1.02 (0.93, 1.12)	1.03 (0.93, 1.14)			
	>141-30341	4	0.60 (0.19, 1.92)	28	0.87 (0.68, 1.12)	1.02 (0.92, 1.13)	0.98 (0.88, 1.09)			
	P _{trend}		0.39		0.27	0.72	0.67			

TSH: Thyroid stimulating hormone; T4: thyroxine; T3: triiodothyronine

^a Adjusted for age, state, body mass index, smoking, correlated pesticides

^b Exposed cases

^c Total exposed

Detailed information for these chemicals was collected on the take-home questionnaire at enrollment
 $p < 0.05$ compared with never users

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript