



HHS Public Access

Author manuscript

Eur J Oncol Nurs. Author manuscript; available in PMC 2019 February 01.

Published in final edited form as:

Eur J Oncol Nurs. 2018 February ; 32: 63–72. doi:10.1016/j.ejon.2017.12.003.

Differences in Symptom Clusters Before and Twelve Months After Breast Cancer Surgery

Melissa Mazor, RN, PhD,

School of Nursing, University of California at San Francisco, San Francisco, California

Janine K. Cataldo, RN, PhD,

School of Nursing, University of California at San Francisco, San Francisco, California

Kathryn Lee, RN, PhD,

School of Nursing, University of California at San Francisco, San Francisco, California

Anand Dhruva, MD,

School of Medicine, University of California at San Francisco, San Francisco, California

Bruce Cooper, PhD,

School of Nursing, University of California at San Francisco, San Francisco, California

Steven M. Paul, PhD,

School of Nursing, University of California at San Francisco, San Francisco, California

Kimberly Topp, PT, PhD, FAAA,

School of Medicine, University of California at San Francisco, San Francisco, California

Betty J. Smoot, PT, DPTSc, MAS,

School of Medicine, University of California at San Francisco, San Francisco, California

Laura B. Dunn, MD,

School of Medicine, Stanford University, Stanford, California

Jon D. Levine, MD, PhD,

School of Medicine, University of California at San Francisco, San Francisco, California

Yvette P. Conley, PhD, and

School of Nursing, University of Pittsburgh, Pittsburgh, Pennsylvania

Christine Miaskowski, RN, PhD

School of Nursing, University of California at San Francisco, San Francisco, California

Abstract

Address correspondence to: Christine Miaskowski, RN, PhD, FAAN, Professor, Department of Physiological Nursing, University of California, 2 Koret Way – N631Y, San Francisco, CA 94143-0610, 415-476-9407 (phone), 415-476-8899 (fax), chris.miaskowski@ucsf.edu.

Conflict of interest: The authors have no conflicts of interest to declare.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Purpose—Given the inter-relatedness among symptoms, research efforts are focused on an evaluation of symptom clusters. The purposes of this study were to evaluate for differences in the number and types of menopausal-related symptom clusters assessed prior to and at 12-months after surgery using ratings of occurrence and severity and to evaluate for changes in these symptom clusters over time.

Methods—Prior to and at 12 months after surgery, 392 women with breast cancer completed the Menopausal Symptoms Scale. Exploratory factor analyses were used to identify the symptom clusters.

Results—Of the 392 women evaluated, the mean number of symptoms (out of 46) was 13.2 (± 8.5) at enrollment and 10.9 (± 8.2) at 12 months after surgery. Using occurrence and severity, three symptom clusters were identified prior to surgery. Five symptom clusters were identified at 12 months following surgery. Two symptom clusters (i.e., pain/discomfort and hormonal) were relatively stable across both dimensions and time points. Two symptom clusters were relatively stable across both dimensions either prior to surgery (i.e., sleep/psychological/cognitive) or at 12 months after surgery (i.e., sleep). The other four clusters (i.e., irritability, psychological/cognitive, cognitive, psychological) were identified at one time point using a single dimension.

Conclusions—While some menopausal-related symptom clusters were consistent across time and dimensions, the majority of symptoms clustered together differently depending on whether they were evaluated prior to or at 12 months after breast cancer surgery. An increased understanding of how symptom clusters change over time may assist clinicians to focus their symptom assessments and management strategies.

Keywords

symptoms; symptom clusters; menopause; breast cancer; factor analysis

Introduction

Prior to and following breast cancer treatment women experience multiple co-occurring menopausal-related symptoms (Barton and Ganz, 2015; Howard-Anderson et al., 2012). Most of this research has focused on descriptions of single menopausal-related symptoms (e.g., hot flashes) during or after chemotherapy (CTX) and/or endocrine therapy (ET) in breast cancer survivors. Given the inter-relatedness among symptoms, current research efforts are focused on an evaluation of symptom clusters (Glaus et al., 2006; Marshall et al., 2016; Seib et al., 2017).

A symptom cluster is defined as a group of two (Kim et al., 2005) or more (Dodd et al., 2001) concurrent symptoms that are related to one another through a common etiology, mechanism, variance, or outcome (Barsevick, 2016; Miaskowski et al., 2007; Miaskowski et al., 2017). The identification of differences in the number and types of menopausal-related symptom clusters before and after breast cancer treatment may assist clinicians to focus both their assessments and management strategies. For example, rather than treating a single symptom, clinicians may be able to target several symptoms within a cluster (Kwekkeboom et al., 2012) and minimize the need for women to take multiple medications. For example, in a recent study (Lengacher et al., 2017), a mindfulness-based stress reduction intervention

improved the severity of several symptoms within a psychological/cognitive symptom cluster. Given that relative to single symptoms, symptoms clusters are associated with poorer functional status and quality of life (QOL) (Kim et al., 2012), management of several symptoms within a cluster may improve patient outcomes. Moreover, the identification of menopausal-related symptom clusters may suggest a common etiology for symptoms within a cluster.

In women with breast cancer, the majority of the research on symptom clusters has evaluated common symptoms associated with cancer treatment using instruments like the Memorial Symptom Assessment Scale (MSAS) (Portenoy et al., 1994) or the MD Anderson Symptom Inventory (MDASI) (Cleeland et al., 2000) (for review see (Nguyen et al., 2011)). While both the MSAS and MDASI include some symptoms that would be found on a menopausal symptom inventory like the Menopausal Symptom Scale (MSS) (Radtke et al., 2011), neither instrument assess for hot flashes, the most commonly reported menopausal-related symptom in women with (Gupta et al., 2006) and without (Woods and Mitchell, 2005) breast cancer.

In studies of individual symptoms, breast cancer patients reported bothersome symptoms prior to and following surgery and that these symptoms had a negative impact on their QOL (Denieffe et al., 2014). Yet, with the major focus on the identification of symptom clusters during treatment, no studies were identified that evaluated for menopausal-related symptom clusters prior to breast cancer surgery. Only three cross-sectional studies (Glaus et al., 2006; Marshall et al., 2016; Seib et al., 2017) evaluated for menopausal-related symptom clusters following breast cancer treatment (Supplementary Table 1). In the first study that evaluated breast cancer patients on ET, a single symptom cluster was found using symptom occurrence rates from the Clinical Checklist for Patients with Endocrine Therapy (C-PET) (Glaus et al., 2006). This single 'menopausal' cluster included: hot flashes, weight-gain, tiredness, reduced sexual interest, and vaginal dryness.

In the second study that evaluated breast cancer survivors 8 months after their cancer diagnosis (Marshall et al., 2016), menopausal-related symptom clusters were derived from the Women's Health Initiative Checklist. Five clusters were identified using dichotomous ratings of severity. These five clusters varied slightly depending on whether moderate and severe (i.e., menopausal, pain, fatigue/sleep/gastrointestinal (GI), psychological, increased weight/appetite) or severe (i.e., menopausal, pain, fatigue/psychological/GI, GI, increased weight/appetite) symptoms were evaluated. In addition, menopausal-related symptom clusters were evaluated using symptom data derived from messages on a breast cancer forum. The four clusters identified from the social media data were: pain/fatigue, menopausal/psychological, GI, and miscellaneous.

The third study compared menopausal-related symptom clusters in women with and without breast cancer (Seib et al., 2017). The symptom clusters were using the severity scores from the Greene Climacteric Scale. In the women with breast cancer, the following symptom clusters were identified: psychological, vasoactive, sensory somatic, peripheral somatic, nervous tension, and general somatic. With the exception of the general somatic symptom cluster, the same clusters were identified in women without breast cancer. However, the

specific symptoms within each of the five clusters varied between these two groups of women. Across these three studies of breast cancer survivors (Glaus et al., 2006; Marshall et al., 2016; Seib et al., 2017), the menopausal-related cluster was the only consistent symptom cluster identified in patients with breast cancer. Within this cluster, hot flashes was the only consistent symptom.

While these three studies provide preliminary evidence of menopausal-related symptom clusters in breast cancer survivors (Glaus et al., 2006; Marshall et al., 2016; Seib et al., 2017), several limitations warrant consideration. The instruments and dimensions used to evaluate for symptom clusters were not consistent. In addition, all three studies evaluated for symptom clusters using only a single dimension of the symptom experience (i.e., occurrence (Glaus et al., 2006) or severity (Marshall et al., 2016; Seib et al., 2017)). Finally, time since cancer diagnosis (Glaus et al., 2006; Marshall et al., 2016; Seib et al., 2017), demographic and clinical characteristics (Marshall et al., 2016), and specific cancer treatments (Seib et al., 2017) were not reported. These limitations make it difficult to compare findings across these three studies.

Given these limitations and the paucity of research on menopausal-related symptom clusters in women prior to and following breast cancer surgery, the purposes of this study, in a sample of breast cancer patients, were to: evaluate for differences in the number and types of menopausal-related symptom clusters assessed prior to and at 12-months after surgery using ratings of occurrence and severity and to evaluate for changes in these symptom clusters over time. Given that the occurrence and severity of menopausal-related symptoms vary over the course of treatment (Ganz et al., 2011), we hypothesized that the number and types of symptom clusters would differ over time but not by dimension.

Methods

This study is part of a larger descriptive, longitudinal study that evaluated neuropathic pain and lymphedema in women who underwent breast cancer surgery. The methods for this study are described in detail elsewhere (Doong et al., 2015; Kyranou et al., 2013; Langford et al., 2014; McCann et al., 2012; Van Onselen et al., 2013). In brief, patients were recruited from Breast Cancer Centers located in a Comprehensive Cancer Center, two public hospitals, and four community practices. Eligibility criteria included: adult women (18 years) who were scheduled for unilateral breast cancer surgery; were able to read, write, and understand English; agreed to participate; and provided written informed consent. Patients were excluded if they had bilateral breast surgery and/or had distant metastases at the time of diagnosis.

Instruments

The demographic questionnaire obtained information on age, education, ethnicity, marital status, employment status, living situation, and financial status. Menopausal status was determined by the patient's response (yes/no) at the time of enrollment to the question "Have you gone through menopause yet (stopped having your menstrual cycle)?" Patients were asked to indicate if they exercised on a regular basis (yes/no). The Karnofsky Performance Status (KPS) scale was used to evaluate functional status. Patients rated their functional

status using the KPS scale that ranged from 30 (I feel severely disabled and need to be hospitalized) to 100 (I feel normal; I have no complaints or symptoms). The KPS scale has well established validity and reliability (Karnofsky et al., 1948).

The Self-Administered Comorbidity Questionnaire (SCQ) is a short and easily understood instrument that was developed to measure comorbidity in clinical and health service research settings (Sangha et al., 2003). The questionnaire consists of 13 common medical conditions that were simplified into language that could be understood without any prior medical knowledge. Patients indicated if they had the condition; if they received treatment for it (proxy for disease severity); and if it limited their activities (indication of functional limitations). SCQ scores can range from 0 to 39. The SCQ has well established validity and reliability and has been used in studies of patients with a variety of chronic conditions (Brunner et al., 2008; Cieza et al., 2006).

The Menopausal Symptoms Scale (MSS) was modified from the Seattle Women's Health Study questionnaire (Woods et al., 1999). The MSS evaluated the occurrence, severity, and distress of 46 menopausal-related symptoms. Patients were asked to indicate whether they experienced each symptom during the past week (i.e., symptom occurrence). If they experienced the symptom, they were asked to rate its severity and distress. Symptom severity was rated using a 0 (none) to 10 (intolerable) numeric rating scale (NRS). The MSS has well established validity and reliability (Woods et al., 2014).

Study Procedures

The study was approved by the Committee on Human Research at the University of California, San Francisco and by the Institutional Review Boards at each of the study sites. A clinician explained the study and determined patient's willingness to participate during her scheduled preoperative visit. After the visit, the clinician introduced the patient to the research nurse who met with the woman, determined eligibility, and obtained written informed consent. Then, patients completed the enrollment questionnaires an average of four days prior to surgery. For the current study, data from prior to and 12 months after surgery were analyzed. Medical records were reviewed for disease and treatment information.

Statistical Analysis

Data were analyzed using the Statistical Package for the Social Sciences 23 (SPSS, 2015) and MPlus Version 7.3 (Muthén, 1989; Muthén and Muthén, 1998-2015). Descriptive statistics and frequency distributions were calculated for the demographic and clinical characteristics. As part of the evaluation of symptom occurrence and severity, occurrence rates were generated for each symptom and mean severity scores were calculated for patients who reported the symptom (without zeros) and for all of the patients (with zeros included).

Creation of symptom clusters using exploratory factor analysis (EFA)—Four separate EFAs were done to evaluate for symptom clusters using the dichotomous ratings of symptom occurrence and ordinal ratings of symptom severity obtained prior to and at 12 months after breast cancer surgery. Factor analysis aims to identify whether correlations between a set of observed variables can be explained by a few latent, unobserved variables

(i.e., factors) (Brown, 2015). The “factors” in the current study are referred to as symptom clusters (Kim et al., 2009; Miaskowski et al., 2004).

For each EFA, factor loadings were considered meaningful if the loading had an absolute value of ≥ 0.40 (Browne, 2001; Muthén, 1989; Muthén and Muthén, 1998-2015). While it is common to require that each item load strongly on only one factor, in this study, items that loaded with an absolute value of ≥ 0.40 on two factors (i.e., cross loaded), were retained and used to define both factors. The cross loading of symptoms on more than one factor may be beneficial in the interpretation of potential causal mechanisms, especially when oblique rotation is employed (Brown, 2015; Browne, 2001; Miaskowski and Aouizerat, 2007).

EFA was used to identify symptom clusters from the occurrence rates and the severity ratings of 24 out of the 46 MSS symptoms. In order to have sufficient variation and covariation to perform the EFAs, only symptoms that were present in $>18\%$ and $<80\%$ of the patients at both time points were included in these analyses. The Cronbach's alphas ranged from good to excellent with all 46 symptoms after and before surgery (i.e., $\alpha=.909$ and $.920$) and with 24 symptoms before and after surgery (i.e., $\alpha=.896$ and $.880$). Twenty-two symptoms on the MSS were excluded from the analyses due to insufficient variation in the occurrence of these symptoms. These excluded symptoms were tearful/crying spells, nervousness, panic feelings, lost sexual interest, constipation, urinary frequency, loss of interest in things, loss of appetite, heart races/pounds, abdominal bloating, diarrhea, nausea/upset stomach, swollen hands/feet, eating more than usual, indigestion, shortness of breath, skin breakout/acne, hostility, cramps, dizziness, alcohol cravings, increased sexual desire.

The occurrence items were evaluated as dichotomous variables (i.e., had versus did not have the symptom). For these EFAs, tetrachoric correlations were used to create the matrix of associations. In order to provide better estimation of the results, the 0 to 10 severity scores were recoded into: 0 (none), 1 (mild = 1 to 3), 2 (moderate = 4 to 6), and 3 (severe > 7). This approach was used because the distribution of the scores on the 0 to 10 continuous scale were positively skewed, violating the bivariate normality assumption. Reducing the number of ordinal points through recoding better captured the meaning of the scale and allowed for better estimation of the covariance structure for the EFAs. For the rescored severity EFAs, polychoric correlations were used to create the matrix of associations (Muthén, 1989; Muthén and Muthén, 1998-2015).

The simple structures for the occurrence and severity EFAs were estimated using the method of unweighted least squares with geomin (i.e., oblique) rotation. The geomin rotation method was used to identify the model with the best fit (i.e., optimum number of factors using the criteria for simple structure described above). Adopting this rotational method provided an improved representation of how the factors were correlated and improved the interpretability of each factor solution (Muthén, 1989; Muthén and Muthén, 1998-2015). The unweighted least squares estimator (ulsmv: unweighted least squares parameter estimates with standard errors and a mean and variance adjusted (chi-square test using a full weight matrix (Muthén, 1989; Muthén and Muthén, 1998-2015)) was selected in order to achieve more reliable results because the scales for the MSS items were dichotomous (i.e., occurrence) and ordinal (i.e., severity).

Factor solutions were estimated for two through six factors. After examining all of the factor solutions, the factor solution with the greatest interpretability and clinical meaningfulness was selected, given that it met the criteria set for evaluating simple structure (i.e., size of item loadings, number of items on a factor).

Differences in number and types of symptom clusters—To evaluate the agreement among the symptoms within the same cluster using occurrence and severity ratings, within and across each assessment, we used the criteria proposed by Kirkova and Walsh (Kirkova and Walsh, 2007). In their paper, they suggested that to be in agreement with each other, at least 75% of the symptoms in the clusters should be present including the prominent and important symptom, namely the symptom with the greatest weight from the EFAs.

Results

Demographic and Clinical Characteristics

The demographic and clinical characteristics of the patients are summarized in Table 1. The mean age of the women was 54.9 ± 11.6 years (range: 29 to 91 years), 41.9% were married/partnered, 24.2% lived alone, and 64.1% were postmenopausal prior to surgery. The majority of the patients was White (64.4%) and well educated (15.7 ± 2.7 years).

At enrollment, women were 9.5 ± 11.0 weeks from their cancer diagnosis. They had a mean SCQ score of 4.3 ± 2.8 , with high blood pressure (30.9%), back pain (28.1%), and depression (21.9%) being the most common comorbidities. The mean KPS score was 93.2 ± 10.3 . Approximately 19.9% of the patients had received neoadjuvant CTX. The majority of patients had breast conservation surgery (79.9%), 82.4% had a sentinel lymph node biopsy, and 21.7% had breast reconstruction at time of surgery. During the first 12 months after surgery, 33.5% received adjuvant CTX, 72.5% received external beam radiation therapy (RT), and 63.2% were taking ET.

Symptom Occurrence and Severity

The mean number of symptoms (out of 46) was 13.2 ± 8.5 at enrollment and 10.9 ± 8.2 at 12 months after surgery (Table 1). The occurrence and severity scores for the 46 symptoms are summarized in Supplementary Table 2. The severity scores are reported as: ordinal with zero including women who did not report the occurrence of the symptom; ordinal including only women who reported the occurrence of the symptom; and continuous (i.e., 0 to 10 NRS) including only women who reported the occurrence of the symptom. Across the two assessments, the most common and the most severe symptom was wake during the night and lost sexual interest, respectively.

Symptom Clusters Based on Occurrence Rates

As shown in Table 2, prior to surgery, a four factor solution indicated the best fit for the occurrence data. Factor 1 that included five symptoms (i.e., anger, impatience, irritability, mood swings, tension) was named the irritability symptom cluster. Factor 2 that consisted of six symptoms (i.e., backache or neckache, general body aches, joint pain or stiffness, numbness or tingling, painful/tender breasts, weight gain) was named the pain/discomfort

symptom cluster. Factor 3 that included eight symptoms (i.e., tension, anxiety, depression, difficulty concentrating, difficulty falling asleep, fatigue or tiredness, wake during the night, waking too early) was named the psychological/cognitive/sleep symptom cluster. Factor 4 that consisted of four symptoms (i.e., daytime sweats, hot flashes, night sweats, vaginal dryness) was named the hormonal symptom cluster. Two symptoms (i.e., forgetfulness and headache) did not load on any factor.

As shown in Table 3, at 12 months after surgery, a four factor solution was the best fit for the occurrence data. Factor 1 that included ten symptoms (i.e., anger, anxiety, depression, difficulty concentrating, fatigue or tiredness, forgetfulness, impatience, irritability, mood swings, tension) was named the psychological/cognitive symptom cluster. Factor 2 that included three symptoms (i.e., daytime sweats, hot flashes, night sweats) was named the hormonal symptom cluster. Factor 3 that included three symptoms (i.e., backache or neckache, general body aches, joint pain or stiffness) was named the pain/discomfort symptom cluster. Factor 4 that included three symptoms (i.e., difficulty falling asleep, wake during the night, waking too early) was named the sleep symptom cluster. Five symptoms (i.e., headache, numbness or tingling, painful/tender breasts, vaginal dryness, weight gain) did not load on any factor.

Symptom Clusters Based on Severity Ratings

As shown in Table 4, prior to surgery, a three factor solution was the best fit for the severity data. Factor 1 that included fourteen symptoms (i.e., anger, anxiety, depression, difficulty concentrating, difficulty falling asleep, fatigue or tiredness, forgetfulness, headache, impatience, irritability, mood swings, tension, wake during the night, waking too early) was named the psychological/cognitive/sleep symptom cluster. Factor 2 that included seven symptoms (i.e., daytime sweats, general body aches, hot flashes, night sweats, numbness or tingling, vaginal dryness, weight gain) was named the hormonal symptom cluster. Factor 3 that consisted of four symptoms (i.e., general body aches, numbness or tingling, backache or neckache, joint pain or stiffness) was named the pain/discomfort symptom cluster. Only painful/tender breasts did not load on any factor.

As shown in Table 5, at 12 months after surgery, a five factor solution was the best fit for the severity data. Factor 1 that included seven symptoms (i.e., anger, anxiety, depression, impatience, irritability, mood swings, tension) was named the psychological cluster. Factor 2 that included four symptoms (difficulty concentrating, fatigue or tiredness, forgetfulness, painful/tender breasts) was named the cognitive symptom cluster. Factor 3 that included three symptoms (i.e., daytime sweats, hot flashes, night sweats) was named the hormonal symptom cluster. Factor 4 that included four symptoms (i.e., backache or neckache, general body aches, headache, joint pain or stiffness) was named the pain/discomfort symptom cluster. Factor 5 that included three symptoms (i.e., difficulty falling asleep, wake during the night, wake too early) was named the sleep symptom cluster. Three symptoms (i.e., numbness/tingling, vaginal dryness, and weight gain) did not load on any factor.

Agreement in the Number and Types of Symptom Clusters

Table 6 presents a summary of the factor loadings and percentage agreement among the symptoms within each cluster that were identified regardless of dimension (i.e., occurrence and severity) and time (i.e., prior to surgery, 12 months after surgery). For the pain/discomfort symptom cluster, that was identified for both dimensions and at both time points, the total number of symptoms ranged from three to six and the percent agreement ranged from 42.9% to 85.7%. Across both dimensions and assessments, the three symptoms that were included in the pain/discomfort cluster were: backache or neckache, general body aches, and joint pain or stiffness.

For the hormonal symptom cluster, that was identified for both dimensions and at both time points, the total number of symptoms ranged from 3 to 7 and the percent agreement ranged from 42.9% to 100.0%. Across both dimensions and assessments, the three symptoms that were included in the hormonal cluster were: daytime sweats, hot flashes, and night sweats.

Differences in the Number and Types of Symptom Clusters

Table 7 presents a summary of the factor loadings and symptoms within each cluster that were different across time (i.e., prior to surgery, 12 months after surgery). For the psychological/cognitive/sleep symptom cluster that was identified prior to surgery using both dimensions, the total number of symptoms ranged from 8 to 14. The eight symptoms included in this cluster were: tension, anxiety, depression, difficulty concentrating, difficulty falling asleep, fatigue or tiredness, wake during the night, and waking too early.

The irritability cluster was only identified for the occurrence dimension assessed prior to surgery. The five symptoms included in this cluster were: anger, impatience, irritability, mood swings, and tension.

The sleep symptom cluster was identified using both occurrence and severity dimensions assessed at 12 months after surgery. The three symptoms in this cluster were: difficulty falling asleep, wake during the night, and waking too early.

The psychological/cognitive symptom cluster was identified only for the occurrence dimension assessed at 12 months after surgery. The ten symptoms included in this cluster were: anger, anxiety, depression, difficulty concentrating, fatigue or tiredness, forgetfulness, impatience, irritability, mood swings, and tension.

The psychological symptom cluster was identified only for the severity dimension assessed at 12 months after surgery. The seven symptoms included in this cluster were: anger, anxiety, depression, impatience, irritability, mood swings, and tension.

The cognitive symptom cluster was identified only for the severity dimension assessed at 12 months after surgery. The four symptoms included in this cluster were: difficulty concentrating, fatigue or tiredness, forgetfulness, and painful/tender breasts.

Discussion

This study is the first to assess for differences in the number and types of menopausal-related symptom clusters in women before and at 12 months after breast cancer surgery using occurrence rates and severity ratings. Our hypothesis, regarding the consistency in number and types of symptom clusters across dimensions and time points, was only partially supported. While eight distinct symptom clusters were identified across the two dimensions and time points, only two (i.e., pain/discomfort and hormonal) were relatively stable across both dimensions and time points. In addition, two symptom clusters were relatively stable across both dimensions, either prior to surgery (i.e., sleep/psychological/cognitive) or at 12 months after surgery (i.e., sleep). The other four clusters (i.e., irritability, psychological/cognitive, cognitive, psychological) were only identified at one time point using a single dimension. In the remainder of the discussion, the two symptom clusters that were the same across all four EFAs will be discussed first followed by a discussion of the six distinct symptom clusters.

Consistent Symptom Clusters

Pain/Discomfort Symptom Cluster—The pain/discomfort cluster was identified using both dimensions and at both assessment times. While the number of symptoms in this cluster ranged from three to six, backache/neckache, general body ache, and joint pain or stiffness were the common symptoms across all four EFAs. This finding is consistent with previous studies of symptom clusters in healthy women (Cray et al., 2013) as well as in women during (Phligbua W, 2013; Suwith, 2008) and after (Fu et al., 2009; Marshall et al., 2016; Roiland and Heidrich, 2011) breast cancer treatment.

While this ‘pain-like’ cluster had various names (i.e., pain (Cray et al., 2013; Fu et al., 2009; Marshall et al., 2016), pain/discomfort (Suwith, 2008), discomfort (Phligbua W, 2013), musculoskeletal (Roiland and Heidrich, 2011)), backache, joint pain, and headache were the consistent symptoms regardless of whether healthy women (Cray et al., 2013) or women after breast cancer treatment were assessed and regardless of the symptom dimension evaluated (i.e., occurrence (Marshall et al., 2016), severity (Fu et al., 2009), or both (Roiland and Heidrich, 2011)). In contrast, in studies of patients with breast cancer undergoing CTX or RT, while joint pain (Phligbua W, 2013) and numbness and tingling (Phligbua W, 2013; Suwith, 2008) were found in the pain cluster, other symptoms in this cluster included: dry mouth, feeling irritable, dizziness, difficulty concentrating, vaginal dryness, worrying, skin changes, and lack of energy. These inconsistent findings may be related to the symptom assessment instrument used and the timing of the assessments.

Despite these inconsistencies, a growing body of evidence suggests that a pain cluster (i.e., joint pain, back pain, headache) occurs prior to breast cancer surgery and appears to persist well into survivorship. Of note, in our study, all three symptoms occurred in over 20% of the women and were in the moderate to severe range at both assessments. Given the interrelationships among symptoms and the high occurrence of joint pain with aromatase inhibitors (Glaus et al., 2006), future studies need to evaluate the underlying mechanisms for the symptoms within this cluster. Clinicians need to assess for the specific types and causes of pain throughout the course of breast cancer treatment.

Hormonal Symptom Cluster—A hormonal symptom cluster was identified across all four EFAs. The total number of symptoms in this cluster ranged from three to seven. Regardless of dimension or time, daytime sweats, hot flashes, and night sweats were the three vasomotor symptoms included in this cluster. While the names for this cluster varied (i.e., menopausal (Glaus et al., 2006; Huang et al., 2016; Marshall et al., 2016), hormonal (Fu et al., 2009; Roiland and Heidrich, 2011; Yates et al., 2015), vasoactive (Seib et al., 2017), and vasomotor (Cray et al., 2013; Mitchell and Woods, 1996)), similar clusters were identified in healthy women (Avis et al., 2001; Cray et al., 2013) and in women with breast (Cray et al., 2013; Fu et al., 2009; Glaus et al., 2006; Marshall et al., 2016; Mitchell and Woods, 1996; Roiland and Heidrich, 2011; Seib et al., 2017), ovarian (Huang et al., 2016), or heterogeneous (Yates et al., 2015) cancer diagnoses.

Across these studies, the consistent symptoms included hot flashes (Avis et al., 2001; Cray et al., 2013; Fu et al., 2009; Glaus et al., 2006; Marshall et al., 2016; Mitchell and Woods, 1996; Phligbua W, 2013; Roiland and Heidrich, 2011; Seib et al., 2017), night sweats (Avis et al., 2001; Marshall et al., 2016; Mitchell and Woods, 1996; Phligbua W, 2013; Seib et al., 2017), vaginal dryness (Avis et al., 2001; Glaus et al., 2006; Marshall et al., 2016; Roiland and Heidrich, 2011), lost sexual interest (Fu et al., 2009; Glaus et al., 2006; Seib et al., 2017), and weight gain (Glaus et al., 2006; Marshall et al., 2016; Roiland and Heidrich, 2011). While not found in our hormonal cluster, psychological symptoms (i.e., mood swings (Marshall et al., 2016; Phligbua W, 2013; Roiland and Heidrich, 2011), depression (Marshall et al., 2016; Roiland and Heidrich, 2011), anxiety (Roiland and Heidrich, 2011), irritated (Huang et al., 2016)) were included in this cluster in previous studies. These variable associations between vasomotor and psychological symptoms are well established and may be related to the multiple and complex etiologies for hormonal symptoms (for review see Avis et al., 2005) and differences in the symptoms included on the various assessment instruments.

Consistent with a previous study (Savard et al., 2009), in our study, occurrence rates of and severity ratings for hot flashes and daytime and night sweats increased from before to 12 months after surgery (see Supplemental Table 2). Given that hormonal symptoms are common and severe in women treated for breast cancer, these symptoms need to be added to multidimensional instruments like the MSAS (Portenoy et al., 1994) and MDASI (Cleeland et al., 2000).

Variable Symptom Clusters

In our study, psychological, cognitive, and sleep symptoms clustered together differently depending on whether the EFAs were conducted before or at 12 months after surgery. Prior to surgery, regardless of the symptom dimension considered, these symptoms loaded on one factor that was named the psychological/cognitive/sleep cluster. In addition, five psychological symptoms formed a distinct irritation cluster using occurrence ratings prior to surgery. Of note, at 12 months after surgery, these same symptoms formed two (i.e., psychological/cognitive, sleep disturbance) or three (i.e., psychological, cognitive, sleep disturbance) distinct clusters depending on whether occurrence rates or severity ratings were used in the EFAs, respectively.

Clusters with a combination of psychological, cognitive, and sleep symptoms

—Symptom clusters that include psychological, cognitive, and sleep symptoms are common in patients with (Bender et al., 2005; Evangelista and Santos, 2012; Fu et al., 2009; Huang et al., 2016; Hwang et al., 2016; Kim et al., 2008; Marshall et al., 2016; Phligbua W, 2013; Reich et al., 2017; Seib et al., 2017; Suwith, 2008) and without (Avis et al., 2001; Cray et al., 2013) cancer. For example, in healthy women across four menopausal stages (Cray et al., 2013), a mood/cognitive/nervous cluster was identified that included six symptoms that were similar to our psychological/cognitive/sleep cluster (i.e., depression, difficulty concentrating, tiredness, forgetfulness, irritability, tension).

Moreover, most ‘psychological’ clusters found in studies of women during treatment for breast (Kim et al., 2008; Phligbua W, 2013; Suwith, 2008) or ovarian (Huang et al., 2016; Hwang et al., 2016) cancer or after breast cancer treatment (Bender et al., 2005; Evangelista and Santos, 2012; Fu et al., 2009; Marshall et al., 2016; Reich et al., 2017; Seib et al., 2017), included cognitive and/or sleep symptoms. For example, a psychoneurological cluster was found in breast cancer patients before, during, and after RT (Kim et al., 2008). Although this cluster consisted of several symptoms that were similar to our psychological/cognitive/sleep cluster (i.e., depressed mood, fatigue, and insomnia (Kim et al., 2008)), it did not vary over time. These inconsistent findings are likely due to differences in the instruments used and the timing of assessments (Kim et al., 2008).

To date, only a limited number of studies have evaluated underlying mechanisms that may contribute to a psychoneurological symptom cluster in cancer patients (for reviews see Miaskowski et al., 2017; Starkweather et al., 2013b). In the two studies that evaluated associations between cytokine levels and this cluster in breast cancer patients, higher symptom burden was associated with several cytokines (e.g., interleukin (IL)3, IL5, IL6, IL7) (Starkweather et al., 2013a) and C-reactive protein (Starkweather et al., 2017). Additional mechanisms that may underlie this cluster include: dysregulation of the hypothalamic-pituitary-adrenocortical (HPA) axis (Thornton et al., 2010) or stress-induced epigenetic changes in neuroendocrine-immune signaling pathways (Lutgendorf and Sood, 2011). Further research is warranted to confirm these preliminary findings.

Clusters of distinct psychological, cognitive, and sleep symptoms—Symptom clusters that contain distinct psychological, cognitive, or sleep symptoms, are less common in the literature. To our knowledge, only four studies identified distinct psychological (Bender et al., 2005; Fu et al., 2009; Reich et al., 2017), cognitive (Bender et al., 2005; Reich et al., 2017; Roiland and Heidrich, 2011), and/or sleep (Roiland and Heidrich, 2011) clusters in women with breast cancer. The consistent symptoms across these three distinct clusters were: anxiety and depression for psychological (Bender et al., 2005; Fu et al., 2009; Reich et al., 2017); memory problems (Bender et al., 2005; Reich et al., 2017; Roiland and Heidrich, 2011) and difficulty concentrating (Bender et al., 2005; Roiland and Heidrich, 2011) for cognitive; and difficulty falling asleep, wake during the night, and waking too early (Roiland and Heidrich, 2011) for sleep.

Of interest and consistent with our study, all of these distinct symptom clusters were identified in women after completion of primary breast cancer treatment (i.e., surgery and

adjuvant CTX and/or RT). Moreover, findings from a randomized controlled trial that evaluated the impact of mindfulness based stress reduction (MBSR) on symptom cluster burden in breast cancer survivors (Reich et al., 2017), suggested improvements in the psychological cluster after six weeks of MBSR. Of note, symptom burden associated with the cognitive and physical clusters did not improve. These findings provide initial evidence of distinct cognitive, psychological, and sleep symptom clusters after breast cancer treatment (Bender et al., 2005; Fu et al., 2009; Reich et al., 2017; Roiland and Heidrich, 2011). Additional research is warranted to confirm these findings and to evaluate for common and distinct underlying mechanisms.

Hypotheses to explain these variable symptom clusters—Similar to our study, previous findings suggest that associations between symptoms vary over time. For example, Sanford and colleagues (Sanford et al., 2014) found that while fatigue, anxiety, and depression were strongly correlated before CTX, fatigue and sleep were highly correlated after CTX. In addition, in a study of a pre-specified symptom cluster (i.e., sleep, depression, and fatigue), correlations among these three symptoms varied over the three assessments that were done before and after adjuvant CTX (Ho et al., 2015).

One potential explanation for the differences in the number and types of symptom clusters is that the etiologies and associated mechanisms for these symptoms at the two assessment times are distinct. For example, prior to surgery, women reported a high level of distress and co-occurring psychoneurological symptoms, which negatively impacted their social and emotional functioning and overall QOL (Denieffe et al., 2014; Doong et al., 2015; Kyranou et al., 2013; McCann et al., 2012; Van Onselen et al., 2013). Consistent with these findings, women in our study reported relatively higher occurrence rates and severity scores for these types of symptoms (i.e., fatigue, anxiety, irritability, difficulty concentrating, tension, depression, anger, mood swings, nervousness) before as compared to after surgery. For example, from before to after surgery, the occurrence rates for anxiety, depression, and difficulty concentrating decreased by 41%, 27%, and 37%, respectively. It is plausible that the increased stress associated with a cancer diagnosis and the impending surgery may contribute to the higher occurrence rates for these co-occurring symptoms as well as to the psychological/cognitive/sleep and irritability clusters found in our study prior to surgery.

However, at one year after surgery, most if not all of these women were cancer survivors. At this time, the majority of the women in our study had completed their adjuvant CTX and/or RT and was on ET. Therefore, it is possible that the etiologies and mechanisms for these symptoms were treatment-related and/or associated with survivorship concerns (e.g., fear of recurrence). For example, cognitive changes after cancer treatment are well documented and may be due to a variety of treatment-related toxicities (e.g., inflammation, peripheral tissue damage, estrogen deprivation) (Merriman et al., 2013). Hence, in this example, the mechanisms that contribute to treatment-related cognitive changes may be different than those that contribute to stress-induced changes prior to surgery. Given that this study is the first to evaluate for differences in the number and types of symptom clusters from before to 12 months after surgery, these findings and hypotheses warrant confirmation in future studies.

Limitations

Several limitations warrant consideration. Given that the majority of women in our study were diagnosed with stage I and II breast cancer, our findings may not generalize to women with advanced stage disease. Because we used EFA to create the symptom clusters, symptoms with lower occurrence rates could not be included in the analysis. Therefore, it is possible that different clusters would be identified, if these symptoms were included in the EFAs. Of note, in a previous study (Ho et al., 2015), the pre-specified symptom cluster varied based on patients' age, menopausal status, and cancer treatments. Future studies need to evaluate for differences in symptom clusters using a variety of demographic and clinical characteristics. In addition, given that the MSS assesses menopausal-related symptoms, more common symptom clusters (e.g., sickness behavior) may have been missed. While two time points were evaluated, the inclusion of a third assessment, perhaps during treatment would provide additional insights into changes in symptom clusters over time. The primary reason for refusal was being overwhelmed with the cancer diagnosis. Therefore, our findings may underestimate the symptom burden of these women. Lastly, the majority of our sample was White and well educated, which limits the generalizability of our findings.

Implications for Clinical Practice and Research

Findings from this study confirm that women with breast cancer experience multiple co-occurring menopausal-related symptoms at two distinct points in their cancer treatment trajectory. Of note, the most common and severe symptoms varied depending on whether symptoms were assessed prior to or at 12 months after surgery. In addition, while two clusters were consistent at both assessments, two clusters were time but not dimension dependent. An increased understanding of how symptom clusters change over time may assist clinicians to better focus their symptom assessment and management strategies. For example, in response to a patient's report of a single symptom (e.g., depression), clinicians may assess the occurrence and severity of additional symptoms within the cluster (e.g., difficulty concentrating and difficulty falling asleep). In addition, our findings suggest that comprehensive evaluations of patient-reported outcomes are warranted as part of survivorship care plans. These evaluations should assess multiple dimensions of the symptom experience (e.g., occurrence, severity) and be done at multiple time points across the cancer treatment trajectory. Given the high occurrence rates and severity scores for the psychological, sleep, and cognitive symptoms prior to surgery, referrals to mental health clinicians or social workers may help women better manage these symptoms.

Given that this study is the first to evaluate for menopausal-related symptom clusters in women before and after breast cancer surgery, additional studies are needed to confirm our findings. Research should focus on the identification of symptom clusters at different points across the treatment trajectory. In addition, future studies should evaluate for differences in the number and types of symptom clusters based on individual factors (e.g., menopausal status, age, types of treatment). For example, the occurrence and severity of hot flashes vary across menopausal stages and cancer treatments. Therefore, future studies should evaluate for these characteristics. Finally, associations between phenotypic and genetic characteristics and symptom clusters warrant evaluation. These investigations may support intervention studies that target multiple co-occurring symptoms and symptom clusters.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was funded by grants from the National Cancer Institute (NCI, CA107091 and CA118658). Dr. Miaskowski is an American Cancer Society Clinical Research Professor and is funded by a K05 award from the NCI (CA168960). This project was supported by NIH/NCRR UCSF-CTSI Grant Number UL1 RR024131. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH. Dr. Mazor was funded by the American Cancer Society and by the National Institute of Nursing Research (T32 NR007088).

References

- Avis NE, Brockwell S, Colvin A. A universal menopausal syndrome? *The American journal of medicine*. 2005; 118(12B):37–46. [PubMed: 16414325]
- Avis NE, Stellato R, Crawford S, Bromberger J, Ganz P, Cain V, Kagawa-Singer M. Is there a menopausal syndrome? Menopausal status and symptoms across racial/ethnic groups. *Social Science & Medicine*. 2001; 52:345–356. [PubMed: 11330770]
- Barsevick A. Defining the symptom cluster: How far have we come? *Seminars in Oncology Nursing*. 2016; 32:334–350.
- Barton DL, Ganz PA. Symptoms: Menopause, Infertility, and Sexual Health. *Advances in Experimental Medicine and Biology*. 2015; 862:115–141. [PubMed: 26059933]
- Bender CM, Ergyn FS, Rosenzweig MQ, Cohen SM, Sereika SM. Symptom clusters in breast cancer across 3 phases of the disease. *Cancer Nursing*. 2005; 28:219–225. [PubMed: 15915067]
- Brown, TA. *The Common Factor Model and Exploratory Factor Analysis*. In: Litte, T., editor. *Confirmatory factor analysis for applied research*. Second. The Guilford Press; New York; London: 2015.
- Browne MW. An overview of analytic rotation in exploratory factor analysis. *Multivariate Behavioral Research*. 2001; 36:111–150.
- Brunner F, Bachmann LM, Weber U, Kessels AG, Perez RS, Marinus J, Kissling R. Complex regional pain syndrome 1--the Swiss cohort study. *BMC Musculoskeletal Disorders*. 2008; 9:92.
- Cieza A, Geyh S, Chatterji S, Kostanjsek N, Ustun BT, Stucki G. Identification of candidate categories of the International Classification of Functioning Disability and Health (ICF) for a Generic ICF Core Set based on regression modelling. *BMC Medical Research Methodology*. 2006; 6:36. [PubMed: 16872536]
- Cleland CS, Mendoza TR, Wang XS, Chou C, Harle MT, Morrissey M, Engstrom MC. Assessing symptom distress in cancer patients: the M.D. Anderson Symptom Inventory. *Cancer*. 2000; 89:1634–1646. [PubMed: 11013380]
- Cray LA, Woods NF, Mitchell ES. Identifying symptom clusters during the menopausal transition: observations from the Seattle Midlife Women's Health Study. *Climacteric*. 2013; 16:539–549. [PubMed: 23153001]
- Denieffe S, Cowman S, Gooney M. Symptoms, clusters and quality of life prior to surgery for breast cancer. *Journal of Clinical Nursing*. 2014; 23:2491–2502. [PubMed: 24329603]
- Dodd MJ, Miaskowski C, Paul SM. Symptom clusters and their effect on the functional status of patients with cancer. *Oncology Nursing Forum*. 2001; 28:465–470. [PubMed: 11338755]
- Doong SH, Dhruva A, Dunn LB, West C, Paul SM, Cooper BA, Elboim C, Abrams G, Merriman JD, Langford DJ, Leutwyler H, Baggott C, Kober K, Aouizerat BE, Miaskowski C. Associations between cytokine genes and a symptom cluster of pain, fatigue, sleep disturbance, and depression in patients prior to breast cancer surgery. *Biological Research in Nursing*. 2015; 17:237–247.
- Evangelista AL, Santos EM. Cluster of symptoms in women with breast cancer treated with curative intent. *Supportive Care in Cancer*. 2012; 20:1499–1506. [PubMed: 21845454]

- Fu OS, Crew KD, Jacobson JS, Greenlee H, Yu G, Campbell J, Ortiz Y, Hershman DL. Ethnicity and persistent symptom burden in breast cancer survivors. *Journal of Cancer Survivorship*. 2009; 3:241–250. [PubMed: 19859813]
- Ganz PA, Kwan L, Stanton AL, Bower JE, Belin TR. Physical and psychosocial recovery in the year after primary treatment of breast cancer. *Journal of Clinical Oncology*. 2011; 29:1101–1109. [PubMed: 21300931]
- Glaus A, Boehme C, Thurlimann B, Ruhstaller T, Hsu Schmitz SF, Morant R, Senn HJ, von Moos R. Fatigue and menopausal symptoms in women with breast cancer undergoing hormonal cancer treatment. *Annals of Oncology*. 2006; 17:801–806. [PubMed: 16507565]
- Gupta P, Sturdee DW, Palin SL, Majumder K, Fear R, Marshall T, Paterson I. Menopausal symptoms in women treated for breast cancer: the prevalence and severity of symptoms and their perceived effects on quality of life. *Climacteric*. 2006; 9:49–58. [PubMed: 16428125]
- Ho SY, Rohan KJ, Parent J, Tager FA, McKinley PS. A longitudinal study of depression, fatigue, and sleep disturbances as a symptom cluster in women with breast cancer. *Journal of Pain and Symptom Management*. 2015; 49:707–715. [PubMed: 25461671]
- Howard-Anderson J, Ganz PA, Bower JE, Stanton AL. Quality of life, fertility concerns, and behavioral health outcomes in younger breast cancer survivors: a systematic review. *Journal of the National Cancer Institute*. 2012; 104:386–405. [PubMed: 22271773]
- Huang J, Gu L, Zhang L, Lu X, Zhuang W, Yang Y. Symptom clusters in ovarian cancer patients with chemotherapy after surgery: A longitudinal survey. *Cancer Nursing*. 2016; 39:106–116. [PubMed: 25837811]
- Hwang KH, Cho OH, Yoo YS. Symptom clusters of ovarian cancer patients undergoing chemotherapy, and their emotional status and quality of life. *European Journal of Oncology Nursing*. 2016; 21:215–222. [PubMed: 26645947]
- Karnofsky D, Abelmann WH, Craver LV, Burchenal JH. The use of nitrogen mustards in the palliative treatment of carcinoma. *Cancer*. 1948; 1:634–656.
- Kim E, Jahan T, Aouizerat BE, Dodd MJ, Cooper BA, Paul SM, West C, Lee K, Swift PS, Wara W, Miaskowski C. Differences in symptom clusters identified using occurrence rates versus symptom severity ratings in patients at the end of radiation therapy. *Cancer Nursing*. 2009; 32:429–436. [PubMed: 19816162]
- Kim HJ, Barsevick AM, Beck SL, Dudley W. Clinical subgroups of a psychoneurologic symptom cluster in women receiving treatment for breast cancer: a secondary analysis. *Oncology Nursing Forum*. 2012; 39:E20–30. [PubMed: 22201665]
- Kim HJ, Barsevick AM, Tulman L, McDermott PA. Treatment-related symptom clusters in breast cancer: a secondary analysis. *Journal of Pain and Symptom Management*. 2008; 36:468–479. [PubMed: 18718735]
- Kim HJ, McGuire DB, Tulman L, Barsevick AM. Symptom clusters: concept analysis and clinical implications for cancer nursing. *Cancer Nursing*. 2005; 28:270–282. [PubMed: 16046888]
- Kirkova J, Walsh D. Cancer symptom clusters—a dynamic construct. *Supportive Care in Cancer*. 2007; 15:1011–1013. [PubMed: 17479300]
- Kwekkeboom KL, Abbott-Anderson K, Cherwin C, Roiland R, Serlin RC, Ward SE. Pilot randomized controlled trial of a patient-controlled cognitive-behavioral intervention for the pain, fatigue, and sleep disturbance symptom cluster in cancer. *Journal of Pain and Symptom Management*. 2012; 44:810–822. [PubMed: 22771125]
- Kyranou M, Paul SM, Dunn LB, Puntillo K, Aouizerat BE, Abrams G, Hamolsky D, West C, Neuhaus J, Cooper B, Miaskowski C. Differences in depression, anxiety, and quality of life between women with and without breast pain prior to breast cancer surgery. *European Journal of Oncology Nursing*. 2013; 17:190–195. [PubMed: 22892272]
- Langford DJ, West C, Elboim C, Cooper BA, Abrams G, Paul SM, Schmidt BL, Levine JD, Merriman JD, Dhruva A, Neuhaus J, Leutwyler H, Baggott C, Sullivan CW, Aouizerat BE, Miaskowski C. Variations in potassium channel genes are associated with breast pain in women prior to breast cancer surgery. *Journal of Neurogenetics*. 2014; 28:122–135. [PubMed: 24392765]
- Lengacher CA, Reich RR, Ramesar S, Alinat CB, Moscoso M, Cousin L, Marino VR, Elias MN, Paterson CL, Pleasant ML, Rodriguez CS, Wang HL, Kip KE, Meng H, Park JY. Feasibility of the

mobile mindfulness-based stress reduction for breast cancer (mMBSR(BC)) program for symptom improvement among breast cancer survivors. *Psychooncology*. 2017

- Lutgendorf SK, Sood AK. Biobehavioral factors and cancer progression: physiological pathways and mechanisms. *Psychosomatic Medicine*. 2011; 73:724–730. [PubMed: 22021459]
- Marshall SA, Yang CC, Ping Q, Zhao M, Avis NE, Ip EH. Symptom clusters in women with breast cancer: an analysis of data from social media and a research study. *Quality of Life Research*. 2016; 25:547–557. [PubMed: 26476836]
- McCann B, Miaskowski C, Koettters T, Baggott C, West C, Levine JD, Elboim C, Abrams G, Hamolsky D, Dunn L, Rugo H, Dodd M, Paul SM, Neuhaus J, Cooper B, Schmidt B, Langford D, Cataldo J, Aouizerat BE. Associations between pro- and anti-inflammatory cytokine genes and breast pain in women prior to breast cancer surgery. *Journal of Pain*. 2012; 13:425–437. [PubMed: 22515947]
- Merriman JD, Von Ah D, Miaskowski C, Aouizerat BE. Proposed mechanisms for cancer- and treatment-related cognitive changes. *Seminars in Oncology Nursing*. 2013; 29:260–269. [PubMed: 24183157]
- Miaskowski C, Aouizerat BE. Is there a biological basis for the clustering of symptoms? *Seminars in Oncology Nursing*. 2007; 23:99–105. [PubMed: 17512436]
- Miaskowski C, Aouizerat BE, Dodd M, Cooper B. Conceptual issues in symptom clusters research and their implications for quality-of-life assessment in patients with cancer. *Journal of the National Cancer Institute*. 2007; 37:39–46.
- Miaskowski C, Barsevick A, Berger A, Casagrande R, Grady PA, Jacobsen P, Kutner J, Patrick D, Zimmerman L, Xiao C, Matocha M, Marden S. Advancing symptom science through symptom cluster research: Expert panel proceedings and recommendations. *Journal of the National Cancer Institute*. 2017; 109
- Miaskowski C, Dodd M, Lee K. Symptom clusters: the new frontier in symptom management research. *Journal of the National Cancer Institute*. 2004; 37:17–21.
- Mitchell ES, Woods NF. Symptom experiences of midlife women: observations from the Seattle Midlife Women's Health Study. *Maturitas*. 1996; 25:1–10. [PubMed: 8887303]
- Muthén BO. Dichotomous factor analysis of symptom data. *Sociological Methods & Research*. 1989; 18:19–65.
- Muthén, LK., Muthén, BO. *Mplus User's Guide*. 7th. Muthén & Muthén; Los Angeles, CA: 1998-2015.
- Nguyen J, Cramarossa G, Bruner D, Chen E, Khan L, Leung A, Lutz S, Chow E. A literature review of symptom clusters in patients with breast cancer. *Expert Review in Pharmacoeconomics and Outcomes Research*. 2011; 11:533–539.
- Phligbua WP, K, Knobf TM, Junda T, Viwatwongkasem C, Srimuninnimit V. Symptom clusters and quality of life in women with breast cancer receiving adjuvant chemotherapy. *Pacific Rim International Journal of Nursing Research*. 2013; 17:249–266.
- Portenoy RK, Thaler HT, Kornblith AB, Lepore JM, Friedlander-Klar H, Kiyasu E, Sobel K, Coyle N, Kemeny N, Norton L, et al. The Memorial Symptom Assessment Scale: an instrument for the evaluation of symptom prevalence, characteristics and distress. *European Journal of Cancer*. 1994; 30A:1326–1336. [PubMed: 7999421]
- Radtke JV, Terhorst L, Cohen SM. The Menopause-Specific Quality of Life Questionnaire: psychometric evaluation among breast cancer survivors. *Menopause*. 2011; 18:289–295. [PubMed: 20881889]
- Reich RR, Lengacher CA, Alinat CB, Kip KE, Paterson C, Ramesar S, Han HS, Ismail-Khan R, Johnson-Mallard V, Moscoso M, Budhrani-Shani P, Shivers S, Cox CE, Goodman M, Park J. Mindfulness-based stress reduction in post-treatment breast cancer patients: Immediate and sustained effects across multiple symptom clusters. *Journal of Pain and Symptom Management*. 2017; 53:85–95. [PubMed: 27720794]
- Roiland RA, Heidrich SM. Symptom clusters and quality of life in older adult breast cancer survivors. *Oncology Nursing Forum*. 2011; 38:672–680. [PubMed: 22037330]

- Sanford SD, Beaumont JL, Butt Z, Sweet JJ, Cella D, Wagner LI. Prospective longitudinal evaluation of a symptom cluster in breast cancer. *Journal of Pain and Symptom Management*. 2014; 47:721–730. [PubMed: 23880589]
- Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis and Rheumatism*. 2003; 49:156–163. [PubMed: 12687505]
- Savard MH, Savard J, Quesnel C, Ivers H. The influence of breast cancer treatment on the occurrence of hot flashes. *Journal of Pain and Symptom Management*. 2009; 37:687–697. [PubMed: 18973980]
- Seib C, Porter-Steele J, McGuire A, McCarthy A, Balaam S, Anderson DJ. Menopausal symptom clusters and their correlates in women with and without a history of breast cancer: a pooled data analysis from the Women's Wellness Research Program. *Menopause*. 2017
- SPSS. IBM SPSS for Windows. 23. SPSS, Inc; Chicago, IL: 2015.
- Starkweather A, Kelly DL, Thacker L, Wright ML, Jackson-Cook CK, Lyon DE. Relationships among psychoneurological symptoms and levels of C-reactive protein over 2 years in women with early-stage breast cancer. *Supportive Care in Cancer*. 2017; 25:167–176. [PubMed: 27599815]
- Starkweather AR, Lyon DE, Elswick RK Jr, Montpetit A, Conley Y, McCain NL. Symptom cluster research in women with breast cancer: A comparison of three subgrouping techniques. *Advances in Breast Cancer Research*. 2013a; 2:107–113. [PubMed: 24498579]
- Starkweather AR, Lyon DE, Elswick RK Jr, Montpetit AJ, Conley Y, McCain NL. A conceptual model of psychoneurological symptom cluster variation in women with breast cancer: Bringing nursing research to personalized medicine. *Current Pharmacogenomics and Personalized Medicine*. 2013b; 11:224–230. [PubMed: 24497894]
- Suwith N, Hanucharurnkul S, Dodd M, Vorapongsathron T, Pongthavorakamol K, Asavemetha N. Symptom clusters and functional status of women with breast cancer. *Thai Journal of Nursing Research*. 2008; 12:153–165.
- Thornton LM, Andersen BL, Blakely WP. The pain, depression, and fatigue symptom cluster in advanced breast cancer: covariation with the hypothalamic-pituitary-adrenal axis and the sympathetic nervous system. *Health Psychology*. 2010; 29:333–337. [PubMed: 20496988]
- Van Onselen C, Aouizerat BE, Dunn LB, Paul SM, West C, Hamolsky D, Lee K, Melisko M, Neuhaus J, Miaskowski C. Differences in sleep disturbance, fatigue and energy levels between women with and without breast pain prior to breast cancer surgery. *Breast*. 2013; 22:273–276. [PubMed: 22858121]
- Woods NF, Mitchell ES. Symptoms during the perimenopause: prevalence, severity, trajectory, and significance in women's lives. *American Journal of Medicine*. 2005; 118(12B):14–24. [PubMed: 16414323]
- Woods NF, Mitchell ES, Lentz M. Premenstrual symptoms: delineating symptom clusters. *Journal of Womens Health and Gender Based Medicine*. 1999; 8:1053–1062.
- Woods NF, Mitchell ES, Schnell JG, Cray L, Ismail R, Taylor-Swanson L, Thomas A. Effects of mind-body therapies on symptom clusters during the menopausal transition. *Climacteric*. 2014; 17:10–22. [PubMed: 23937432]
- Yates P, Miaskowski C, Cataldo JK, Paul SM, Cooper BA, Alexander K, Aouizerat B, Dunn L, Ritchie C, McCarthy A, Skerman H. Differences in composition of symptom clusters between older and younger oncology patients. *Journal of Pain and Symptom Management*. 2015; 49:1025–1034. [PubMed: 25582681]

Highlights

- Prior to and at 12 months after surgery, patients reported an average of 11 symptoms.
- Hormonal and pain/discomfort symptom clusters were identified regardless of time and dimension.
- Clusters that included psychological, cognitive and sleep symptoms varied over time.
- Identification of symptom clusters can guide symptom management interventions.

Table 1
Demographic and clinical characteristics of the sample at enrollment (n=398)

Demographic characteristics	Mean (SD)
Age (years)	54.9 (11.6)
Education (years)	15.7 (2.7)
	% (n)
Ethnicity	
White	64.4 (255)
Non-white	35.6 (141)
Lives alone (% yes)	24.2 (95)
Married/partnered (% yes)	41.9 (165)
Currently working for pay (% yes)	47.8 (189)
Total annual household income	
< \$10,000 to \$19,999	9.7 (32)
\$20,000 to \$99,000	52.3 (172)
\$100,000	38.0 (125)
	% (n)
Clinical characteristics	
	Mean (SD)
Body mass index (kg/m ²)	26.8 (6.2)
Karnofsky Performance Status score	93.2 (10.3)
Self-Administered Comorbidity Questionnaire score	4.3 (2.8)
Mean time since diagnosis (weeks)	9.5 (11.0)
Number of menopausal symptoms prior to surgery (out of 46)	13.2 (8.5)
Number of menopausal symptoms 12 months after surgery (out of 46)	10.9 (8.2)
	% (n)
Occurrence of comorbid conditions (% and number of women who reported each comorbid condition from the Self-Administered Comorbidity Questionnaire)	
Heart disease	3.8 (15)
High blood pressure	30.9 (123)
Lung disease	3.0 (12)
Diabetes	7.8 (31)
Ulcer	3.8 (15)
Kidney disease	0.8 (3)
Liver disease	2.5 (10)

Demographic characteristics	Mean (SD)
Anemia	8.0 (32)
Depression	21.9 (87)
Osteoarthritis	17.3 (69)
Back pain	28.1 (112)
Rheumatoid arthritis	3.5 (14)
Exercise on a regular basis (% yes)	69.6 (275)
Postmenopausal prior to surgery (% yes)	64.1 (248)
Diagnosed with mastitis (% yes)	12.0 (47)
Diagnosed with fibrocystic disease (% yes)	19.1 (73)
Ever breast fed (% yes)	47.0 (186)
Prior hysterectomy (% yes)	13.6 (54)
Prior oophorectomy (% yes)	10.8 (43)
On HRT prior to surgery (% yes)	16.9 (67)
Stage of disease	
Stage 0	16.9 (64)
Stage I	37.7 (143)
Stage IIA and IIB	36.4 (138)
Stage IIIA, IIIB, IIIC, and IV	9.0 (34)
Estrogen receptor positive (% yes)	77.3 (307)
Progesterone receptor positive (% yes)	70.3 (279)
HER2/neu receptor positive (% yes)	16.4 (59)
Received neoadjuvant chemotherapy (% yes)	19.9 (79)
Type of surgery	
Breast conservation	79.9 (318)
Mastectomy	20.1 (80)
Reconstruction at the time of surgery (% yes)	21.7 (86)
Sentinel lymph node biopsy (% yes)	82.4 (328)
Axillary lymph node dissection (% yes)	37.5 (149)
Received adjuvant chemotherapy during the 12 months (% yes)	33.5 (112)
Received external beam radiation therapy during the 12 months (% yes)	72.5 (242)
On endocrine therapy during the 12 months (% yes)	63.2 (211)
BRCA1 and BRCA2 genetic testing	

Demographic characteristics	Mean (SD)
Positive	2.0 (8)
Negative	10.7 (42)
Not done	87.3 (344)

Abbreviations: BRCA = breast cancer; HER2/neu = human epidermal growth factor receptor 2; HRT = hormone replacement therapy; kg = kilogram; m² = meters squared; SD = standard deviation

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2
Exploratory Factor Analysis Using Ratings of Symptom Occurrence Prior to Breast Cancer Surgery^a

Symptom	Factor 1	Factor 2	Factor 3	Factor 4
	Irritability Symptom Cluster	Pain/Discomfort Symptom Cluster	Psychological/Cognitive/Sleep Symptom Cluster	Hormonal Symptom Cluster
Anger	0.776	-0.028	0.021	0.150
Impatience	0.738	0.046	0.023	0.291
Irritability	0.797	0.020	0.011	0.314
Mood swings	0.565	0.106	0.086	0.248
Tension	0.566	0.036	0.453	-0.121
Backache or neckache	0.332	0.578	-0.040	-0.059
General body aches	0.059	0.973	-0.111	-0.007
Joint pain or stiffness	-0.064	0.745	0.070	0.023
Numbness or tingling	-0.285	0.675	0.221	0.164
Painful/tender breasts	0.174	0.520	0.090	-0.267
Weight gain	0.061	0.409	-0.129	0.302
Anxiety	0.390	-0.050	0.644	-0.128
Depression	0.392	-0.001	0.463	0.012
Difficulty concentrating	0.385	-0.072	0.586	0.123
Difficulty falling asleep	0.022	0.077	0.737	0.023
Fatigue or tiredness	0.101	0.373	0.535	0.027
Wake during the night	-0.088	0.357	0.708	0.076
Waking too early	0.043	0.324	0.567	-0.070
Daytime sweats	0.145	-0.022	-0.020	0.825
Hot flashes	-0.043	0.008	0.094	0.939
Night sweats	0.091	0.043	0.024	0.839
Vaginal dryness	-0.044	0.260	0.017	0.431
Forgetfulness	0.241	0.145	0.388	0.197
Headache	0.375	0.389	-0.011	0.009
Total number of symptoms in the cluster	5	6	8	4

Extraction method: unweighted least squares.
Rotation method: Geomin (oblique) rotation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3
Exploratory Factor Analysis Using Ratings of Symptom Occurrence 12 Months After Breast Cancer Surgery^a

Symptom	Factor 1	Factor 2	Factor 3	Factor 4
	Psychological/Cognitive Symptom Cluster	Hormonal Symptom Cluster	Pain/Discomfort Symptom Cluster	Sleep Symptom Cluster
Anger	0.823	0.205	-0.006	-0.183
Anxiety	0.913	-0.122	-0.129	0.028
Depression	0.873	0.035	-0.095	0.024
Difficulty concentrating	0.687	0.091	0.054	0.090
Fatigue or tiredness	0.510	-0.033	0.182	0.239
Forgetfulness	0.587	0.020	0.189	0.096
Impatience	0.889	-0.044	-0.049	-0.024
Irritability	0.995	0.030	0.038	-0.268
Mood swings	0.795	-0.067	0.011	0.093
Tension	0.823	-0.002	0.042	-0.029
Daytime sweats	0.044	0.922	0.017	-0.093
Hot flashes	0.003	0.885	0.008	0.093
Night sweats	-0.025	0.872	-0.040	0.159
Backache or neckache	0.058	0.000	0.536	0.113
General body aches	-0.022	0.020	0.902	-0.003
Joint pain or stiffness	0.011	-0.032	0.751	-0.070
Difficulty falling asleep	0.249	0.130	0.111	0.505
Wake during the night	-0.011	0.182	0.080	0.822
Waking too early	0.035	-0.013	-0.129	0.966
Headache	0.121	0.085	0.383	0.116
Numbness or tingling	0.317	0.145	0.121	0.021
Painful/tender breasts	0.043	0.020	0.238	0.182
Vaginal dryness	0.043	0.020	0.238	0.182
Weight gain	0.143	0.192	0.14	0.174
Total number of symptoms in the cluster	10	3	3	3

Extraction method: unweighted least squares.
Rotation method: Geomin (oblique) rotation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4
Exploratory Factor Analysis Using Ratings of Symptom Severity Prior to Breast Cancer Surgery^a

Symptom	Factor 1	Factor 2	Factor 3
	Psychological/Cognitive/Sleep Symptom Cluster	Hormonal Symptom Cluster	Pain/Discomfort Symptom Cluster
Anger	0.713	-0.110	0.017
Anxiety	0.925	-0.272	-0.125
Depression	0.773	-0.097	-0.120
Difficulty concentrating	0.837	0.027	-0.132
Difficulty falling asleep	0.655	0.068	0.013
Fatigue or tiredness	0.600	0.152	0.161
Forgetfulness	0.557	0.251	0.002
Headache	0.411	0.024	0.280
Impatience	0.754	0.015	-0.004
Irritability	0.729	0.070	0.019
Mood swings	0.692	0.075	0.024
Tension	0.977	-0.309	-0.002
Wake during the night	0.535	0.304	0.056
Waking too early	0.569	0.075	0.089
Daytime sweats	0.036	0.835	-0.140
General body aches	0.003	0.427	0.740
Hot flashes	-0.016	0.962	-0.100
Night sweats	0.065	0.849	-0.096
Numbness or tingling	-0.048	0.506	0.443
Vaginal dryness	-0.043	0.485	0.118
Weight gain	0.035	0.413	0.283
Backache or neckache	0.342	-0.011	0.507
Joint pain or stiffness	0.018	0.377	0.569
Painful/tender breasts	0.310	-0.092	0.362
Total number of symptoms in the cluster	14	7	4

^aExtraction method: unweighted least squares.

Rotation method: Geomin (oblique) rotation.

Table 5
Exploratory Factor Analysis Using Ratings of Symptom Severity 12 Months After Breast Cancer Surgery^a

Symptom	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5
	Psychological Symptom Cluster	Cognitive Symptom Cluster	Hormonal Symptom Cluster	Pain/Discomfort Symptom Cluster	Sleep Symptom Cluster
Anger	0.881	-0.070	0.180	0.055	-0.132
Anxiety	0.879	-0.028	-0.149	-0.072	0.082
Depression	0.706	0.095	0.026	0.001	0.036
Impatience	0.793	0.103	0.020	-0.115	0.010
Irritability	0.946	0.035	0.026	0.030	-0.137
Mood swings	0.676	0.109	-0.016	0.017	0.120
Tension	0.724	0.106	-0.049	0.030	0.028
Difficulty concentrating	0.207	0.678	0.032	-0.014	0.044
Fatigue or tiredness	0.231	0.420	-0.078	0.146	0.199
Forgetfulness	-0.005	0.981	0.069	-0.004	-0.065
Painful/tender breasts	0.063	0.411	-0.107	0.204	0.004
Daytime sweats	0.111	-0.036	0.897	0.059	-0.062
Hot flashes	-0.026	0.150	0.848	-0.011	0.106
Night sweats	-0.008	0.018	0.832	-0.035	0.189
Backache or neckache	0.123	0.015	-0.042	0.618	0.076
General body aches	-0.058	0.016	0.028	0.897	0.006
Headache	0.260	-0.069	0.045	0.466	0.012
Joint pain or stiffness	0.008	0.118	0.023	0.625	-0.020
Difficulty falling asleep	0.132	0.154	0.022	0.084	0.518
Wake during the night	0.051	-0.062	0.138	0.021	0.883
Waking too early	-0.065	0.043	-0.007	-0.002	0.891
Numbness/tingling	0.299	0.077	0.108	0.097	0.071
Vaginal dryness	0.016	0.092	0.037	0.218	0.160
Weight gain	0.019	0.301	0.163	0.131	0.055
Total number of symptoms in the cluster	7	4	3	4	3

Extraction method: unweighted least squares.
Rotation method: Geomin (oblique) rotation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 6
Summary of Symptom Clusters that Were Relatively Consistent Across Dimensions and Time

Symptom Cluster	Symptoms Within the Cluster		Occurrence		Severity	
	Time 1	Time 2	Time 1	Time 2	Time 1	Time 2
<i>Pain/Discomfort Symptom Cluster</i>	Backache/neckache	0.578	0.536	0.578	0.507	0.618
	General body aches	0.973	0.902	0.740	0.897	0.625
	Joint pain or stiffness	0.745	0.751	0.569	0.443	
	Numbness or tingling	0.675				
	Painful/tender breasts	0.520				
	Weight gain	0.409				
	Headache					0.466
	Percent agreement	85.7	42.9	57.1	57.1	57.1
	Daytime sweats	0.825	0.922	0.835	0.897	0.897
	Hot flashes	0.939	0.885	0.962	0.848	0.848
<i>Hormonal Symptom Cluster</i>	Night sweats	0.839	0.872	0.849	0.832	0.832
	Vaginal dryness	0.431		0.485		
	General body aches			0.427		
	Numbness or tingling			0.506		
	Weight gain			0.413		
	Percent agreement	57.1	42.9	100.0	42.9	42.9

Table 7

Summary of Symptom Clusters that Varied Across Time

Symptom Cluster	Occurrence		Severity	
	Symptom	Factor Loading	Symptom	Factor Loading
<i>Psychological/Cognitive/Sleep Symptom Cluster</i>	Time 1 – Prior to Surgery			
	Tension	0.435	Tension	0.977
	Anxiety	0.644	Anxiety	0.935
	Depression	0.463	Depression	0.733
	Difficulty concentrating	0.586	Difficulty concentrating	0.837
	Difficulty falling asleep	0.737	Difficulty falling asleep	0.655
	Fatigue or tiredness	0.535	Fatigue or tiredness	0.600
	Wake during the night	0.708	Wake during the night	0.535
	Waking too early	0.567	Waking too early	0.569
			Anger	0.713
			Forgetfulness	0.557
			Headache	0.411
			Impatience	0.754
			Irritability	0.729
		Mood swings	0.692	
<i>Irritability Symptom Cluster</i>	Time 2 – 12 Months After Surgery			
	Anger	0.778	<i>Not Identified</i>	
	Impatience	0.738		
	Irritability	0.797		
	Mood swings	0.565		
	0.566			
<i>Sleep Symptom Cluster</i>	Time 2 – 12 Months After Surgery			
	Difficulty falling asleep	0.505	Difficulty falling asleep	0.518
	Wake during the night	0.822	Wake during the night	0.883
<i>Psychological/Cognitive Symptom Cluster</i>	Waking too early	0.966	Waking too early	0.891
	Anger	0.823	<i>Not Identified</i>	
	Anxiety	0.913		

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Symptom Cluster	Occurrence		Severity		
	Symptom	Factor Loading	Symptom	Factor Loading	
	Depression	0.873			
	Difficulty concentrating	0.687			
	Fatigue or tiredness	0.510			
	Forgetfulness	0.587			
	Impatience	0.889			
	Irritability	0.995			
	Mood swings	0.795			
	Tension	0.823			
	<i>Psychological Symptom Cluster</i>	<i>Not Identified</i>		Anger	0.881
		<i>Not Identified</i>		Anxiety	0.879
<i>Not Identified</i>		Depression	0.706		
<i>Not Identified</i>		Impatience	0.793		
<i>Not Identified</i>		Irritability	0.946		
<i>Not Identified</i>		Mood swings	0.676		
<i>Not Identified</i>		Tension	0.724		
<i>Not Identified</i>		Difficulty concentrating	0.678		
<i>Not Identified</i>		Fatigue or tiredness	0.420		
<i>Not Identified</i>		Forgetfulness	0.981		
<i>Cognitive Symptom Cluster</i>	<i>Not Identified</i>		Painful/tender breasts	0.411	