

Original Article
Musculoskeletal Disorders



Incidence and Seasonal Variation of Distal Radius Fractures in Korea: a Population-based Study

Young-Hoon Jo ,^{1,*} Bong-Gun Lee ,^{1,*} Hee-Soo Kim ,¹ Joo-Hak Kim ,² Chang-Hun Lee ,³ Sung-Jae Kim ,⁴ Wan-Sun Choi ,⁵ Jae-Ho Lee ,¹ and Kwang-Hyun Lee ¹

¹Department of Orthopaedic Surgery, Hanyang University College of Medicine, Seoul, Korea

²Department of Orthopaedic Surgery, Myongji Hospital, Goyang, Korea

³Department of Orthopaedic Surgery, Eulji University College of Medicine, Seoul, Korea

⁴Department of Orthopaedic Surgery, Hallym University Dongtan Sacred Heart Hospital, Hallym University College of Medicine, Hwaseong, Korea

⁵Department of Orthopaedic Surgery, Ajou University School of Medicine, Suwon, Korea

OPEN ACCESS

Received: Aug 25, 2017

Accepted: Nov 28, 2017

Address for Correspondence:

Kwang-Hyun Lee, MD

Department of Orthopaedic Surgery, Hanyang University College of Medicine, 222-1 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea.
E-mail: leegh@hanyang.ac.kr

*Young-Hoon Jo and Bong-Gun Lee contributed equally to this work.

© 2018 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Young-Hoon Jo

<https://orcid.org/0000-0002-4299-2496>

Bong-Gun Lee

<https://orcid.org/0000-0002-4003-5529>

Hee-Soo Kim

<https://orcid.org/0000-0002-2643-1402>

Joo-Hak Kim

<https://orcid.org/0000-0003-3330-1224>

Chang-Hun Lee

<https://orcid.org/0000-0003-4330-7726>

ABSTRACT

Background: The present study aimed to investigate the incidence and seasonal variation of distal radius fractures (DRFs) in Korea.

Methods: We analyzed a nationwide database acquired from the Korean Health Insurance Review and Assessment Service from 2011 to 2015. We used International Classification of Diseases, 10th revision codes and procedure codes to identify patients of all ages with newly diagnosed DRFs.

Results: An average of about 130,000 DRFs occurred annually in Korea. The incidence of DRF, by age group, was highest in the 10 to 14-year-old age group for males and the highest in the 70s age group for females, with a rapid increase of incidence after 50 years. The peak incidence of DRF occurred during winter; however, the incidence greatly varied annually when compared with that of other seasons. The incidence of DRFs during the winter season was correlated with the average temperature.

Conclusion: The annual incidence of DRF was 130,000 in Korea. The incidence increased under an intense cold surge during winter. Active preventive measures are recommended especially in women exceeding 50 years considering the higher incidence in this age group.

Keywords: Distal Radius Fracture; Epidemiology; Incidence; Seasonal Variation

INTRODUCTION


Distal radius fractures (DRFs) are one of the most common fractures not only among adults but also among children and adolescents.^{1,2} More than 640,000 cases of DRFs are reported annually in the United States,³ and almost 71,000 cases are reported in the British adult population.⁴ The high incidence of DRF causes a substantial economic burden and other social problems including decreased school attendance, lost work hours, loss of independence, and lasting disability.⁵ Therefore, DRF prevention is important, and it is essential to understand the epidemiology of DRFs to achieve this.

Sung-Jae Kim 

<https://orcid.org/0000-0003-4785-8154>

Wan-Sun Choi 

<https://orcid.org/0000-0002-2164-6965>

Jae-Ho Lee 

<https://orcid.org/0000-0003-1224-7038>

Kwang-Hyun Lee 

<https://orcid.org/0000-0002-3060-3916>

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Jo YH, Lee KH. Data

curation: Jo YH, Lee JH. Formal analysis:

Jo YH, Kim SJ. Investigation: Jo YH, Lee KH.

Writing - original draft: Jo YH, Lee BG, Kim HS.

Writing - review & editing: Jo YH, Lee BG, Kim

JH, Lee CH, Choi WS, Lee KH.

Since most DRFs are caused by low-energy trauma, such as a fall from a standing height,⁶ slippery weather conditions may cause fracture epidemics.⁷ Thus, the incidence of DRF, which shows a clear seasonal variation, is known to increase in winter.^{8,9} However, most previous epidemiological studies of the incidence and seasonal variation of DRFs have been confined to European countries and North America.^{7,10} Few studies have examined seasonal variation in DRFs in Asia.¹¹ Lee et al.¹² reported seasonal variations in DRF incidence in Korea; however, their study had some limitations; it had a short-term study period and was conducted in a single center.

Thus, our aim in the present study was to examine the incidence and seasonal variation of DRFs in Korea based on an analysis of nationwide data acquired from the Korean Health Insurance Review and Assessment Service (HIRA).

METHODS

Data source

The authors analyzed nationwide data obtained from the HIRA from 2011 to 2015. In Korea, 97% of the entire population is legally obligated to enroll in the National Health Insurance (NHI) program. Patients only pay about 30% of the total medical cost to healthcare institutions, and all healthcare providers submit claims data for inpatient and outpatient management, including diagnostic codes, which are classified according to the International Classification of Diseases, 10th revision (ICD-10), procedure codes, prescription records, demographic information, and direct medical costs to the HIRA to request reimbursement for the remaining 70% of the medical cost from the NHI service. Of the remaining 3% of the population not registered in the NHI program, excluding illegal residents, most receive healthcare coverage through the Medical Aid Program. The claims data for patients covered by the Medical Aid Program are also reviewed by the HIRA. Hence, the medical records of almost all newly admitted or hospitalized patients in healthcare institutions in Korea are prospectively recorded by the HIRA.

Data collection

The authors used ICD-10 codes and procedure codes to identify patients of all ages with newly diagnosed DRFs in Korea between 2011 and 2015 (**Table 1**). Although HIRA data provide patient identifiers, if a patient with a DRF makes multiple visits to a healthcare institution, claims data for the number of visits are generated. To avoid the risk of multiple counting, the authors used a previously reported method,¹³ as follows. First, patients with surgically treated DRFs were identified with ICD-10 codes (S52.5 and S52.6) and operation codes (N0603, N0613, N0607, N0617, N0993, N0994, and N0983) (**Table 1**). Each operation code was counted as a single case. However, percutaneous pinning and external fixation are often performed simultaneously. Therefore, if a patient received the two operations on the same day, they were considered to be performed on a single DRF. The same criterion was applied when a patient received open reduction with internal fixation and external fixation on the same day. Subsequently, in order to identify patients with DRFs who received conservative treatment, those who underwent surgical treatment were excluded from the HIRA data using patient identifiers, and those with splint or cast codes (T6020, T6030, T6151, and T6152) for DRF ICD-10 codes (S52.5 and S52.6) were included (**Table 1**). For conservative treatment, multiple splint or cast codes are commonly entered for a single case of DRF because an

Table 1. ICD-10 diagnosis codes and procedure codes of distal radius fractures

Codes	Description
ICD-10 code	
S52.5	Fracture of lower end of radius
S52.6	Fracture of lower end of both ulna and radius
Procedure code in Korea	
Operation code	
N0607 or N0617	ORIF for radius
N0603 or N0613	ORIF for both radius and ulna
N0993	Percutaneous pinning for radius
N0994	Percutaneous pinning for radius and ulna
N0983	External fixation
Splint or cast code	
T6020	Long-arm cast
T6030	Short-arm cast
T6151	Long-arm splint
T6152	Short-arm splint

ICD-10 = International Classification of Diseases, 10th revision, ORIF = open reduction with internal fixation.

initially applied splint can be substituted by a cast at a later stage, or a long-arm cast can be changed to a short-arm cast, etc. For this reason, additional codes entered after a period of 6 months following the initial entry of splint or cast codes were recounted.¹⁴

We examined patient data to identify the year and month in which the fracture occurred, and the patient's age and sex. We first counted the number of DRFs for each year and then calculated age-adjusted and sex-specific incidence rates per 100,000 persons with DRFs using the 2010 United States population as the standard population. Estimated year-specific, age-specific, and sex-specific populations in Korea were obtained from the "Statistics Korea" website (<http://www.kosis.kr>). Patients were divided into groups according to age (5-year intervals), and then the incidence of DRFs per age group per study year was calculated according to the population of each age group in Korea for a given study year (per 100,000 persons).

We divided the year into 4 seasons to examine the seasonal variation in the incidence of DRFs; March through May was defined as spring, June through August as summer, September through November as autumn, and December through February as winter. Seasonal variations in the incidence of DRFs were examined according to year and age group (15-year intervals). Lastly, the relationship between the monthly incidence of DRFs and the monthly national average temperature in Korea during winter was investigated.

Statistical analysis

The annual percentage changes in the age-adjusted incidence rate of DRFs were calculated from 2011 to 2015 using joinpoint regression analysis (Joinpoint Regression Program, version 4.3.1.0; National Cancer Institute, Bethesda, MD, USA). All other data sets were analyzed using SAS statistical software version 9.13 (SAS Institute, Cary, NC, USA). Univariate analysis was conducted using the t-test. The correlation between the monthly number of DRFs and the national monthly mean temperatures in winter (<https://data.kma.go.kr>) was analyzed using the Spearman rank test. Values of $P < 0.05$ were considered significant.

Ethics statement

This study protocol was exempted for review by the Institutional Review Board of the Hanyang University Hospital (HYUH 2017-06-027) in accordance with the exemption criteria.

RESULTS

A total of 647,655 DRFs (621,451 patients) occurred from 2011 to 2015; thus, an average of about 130,000 DRFs occurred annually in Korea. Of these, 259,453 cases (249,457 patients) involved male patients, and 388,202 cases (371,994 patients) involved female patients, with a male to female ratio of 1.50. The mean age of patients with DRFs was 47.2 years (standard deviation [SD] \pm 25.5). The mean age of men was 30.2 years (SD \pm 23.2), while that of women was significantly higher at 58.6 years (SD \pm 20.2, $P < 0.001$). The total number of DRFs increased from 2011 to 2013 (128,000 and 137,054, respectively), and then decreased after 2013 (129,687 in 2014 and 116,750 in 2015) (Table 2). The age-adjusted incidence rate per 100,000 persons increased between 2011 and 2012 (278.0 and 291.7, respectively), and then decreased after 2012 (288.33 in 2013 and 246.00 in 2015) (Fig. 1, Table 2). The annual percentage change in the age-adjusted rate for 5 years was calculated as -2.9% (95% confidence interval [CI], -8.2% – 2.6%), which was not statistically significant ($P = 0.190$).

The 5-year age group-specific number of DRFs from 2011 to 2015 was highest in the 10 to 14-year-old age group (124,591 cases, 19%), followed by the 55 to 59-year-old age group (70,978 cases, 11%), and the 60 to 64 year-old age group (61,608 cases, 10%) (Fig. 2). Among those under 45-year-old, the incidence was higher in male than in female individuals; in particular, boys accounted for 82% (101,796 cases) of DRF cases in the 10 to 14-year-old age group. The age group of 45-year-old women had a higher incidence of DRFs than their male counterparts, and 82% (317,687 DRFs) of 387,607 DRFs in persons over 50 years occurred in the female population (Fig. 2, Table 3). The incidence rate per 100,000 persons by age




Fig. 1. Age-adjusted and sex-specific rate of distal radius fractures per 100,000 persons in Korea from 2011 to 2015.

Table 2. Number of cases and age-adjusted rates of distal radius fractures from 2011 to 2015

Years	No. of cases			Age-adjusted rates per 100,000 persons-year ^a		
	Total	Men	Women	Total	Men	Women
2011	128,000	52,822	75,178	277.99	224.20	307.59
2012	136,164	53,480	82,684	291.65	231.29	327.86
2013	137,054	51,086	85,968	288.33	223.93	330.16
2014	129,687	53,525	76,162	274.16	239.50	287.21
2015	116,750	48,540	68,210	246.00	221.25	252.33
Overall	647,655	259,453	388,202			

^aUse of the United States population in 2010 as the control.




Fig. 2. The total number of cases with distal radius fractures, by age group, from 2011 to 2015.

Table 3. Number of distal radius fractures divided by the age of 50 from 2011 to 2015

Years	Men		Women		Total	
	< 50 yr	≥ 50 yr	< 50 yr	≥ 50 yr	< 50 yr	≥ 50 yr
2011	40,067	12,755	15,043	60,135	55,110	72,890
2012	39,472	14,008	14,998	67,686	54,470	81,694
2013	35,691	15,395	14,470	71,498	50,161	86,893
2014	39,434	14,091	13,676	62,486	53,110	76,577
2015	34,869	13,671	12,328	55,882	47,197	69,553
Overall	189,533	69,920	70,515	317,687	260,048	387,607

group peaked in the 10 to 14-year-old age group, then decreased, and increased again as they aged (Fig. 3A). In the female population, the annual incidence rate per 100,000 persons was highest in the 70 to 79-year-old age group, while the 10 to 14-year-old age group showed the highest incidence rate in the male population (Fig. 3B and C).

During the study period, the number of DRFs was highest in winter (33%), followed by autumn (23%), spring (23%), and summer (22%) (Table 4). The annual change in the number of DRFs in winter was more prominent than in other seasons (Fig. 4, Table 4). While the incidence proportion of DRFs in children, adolescents and young men under 30-year-old peaked during spring, females over 15-year-old and men over 30-year-old had the highest incidence proportion during winter (Fig. 5). During the study period, the highest mean monthly number of DRFs occurred in January (15,722 cases, 12%), followed by December (15,508 cases, 12%), and February (11,321 cases, 9%) (Fig. 6). As the national monthly mean winter temperature increased, the monthly number of DRFs tended to decrease, and the correlation was significant ($P = 0.003$, $r = -0.704$) (Fig. 7).

DISCUSSION

An average of 130,000 cases of DRFs occurred annually from 2011 to 2015 in Korea. The annual age-adjusted incidence rate per 100,000 persons was between 240 and 300, but no statistically significant annual change was observed. The incidence of DRFs in males was highest in children and adolescents, while in females it was highest in those in their 70s. Further, females showed a markedly increased DRF incidence after the age of 50 years. DRFs reached peak incidence during winter. However, the annual incidence of DRFs occurring in




Fig. 3. Incidence rate of distal radius fractures per 100,000 persons, by age group — (A) Total population, (B) Men, and (C) Women.

Table 4. Seasonal variation of number of distal radius fractures from 2011 to 2015

Years	Spring	Summer	Autumn	Winter
2011	28,465	27,408	29,247	42,880
2012	29,062	28,403	29,421	49,278
2013	28,934	27,009	28,744	52,367
2014	30,210	29,873	30,407	39,197
2015	29,801	28,542	29,369	29,038
Overall	146,472 (22.6)	141,235 (21.8)	147,188 (22.7)	212,760 (32.9)

Values are presented as number (%).




Fig. 4. Seasonal variation in the incidence of distal radius fractures by year in the total population.

winter greatly varied compared with that occurring in other seasons and correlated with the average temperature during the winter season. Children, adolescents and young men under 30-year-old had a peak incidence proportion of DRFs during spring, and females over 15-year-old and males over 30-year-old had the highest incidence proportion of DRFs during winter.

In studies using national claims data, the number of DRFs may vary slightly from study to study depending on their assessment methods.^{14,15} Kwon et al.¹⁴ examined the DRF incidence and related mortality in patients aged ≥ 50 years with between 2008 and 2012, and reported 65,654 and 74,720 DRFs in 2011 and 2012, respectively. In our study, however, the numbers of DRFs in patients aged ≥ 50 years were 72,890 and 81,694 in 2011 and 2012 respectively, and higher than those of the previous study (Table 3). We considered conservatively-treated patients with multiple procedure codes for splints or casts within 6 months as a single DRF, similar to the method used by Kwon et al.¹⁴ However, patients who received two operations were counted as two DRFs, assuming that they had bilateral DRFs. If the primary surgery was not successful, the patients needed reoperation. In our study, if reoperation was performed, we counted these cases as 2 DRFs. Our method has the potential for overestimating the number of DRFs, but we consider these are very rare cases. Thus, the method we used to assess the number of DRFs produced slightly different results than those reported by the previous study, but this difference was less than 10%.¹⁴

In our study, the incidence rates of DRFs in children and adolescents between 10 and 14 years of age were high, especially in the males, similar to those of previous studies.¹⁶ The cause of a peak incidence rate is known to have a correlation with pubertal growth spurt;¹⁷ the process of bone mineralization cannot keep up with the abrupt increase in new bone development,




Fig. 5. Seasonal variation in the incidence of distal radius fractures by age group from 2011 to 2015 — **(A)** Men and **(B)** Women.

resulting in bones that are particularly susceptible to fracture. It is known that physical activities among these age groups are vigorous¹⁸ and most pediatric forearm fractures are reported to correlate with playing or sports activities.^{19,20} In particular, the incidence proportion of the DRFs in children and adolescents was previously reported to be high in spring.^{19,21} This observation is consistent with our study that children, adolescents and males younger than 30-year-old engaging in vigorous physical activities showed a high incidence proportion of DRFs in spring. Such results are considered due in part to the nature of spring season being appropriate for playing and sports activities with better weather conditions.

We also found that the incidence rate of DRFs notably increased in individuals over 50 years of age, particularly in women, which was similar to the findings of previous studies.^{4,22} The incidence of DRFs increased in older women due to various risk factors, including decreased bone mineral density (BMD), low vitamin D levels, and postural balance decline.²³⁻²⁵ Postural balance decline in older women is associated with a decrease in lower limb strength, sensory




Fig. 6. The mean monthly number of distal radius fractures from 2011 to 2015.




Fig. 7. The correlation between the monthly number of distal radius fractures and the national monthly mean winter temperatures.

system function (vision, vestibular, and proprioception), and flexibility.²⁶ Fu et al.²⁷ reported that sedentary women aged 40 to 60 years improved postural stability through a balance training program. However, the importance of such balance training is often overlooked compared to BMD evaluation.²³ The prevention of osteoporosis and balance decline is needed to reduce the incidence of DRFs in women 50-year-old and older.

Several previous studies in Europe and North America have confirmed that the incidence of DRFs increases in the winter, and this pattern is reported to be associated with snowy and icy conditions.^{7,8,10} However, in East Asia, including China, Japan, and Korea, there have been few studies investigating the seasonal variation in the incidence of DRFs. The marked decrease in temperature during winter in East Asia is accompanied by widespread inflows of cold continental air, called cold surges or cold waves.²⁸ A cold surge is one of the most

important characteristic weather phenomena of the East Asian winter monsoon on synoptic time scales.²⁹ Therefore, seasonal variations in the incidence of DRFs may occur due to such seasonal characteristics in East Asia. All age groups, excluding those usually associated with high-level physical activities, showed a high incidence proportion of DRF during winter.

Bischoff-Ferrari et al.⁹ reported that higher winter temperatures were inversely related to the risk for DRFs. In this study, the incidence of DRFs tended to decrease as the national mean temperature in winter increased. Such effects from weather conditions resulted in annual variations of the DRF incidence rate in winter. During winter, when under an intense cold surge, proper maintenance of roads is necessary in order to reduce the incidence of DRFs.

Although this study employed a large sample size, based on a nationwide database, it also has some limitations. First, the correlation between the incidence of DRFs and snowfall could not be analyzed. Although the Korean Meteorological Administration (<https://data.kma.go.kr>) provides precipitation data, which covers both rainfall and snowfall, it does not distinguish between these 2 parameters. Thus, we could not analyze the correlation between the incidence of DRFs and snowfall. Second, since the HIRA data did not provide the cause of injury, an analysis on the injury mechanism was not available. Third, primary surgery and reoperations could not be distinguished in the coding system. As such, when 2 operation codes were entered for a single patient, it was assumed that the second operation code was entered for the opposite side, rather than for reoperation. With this method, the patients with reoperation were counted as two DRFs, and thus the number of DRFs may have been overestimated. Finally, there is the possibility of some code errors in such a large database.

In conclusion, the annual incidence of DRF was 130,000 in Korea. Children, adolescents, and men under 30-year-old had a high incidence of DRFs in spring. However, the incidence in females over 15-year-old and men over 30-year-old was highest in winter. The incidence of DRFs in winter varied notably from year to year. In winter, the DRF incidence increased under an intense cold surge. Active preventive measures are essential, especially in women 50 years or older given the high DRF incidence in this group.

REFERENCES

1. Court-Brown CM, Caesar B. Epidemiology of adult fractures: a review. *Injury* 2006;37(8):691-7. [PUBMED](#) | [CROSSREF](#)
2. Rennie L, Court-Brown CM, Mok JY, Beattie TF. The epidemiology of fractures in children. *Injury* 2007;38(8):913-22. [PUBMED](#) | [CROSSREF](#)
3. Chung KC, Spilson SV. The frequency and epidemiology of hand and forearm fractures in the United States. *J Hand Surg Am* 2001;26(5):908-15. [PUBMED](#) | [CROSSREF](#)
4. O'Neill TW, Cooper C, Finn JD, Lunt M, Purdie D, Reid DM, et al. Incidence of distal forearm fracture in British men and women. *Osteoporos Int* 2001;12(7):555-8. [PUBMED](#) | [CROSSREF](#)
5. Shauver MJ, Yin H, Banerjee M, Chung KC. Current and future national costs to medicare for the treatment of distal radius fracture in the elderly. *J Hand Surg Am* 2011;36(8):1282-7. [PUBMED](#) | [CROSSREF](#)
6. Diamantopoulos AP, Rohde G, Johnsrud I, Skoie IM, Hochberg M, Haugeberg G. The epidemiology of low- and high-energy distal radius fracture in middle-aged and elderly men and women in Southern Norway. *PLoS One* 2012;7(8):e43367. [PUBMED](#) | [CROSSREF](#)

7. Flinkkilä T, Sirniö K, Hippi M, Hartonen S, Ruuhela R, Ohtonen P, et al. Epidemiology and seasonal variation of distal radius fractures in Oulu, Finland. *Osteoporos Int* 2011;22(8):2307-12.
[PUBMED](#) | [CROSSREF](#)
8. Øyen J, Rohde GE, Hochberg M, Johnsen V, Haugeberg G. Low-energy distal radius fractures in middle-aged and elderly women-seasonal variations, prevalence of osteoporosis, and associates with fractures. *Osteoporos Int* 2010;21(7):1247-55.
[PUBMED](#) | [CROSSREF](#)
9. Bischoff-Ferrari HA, Orav JE, Barrett JA, Baron JA. Effect of seasonality and weather on fracture risk in individuals 65 years and older. *Osteoporos Int* 2007;18(9):1225-33.
[PUBMED](#) | [CROSSREF](#)
10. Giladi AM, Shauver MJ, Ho A, Zhong L, Kim HM, Chung KC. Variation in the incidence of distal radius fractures in the U.S. elderly as related to slippery weather conditions. *Plast Reconstr Surg* 2014;133(2):321-32.
[PUBMED](#) | [CROSSREF](#)
11. Sebastin SJ, Chung KC. An Asian perspective on the management of distal radius fractures. *Hand Clin* 2012;28(2):151-6.
[PUBMED](#) | [CROSSREF](#)
12. Lee HJ, Han SH, Jo YG, Lee SH, Bong ST. Epidemiology and seasonal variance of incidence of distal radius fractures. *J Korean Orthop Assoc* 2015;50(4):307-12.
[CROSSREF](#)
13. Jo YH, Lee BG, Kim JH, Lee CH, Kim SJ, Choi WS, et al. National surgical trends for distal radius fractures in Korea. *J Korean Med Sci* 2017;32(7):1181-6.
[PUBMED](#) | [CROSSREF](#)
14. Kwon GD, Jang S, Lee A, Park CM, Lee YK, Kim TY, et al. Incidence and mortality after distal radius fractures in adults aged 50 years and older in Korea. *J Korean Med Sci* 2016;31(4):630-4.
[PUBMED](#) | [CROSSREF](#)
15. Park C, Ha YC, Jang S, Jang S, Yoon HK, Lee YK. The incidence and residual lifetime risk of osteoporosis-related fractures in Korea. *J Bone Miner Metab* 2011;29(6):744-51.
[PUBMED](#) | [CROSSREF](#)
16. Khosla S, Melton LJ 3rd, Dekutoski MB, Achenbach SJ, Oberg AL, Riggs BL. Incidence of childhood distal forearm fractures over 30 years: a population-based study. *JAMA* 2003;290(11):1479-85.
[PUBMED](#) | [CROSSREF](#)
17. Parfitt AM. The two faces of growth: benefits and risks to bone integrity. *Osteoporos Int* 1994;4(6):382-98.
[PUBMED](#) | [CROSSREF](#)
18. Caspersen CJ, Pereira MA, Curran KM. Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Med Sci Sports Exerc* 2000;32(9):1601-9.
[PUBMED](#) | [CROSSREF](#)
19. Ryan LM, Teach SJ, Searcy K, Singer SA, Wood R, Wright JL, et al. Epidemiology of pediatric forearm fractures in Washington, DC. *J Trauma* 2010;69(4 Suppl):S200-5.
[PUBMED](#) | [CROSSREF](#)
20. Shah NS, Buzas D, Zinberg EM. Epidemiologic dynamics contributing to pediatric wrist fractures in the United States. *Hand (N Y)* 2015;10(2):266-71.
[PUBMED](#) | [CROSSREF](#)
21. Wareham K, Johansen A, Stone MD, Saunders J, Jones S, Lyons RA. Seasonal variation in the incidence of wrist and forearm fractures, and its consequences. *Injury* 2003;34(3):219-22.
[PUBMED](#) | [CROSSREF](#)
22. Tsai CH, Muo CH, Fong YC, Lo WY, Chen YJ, Hsu HC, et al. A population-based study on trend in incidence of distal radial fractures in adults in Taiwan in 2000–2007. *Osteoporos Int* 2011;22(11):2809-15.
[PUBMED](#) | [CROSSREF](#)
23. Louer CR, Boone SL, Guthrie AK, Motley JR, Calfee RP, Wall LB. Postural stability in older adults with a distal radial fracture. *J Bone Joint Surg Am* 2016;98(14):1176-82.
[PUBMED](#) | [CROSSREF](#)
24. Melton LJ 3rd, Christen D, Riggs BL, Achenbach SJ, Müller R, van Lenthe GH, et al. Assessing forearm fracture risk in postmenopausal women. *Osteoporos Int* 2010;21(7):1161-9.
[PUBMED](#) | [CROSSREF](#)
25. Jang WY, Chung MS, Baek GH, Song CH, Cho HE, Gong HS. Vitamin D levels in post-menopausal Korean women with a distal radius fracture. *Injury* 2012;43(2):237-41.
[PUBMED](#) | [CROSSREF](#)
26. Low Choy NL, Brauer SG, Nitz JC. Age-related changes in strength and somatosensation during midlife: rationale for targeted preventive intervention programs. *Ann N Y Acad Sci* 2007;1114(1):180-93.
[PUBMED](#) | [CROSSREF](#)

27. Fu S, Choy NL, Nitz J. Controlling balance decline across the menopause using a balance-strategy training program: a randomized, controlled trial. *Climacteric* 2009;12(2):165-76.
[PUBMED](#) | [CROSSREF](#)
28. Ryoo SB, Kwon WT, Jhun JG. Characteristics of wintertime daily and extreme minimum temperature over South Korea. *Int J Climatol* 2004;24(2):145-60.
[CROSSREF](#)
29. Zhang Y, Sperber KR, Boyle JS. Climatology and interannual variation of the East Asian winter monsoon: Results from the 1979–95 NCEP/NCAR reanalysis. *Mon Weather Rev* 1997;125(10):2605-19.
[CROSSREF](#)