

Loss of *Drosophila* Mei-41/ATR Alters Meiotic Crossover Patterning

Morgan M. Brady,* Susan McMahan,^{†,‡} and Jeff Sekelsky*^{†,‡,1}

*Curriculum in Genetics and Molecular Biology, [†]Department of Biology, and [‡]Integrative Program in Biological and Genome Sciences, University of North Carolina, Chapel Hill, North Carolina 27599

ORCID ID: 0000-0002-4424-677X (J.S.)

ABSTRACT Meiotic crossovers must be properly patterned to ensure accurate disjunction of homologous chromosomes during meiosis I. Disruption of the spatial distribution of crossovers can lead to nondisjunction, aneuploidy, gamete dysfunction, miscarriage, or birth defects. One of the earliest identified genes involved in proper crossover patterning is *Drosophila* *mei-41*, which encodes the ortholog of the checkpoint kinase ATR. Analysis of hypomorphic mutants suggested the existence of crossover patterning defects, but it was not possible to assess this in null mutants because of maternal-effect embryonic lethality. To overcome this lethality, we constructed *mei-41* null mutants in which we expressed wild-type *Mei-41* in the germline after completion of meiotic recombination, allowing progeny to survive. We find that crossovers are decreased to about one-third of wild-type levels, but the reduction is not uniform, being less severe in the proximal regions of chromosome 2L than in medial or distal 2L or on the X chromosome. None of the crossovers formed in the absence of *Mei-41* require *Mei-9*, the presumptive meiotic resolvase, suggesting that *Mei-41* functions everywhere, despite the differential effects on crossover frequency. Interference appears to be significantly reduced or absent in *mei-41* mutants, but the reduction in crossover density in centromere-proximal regions is largely intact. We propose that crossover patterning is achieved in a stepwise manner, with the crossover suppression related to proximity to the centromere occurring prior to and independently of crossover designation and enforcement of interference. In this model, *Mei-41* has an essential function in meiotic recombination after the centromere effect is established but before crossover designation and interference occur.

KEYWORDS meiotic recombination; interference; centromere effect; *Drosophila*; ATR kinase

MEIOTIC crossovers are subject to numerous mechanisms of spatial control to ensure proper disjunction of homologous chromosomes and the generation of genetic diversity. Sturtevant (1913) described the phenomenon of crossover interference, where the presence of one crossover reduces the probability of crossovers nearby (reviewed in Berchowitz and Copenhaver 2010). Mather (1937) pointed out that for small chromosomes “the chiasma frequency equals one, no matter what the size”; Owen (1949) referred to this as the “obligate chiasma.” The phenomenon in which every pair of homologous chromosomes has at least one

crossover that generates a chiasma to promote disjunction is commonly called crossover assurance (reviewed in Wang *et al.* 2015). Together with crossover homeostasis, which buffers crossover formation from increases or decreases in potential crossover precursors (Martini *et al.* 2006), assurance and interference demarcate the minimum and maximum number of crossovers per meiosis. Modeling suggests that crossover assurance, interference, and homeostasis are the result of a single patterning process (Wang *et al.* 2015). The mechanisms that achieve assurance, interference, and homeostasis remain obscure.

Less attention has been paid to the centromere effect; a spatial crossover patterning phenomenon first described by Beadle (1932). Crossovers are excluded from the vicinity of centromeres in many organisms, presumably because very proximal crossovers can interfere with homolog disjunction (Koehler *et al.* 1996; Lamb *et al.* 1996). There are two components to the reduction in crossovers near the centromere. First, Muller and Painter (1932) reported that crossing over is

Copyright © 2018 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.117.300634>

Manuscript received May 30, 2017; accepted for publication December 14, 2017; published Early Online December 15, 2017.

Supplemental material is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.117.300634/-/DC1.

¹Corresponding author: CB #3280, 303 Fordham Hall, Department of Biology, University of North Carolina, Chapel Hill, NC 27599-3280. E-mail: sekelsky@unc.edu

absent or extremely rare within the “inert regions,” now known to comprise heterochromatic, pericentromeric satellite sequence. The second component, which we refer to as the centromere effect, is the phenomenon Beadle described: the reduction in crossing over within crossover-competent regions of the genome as a function of proximity to the centromere. Beadle noticed that when regions with high crossover density were moved closer to the centromere by chromosome rearrangement, crossover frequency decreased. The converse—increased crossover density when centromere-proximal regions are moved away from the centromere—was shown by Mather (1939). The mechanisms underlying the centromere effect are also unknown.

Meiotic recombination is initiated by the formation of DNA double-strand breaks (DSBs). Each DSB can be repaired into a crossover or a noncrossover; the latter can be detected when they result in gene conversion, the unidirectional transfer of sequence from a donor (a homologous chromosome) to a recipient (the chromatid that received the DSB). In *Drosophila*, DSBs appear to be excluded from the pericentric heterochromatin, explaining the absence of crossovers in those regions (Mehrotra and McKim 2006). The centromere effect could, in principle, be explained by decreased DSB density in proximal regions. However, recent whole-genome sequencing reveals that the density of noncrossover gene conversion is relatively constant across the assembled genome on each arm (Comeron *et al.* 2012; Miller *et al.* 2016). This suggests that DSB density is also fairly constant across the chromosome arm and that the centromere effect is exerted by regulating the outcome of DSB repair (crossover or noncrossover) in a manner that is dependent on distance to the centromere.

Mutations in the *Drosophila mei-41* gene were first described by Baker and Carpenter (Baker and Carpenter 1972; Hari *et al.* 1995) who reported a polar reduction in crossovers, with a less severe effect on crossovers in proximal regions, and a possible decrease in interference. These observations suggest a potential role for *Mei-41* in crossover patterning. *Mei-41* is the *Drosophila* ortholog of ATR kinase, best known for regulating DNA damage-dependent cell cycle checkpoints (Hari *et al.* 1995). Consistent with this role, *Mei-41* establishes a checkpoint that monitors progression of meiotic recombination (Ghabrial and Schüpbach 1999; Abdu *et al.* 2002). In addition, *Mei-41* acts redundantly with ATM kinase to promote phosphorylation of histone H2AV at sites of meiotic DSBs (Joyce *et al.* 2011). However, it is unlikely that either of these functions explains the effects of crossover number or position noted by Baker and Carpenter.

Understanding this role is further complicated by the finding that *Mei-41* has an essential function in slowing the rapid nuclear cycles at the midblastula transition in embryonic development (Sibon *et al.* 1999). Females with null mutations in *mei-41* are sterile because this function is lost, and thus the mutations used in previous studies of meiotic recombination are either hypomorphic or separation-of-function alleles (Laurençon *et al.* 2003).

We sought to investigate the possible function for *Mei-41* in crossover patterning by analyzing crossover distribution in *mei-41* null mutants. To overcome the requirement for maternal *Mei-41*, we used a transgene in which *mei-41* expression is under control of a promoter that turns on only after recombination has been completed, thereby generating a fertile *mei-41* “meiotic recombination null” mutant. We find that crossover and nondisjunction phenotypes are more severe in this mutant than in previously reported hypomorphic mutants. We observe a polar effect on chromosome 2L but not on the X; we suggest that this is due to retention of the centromere effect, which is weak on the X. However, interference and assurance are greatly decreased or lost. We propose that the centromere effect is established early in the meiotic recombination pathway and that *Mei-41* has a recombination role after this establishment but before interference and assurance are achieved. Loss of *Mei-41* leads to exit from the meiotic recombination pathway after establishment of the centromere effect and repair is then completed by alternative mechanisms that lack interference and assurance. These findings provide insight into the establishment of crossover patterning.

Materials and Methods

Drosophila stocks

Flies were maintained at 25° on standard medium. To overcome the maternal-effect embryonic lethality of *mei-41^{29D}* null mutation (Sibon *et al.* 1999; Laurençon *et al.* 2003), wild-type genomic *mei-41* was cloned into the *P{attB, UASp::, w⁺m}* vector (courtesy of Steve Rogers) via In-Fusion HD (Takara Bio, Mountain View, CA) and transformed into XL10-Gold Ultracompetent Cells (Agilent Technologies, Santa Clara, CA). This construct was injected via phiC31 integrase-mediated transgenesis into the X chromosome landing site *M{3xP3-RFP.attP}*ZH-2A (BestGene, Chino Hills, CA). The resulting integrants, abbreviated herein as *M{UASp::mei-41}*, were crossed into a *P{mata4::GAL4-VPI6}* background. All *mei-41* null assays used the genotype:

$$\frac{w \text{ mei} - 41^{29D}}{y \text{ M}\{UASp::\text{mei} - 41\} w \text{ mei} - 41^{29D}}, \frac{P\{\text{mata}\alpha::GAL4 - VPI6\}}{+}$$

The *mei-41 mei-P22* double mutant genotype was as above except the third chromosomes were *mei-P22¹⁰³ st/mei-P22¹⁰³ Blm^{D2} Sb P{mata4::GAL4-VP16}*. The *mei-9 mei-41* double mutant genotype was as above except the *X* chromosomes were *y mei-9^a mei-41^{29D}/y M{UASp::mei-41} w mei-9^a mei-41^{29D}*. The presence of the *mei-9^a* mutation was confirmed by allele-specific PCR and by genetic tests (see Supplemental Material, Table S5 in File S1).

Hatch rates

To test *M{UASp::mei-41}* rescue efficiency, 60 virgin females of appropriate genotypes were crossed to 20 isogenized Oregon-Rm males (courtesy of Scott Hawley). Adults were mated in grape-juice agar cages containing yeast paste for 2 days prior to collection. Embryos were collected on grape-juice agar plates for 5 hr and scored for hatching 48 hr later.

Crossover assays and analyses

Meiotic crossovers on chromosome 2L were quantified by crossing *net dpp^{d-ho} dp b pr cn/+* virgin females of the appropriate mutant background to *net dpp^{d-ho} dp b pr cn* males. All six markers were scored in progeny from each genotype, with the of exception *mei-41; mei-P22*. In that case, 731 XX females were scored for all six markers and an additional 1023 XXY females and XY males were scored for *net-b*; eye color markers *pr* and *cn* were excluded because of the presence of a *w* mutation in the mothers. These data were pooled for a final number of 1754 progeny scored.

Meiotic crossovers on *X* were quantified by crossing *y sc cv v gf · y⁺* virgin females of the appropriate background to *y sc cv v gf* male. “*y⁺*” is *Dp(1;1)sc^{V1}*, a duplication of the left end of the *X*, carrying *y⁺*, onto XR. All six markers were scored in all progeny.

To measure chromosome 4 crossovers, the *mei-41* rescue genotype given above was made heterozygous for *PBac{y⁺ w^{+m}}* (101F) and *sv^{spa-pol}*, which are near opposite ends of the assembled region of chromosome 4. These females were crossed with *w¹¹¹⁸; sv^{spa-pol}* males and the progeny were scored for the poliart eye phenotype associated with *sv^{spa-pol}* homozygosity and the *w^{+m}* of the *PBac* transgene. Although both the *M{UASp::mei-41}* and *P{mata4::GAL4-VP16}* transgenes also carry a *w^{+m}*, both confer only mild eye coloration, so the strong red-eye phenotype of *PBac{y⁺ w^{+m}}* (101F) is easily discerned.

Genetic distances, expressed here in centiMorgans (cM) rather than “map units,” as is traditionally used in *Drosophila*, were calculated using the equations of Stevens (1936). Crossover density was calculated by dividing the centiMorgans by the distance between markers (rounded to nearest 10 kb), using *Drosophila melanogaster* reference genome release 6.12 with transposable elements excluded, as described in Hatkevich *et al.* (2017). Including transposable elements in distances did not change any conclusions (see Table S4, a and b, in File S1).

The coefficient of coincidence (*c*) is calculated as $c = (d)(n)/(a)(b)$, where *a* and *b* are the number of single-

crossover progeny in two intervals being compared, *d* is the number of double-crossover (DCO) progeny, and *n* is the total progeny scored. This is equivalent to observed DCOs divided by expected DCOs if the two intervals are independent (no interference). Interference (*I*) is $1 - c$. Thus, $I = 0$ in the absence of interference and $I = 1$ if there is complete positive interference (no DCOs observed).

The centromere effect was quantified as in Hatkevich *et al.* (2017). The definition parallels that of *I*: $CE = 1 - (O/E)$, where *O* is the number of crossovers observed and *E* is the number expected based on the average crossover density across the entire region assayed. *CE* therefore describes the deviation in crossover density in any interval from the mean density across all intervals.

For crossover assurance, we obtained the expected number of meioses in which a given region of the genome (*X* or *net-cn*) had no crossovers (*E₀*) from the Poisson distribution, using mean number of crossovers in that region as the average rate of success. To convert observed crossover classes (parental, single, double, and triple crossover) to bivalent exchange classes (*E₀*, *E₁*, *E₂*, and *E₃*) we used the method of Weinstein (1936). This method accounts for the fact that an *E₁* “tetrad” gives two crossover chromatids and two parental chromatids, so the probability of recovering the crossover in the progeny is 0.5. Weinstein tested models with and without sister chromatid exchange and with and without chromatid interference (*i.e.*, whether the chromatids involved in the two crossovers of a DCO are independent of one another). We used the model that he found to be the best fit to two large *Drosophila* data sets: no sister chromatid exchange and no chromatid interference.

X nondisjunction was scored by crossing virgin mutant females of the appropriate genotypes to *y sc cv v gf/Dp(1:Y)B^S* males. Exceptional progeny for *X* nondisjunction events originate from diplo-*X* and nullo-*X* ova, resulting in XXY (bar-eyed females) and XO (wild-eyed males) progeny, respectively. Numbers of exceptional progeny were doubled to account for those that do not survive to adulthood (XXX and YO).

Statistical analyses

For *cM* and *c* (and therefore *I*), 95% confidence intervals were calculated as $\pm 1.96\sqrt{V(x)}$, where $V(x)$ is the variance of parameter *x*. $V(cM) = (cM)(1 - cM)/n$ and $V(c) = (c/n)(1 - ca - cb - cab + 2c^2ab/ab)$ (Stevens 1936). For nondisjunction, 95% confidence intervals and comparisons of rates across genotypes followed the statistical methods developed by Zeng *et al.* (2010).

For statistical analyses of interference, we conducted χ^2 tests on two-by-two contingency tables of observed and expected DCOs for each genotype. A two-by-two table is appropriate for counts of events that are positive integer values and for which there is an expectation under the null hypothesis that mutant and wild type have the same levels of interference given their levels of recombination. This expected number of DCOs is derived by applying a model of the frequency of DCOs under no interference. Since the data do not

Table 1 Hatch rates for embryos from *mei-41* mutants

Maternal genotype	Hatched (%)	Total (n)
Wild type	73.1 ^a	2035
<i>mei-41</i> ^{29D}	0	527
<i>P{UASp::mei-41} mei-41</i> ^{29D}	0	837
<i>P{UASp::mei-41} mei-41</i> ^{29D} ; <i>P{matα4::GAL4-VP16}/+</i>	52.8 ^b	1187

^a This number is lower than expected for wild type. The cause of this is unknown.

^b The apparent lack of complete rescue may be the result of a high frequency of aneuploidy resulting from the absence of *mei-41* during meiotic recombination.

have covariates or repeated measures, a χ^2 test is the most straightforward. We applied Yates' continuity correction because of low counts in some categories. χ^2 tests were conducted using the GraphPad QuickCalcs online tool (<https://www.graphpad.com/quickcalcs/contingency1.cfm>).

For within-genotype analyses of interference, contingency tables were constructed in two ways. In the first method, columns had numbers with and without crossovers in the first interval, and rows had numbers with and without crossovers in the second interval. These *P*-values are reported in Figure 2. In the second method, columns had number of progeny with and without a DCO in the intervals being compared, and rows were observed and expected. Although *P*-values varied slightly between the two methods, none of the differences affected interpretations concerning statistical significance. For between-genotype comparisons, columns had observed and expected DCOs and rows had wild-type and mutant flies.

A similar argument holds for assurance and *CE*. For assurance, columns were E_0 and $E_{>0}$, and rows had observed and expected counts. For *CE*, within-genotype comparisons had columns for with and without a crossover in the proximal interval and rows were observed and expected. For between-genotype comparisons, columns were observed and expected and rows were wild-type and mutant genotypes.

Data availability

All data necessary for confirming the conclusions presented in the article are presented within the article and the supplemental tables in File S1. *Drosophila* stocks are available upon request.

Results

Postgermarium expression of *Mei-41* rescues embryonic lethality and creates a meiotic recombination null

Drosophila females homozygous for null mutations in *mei-41* produce no viable progeny due to a requirement for maternally deposited *Mei-41* at the midblastula transition (Sibon *et al.* 1999). *Blm* null mutants also exhibit maternal-effect embryonic lethality (McVey *et al.* 2007). To study meiotic recombination in *Blm* null mutants, Kohl *et al.* (2012) expressed wild-type *Blm* under indirect control of the α -tubulin 67C (*matα*) promoter via the Gal4>*UASp* system. This promoter is specific to the female germline, with expression initiating in the early vitellarium (Sanghavi *et al.* 2013), by which time recombination should be complete. In support of this expectation, crossover and nondisjunction assays on the occasional surviving progeny of *Blm* mutant females give similar results

to those from embryos rescued by expressing *UASp::Blm* with the *matα4::GAL4-VP16* driver in *Blm* null mothers (McVey *et al.* 2007; Kohl *et al.* 2012; Hatkevich *et al.* 2017).

We used the same system to overcome the maternal-effect inviability of embryos from *mei-41*^{29D} homozygous null females (see *Materials and Methods*). To quantify the extent of maternal *M{UASp::mei-41}* rescue, we compared hatch rates of embryos from wild-type, *mei-41*^{29D}, and *P{UASp::mei-41} mei-41*^{29D} with and without *P{matα4::GAL4-VP16}* (Table 1). Embryos from females homozygous for *mei-41*^{29D} with or without *M{UASp::mei-41}* but lacking *P{matα4::GAL4-VP16}* did not survive to hatching, whereas embryos from females with both components of the Gal4>*UASp* rescue system had a hatch rate of 52.8%. Most or all of the residual lethality is likely due to aneuploidy resulting from high nondisjunction in *mei-41* mutants (13.6% X nondisjunction among progeny surviving to adulthood; Table S5 in File S1). Larvae that did hatch survived to adulthood, allowing for analysis of the crossover patterning landscape in a *mei-41* null mutant. For simplicity, flies carrying this transgene system are denoted below as *mei-41*^{29D} or *mei-41* null mutants.

Crossover reduction in *mei-41* null mutants

Drosophila mei-41 was initially characterized as a meiotic mutant by Baker and Carpenter (1972). Hypomorphic *mei-41* alleles resulted in an overall 46% decrease in crossovers relative to wild-type controls, and was measured in five adjacent intervals spanning the entirety of chromosome 2L and proximal 2R (~20% of the euchromatic genome). We measured crossovers in this same region in *mei-41* null mutant females and found a significantly more severe reduction of 67% ($P < 0.0001$; Figure 1, A and B). Given the many functions of *Mei-41* in mitotically proliferating cells, we wanted to determine whether the remaining crossovers were meiotic or whether they possibly resulted from DNA damage within the premeiotic germline. As *Mei-P22* is required to generate meiotic DSBs (Liu *et al.* 2002; Robert *et al.* 2016), any crossovers that are independent of *Mei-P22* most likely result from damage occurring in premeiotic mitotic cell cycles or premeiotic S phase. Crossovers were completely abolished in *mei-41*^{29D}; *mei-P22*¹⁰³ double mutants ($n = 1754$). One vial had two female progeny that were mutant for all markers on the *net-cn* chromosome except *pr*. These may have arisen from a DCO in the adjacent *b-pr* and *pr-cn* regions, gene conversion of the *pr* mutation, or reversion of this mutation (an insertion of a 412 transposable element). Since these were in the same vial, they likely represent a single premeiotic




Figure 1 Reduction of crossing over in *mei-41* null mutants. (A and B) Crossover distribution on chromosomes 2L (A) and X (B) in *mei-41*^{29D} mutants compared to wild type. Marker location indicated at top based on genome assembly position (megabase), excluding the centromere, unassembled pericentromeric satellite sequences, and transposable elements. Crossover density (solid lines) was determined for wild-type and *mei-41* mutant females. Dotted lines show mean crossover density across the entire region. (C and D) Crossing over on chromosomes 2L (C) and X (D) in *mei-41*^{29D} mutants as a percentage of wild type. The x-axis is scaled to genetic distance (centiMorgan) in wild-type females. Bars are 95% confidence intervals. (E) Crossover density in *mei-9* and *mei-41* single and double mutants. Note scale difference compared to (A). Wild-type chromosome 2L: $n = 4222$ progeny, 1943 crossovers. *mei-41* chromosome 2L: $n = 7801$ progeny, 1175 crossovers. Wild-type X: $n = 2179$ progeny, 1367 crossovers. *mei-41* X: $n = 5174$ progeny, 1396 crossovers. *mei-9*: $n = 2433$ progeny, 67 crossovers. *mei-9 mei-41*: $n = 1059$ progeny, 165 crossovers. Wild-type and *mei-9* single mutant data are from Hatkevich *et al.* (2017), used with permission. Full data sets are in Tables S1 and S2 in File S1. cM and cM/Mb, with 95% confidence intervals, are in Table S3 in File S1.

event. We conclude that the vast majority of crossovers observed in the *mei-41* null mutant females are meiotic in origin.

Baker and Carpenter (1972) described crossover reduction in *mei-41* hypomorphic mutants as polar, with a more severe decrease in medial and distal regions of the chromosome than in proximal regions. This is also true in our null mutant: Although crossovers are significantly reduced in every interval, the average reduction in the three distal intervals is 75%, while in the two proximal intervals the decrease averages only 16% (Figure 1B). We also assayed crossing over across the entire X chromosome. Crossovers were reduced by an average of 57% on this chromosome; notably, the decrease was uniform across the entire chromosome, with no apparent polar effect (Figure 1, C and D).

One hypothesis to explain the polar effect on recombination on chromosome 2L is that there are region-specific requirements for *Mei-41*, with the protein being less important in proximal 2L. We tested this hypothesis by assessing the dependence of crossovers on *Mei-9*, the catalytic subunit of the putative meiotic resolvase (Sekelsky *et al.* 1995). Meiotic crossovers are reduced by ~90% in *mei-9* mutants, suggesting that most or all crossovers generated in wild-type flies require *Mei-9* (Baker and Carpenter 1972; Sekelsky *et al.*

1995). However, in many mutants that affect meiotic recombination, including *Blm*, *mei-218*, and *rec*, crossovers are independent of *Mei-9* (Sekelsky *et al.* 1995; Blanton *et al.* 2005; Hatkevich *et al.* 2017). Our interpretation is that when the meiotic crossover pathway is blocked because of loss of a critical component, repair is completed by alternative pathways that are independent of *Mei-9* and other downstream meiotic recombination proteins. If *Mei-41* is less important in proximal chromosome 2L, then crossovers in these regions may remain dependent on *Mei-9*. We scored crossovers along chromosome 2L in *mei-9*^a *mei-41*^{29D} double mutants (Figure 1E; we did not score the X chromosome because of the difficulty of recombining the *mei-9* and *mei-41* mutations and the *UAS::mei-41* transgene onto the multiple-marked chromosome, and because the requirement for *Mei-41* appeared to be similar across the X). The total genetic map length was similar between *mei-9 mei-41* double mutants and *mei-41* single mutants (15.58 cM vs. 15.06 cM; $P = 0.6679$ by χ^2 test comparing total crossovers and number of progeny scored), but significantly greater than that of *mei-9* mutants (2.8 cM; $P < 0.0001$). There was no apparent difference in requirement for *Mei-9* between the proximal and distal intervals. We conclude that all crossovers generated in *mei-41* mutants are independent of *mei-9*, regardless of chromosomal

location. This suggests that loss of *Mei-41* disrupts progression through the meiotic crossover pathway at all sites along the chromosome.

The apparent polar effect on crossing over in *mei-41* mutants can be explained by retention of the centromere effect

Compared to wild-type crossing over, the effects of loss of *Mei-41* on meiotic crossing over is puzzling, as there seem to be substantially stronger effects in some regions of the genome than others, yet all crossovers in the mutant are independent of *Mei-9*. The conclusion that there is a polar effect on crossing over is based on comparing crossover frequencies in the mutant to those in wild-type females. Insight can also be gleaned by analyzing crossover distribution in the mutant in isolation. For example, Hatkevich *et al.* (2017) noted an apparently flat distribution of crossover in mutants lacking the *Blm* helicase. Their interpretation was that all crossover patterning is lost in *Blm* mutants, resulting in a distribution that reflects the DSB distribution.

Crossover distribution in *mei-41* mutants does not mimic that of *Blm* mutants, at least in proximal chromosome 2L, suggesting that crossover patterning is not entirely lost in *mei-41* mutants. In wild-type flies, crossover density is substantially lower in the *pr-cn* interval than in any of the other nine intervals that we assayed (0.11 ± 0.03 cM/Mb vs. 0.56 ± 0.09 cM/Mb in the next lowest interval, *net-ho*). The *pr-cn* interval is noteworthy because it spans the centromere, so recombination is strongly influenced by the centromere effect. To determine whether this phenomenon is affected by loss of *Mei-41*, we calculated *CE* as a measure of the centromere effect (Hatkevich *et al.* 2017). *CE* quantifies the difference in crossover density in a given interval relative to mean crossover density. It is thought that the significant decrease in the *pr-cn* interval is primarily due to the centromere effect. In wild-type females, if crossover density in the *pr-cn* interval were equal to the mean density across the entire region assayed, 643 crossovers would be expected; only 73 were observed ($P < 0.0001$), giving a *CE* value of 0.89. In *mei-41*^{29D} mutants, 390 were expected but only 82 were observed ($P < 0.0001$), yielding a *CE* value of 0.79. This high value of *CE* is consistent with most or all of the centromere effect being intact in *mei-41*^{29D} mutants, although the significant difference between *mei-41*^{29D} and wild-type females ($P = 0.0004$) suggests that there may be mild amelioration in *mei-41* mutants.

The decrease in crossing over on the X chromosome does not appear to be polar (Figure 1, C and D). The pericentric heterochromatin of the X chromosome spans ~19 Mb, compared to ~5 Mb on chromosome 2L and 7 Mb on 2R. This results in a much weaker centromere effect in the most proximal euchromatin of the X (Yamamoto and Miklos 1978). Thus, the lack of a polar decrease in crossing over on the X in *mei-41* null mutants may be because the entire region being analyzed is, with respect to distance from the centromere, equivalent to the distal half of chromosome 2L.

The small chromosome 4 of *D. melanogaster* never has meiotic crossovers in wild-type females (reviewed in Hartmann and Sekelsky 2017), but does have crossovers in *Blm* mutants (Hatkevich *et al.* 2017). Hatkevich *et al.* argued that the absence of crossovers on chromosome 4 is due in large part to the centromere effect (~4 Mb of heterochromatic satellite sequence between the centromere and the gene-containing region), and that it is loss of the centromere effect that permits crossing over on chromosome 4 in *Blm* mutants. We measured crossing over on chromosome 4 in *mei-41* mutants. We recovered no crossovers between markers at opposite ends of the gene-containing region of chromosome 4 ($n = 5555$; $P < 0.0001$ compared to *Blm*), consistent with our interpretation that the centromere effect is not lost in *mei-41* mutants.

Effects of loss of *Mei-41* on crossover interference and assurance

Given the apparent retention of the centromere effect on crossing over, we asked whether the crossover patterning phenomena interference and assurance are affected by loss of *Mei-41*. We calculated interference (*I*) using the method of Stevens (1936). Stevens defined *I* as $1 - (O/E)$, where *O* is the number of DCOs observed and *E* is the number of DCO expected if the two intervals are independent of one another (see *Materials and Methods*). Thus, $I = 1$ indicates complete positive interference (no DCOs observed) and $I = 0$ indicates no interference (the two intervals are independent of one another).

Values for *O*, *E*, and *I* are given in Figure 2, A and B. On chromosome 2L, the only pairs of adjacent intervals that have enough DCOs to analyze interference are *ho-dp/dp-b* and *dp-b/b-pr*. In wild-type females, *I* was 0.93 ± 0.05 between the first pair and 0.64 ± 0.15 between the second pair (Figure 2A). In *mei-41* mutants, we did not detect significant interference between the first pair ($I = 0.26 \pm 0.52$, $P = 0.4977$), indicating a significant difference from the wild type ($P < 0.0001$). However, we did detect interference between the second pair of intervals in the *mei-41* mutant ($I = 0.63 \pm 0.25$, $P = 0.0019$), suggesting that interference is intact in this region ($P = 0.9922$ compared to the wild type). The *dp-b* interval is typically not used for measuring interference because of its large size (>27 cM in wild-type flies). Therefore, we reexamined interference within this region by subdividing it with another marker, *wg*^{Sp-1}. Again, interference was strong in wild-type females ($I = 0.92 \pm 0.07$) but absent from *mei-41* mutants ($I = 0.04 \pm 0.38$; Figure 2A, bottom). Using the same analysis of interference across the X chromosome, we found significant positive interference between every pair of adjacent intervals in wild-type females, but no detectable interference in *mei-41* mutants (Figure 2B; the *P*-value for *cv-v* and *v-g* is 0.0131, which would be considered significant, but this is because there were significantly more DCOs observed than expected, resulting in a negative value for *I*).

We used one additional method to assess the distribution of crossovers relative to one another. In many species, crossovers are distributed among bivalents such that the probability that any pair of homologous chromosomes does not receive




Figure 2 Interference and assurance in *mei-41* null mutants. (A) Interference on chromosome 2L. Black line represents genetic map of markers used, with size of each interval (in centiMorgan) listed above the line for wild-type females and below the line for *mei-41* mutants. The *pr-cn* interval was omitted because it spans the centromere and because of low numbers of DCOs between this and the adjacent interval. Arcs represent pairs of adjacent intervals in which interference was assessed. Above (for wild type, brown) or below (*mei-41*, blue) each arc is listed the number of observed DCOs (O) and the number expected (E) if the two intervals are independent (no interference). P -values indicate the probability of observing these data under the null hypothesis, which is that the two intervals are independent (see *Materials and Methods*). Stevens' interference (I), which equals $1 - (O/E)$, is also given. Red * below map lines are from Chi-squared analysis comparing O and E between wild type and mutant (see *Materials and Meth-*

ods). In a separate experiment, the large *dp-b* interval was further divided by the addition of *Sp* (*wg^{Sp-1}*). (B) Similar analysis of interference on the X chromosome. The *f-y+* region spans the centromere, but since the marker on the right arm (*y+*) is hemizygous (*i.e.*, a duplication of the tip of chromosome XL onto XR on one homolog), all crossovers must be to the left of the centromere. (C) Crossover assurance assessed by comparing frequencies of E_0 bivalents. Expected E_0 frequency is based on Poisson distribution from the average number of crossovers per meiosis; observed frequencies were calculated using the method of Weinstein (1936). Statistical significance between expected and observed E_0 frequencies determined via Chi-square tests. Bars show 95% confidence intervals. Sample sizes for chromosomes 2L and X are given in Figure 1. For the *dp-Sp-b* experiment, $n = 3325$ flies, 928 crossovers for wild-type; $n = 9740$ flies, 972 crossovers for mutants. P -values reported were not corrected for multiple comparisons. Nonsignificance (ns) is for $P > 0.05$. * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$.

a crossover is significantly lower than expected by chance, a phenomenon known as crossover assurance. It has been proposed that if there are sufficient well-spaced, crossover-eligible intermediates, then coupling interference with a mechanism to achieve a specific number of crossovers per meiosis (within a narrow range) will produce crossover assurance (Zhang *et al.* 2014; Wang *et al.* 2015). In this model, assurance is merely an outcome of interference and homeostasis.

Extrapolating from our measurements of crossovers on chromosomes X and 2L, we estimate approximately two crossovers per meiosis in *mei-41* mutants. True assurance requires a minimum of three or five crossovers (one per major chromosome or arm, excluding chromosome 4); however, assurance among the residual crossovers could manifest as the two crossovers being on different chromosomes (or chromosome arms) more often than expected by chance. We compared the expected and observed frequency of meioses in which there were no crossovers (E_0 , for zero-exchange bivalent, frequency) on the X chromosome or on chromosome 2L. For expected E_0 frequency we used the Poisson distribution expectation based on the average number of crossovers per meiosis. We used the method of Weinstein (1936) to transform counts of progeny that inherited parental, single crossover, DCO, *etc.*, chromatids to observed bivalent exchange

classes (see *Materials and Methods*). In wild-type flies, the expected E_0 frequency for the X chromosome is 0.285, but the observed frequency was 0.112 (Figure 2C). This demonstrates crossover assurance that is significant ($P < 0.0001$) but incomplete (11% of meioses have no crossovers between the X chromosomes), as has been observed in previous studies (*e.g.*, Weinstein 1936; Koehler *et al.* 1996). In *mei-41* mutants, reduced crossing over results in a higher expected E_0 frequency (0.582), but unlike the case in wild-type flies, the observed frequency (0.572) was not significantly different ($P = 0.3008$). Similar results were obtained with the chromosome 2L data (Figure 2C). For 2L, the difference between observed and expected in *mei-41* mutants was significant ($P = 0.0046$), but given the small magnitude of the difference (0.740 expected, 0.720 observed) this may not be biologically meaningful.

Together, our data indicate that interference and assurance are significantly decreased or lost in *mei-41* mutants, though it is possible that crossovers in proximal chromosome 2L retain interference.

Discussion

We have demonstrated that the *Gal4>UASp* rescue successfully overcomes maternal-effect embryonic lethality of *mei-41* mutants, allowing us to perform meiotic crossover patterning

analysis in *mei-41* null mutants. The crossover reduction in null mutants is more severe than that previously reported for hypomorphic mutants (Baker and Carpenter 1972), but the nonuniform reduction in crossing over on chromosome 2 is still present (Figure 1B).

We considered the hypothesis that the polar effect stems from differential requirement for *Mei-41* in proximal and distal regions of the chromosome. However, in *mei-41* mutants, crossovers in all regions are independent of the presumptive resolvase *Mei-9* (Figure 1C). Our interpretation is that this reveals an essential role for *Mei-41* in carrying out meiotic recombination throughout the genome. In the absence of *Mei-41*, the meiotic pathway is disrupted and repair is completed by alternative pathways that neither require functions specific to the meiotic pathway nor result in properties normally associated with meiotic recombination, such as crossover patterning.

Since the apparent polar effect is observed on chromosome 2 but not on the *X*, we hypothesized that the centromere effect is retained in *mei-41* mutants. We calculated *CE*, a measure of how much crossover density in an interval deviates from the mean crossover density (Hatkevich *et al.* 2017), to compare the centromere effect between wild-type females and *mei-41* mutants. Although every interval deviates significantly from the mean in wild-type flies, the very strong deviation in the *pr-cn* region (*CE* = 0.89) is probably due primarily to the suppression of crossovers associated with proximity to the centromere. Direct confirmation of the presence of a centromere effect requires moving the sequences to be analyzed away from the centromere through chromosome rearrangement, a difficult experiment because of the need to have structural homozygosity combined with heterozygosity for markers. Nonetheless, our data suggest that a strong centromere effect is retained in the absence of *Mei-41*.

In contrast to the absence of a strong impact on the centromere effect, our analysis suggests that interference and assurance are significantly disrupted when *Mei-41* is absent. We did not detect any significant interference across the *X* chromosome (Figure 2B). On chromosome 2L, interference was lost between *dp* and *b* (when divided into two intervals), but was retained between this interval and *b-pr* (Figure 2A). It is notable that the only interval that appears to retain interference is *b-pr*, which is the closest interval to the centromere. This could indicate that *Mei-41* does have different functions in proximal regions than in other parts of the genome. However, given that there appears to be no interference within the adjacent *dp-b* interval, the presence of interference between these intervals would require that crossover-eligible intermediates between *dp* and *b* should be subject to interference exerted by crossover designations in the proximal interval, while at the same time any crossovers designated between *dp* and *b* should not signal interference themselves. This seems unlikely, and perhaps indicates that there is some other effect or some idiosyncrasy associated with the *b-pr* interval.

Another argument regarding the reduced or absent interference in *mei-41* mutants is that the strength of interference




Figure 3 Model for progressive enforcement of crossover patterning. The drawing at the top represents a chromosome arm. Thick solid line is pericentric satellite DNA, circle is centromere. (A) Based on whole-genome sequencing, the initial DSB distribution (dotted line) is flat at large scales (DSBs are excluded from the heterochromatic satellite DNA). The centromere effect revokes the eligibility of intermediates near the centromere to become crossovers. (B) This results in a distribution of crossover-eligible intermediates that is flat across much of the arm, then tailed as the centromere is approached. The shape of the tailing is unknown; a sigmoidal drop is shown here for illustrative purposes. Later, some intermediates are designated to become crossovers, and the resultant interference discourages other intermediates over large distances from achieving crossover designation. The resultant distribution of crossover-designated intermediates (C) and crossovers (D) is approximately skew normal, with the degree of skew being proportional to the length of satellite sequence that separates DSB-competent regions from the centromere. *Blm*, *Mei-41*, and *Mei-9* are essential at different times in the crossover pathway, so crossovers generated in mutants that lack these proteins are made outside the normal meiotic pathway, and are therefore *Mei-9* independent, and are either unpatterned (*Blm* mutants, distribution similar to A), are partially patterned (*mei-41* mutant, resembles B), or are fully patterned (*mei-9* mutant, resembles C and D).

is inversely proportional to the genetic size of the interval in which it is measured. Therefore, since genetic intervals become shorter in *mei-41* mutants, interference might be expected to become stronger. This expectation would not hold for the recombination proteins required to generate crossovers after interference has occurred, such as the proteins that resolve crossover-designated intermediates into crossovers [the “crossover maturation” step in the models of Zhang *et al.* (2014)]. *Mei-9* and associated proteins are thought to be required for resolution (Baker and Carpenter 1972; Sekelsky *et al.* 1995; Yildiz *et al.* 2002). It is not meaningful to discuss interference in *mei-9* mutants, since the number of crossovers per meiosis is well below one (0.06), but the uniform decrease in crossovers across chromosome 2L led Baker and Carpenter (1972) to conclude that *Mei-41* acts earlier in crossover generation than *Mei-9*.

We believe the most parsimonious interpretation of our data is that loss of *Mei-41* has little or no impact on the centromere effect but reduces or eliminates interference and assurance. We propose that crossover patterning in *Drosophila* occurs in a stepwise manner (Figure 3). Analysis of noncrossover gene conversion events mapped through whole-genome sequencing suggests that DSBs are, at a large scale, spread evenly throughout the assembled genome (Comeron *et al.* 2012; Miller *et al.* 2016; Hatkevich *et al.* 2017). The centromere effect is applied early by making some intermediates ineligible to enter the crossover pathway, with the probability of being affected in this way being related to distance to the centromere (Figure 3B). Subsequently, when any remaining crossover-eligible intermediate becomes designated for crossing over, interference precludes nearby intermediates from also adopting this fate (Figure 3C).

Given a uniform distribution of DSBs and the fact that each of the chromosome arms in *Drosophila* has ~1.0–1.3 crossovers per meiosis, interference alone will produce a crossover density that resembles a normal distribution (*e.g.*, simulations in Zhang *et al.* 2014). The combination of a strong centromere effect and interference will yield a crossover density map that approximates a skew normal distribution (Figure 3D). Crossover distribution maps in *Drosophila* do resemble skew normal distributions, with much more skew on the major autosome arms than on the *X* chromosome, which also lacks a strong centromere effect (see figure S2 in Comeron *et al.* 2012).

Blm helicase has been proposed to have an essential function early in the meiotic recombination pathway (reviewed in Hatkevich and Sekelsky 2017). Loss of *Blm* results in an early exit from the meiotic pathway and completion of repair by alternative mechanisms. Since these alternative mechanisms do not involve patterning, the probability of becoming a crossover is the same for each intermediate, resulting in crossovers being evenly distributed across each chromosome arm. We propose that *Mei-41* has some critical function after the centromere effect has been at least partially established. Loss of *Mei-41* leads to exit from the meiotic pathway at this

point. As with *Blm* mutants, every remaining intermediate has the same probability of becoming a crossover, so the crossover distribution in *mei-41* mutants is similar to that in Figure 3B. *Mei-9* is required only for maturation of crossover-designated intermediates into crossovers. Since this occurs after crossover designation, residual crossovers in a *mei-9* mutant are patterned like crossovers in wild-type flies; but there are far fewer crossovers because most intermediates that had been designated to become crossovers are instead processed into noncrossover products.

In many model organisms, a subset of crossovers do not participate in interference and are generated by different resolvases than those that generate interfering crossovers (reviewed in Kohl and Sekelsky 2013). These “class II” crossovers are sometimes defined as lacking interference or being unpatterned. In the model discussed above, this distinction is not always appropriate, at least in mutant situations. Rather, crossovers generated outside of the primary pathway may be unpatterned (as in *Blm* mutants), partially patterned (as in *mei-41* mutants), or patterned (as in *mei-9* mutants). The only features that these crossovers have in common is that they are generated outside of the normal meiotic crossover pathway, presumably through general DSB repair pathways that act to ensure there are no unrepaired DNA structures persisting until the meiotic divisions begin.

Our data provide little insight into the molecular function of *Mei-41* in the meiotic DSB repair pathway. In mitotic DSB repair, *mei-41* mutants have no observable defects in the early steps of homologous repair of DSBs (*e.g.*, resection, strand invasion, and repair synthesis), but *Mei-41* appears to be required for the annealing or ligation steps of synthesis-dependent strand annealing (LaRocque *et al.* 2007). Korda Holsclaw and Sekelsky (2017) hypothesized that *Mei-41* activates Marcal1, which then catalyzes annealing of complementary sequences. Synthesis-dependent strand annealing promotes formation of noncrossover products, in contrast to the apparent role for *Mei-41* in promoting crossovers during meiotic recombination. Nonetheless, *Mei-41* might have similar functions in mitotic and meiotic DSB repair if, in the latter, it activates a protein that catalyzes the annealing required for second-end capture, a process that might occur after an early requirement for *Blm* helicase but prior to crossover designation (*e.g.*, models in Crown *et al.* 2014). Future studies to elucidate the role for *Mei-41* might provide additional insights into meiotic crossover pathways and patterning.

Acknowledgments

We thank Nicole Crown, Michaelyn Hartmann, and Talia Hatkevich for comments on the manuscript and Corbin Jones and Nadia Singh for advice on statistical analyses. This work was supported by a grant from the National Institute of General Medical Sciences to J.S. under award 1R35 GM-118127.

Literature Cited

- Abdu, U., M. Brodsky, and T. Schüpbach, 2002 Activation of a meiotic checkpoint during *Drosophila* oogenesis regulates the translation of Gurken through Chk2/Mnk. *Curr. Biol.* 12: 1645–1651.
- Baker, B. S., and A. T. C. Carpenter, 1972 Genetic analysis of sex chromosomal meiotic mutants in *Drosophila melanogaster*. *Genetics* 71: 255–286.
- Beadle, G. W., 1932 A possible influence of the spindle fibre on crossing-over in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 18: 160–165.
- Berchowitz, L. E., and G. P. Copenhaver, 2010 Genetic interference: don't stand so close to me. *Curr. Genomics* 11: 91–102.
- Blanton, H. L., S. J. Radford, S. McMahan, H. M. Kearney, J. G. Ibrahim *et al.*, 2005 REC, *Drosophila* MCM8, drives formation of meiotic crossovers. *PLoS Genet.* 1: e40.
- Comeron, J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in *Drosophila melanogaster*. *PLoS Genet.* 8: e1002905.
- Crown, K. N., S. McMahan, and J. Sekelsky, 2014 Eliminating both canonical and short-patch mismatch repair in *Drosophila melanogaster* suggests a new meiotic recombination model. *PLoS Genet.* 10: e1004583.
- Ghahrial, A., and T. Schüpbach, 1999 Activation of a meiotic checkpoint regulates translation of Gurken during *Drosophila* oogenesis. *Nat. Cell Biol.* 1: 354–357.
- Hari, K. L., A. Santerre, J. Sekelsky, K. S. McKim, J. B. Boyd *et al.*, 1995 The *mei-41* gene of *D. melanogaster* is a structural and functional homolog of the human ataxia telangiectasia gene. *Cell* 82: 815–821.
- Hartmann, M. A., and J. Sekelsky, 2017 The absence of crossovers on chromosome 4 in *Drosophila melanogaster*: Imperfection or interesting exception? *Fly (Austin)* 11: 253–259.
- Hatkevich, T., and J. Sekelsky, 2017 Bloom syndrome helicase in meiosis: pro-crossover functions of an anti-crossover protein. *Bioessays* 39. DOI: 10.1002/bies.201700073.
- Hatkevich, T., K. P. Kohl, S. McMahan, M. A. Hartmann, A. M. Williams *et al.*, 2017 Bloom syndrome helicase promotes meiotic crossover patterning and homolog disjunction. *Curr. Biol.* 27: 1–5.
- Joyce, E. F., M. Pedersen, S. Tiong, S. K. White-Brown, A. Paul *et al.*, 2011 *Drosophila* ATM and ATR have distinct activities in the regulation of meiotic DNA damage and repair. *J. Cell Biol.* 195: 359–367.
- Koehler, K. E., C. L. Boulton, H. E. Collins, R. L. French, K. C. Herman *et al.*, 1996 Spontaneous X chromosome MI and MII nondisjunction events in *Drosophila melanogaster* oocytes have different recombinational histories. *Nat. Genet.* 14: 406–414.
- Kohl, K. P., and J. Sekelsky, 2013 Meiotic and mitotic recombination in meiosis. *Genetics* 194: 327–334.
- Kohl, K. P., C. D. Jones, and J. Sekelsky, 2012 Evolution of an MCM complex in flies that promotes meiotic crossovers by blocking BLM helicase. *Science* 338: 1363–1365.
- Korda Holsclaw, J., and J. Sekelsky, 2017 Annealing of complementary DNA sequences during double-strand break repair in *Drosophila* is mediated by the ortholog of SMARCAL1. *Genetics* 206: 467–480.
- Lamb, N. E., S. B. Freeman, A. Savage-Austin, D. Pettay, L. Taft *et al.*, 1996 Susceptible chiasmate configurations of chromosome 21 predispose to non-disjunction in both maternal meiosis I and meiosis II. *Nat. Genet.* 14: 400–405.
- LaRocque, J. R., B. Jaklevic, T. T. Su, and J. Sekelsky, 2007 *Drosophila* ATR in double-strand break repair. *Genetics* 175: 1023–1033.
- Laurençon, A., A. Purdy, J. Sekelsky, R. S. Hawley, and T. T. Su, 2003 Phenotypic analysis of separation-of-function alleles of MEI-41, *Drosophila* ATM/ATR. *Genetics* 164: 589–601.
- Liu, H., J. K. Jang, N. Kato, and K. S. McKim, 2002 *Mei-P22* encodes a chromosome-associated protein required for the initiation of meiotic recombination in *Drosophila melanogaster*. *Genetics* 162: 245–258.
- Martini, E., R. L. Diaz, N. Hunter, and S. Keeney, 2006 Crossover homeostasis in yeast meiosis. *Cell* 126: 285–295.
- Mather, K., 1937 The determination of position in crossing-over. II. The chromosome length-chiasma frequency relation. *Cytologia Fujii Jubilee Volume*: 514–526.
- Mather, K., 1939 Crossing over and heterochromatin in the X chromosome of *Drosophila melanogaster*. *Genetics* 24: 413–435.
- McVey, M., S. L. Andersen, Y. Broze, and J. Sekelsky, 2007 Multiple functions of *Drosophila* BLM helicase in maintenance of genome stability. *Genetics* 176: 1979–1992.
- Mehrotra, S., and K. S. McKim, 2006 Temporal analysis of meiotic DNA double-strand break formation and repair in *Drosophila* females. *PLoS Genet.* 2: e200.
- Miller, D. E., C. B. Smith, N. Yeganeh Kazemi, A. J. Cockrell, A. V. Arvanitakis *et al.*, 2016 Whole-genome analysis of individual meiotic events in *Drosophila melanogaster* reveals that noncrossover gene conversions are insensitive to interference and the centromere effect. *Genetics* 203: 159–171.
- Muller, H. J., and T. S. Painter, 1932 The differentiation of the sex chromosomes of *Drosophila* into genetically active and inert regions. *Z. Induct. Abstammungs Vererbungsl.* 62: 316–365.
- Owen, A. R. G., 1949 A possible interpretation of the apparent interference across the centromere found by Callan and Montalenti in *Culex pipiens*. *Heredity (Edinb)* 3: 357–367.
- Robert, T., A. Nore, C. Brun, C. Maffre, B. Crimi *et al.*, 2016 The TopoVIB-Like protein family is required for meiotic DNA double-strand break formation. *Science* 351: 943–949.
- Sanghavi, P., S. Laxani, X. Li, S. L. Bullock, and G. B. Gonsalvez, 2013 Dynein associates with oskar mRNPs and is required for their efficient net plus-end localization in *Drosophila* oocytes. *PLoS One* 8: e80605.
- Sekelsky, J. J., K. S. McKim, G. M. Chin, and R. S. Hawley, 1995 The *Drosophila* meiotic recombination gene *mei-9* encodes a homologue of the yeast excision repair protein Rad1. *Genetics* 141: 619–627.
- Sibon, O. C., A. Laurençon, R. Hawley, and W. E. Theurkauf, 1999 The *Drosophila* ATM homologue Mei-41 has an essential checkpoint function at the midblastula transition. *Curr. Biol.* 9: 302–312.
- Stevens, W. L., 1936 The analysis of interference. *J. Genet.* 32: 51–64.
- Sturtevant, A. H., 1913 The linear arrangement of six sex-linked factors in *Drosophila*, as shown by their mode of association. *J. Exp. Biol.* 14: 43–59.
- Wang, S., D. Zickler, N. Kleckner, and L. Zhang, 2015 Meiotic crossover patterns: obligatory crossover, interference and homeostasis in a single process. *Cell Cycle* 14: 305–314.
- Weinstein, A., 1936 The theory of multiple-strand crossing over. *Genetics* 21: 155–199.
- Yamamoto, M., and G. L. Miklos, 1978 Genetic studies on heterochromatin in *Drosophila melanogaster* and their implications for the functions of satellite DNA. *Chromosoma* 66: 71–98.
- Yıldız, Ö., S. Majumder, B. C. Kramer, and J. Sekelsky, 2002 *Drosophila* MUS312 interacts with the nucleotide excision repair endonuclease MEI-9 to generate meiotic crossovers. *Mol. Cell* 10: 1503–1509.
- Zeng, Y., H. Li, N. M. Schweppe, R. S. Hawley, and W. D. Gilliland, 2010 Statistical analysis of nondisjunction assays in *Drosophila*. *Genetics* 186: 505–513.
- Zhang, L., Z. Liang, J. Hutchinson, and N. Kleckner, 2014 Crossover patterning by the beam-film model: analysis and implications. *PLoS Genet.* 10: e1004042.

Communicating editor: J. Birchler