

## Worldwide study of hematopoietic allogeneic stem cell transplantation in pyruvate kinase deficiency

Pyruvate kinase deficiency (PKD) is the most frequent glycolytic enzyme defect causing hereditary non-spherocytic hemolytic anemia.<sup>1</sup> PKD leads to energy deprivation of the red cell, ultimately resulting in premature red cell death. Premature red cell death causes clinical symptoms of hemolytic anemia. The degree of hemolysis can vary widely, from very mild and fully compensated forms to life-threatening anemia with transfusion dependency.<sup>2</sup> The treatment for PKD is mainly supportive, and consists of regular red blood cell transfusions, splenectomy and chelation therapy for iron overload.<sup>3</sup> Hematopoietic allogeneic stem cell transplantation (HSCT) has the potential to cure PKD. However, there is little experience of applying HSCT in PKD. The current knowledge of HSCT in PKD is predominantly based on animal studies, and guidelines are not available.<sup>4,5</sup> To date, only four human cases of HSCT have been published in the literature.<sup>6-8</sup> The total number of cases transplanted worldwide is unknown.

The aim of this study is to make a worldwide inventory of PKD cases that have been treated by HSCT, and to evaluate indication, procedures employed and outcome as a first step towards the establishment of guidelines for HSCT in PKD. In order to achieve this goal queries were sent to national and international databanks, including the European Society for Blood and Marrow Transplantation (EBMT), the Center for International Blood and Marrow Transplant Research (CIBMTR), and the National Institute of Health (NIH), as well as to physicians known to be involved in HSCT in PKD patients. For each case found, a specifically designed questionnaire was sent to the physician involved. The questionnaire contained questions on disease characteristics, pre-transplant condition, transplant regimen and post-transplant outcome.<sup>9</sup> All data were evaluated by an experienced physician, and institutions were contacted in case of inconsistencies. An adapted EBMT score (*i.e.*, age, donor type and donor-recipient sex combination) was calculated based on the answers provided.<sup>10</sup> In addition, data from two additional cases, published recently, were extracted from the literature and included.<sup>8</sup> To the best of our knowledge, we have included all cases worldwide.

In total, 16 cases were found to be treated by HSCT between 1996 and 2015. Patient characteristics are summarized in Table 1. Patients had all been treated in either European or Asian centers. No cases resulted as being transplanted in the USA. Patient's median age at transplantation was 6.5 years. All patients were transfusion-dependent before transplantation, with median transfusion needs of 13 units of packed red blood cells per year (range: 6 to 34 units).

Conditioning and prophylaxis characteristics are summarized in Table 1. All patients received graft-versus-host disease (GvHD) prophylaxis. *Ex vivo* T-cell depletion was performed in one transplant. In another, red cell depletion was performed. Five transplants were sex-matched, four were female receiver-male donor and four were male receiver-female donor; this information was not available for three cases.

Median follow-up time after transplantation was 2.3 years (range: 2 months to 19 years). Fifteen patients showed engraftment. The sixteenth patient initially showed pancytopenia and mixed chimerism. Following

splenectomy six months post-transplantation, this patient's cell count spontaneously transitioned to normal with full donor chimerism. Two patients suffered from secondary graft loss; in one there was recovery to 91% donor chimerism after donor lymphocyte infusion. The outcome in the second patient was unknown.

Infectious complications and occurrence of GvHD are summarized in Table 1. The most significant infectious complications were aspergillus pneumonia (two patients), suspected aspergillus pneumonia (one patient), suspected fungal pneumonia (one patient), pneumonia (one patient), sepsis (one patient) and bacterial infection *e causa ignota* (one patient). GvHD grade 4 was reported in 6/16 cases (38%). Seven out of 16 cases (44%) did not show symptoms of GvHD. There was no correlation between GvHD prophylaxis or any other clinical factors and the occurrence of GvHD grade 2-4 in these patients.

Five out of 16 patients (31%) did not survive. All died of transplant-related causes. They had a median survival time of 13 months (range: 2-25 months). The two-year cumulative survival was 74%. Two patients had not yet reached the two-year milestone at the time of the questionnaire. The three-year cumulative survival rate was 65% (Figure 1); seven patients had not yet reached the three-year milestone.

Patients who did not survive differed significantly from surviving patients. (Figure 1, Table 2). They were significantly older ( $P=0.036$ ). Nine out of ten patients (90%) <10 years of age survived transplantation, whereas two out of six (33%)  $\geq 10$  survived. Patients <10 years were less often splenectomized ( $P=0.001$ ) and had lower pre-transfusion hemoglobin levels prior to HSCT ( $P=0.04$ ). Patients who did not survive had all been treated in European centers. All patients treated in Asian centers survived transplantation (8/8). Patients treated in Asian centers were younger ( $P=0.001$ ), less often splenectomized ( $P=0.041$ ), and had lower ferritin levels prior to HSCT ( $P=0.048$ ). In addition, they were more often transplanted using peripheral blood stem cells as a source ( $P=0.014$ ) and more often conditioned on a cyclophosphamide regimen ( $P=0.007$ ). Furthermore, patients who did not survive had frequently suffered from GvHD grade 2-4 ( $P=0.031$ ). Notably, four out of five deceased patients had suffered from both GvHD grade 3-4 and infection or viral reactivation.

There were no significant differences in sex, plasma ferritin level, use of pre-transplant chelation therapy, transfusion burden in the 12 months prior to HSCT, adapted EBMT-score, conditioning regimen, relation to donor, graft type, donor-recipient sex combination, or transplant source.

In conclusion, herein we discuss the first global study on the outcome of all patients known to have undergone HSCT in PKD. Since guidelines for HSCT in PKD are lacking, this report may be a helpful first step toward future protocols. Compared to published survival rates for other forms of hereditary anemias, cohorts that are otherwise comparable in age, time period and transplant hospital, the overall survival rate after HSCT in PKD is relatively low.<sup>11-13</sup> The present analysis of all 16 PKD patients known to be transplanted to date showed a three-year overall survival of 65%. Significantly better survival was observed for patients transplanted before the age of ten. A negative effect of age on survival is also reported for other forms of hereditary anemia.<sup>11,12</sup> Concurrently, we noticed a striking difference in survival between patients treated in Asian and European centers, which could possibly be explained by the difference in age at which patients were transplanted. In addition,

Table 1. Patient characteristics.

Sex	Center	PKLR genotype	Splenectomy	N. Transf. 12mo	aEBMT	Age at HSCT	Max Ferritin	Pre. Trans. Ferritin	Trans. year	Donor	Regimen	Matching	Stem cell source	Conditioning regimen	CvHD	Infection	Outcome	Follow-up time (Mo)	
M	Asia	Unknown	No	14	Good (0)	5y	-	<b>950</b>	1996	MSD	Myeloablation	<b>8/8</b>	Bone marrow	Cyph 200 mg/kg Bu 16 mg/kg p.o.	No	Febrile neutropenia unknown origin	Alive*	235	
<i>Pt 1: Severe chronic anemia and progressive splenomegaly</i>																			
F	EU	<i>c.[721G&gt;T;1594C&gt;T] p. [(Glu241*); (Arg537Trp)]</i>	Yes	8	Intermediate (2)	15y	-	596	2002	MFD	Myeloablation	<b>10/10</b>	Bone marrow	ATG 20 mg/kg Cyph 90 mg/kg Flu 100 mg/m <sup>2</sup> Bu 16 mg/kg p.o.	Grade 4 (S,G,L)	Primary CMV infection, asperg. pneum	Deceased	15	
<i>Pt 2: Concerns regarding progressive liver and heart hemosiderosis</i>																			
F	Asia	<i>c.[1044G&gt;T;1076 G&gt;A] p. [(Lys348Asn); (Arg359His)]</i>	No	14	Intermediate (2)	1y, 7mo	3357	206	2009	Cord	Myeloablation	<b>7/8</b>	Cord blood	ATG 7.5 mg/kg Cyph 200 mg/kg Bu 19.2 mg/kg	Grade 1 (S)	Bacterial infection	Alive	72	
<i>Pt 3: Transfusion dependency</i>																			
F	EU	<i>c.[721G&gt;T;1463G&gt;A] p. [(Glu241*); (Arg488Gln)]</i>	No	10	Intermediate (2)	3y	<b>2444</b>	<b>1161</b>	2009	MUD	Non-myeloablation/ RIST		Bone marrow	ATG 30 mg/kg Flu 160 mg/m <sup>2</sup> Thio 8 mg/kg Treo 42 mg/kg	No	No	Alive	65	
<i>Pt 4: High transfusion dependency and secondary hemochromatosis</i>																			
M	Asia	<i>c.[119G&gt;A;1015G&gt;A] p. [(Arg40Gln); (Asp339Asn)]</i>	No	13	Intermediate (1) 2y, 6mo		-	-	2009	MUD	Non-myeloablation/ RIST	<b>8/10</b>	Peripheral blood	ATG 15 mg/kg Cyph 200 mg/kg Flu 120-160 mg/m <sup>2</sup> Bu 3.2-4.8 mg/kg	Grade 2 (S)	Fever unknown origin	Alive	8	
<i>Pt 5: Transfusion dependency</i>																			
F	EU	<i>c.[1123_1133dup11; 1123_1133dup11] p. [(Met377fs); (Met377fs)]</i>	Yes	20	Intermediate (1)	17y	1888	1888	2010	MFD	Myeloablation	<b>8/8</b>	Peripheral blood	Cyph 120 mg/kg Bu 12.8 mg/kg	Grade 4 (S, G)	E. faecium sepsis, susp. fungal pneumonia	Deceased	5	
<i>Pt 6: Progressive transfusion dependency, decreasing quality of life</i>																			
F	EU	<i>c.[494G&gt;T;1529G&gt;A] p. [(Gly165Val); (Arg510Gln)]</i>	Yes	12	Intermediate (2)	39y	<b>1311</b>	<b>650</b>	2011	MUD	Non-myeloablation/ RIST	<b>8/8</b>	Bone marrow	ATG 600 mg/m <sup>2</sup> Flu 120 mg/m <sup>2</sup> Bu 10.8 mg/m <sup>2</sup>	Grade 4 (S,G,L)	No	Deceased	25	
<i>Pt 7: Progressive transfusion dependency</i>																			
F	EU	<i>c.[1532G&gt;A;1612G&gt;T] p. [(Cly511Glu); (Glu538*)]</i>	Yes	8	Intermediate (2)	7y	<b>771</b>	<b>771</b>	2013	MFD	Non-myeloablation/ RIST	<b>10/10</b>	Bone marrow	ATG 4 mg/kg Flu 160 mg/m <sup>2</sup> Thio 8 mg/kg Treo 42 mg/m <sup>2</sup>	No	CMV reactivation	Alive	29	

continued on next page

continued from previous page

**Pt 9: Transfusion dependency**  
 M EU c.[1481T>C;1675C>T] 12 Intermediate (1) 6 y **675** **675** 2013 MUD Non- 9/10 Bone ATG 6 mg/kg Grade 4 CMV and Deceased 2  
 p.[(11e494Thr);(Arg559\*)] myeloablation/ RIST marrow Flu 160 mg/m<sup>2</sup> EBV reactivation  
 Thio 8 mg/kg  
 Treo 42 mg/m<sup>2</sup>

**Pt 10: Transfusion dependency**  
 M Asia c.[848T>C;941T>C] 13 Intermediate (1) 6 mo 593.5 2013 MUD Myeloablation 9/10 Peripheral ATG 15 mg/kg Grade 4 No Alive 34  
 p.[(Val283Ala);(Ile314Thr)] blood Cycph 200 mg/kg (G)  
 Flu 120-160 mg/m<sup>2</sup>  
 Bu 3.2-4.8 mg/kg

**Pt 11: Transfusion dependency, problems with iron overload treatment due to compromised renal function**

M EU c.[1618+37\_2064del; 1618+37\_2064del] 6 Intermediate (1) 10y 4149 **7026** 2014 MUD Myeloablation 9/10 Bone ATG 8 mg/kg Grade 3 Asperg. Deceased 13  
 p.[(Lys541fs);(Lys541fs)] marrow Flu 160 mg/m<sup>2</sup> (S,G) pneum.  
 Bu Targeted dose, target AUC 90

**Pt 12: Transfusion dependency**  
 M Asia c.[661G>T;941T>C] 9 Intermediate (1) 9mo - - 2014 Cord Myeloablation 7/10 Cord blood ATG 15 mg/kg Grade 4 Pneumonia Alive 24  
 p.[(Asp221Tyr);(Ile314Thr)]

**Pt 13: Transfusion dependency**  
 M Asia c.[848T>C;848T>C] 13 Intermediate (1) 2mo 297.3 2015 MUD Non- 10/10 Peripheral ATG 15 mg/kg No No Alive 12  
 p.[(Val283Ala);(Val283Ala)] myeloablation/ blood Cycph 200 mg/kg  
 Flu 120-160 mg/m<sup>2</sup>  
 Bu 3.2-4.8 mg/kg

**Pt 14: Transfusion dependency, secondary hemochromatosis and hepatocarcinoma**  
 M EU c.[993C>A;1015G>C] 34 Intermediate (2) 41y - **1650** 2015 MSD Myeloablation Bone ATG 30 mg/kg No Susp. Alive 12  
 p.[(Asp331Glu);(Asp339His)] marrow Flu 160 mg/m<sup>2</sup> asperg.  
 Thio 8 mg/kg pneum.  
 Treo 42 mg/kg

**Pt 15: Transfusion dependency, secondary hemochromatosis, spinal compression fracture due to osteoporosis**

M# Asia c.[1270-3C>A;1618G>T] 11y - - 2000 MUD Myeloablation 9/10 Peripheral ATG 7.5 mg/kg No - Alive 36  
 p.[(?);(Gly540\*)] blood Cycph 200 mg/kg  
 Flu 120 mg/m<sup>2</sup>

**Pt 16: Transfusion dependency**  
 F# Asia c.[1270-3C>A;1618G>T] 8y - - - MUD Myeloablation Peripheral ATG 7.5 mg/kg No - Alive 30  
 p.[(?);(Gly540\*)] blood Cycph 200mg/kg  
 Flu 200 mg/m<sup>2</sup>

#Data retrieved from Kim, 2016<sup>8</sup>

\*At last follow up

M: male; F: female; Y: years; Mo: months; N: Transf. 12mo: estimated number of red blood cell transfusions in 12 months prior to HSCT; Max Ferritin: maximum ferritin reported in ng/mL; Pre-Trans: Ferritin; pre-transplant ferritin level in ng/mL (ferritin levels in bold: under chelation regimen); MUD: matched unrelated donor; MSD: matched sibling donor; MFD: matched family donor; Cord: cord blood; ATG: anti-thymocyte globulin; Flu: fludarabine; Bu: busulfan; Thio: thiothepa; Treo: treosulfan; Cycph: cyclophosphamide; - : unknown; RIST: reduced-intensity hematopoietic stem cell transplantation; S: skin; G: GI tract; L: liver; CMV: cytomegalovirus; EBV: Epstein-Barr virus; sus: suspected; asperg: aspergillus; pneum: pneumonia; E: faecium sepsis; enterococcus faecium sepsis; GVHD: graft-versus-host disease; HSCT: hematopoietic allogeneic stem cell transplantation; aEBMT: adapted European Society for Blood and Marrow Transplantation score; p.o.: per os.

Asian patients were non-splenectomized in many instances, and had lower pre-transplantation ferritin levels, which could also be related to the young age at which HSCT was performed.

Asian patients were more frequently transplanted with peripheral blood stem cells as opposed to bone marrow-derived stem cells. Peripheral blood stem cells are easier to collect from the donor, but reportedly increase the risk of chronic GvHD.<sup>14</sup> Our cohort, however, was too small to analyze the specific effect of stem cell source on the occurrence of chronic GvHD.

An important limitation of this study is its retrospective character, and the fact that the small sample size did

not allow us to perform *post hoc* correction for multiple testing. Therefore, the quantitative analysis of this data should be interpreted with care. Other limitations include the heterogeneity of conditioning regimens, and heterogeneity in the pre-transplant risk classification systems used. However, we did observe a better survival for patients transplanted prior to age ten. This effect of age might also play a role in the observed differences in survival between patients treated in European centers and those treated in Asian centers.

Although HSCT should be considered an investigational treatment, the strong decline in survival of treated patients over the age of ten suggests the need to evalu-


**Table 2.** Statistical differences between surviving and non-surviving patients.

	Survivor	Non-survivor	P value
Age in years	7.5 – 3.0 (0.8-41)	17.4 – 15.2 (6-39)	0.036*
Asian hospital	8/11 (73%)	0/5	0.026*
Splenectomy performed	3/11 (27%)	4/5 (80%)	0.106
Mean Hb (g/dL) (N=13)	6.0 – 5.5 (4.5-7.9)	7.1 – 6.9 (6.0-8.1)	0.112
Pre-transplant ferritin (ng/ml) (n=12)	804 – 771 (206-1650)	2167 – 675 (596-7026)	0.432
Myeloablation	6/11 (55%)	4/5 (80%)	0.588
Graft type			0.507
MSD	2/11 (18%)	0/5	
MUD	6/11 (55%)	3/5 (60%)	
CORD	2/11 (18%)	0/5	
MFD	1/11 (9%)	2/5 (40%)	
Transplant source			0.333
Bone marrow	4/11 (36%)	4/5 (80%)	
Peripheral blood	5/11 (45%)	1/5 (20%)	
Cord blood	2/11 (18%)	0/5	
GvHD			0.015*
None	7/11 (64%)	0/5	
Grade 1	1/11 (9%)	0/5	
Grade 2	1/11 (9%)	0/5	
Grade 3	0/11	1/5 (20%)	
Grade 4	2/11 (18%)	4/5 (80%)	

(descriptive statistics: mean – median (range) (N), frequencies number/total (percentage))

\*P<0.05

Continuous variables were expressed as mean, median and range, and subgroups were compared using Mann-Whitney U tests. Categorical data was compared using Fisher's exact test for binomial and the Fisher-Freeman-Halton exact test for contingency tables larger than 2x2. Statistical significance was considered as P≤0.05. All tests were two-sided. Post hoc multiple comparison correction was not applied. Graft-versus-host disease (GvHD) is defined and graded according to international criteria.<sup>15</sup> Pre-transplant laboratory results from splenectomized patients are from the period after splenectomy. Hb: hemoglobin; MUD: matched unrelated donor; MSD: matched sibling donor; MFD: matched family donor; Cord: cord blood.



**Figure 1.** Overall survival, according to age.

ate HSCT as a treatment option early in life. However, since the rate of grade 3-4 GvHD was relatively high (7/16 = 44%), and death resulting from GvHD was likewise high (5/16 = 31%), transfusion dependency alone should not be an indication for performing HSCT in PKD.

Stephanie van Straaten,<sup>1,2\*</sup> Marc Bierings,<sup>3\*</sup> Paola Bianchi,<sup>4</sup> Kensuke Akiyoshi,<sup>5</sup> Hitoshi Kanno,<sup>6</sup> Isabel Badell Serra,<sup>7</sup> Jing Chen,<sup>8</sup> Xiaohang Huang,<sup>8</sup> Eduard van Beers,<sup>9</sup> Supachai Ekwattanakit,<sup>10</sup> Tayfun Güngör,<sup>11</sup> Wijnanda Adriana Kors,<sup>12</sup> Frans Smiers,<sup>13</sup> Reinier Raymakers,<sup>14</sup> Lucrecia Yanez,<sup>15</sup> Julian Sevilla,<sup>16</sup> Wouter van Solinge,<sup>1</sup> Jose Carlos Segovia<sup>17,18</sup> and Richard van Wijk<sup>1</sup>

<sup>1</sup>Department of Clinical Chemistry and Haematology, University Medical Center Utrecht, Utrecht University, the Netherlands; <sup>2</sup>Van CreveldKliniek, University Medical Center Utrecht, Utrecht University, the Netherlands; <sup>3</sup>Pediatric Blood and Marrow Transplantation Program, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht University, the Netherlands; <sup>4</sup>Hematology Unit, Anemia Physiopathology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; <sup>5</sup>Department of Pediatrics and Child Neurology, Oita University Faculty of Medicine, Hasama, Yufu, Japan; <sup>6</sup>Department of Transfusion Medicine and Cell Processing, Faculty of Medicine, Tokyo Women's Medical University, Japan; <sup>7</sup>Directora Unidad Pediátrica de Trasplante Hematopoyético, Hospital Santa Creu i Sant Pau, Barcelona, Spain; <sup>8</sup>Department of Hematology and Oncology, Shanghai Children's Medical Center, Shanghai Jiao Tong University School of Medicine, China; <sup>9</sup>Van Creveldklinik, Department of Internal Medicine, University Medical Center Utrecht, Utrecht University, the Netherlands; <sup>10</sup>Department of Medicine, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand; <sup>11</sup>Division of Stem Cell Transplantation, University Children's Hospital, Zurich, Switzerland; <sup>12</sup>Department of Pediatric Oncology, VU University Medical Center, Amsterdam, the Netherlands; <sup>13</sup>Department of Pediatric Haematology, Leiden University Hospital, the Netherlands; <sup>14</sup>Department of Internal Medicine, University Medical Center Utrecht, Utrecht University, the Netherlands; <sup>15</sup>Servicio de Hematología, Hospital Universitario Marqués de Valdecilla, Santander, Spain; <sup>16</sup>Servicio Hemato-Oncología Pediátrica, Hospital Infantil Universitario Niño Jesús, Madrid, Spain; <sup>17</sup>Differentiation and Cytometry Unit, Hematopoietic Innovative Therapies Division, Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Centro de Investigaciones Biomédicas en Red de Enfermedades Raras (CIBER-ER), Madrid, Spain and <sup>18</sup>Advance Therapies Mixed Unit, Instituto de Investigación Sanitaria-Fundación Jiménez Díaz (IIS-FJD), Madrid, Spain

\*SvS and MB contributed equally to this work

Correspondence: r.vanwijk@umcutrecht.nl  
doi:10.3324/haematol.2017.177857

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at [www.haematologica.org](http://www.haematologica.org).

## References

- Miwa S, Fujii H. Molecular basis of erythroenzymopathies associated with hereditary hemolytic anemia: tabulation of mutant enzymes. *Am J Hematol.* 1996;51(2):122-132.
- Zanella A, Fermo E, Bianchi P, et al. Red cell pyruvate kinase deficiency: molecular and clinical aspects. *Br J Haematol.* 2005;130(1):11-25.
- Grace RF, Zanella A, Neufeld EJ, et al. Erythrocyte pyruvate kinase deficiency: 2015 Status report. *Am J Hematol.* 2015;90(9):825-830.
- Morimoto M, Kanno H, Asai H, et al. Pyruvate kinase deficiency of mice associated with nonspherocytic hemolytic anemia and cure of the anemia by marrow transplantation without host irradiation. *Blood.* 1995;86(11):4323-4330.
- Weiden PL, Hackman RC, Deeg HJ, et al. Long-term survival and reversal of iron overload after marrow transplantation in dogs with congenital hemolytic anemia. *Blood.* 1981;57(1):66-70.
- Tanphaichitr VS, Suvatve V, Issaragrisil S, et al. Successful bone marrow transplantation in a child with red blood cell pyruvate kinase deficiency. *Bone Marrow Transplant.* 2000;26(6):689-690.
- Akiyoshi K, Sekiguchi K, Okamoto T, et al. Cord blood transplantation in a young child with pyruvate kinase deficiency. *Pediatr Int.* 2016;58(7):634-636.
- Kim M, Park J, Lee J, et al. Hemolytic anemia with null PKLR mutations identified using whole exome sequencing and cured by hematopoietic stem cell transplantation combined with splenectomy. *Bone Marrow Transplant.* 2016;51(12):1605-1608.
- Sorror ML, Maris MB, Storb R, et al. Hematopoietic cell transplantation (HCT)-specific comorbidity index: a new tool for risk assessment before allogeneic HCT. *Blood.* 2005;106(8):2912-2919.
- Gratwohl A, Hermans J, Goldman JM, et al. Risk assessment for patients with chronic myeloid leukaemia before allogeneic blood or marrow transplantation. Chronic Leukemia Working Party of the European Group for Blood and Marrow Transplantation. *Lancet.* 1998;352(9134):1087-1092.
- Baronciani D, Angelucci E, Potschger U, et al. Hemopoietic stem cell transplantation in thalassemia: a report from the European Society for Blood and Bone Marrow Transplantation Hemoglobinopathy Registry, 2000-2010. *Bone Marrow Transplant.* 2016;51(4):536-541.
- Fagioli F, Quarello P, Zecca M, et al. Haematopoietic stem cell transplantation for Diamond Blackfan anaemia: a report from the Italian Association of Paediatric Haematology and Oncology Registry. *Br J Haematol.* 2014;165(5):673-681.
- Smetsers SE, Smiers FJ, Bresters D, et al. Four decades of stem cell transplantation for Fanconi anaemia in the Netherlands. *Br J Haematol.* 2016;174(6):952-961.
- Adhikari J, Sharma P, Bhatt VR. Optimal graft source for allogeneic hematopoietic stem cell transplant: bone marrow or peripheral blood? *Future Oncol.* 2016;12(15):1823-1832.
- Przepiorka D, Weisdorf D, Martin P, et al. 1994 Consensus conference on acute GVHD grading. *Bone Marrow Transplant.* 1995;15(6):825-828.