

RESEARCH ARTICLE

Schistosoma mansoni infection suppresses the growth of *Plasmodium yoelii* parasites in the liver and reduces gametocyte infectivity to mosquitoes

Taeko Moriyasu^{1,2,3}, Risa Nakamura^{1,2,3}, Sharmina Deloer^{1,2,3}, Masachika Senba⁴, Masato Kubo^{5,6}, Megumi Inoue⁷, Richard Culleton^{1,7*}, Shinjiro Hamano^{1,2,3*}

1 Leading Program, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan, **2** Department of Parasitology, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki, Japan, **3** The Joint Usage/Research Center on Tropical Disease, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki, Japan, **4** Pathology Unit, Department of Pathology, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki, Japan, **5** Laboratory for Cytokine Regulation, Research Center for Integrative Medical Science (IMS), RIKEN Yokohama Institute, Yokohama, Japan, **6** Division of Molecular Pathology, Research Institute for Biomedical Science, Tokyo University of Science, Noda, Japan, **7** Malaria Unit, Department of Pathology, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki, Japan

* richard@nagasaki-u.ac.jp (RC); shinjiro@nagasaki-u.ac.jp (SH)



OPEN ACCESS

Citation: Moriyasu T, Nakamura R, Deloer S, Senba M, Kubo M, Inoue M, et al. (2018) *Schistosoma mansoni* infection suppresses the growth of *Plasmodium yoelii* parasites in the liver and reduces gametocyte infectivity to mosquitoes. PLoS Negl Trop Dis 12(1): e0006197. <https://doi.org/10.1371/journal.pntd.0006197>

Editor: John Pius Dalton, Queen's University Belfast, UNITED KINGDOM

Received: October 5, 2017

Accepted: December 28, 2017

Published: January 26, 2018

Copyright: © 2018 Moriyasu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Japan Society for the Promotion of Science (JSPS) (<http://www.jsps.go.jp/english/>) of project numbers JP25870525, JP24255009 and JP16K21233 to RC, JP17H01684 and S2509 to SH; A Royal Society Bilateral Grant for Co-operative Research <https://royalsociety.org/grants-schemes-awards/>

Abstract

Malaria and schistosomiasis are major parasitic diseases causing morbidity and mortality in the tropics. Epidemiological surveys have revealed coinfection rates of up to 30% among children in Sub-Saharan Africa. To investigate the impact of coinfection of these two parasites on disease epidemiology and pathology, we carried out coinfection studies using *Plasmodium yoelii* and *Schistosoma mansoni* in mice. Malaria parasite growth in the liver following sporozoite inoculation is significantly inhibited in mice infected with *S. mansoni*, so that when low numbers of sporozoites are inoculated, there is a large reduction in the percentage of mice that go on to develop blood stage malaria. Furthermore, gametocyte infectivity is much reduced in mice with *S. mansoni* infections. These results have profound implications for understanding the interactions between *Plasmodium* and *Schistosoma* species, and have implications for the control of malaria in schistosome endemic areas.

Author summary

Malaria and schistosomiasis are parasitic infectious diseases that cause severe morbidity and mortality in the tropics. Chronic schistosomiasis causes malnutrition and impaired intellectual development to children while malaria can cause fatal acute infections. Since coinfection of these two parasites is common in the tropics, many studies of both epidemiology and coinfection in animal models have been performed in order to reveal interactions between them. Previous animal studies on the interactions between *Plasmodium* and *Schistosoma* parasites have focused on the blood stage pathology of the malaria infection,

grants/) to RC; a Sasakawa Foundation Butterfield Award (<http://www.gbsf.org.uk/butterfieldawards/>) to RC; the Global Center of Excellence (GCOE) Program at Nagasaki University (<http://www.tm.nagasaki-u.ac.jp/gcoe/>) to SH. TM was supported by a PhD scholarship by the Leading Program, Graduate School of Biomedical Sciences, Nagasaki University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

and have consistently shown that parasitaemia can be enhanced in the presence of the helminth. In contrast, we focused on liver immunopathology in mice during coinfection between with *Schistosoma* and *Plasmodium*. We show that *S. mansoni* infection inhibits *Plasmodium* parasite growth in the liver resulting in a large reduction in the percentage of mice that go on to develop blood stage malaria following inoculation of low numbers of sporozoites. We also demonstrate that gametocyte infectivity is much reduced in mice with *S. mansoni* infections. Our results imply that *S. mansoni* infection can reduce malaria transmission both from mosquitoes to mice, and from mice to mosquitoes.

Introduction

Malaria and schistosomiasis are two of the most important parasitic diseases in the tropics, and together constitute a severe burden to public health and to the economic development of endemic countries. Malaria is estimated to cause 429,000 deaths per year, 70% of those occurring in children aged under five years old [1]. The WHO has estimated that schistosomiasis causes about 200,000 deaths every year in sub-Saharan Africa and 218 million people were required to undergo preventive chemotherapy against the helminth globally in 2015.

The ranges of *Plasmodium* and *Schistosoma* overlap in much of the tropical world, leading to the potential for a great many coinfections of the two parasitic species. It has, for example, been estimated that there may be a greater than 30% coinfection rate among children in Sub-Saharan Africa [2]. Given the importance of such coinfections, interactions between *Plasmodium* and *Schistosoma* have been extensively studied both in nature, and using animal models in the laboratory.

Epidemiological studies on coinfections have often produced contrasting results: some reports contend that *Schistosoma* infection can increase susceptibility to *Plasmodium falciparum* [3–5], whilst others document a protective effect on *P. falciparum* incidence [6–9]. Differences in study design, genetic background of host populations and other environmental factors presumably contribute to these conflicting results.

Most laboratory-based animal studies have shown an exacerbation of malaria parasitaemia in *Schistosoma* infected mice [10–13] whilst others have revealed a protective effect of *Schistosoma* infection against experimental cerebral malaria and associated mortality [14–17]. In experimental *S. mansoni* infections, it is known that eggs deposited in the liver induce a strong Th2 type immune response [18]. Previous work has suggested that the exacerbation of malaria parasitaemia and protection against experimental cerebral malaria were mediated by a polarized Th2 immune environment which down-modulates inflammatory responses [10, 15].

The interactions between *Schistosoma* and *Plasmodium* are mainly mediated via host immune responses [19, 20]. Previous animal studies have investigated inter-species interactions using experimental infection with *Schistosoma* cercariae and *Plasmodium*-parasitized erythrocytes. As both parasites infect the liver at specific stages in their life cycle, we have focused on the immune reactions against those parasites in the liver. The major immunopathology in schistosomiasis is induced by egg-derived antigens in the liver, and hepatic immune cells develop immunity not only against pre-erythrocytic stages but also the blood stages of malaria parasites [21–24]. We therefore investigated whether an ongoing infection with *S. mansoni* could affect the rodent malaria parasite *Plasmodium yoelii* in the livers of mice challenged with sporozoites (SPZ).

We also evaluated whether *S. mansoni* infection affects malaria parasite gametocyte infectivity to mosquitoes, as it has been shown that the infectivity of malaria gametocytes decreases during the early stage of malaria infection due to host serum-mediated immunity [25–27].

Methods

Ethical statement

All mouse experiments were approved by the Institutional Animal Research Committee of Nagasaki University (No.1506181240) and performed according to Japanese law for the Humane Treatment and Management of Animals (Law No. 105 dated 19 October 1973 modified on 2 June 2006).

Mice

Six week-old female BALB/cCrSlc (hereafter referred to as BALB/c) and C57BL/6NCrSlc (hereafter referred to as B6) mice were purchased from Japan SLC, Inc. (Shizuoka, Japan). Six week-old female CBA/J mice were purchased from Charles River Laboratories Japan, Inc. (Kanagawa, Japan). IFN- γ -deficient (IFN- $\gamma^{-/-}$) mice and IL-4-deficient (IL-4 $^{-/-}$) mice were produced at RIKEN Yokohama Institute, Yokohama, Japan. All mice were maintained in the animal facilities of Nagasaki University with environmentally controlled, specific pathogen free conditions. Experiments were conducted with BALB/c mice unless otherwise specified.

Parasites and infections

The rodent malaria parasites *Plasmodium yoelii yoelii* 17x1.1pp [28] (hereafter referred to as *P. yoelii*) and *Plasmodium berghei* ANKA (hereafter referred to as *P. berghei*) were used throughout these experiments. A Puerto Rican strain of *Schistosoma mansoni* was maintained in the animal facilities of Nagasaki University by passage through *Biomphalaria glabrata* snails and ICR mice or *Meriones unguiculatus* (Mongolian jirds). In the coinfection model, experimental mice were percutaneously infected with 50 cercariae. In the intraportal infusion model, eggs were collected from the liver harvested from ICR mice after 7 weeks infection with 250 cercariae and stored at -30°C until use. The mice were anesthetized and inoculated with 3,000 eggs in a total volume of 100 μ L PBS via the portal vein.

Infections with *P. yoelii* and *P. berghei* were performed by i.v. inoculation with SPZ collected from *Anopheles stephensi* mosquitoes, as previously described [29]. 50, 500, or 1,500 SPZ in a total volume of 100 μ L PBS were inoculated for assessment of malaria parasitaemia or malaria parasite liver burden. Malaria challenges were performed 10 weeks after *S. mansoni*-cercariae infection in the coinfection model and 3 weeks after *S. mansoni* egg inoculation in the intraportal infusion model.

In transmission experiments, mice were intravenously infected with one million *P. yoelii*-parasitized erythrocytes. On day 3, 4, and 5 after *P. yoelii* infection, six-day-post emergence female *A. stephensi* mosquitoes were allowed to feed on individual mice for 30 minutes and reared in the insectary at 24°C with 80% humidity for seven days. Midguts were harvested from the mosquitoes and oocysts counted by light microscopy.

Malaria parasite density in the blood and the liver

Ten microliters of blood were collected from the tale-vein of mice at each time point. Livers were harvested 42 hours after malaria challenge infection. DNA was extracted from blood or livers using the EZ1 BioRobot (Qiagen N.V., Hilden, Germany) following the manufacturer's instructions. Real-time quantitative PCR (qPCR) was performed on DNA samples using the

Table 1. Information for PCR primers.

Gene name	Accession No	Primer sequence	Product size
<i>Plasmodium</i> 18s	M14599.1	F: GGAACGATGTGTGTCTAACACAAGGA	80bp
		R: CGCGTGCAGCCTAGTATATCTAAGGACA	
Mouse G3PDH	NM_008084.3	F: CATCTGAGGGCCCACTGAAG	72bp
		R: TGCTGTTGAAGTCGCAGGAG	

<https://doi.org/10.1371/journal.pntd.0006197.t001>

Applied Biosystems 7500 Real Time PCR system (Thermo Fisher Scientific, Inc., Massachusetts, USA). A master mix of the following reaction components was prepared: 4.75 µL water, 6.25 µL Power SYBR Green Master Mix (Qiagen N.V., Hilden, Germany), 0.25 µL forward primer (10 µM), and 0.25 µL reverse primer (10 µM). 5 µL DNA samples were added as PCR template. *P. yoelii* or *P. berghei* 18s gene was amplified using the primers 18s F1 5' GGAACGATGTGTGTCTAACACAAGGA 3' and 18s R1 CGCGTGCAGCCTAGTATATCTAAGGACA 3' (Table 1). Copy numbers of the parasite 18s gene were quantified with reference to a standard curve calculated from known numbers of plasmids containing the same gene sequence, as previously described [30]. The copy number of parasite 18s gene in each sample was standardized by the simultaneous quantification of the mouse glyceraldehyde 3-phosphate dehydrogenase (G3PDH) gene. The mouse G3PDH gene was amplified using the primers MmG3PDHF1 5' CATCTGAGGGCCCACTGAAG 3' and MmG3PDHR1 5' TGCTGTTGAAGTCGCAGGAG 3' (Table 1).

Monitoring of parasitaemia and gametocytaemia

Thin blood films from the tail-vein were prepared daily from day 2 to 8 after SPZ inoculation, and stained with Giemsa's solution. The numbers of infected and non-infected erythrocytes were counted per 10 microscopic fields to calculate parasitaemia and gametocytaemia.

Flow cytometric analysis

Hepatic nonparenchymal cells were isolated after *Schistosoma* inoculation as follows. Livers taken from each experimental mouse were homogenized in 5 mL of culture medium (RPMI-1640 supplemented with 10% FCS and 1% penicillin/streptomycin) using gentleMACS Dissociator (Miltenyi Biotec, Bergisch Gladbach, Germany), filtered through a mesh and suspended in HBSS. To remove liver parenchymal cells, the cells were resuspended in 33% Percoll (GE Healthcare UK Ltd., Buckinghamshire, England) containing 2.5mL of 5000U/5mL Heparin (Mochida Pharmaceutical Co., Ltd., Tokyo, Japan) and were centrifuged at 900 g for 20 min at room temperature. The pellet was resuspended in RBC lysis buffer then washed in HBSS, and resuspended in the culture medium.

The following mAbs were used for flow cytometric analysis using BD FACSVerser (Nippon Becton Dickinson Company, Ltd., Tokyo, Japan): PE anti-mouse CD3e (145-2C11), and PE anti-mouse T-bet (eBio4B10) (Affymetrix, Inc., California, U.S.), PE-Cy7 anti-mouse CD 4 (GK1.5), Percp-Cy5.5 anti-mouseCD3e (145-2C11), PE-Cy7 anti-mouse F4/80 (BM8), APC anti-mouse Gata3 (16E10A23), APC anti-mouse TCRgd (GL3), PE anti-mouse Ly6G (Gr1) (RB6-8C5), FITC anti-mouse CD11b (M1/70), and Biotin anti-mouse F4/80 (BM8) (BioLegend, California, U.S.), FITC anti-mouse CD49b (DX5), and BV421-streptavidin (Becton, Dickinson and Company, New Jersey, U.S.).

Fixation, permeabilization, and staining of the target transcription factors (T-bet and GATA-3) were conducted with Foxp3/Transcription Factor Staining Buffer Set (Affymetrix Japan K.K., Tokyo, Japan) according to the manufacturer's instructions.

Data analysis

Data analyses were performed using Microsoft Excel 2010 (Microsoft Corporation, Washington, USA) and GraphPad Prism6 (GraphPad Software, Inc., California, USA). Significance between control and treatment groups was determined with Student's t-tests. Survival and protection rates were statistically examined using the log-rank test. P values less than 0.05 were considered significant.

Results

S. mansoni infection significantly reduces the number of malaria parasites in the liver following sporozoite inoculation

To investigate the impact of chronic *S. mansoni* infection on the growth of malaria parasites in the liver, BALB/c mice were infected with 50 cercariae subcutaneously 10 weeks prior to i.v. inoculation of 1,500 *P. yoelii* sporozoites (SPZ). The number of copies of the malaria parasite 18s gene present in the liver of mice was measured by qPCR 42 h after SPZ challenge. Parasitaemia was monitored daily from day 2 to 8 post SPZ challenge.

Malaria parasite liver burden was significantly reduced to one-tenth or less in *S. mansoni*-infected mice compared with non-*S. mansoni* infected controls (Fig 1A). This reduction in the numbers of malaria parasites in the liver resulted in a delay in the onset of parasitaemia in the *S. mansoni*-infected group. All mice in both groups developed blood stage malaria parasite infection, and the peak parasitaemia was not affected by *S. mansoni* infection (Fig 1B). There was no difference in mortality between *S. mansoni*-infected and non-infected mice (Fig 1C).

The reduction of liver-stage malaria parasite burden induced by *S. mansoni* infection was not constrained by malaria species or mouse strains

The genetic background of inbred mouse strains can determine the course of experimentally induced malaria parasite infections [31, 32]. We examined the malaria parasite liver burden in three different mouse strains; BALB/c, B6, and CBA/J in coinfections with *S. mansoni*. Female BALB/c, B6, and CBA/J mice were challenged with two different *Plasmodium* species: *P. yoelii* and *P. berghei* in order to investigate whether the reduction of malaria parasite liver burden occurs across mouse and parasite species.




Fig 1. Growth of malaria parasites in the liver and blood of mice following SPZ inoculation of *Plasmodium yoelii* with and without *Schistosoma mansoni* infection. (A) Copy number of *P. yoelii* 18s RNA gene per 1×10^6 mouse G3PDH gene measured at 42 h post SPZ inoculation. Female BALB/c mice (N = 6) infected with 50 *S. mansoni*-cercariae 10 weeks previously were challenged with 1,500 *P. yoelii* SPZ along with *S. mansoni*-non-infected controls. **P < 0.01, Student's two-tailed t-test, t = 4.362, df = 10. (B) Parasitaemia. Blood stage malaria parasites were monitored daily from day 2 to 8 post i.v. inoculation of 500 *P. yoelii* SPZ. (C) Percentage survival. Data from one representative experiment of three independent repeats are shown.

<https://doi.org/10.1371/journal.pntd.0006197.g001>

Both B6 and CBA/J mice showed significant reduction in *P. yoelii* liver burden in *S. mansoni*-infected mice, consistent with the results observed in BALB/c mice (Fig 2A). The liver-stage growth of *P. berghei* was also reduced in *S. mansoni*-infected mice in all mouse strains examined (Fig 2B).

S. mansoni infection significantly reduces liver-stage malaria parasite burden from 2 h following SPZ inoculation

There are several potential mechanisms for the reduction in malaria parasite burden at 42 hours post SPZ inoculation in *S. mansoni* infected mice. It is possible, for example, that antibodies raised against schistosomes may offer protection against sporozoites through cross-reactivity [33]. It is also possible that liver fibrosis caused by schistosomiasis may physically impede the invasion of hepatocytes by sporozoites. Alternatively, there may be suppression of *P. yoelii* hepatocytic stages in the liver mediated by an altered immune environment in the liver caused by the presence of *S. mansoni* eggs in the organ. In order to investigate at which point *S. mansoni*-mediated reduction of malaria parasite in the liver occurs, we measured the number of malaria parasites in both the blood and the liver at time points throughout the 48 h growth of parasites in the liver. Ten microliters of blood were sampled from mouse tail-veins at 10, 30, 60, and 120 min, and livers were harvested at 2, 12, 24, and 42 h after i.v. inoculation of 1,500 SPZ. We found that malaria parasite load in the blood was not significantly different in infected compared with non-infected mice at any time point following SPZ inoculation (Fig 3A). In contrast, the number of malaria parasites in the liver were significantly reduced in *S. mansoni* infected mice from the first 2 h of development in the liver, with the proliferation rate from 2 to 24 h of liver stage parasites also suppressed in these mice (Fig 3B).

Intraportal inoculation of frozen S. mansoni-eggs significantly reduces liver-stage malaria parasite burden

As eggs are the principle cause of immunopathology in schistosomiasis [34], we assessed whether direct inoculation of *S. mansoni* eggs to the portal vein of mice could also reduce




Fig 2. Malaria parasite liver burden in BALB/c, C57BL/6, and CBA/J mice. (A) Copy number of *Plasmodium yoelii* 18s RNA gene per 1×10^6 mouse G3PDH gene measured at 42 h post sporozoite inoculation. Female BALB/c, B6, and CBA/J mice (N = 5) infected with 50 *Schistosoma mansoni*-cercaria 10 weeks previously were challenged with 1,500 SPZ of *P. yoelii* along with *S. mansoni*-non-infected controls. BALB/c: ***P<0.001, t = 6.283, df = 8; B6: *P<0.05, t = 2.511, df = 8; CBA: **P<0.01, t = 4.220, df = 7. (B) Copy number of *Plasmodium berghei* 18s RNA gene per 1×10^6 mouse G3PDH gene measured at 42 h post sporozoite inoculation. Female BALB/c, B6, and CBA/J mice (N = 5) infected with 50 *S. mansoni*-cercaria 10 weeks previously were challenged with 1,500 SPZ of *P. berghei* along with *S. mansoni*-non-infected controls. BALB/c: *P<0.05, t = 2.306, df = 8; B6: ***P<0.001, t = 5.336, df = 8; CBA: *P<0.05, t = 2.846, df = 7. All data were statistically examined using Student's two-tailed t-test.

<https://doi.org/10.1371/journal.pntd.0006197.g002>




Fig 3. *Plasmodium yoelii* parasite density in the blood and liver. Copy number of *P. yoelii* 18s RNA gene per 1×10^6 mouse G3PDH gene measured at 42 h post sporozoite inoculation. Female BALB/c mice (N = 5) infected with 50 *Schistosoma mansoni*-cercaria 10 weeks previously were challenged with 1,500 SPZ of *P. yoelii* along with *S. mansoni*-non-infected controls. (A) *P. yoelii* parasitaemia in the blood. (B) *P. yoelii* parasite density and proliferation in the liver. *** $P < 0.001$, Student's two-tailed t-test, $t = -6.316$, $df = 8$.

<https://doi.org/10.1371/journal.pntd.0006197.g003>

liver-stage malaria parasite growth. Female B6 mice were inoculated with 3,000 or 10,000 frozen *S. mansoni*-eggs in a total volume of 100 μ L PBS into the portal vein. Control mice were inoculated with 100 μ L PBS. Three weeks after egg inoculation, each group was challenged with 1,500 SPZ of *P. yoelii*. Liver-stage malaria parasite burden was significantly reduced in both groups inoculated with *S. mansoni*-eggs (Fig 4A).

S. mansoni infection causes severe liver damage associated with eosinophilic granulomas, collagen deposition, and fibrosis due to immunologic reactions to *Schistosoma* eggs trapped in the tissues. However, livers harvested from the intraportal infusion group did not show conspicuous damage macroscopically. Correspondingly, livers inoculated with *Schistosoma* eggs developed less and smaller granulomas compared with those harvested from the *S. mansoni*-cercariae infection group (Fig 4B and 4C).

Schistosoma infection is known to induce Th2-biased immune responses in mice. However, egg inoculation induced the infiltration of various immune cells into the liver with little pathology. Interferon-mediated innate immune responses are important modulators of malaria parasite growth in the liver [35]. We therefore focused not only on CD4⁺ T-bet⁺ cells and CD4⁺ GATA-3⁺ cells but also NK, NKT, and $\gamma\delta$ T cells which have previously been implicated as important sources of interferon-gamma (IFN- γ) in the liver during *Plasmodium* infection. The representative FACS plots are shown in the supplementary graph (S1 Fig). As shown in Fig 4D and 4E, the upregulation of the immune reaction induced by *S. mansoni* eggs in B6 mice was similar to that induced by *S. mansoni*-cercariae infection in BALB/c mice. Both CD4⁺ T-bet⁺ cells and CD4⁺ GATA-3⁺ cells significantly increased following intraportal inoculation of *S. mansoni*-eggs as well as after *S. mansoni*-cercariae infection. In both cases, the increase of CD4⁺ GATA-3⁺ cells was the largest among immune cells. The numbers of NK cells significantly increased only during the early phase, one week after intraportal inoculation of *S. mansoni*-eggs or 8 weeks after *S. mansoni*-cercariae infection. The numbers of NKT cells also significantly increased in both cases and this increase was greater in the group inoculated with *S. mansoni*-eggs. There was no significant increase in the numbers of $\gamma\delta$ T cells in either group. B6 mice infected with *S. mansoni*-cercariae showed a similar pattern of upregulation of immune cells (S2 Fig).




Fig 4. Liver immunopathology in *Schistosoma mansoni*-cercariae infection and intraportal infusion of frozen *S. mansoni*-eggs. (A) Malaria parasite liver burden with/without intraportal infusion of frozen *S. mansoni*-eggs. Female B6 mice were intraportally inoculated 3,000/10,000 frozen *S. mansoni*-eggs and challenged with 1,500 sporozoites of *Plasmodium yoelii* along with controls inoculated with PBS (control: N = 3; 3,000 frozen eggs: N = 5, ***P<0.001, t = 1.943, df = 6; 10,000 frozen eggs: N = 3, *P<0.05, Student's two-tailed t-test, t = 4.072, df = 4). (B) Macroscopic and microscopic images of liver pathology. The black lines indicate 250 micro meters. (C) The number of granulomas in the liver. Female BALB/c mice and B6 mice were inoculated with 50 *S. mansoni*-cercariae subcutaneously (naive: N = 5; 50 cercariae: N = 3) or inoculated with 3,000 frozen *S. mansoni*-eggs along with controls inoculated with PBS intraportally (PBS: N = 5; 3,000 frozen eggs: N = 5). The numbers of granuloma were counted in 20 microscopic fields at 100 x magnification. *P<0.05, ***P<0.001. All data were statistically examined using Student's two-tailed t-test. (D) The numbers of immune cells induced by intraportal infusion of 3,000 frozen *S. mansoni*-eggs. Female B6 mice (N = 4/group) were intraportally inoculated 3,000 *S. mansoni*-eggs. (E) The numbers of immune cells induced by 50 *S. mansoni*-cercaria infection. Female BALB/c mice (naive: N = 9, 8 weeks: N = 9, 10 weeks N = 6) were infected with 50 *S. mansoni*-cercariae. *P<0.05, **P<0.01, ***P<0.001. All data were statistically examined using Student's two-tailed t-test.

<https://doi.org/10.1371/journal.pntd.0006197.g004>

The reduction of liver-stage malaria parasite burden was reversed in IFN- γ -deficient and IL-4-deficient mice

We hypothesised that the reduction of liver-stage malaria parasite burden is dependent on the immune environment such as IFN- γ production from host immune cells accumulated and activated by the presence *S. mansoni*-eggs in the liver. IFN- γ is a key cytokine for the control of malaria parasites and is induced in the liver during *S. mansoni* infection. Interleukin-4 (IL-4) is a major mediator for the induction of the immune response against *S. mansoni*. Therefore, we examined the impact of IFN- γ and IL-4 on liver-stage malaria parasite burden using IFN- γ -deficient (IFN- $\gamma^{-/-}$) and IL-4-deficient (IL-4 $^{-/-}$) mice.

IFN- $\gamma^{-/-}$ and IL-4 $^{-/-}$ mice along with wild-type B6 (B6 WT) mice were inoculated 3,000 *S. mansoni*-eggs 3 weeks prior to SPZ challenge. Liver-stage malaria parasite burden was measured 42 h after i.v. challenge with 1,500 *P. yoelii* SPZ. Consistent with the result shown in Fig 4A, malaria liver burden was significantly reduced in the intraportal infusion group; however, this reduction was abrogated in IFN- $\gamma^{-/-}$ and IL-4 $^{-/-}$ mice (Fig 5).

S. mansoni infection inhibits the development of blood stage malaria following i.v. inoculation of entomologically relevant numbers (50) of SPZ

Through challenge with 1,500 SPZ of *P. yoelii*, we have demonstrated the impact of *S. mansoni* infection on liver-stage malaria parasite burden; this number of sporozoites, however, would not be expected to be inoculated naturally during a mosquito bite. In order to mimic the numbers inoculated by a mosquito bite, we reduced the challenge dose from 1,500 to 50 SPZ and examined the outcome of *Plasmodium* infection.

BALB/c mice (N = 16) infected with 50 *S. mansoni*-cercariae 10 weeks previously along with non-infected controls were challenged with 50 *P. yoelii* SPZ. Seven out of 16 *S. mansoni*-coinfected mice did not develop malaria parasitaemia, in contrast to all non-*S. mansoni* infected control mice developing blood-stage malaria on day 5 after SPZ challenge (Fig 6A). When blood stage malaria parasite infection became patent in coinfecting mice, it increased rapidly and matched the peak parasitaemia of non-*S. mansoni* infected controls (Fig 6B).

Plasmodium yoelii infections in mice infected with *S. mansoni* were less infectious to mosquitoes

We measured the influence of *S. mansoni* on the infectivity of malaria parasites to mosquitoes. We allowed *Anopheles stephensi* mosquitoes to feed on mice infected with *S. mansoni* + *P. yoelii* or *P. yoelii* alone on days 3 and 4 post-infection, when *P. yoelii* is at its most infectious to mosquitoes. A minimum of eight mosquitoes were allowed to feed on each of 5 mice per group on each day, and the resulting infectivity measured by assessing the proportion of mosquitoes




Fig 5. Malaria parasite liver burden in wild-type B6, IFN-gamma-deficient, and Interleukin-4-deficient mice. Copy number of *Plasmodium yoelii* 18s RNA gene per 1×10^6 mouse G3PDH gene measured at 42 h post sporozoite inoculation. Female IFN- $\gamma^{-/-}$ mice (N = 8), IL-4 $^{-/-}$ mice (N = 8), and B6 WT mice (N = 4) were intraperitoneally inoculated with 3,000 frozen *S. mansoni*-eggs and challenged with 1,500 SPZ of *P. yoelii* along with each control groups inoculated 100 μ L PBS (N = 4). B6 WT mice: *P<0.05, t = 3.017, df = 5; IFN- $\gamma^{-/-}$ mice: Not significant (NS), t = -1.303, df = 10; IL-4 $^{-/-}$ mice: NS, t = -0.016, df = 10.

<https://doi.org/10.1371/journal.pntd.0006197.g005>

that were infected with oocysts eight days post-feeding, and through quantifying the numbers of oocysts per infected mosquito per group on each day of feeding. Prior to mosquito feeding, the parasitaemia and number of gametocytes circulating in the peripheral blood of mice was calculated via microscopy of thin blood films following i.v. inoculation of *P. yoelii*-parasitized erythrocytes.

The number of gametocytes circulating in *S. mansoni*-infected mice was significantly higher than that of mice without *S. mansoni* infection on days 3 and 4 (Fig 7B). Despite this, the




Fig 6. Malaria outcomes with low dose SPZ challenge. Female BALB/c mice (N = 16) infected with 50 *Schistosoma mansoni*-cercariae 10 weeks previously were challenged with 50 sporozoites of *Plasmodium yoelii* along with *S. mansoni*-non-infected controls. (A) The percentage of mice that did not develop blood stage infection following inoculation of low doses SPZ. Data were statistically examined using the log-rank test. ***P<0.001, x2 = 29.8, df = 1. (B) Parasitaemia. Mean parasitaemia in *S. mansoni* infected group was calculated only among blood stage malaria positive mice.

<https://doi.org/10.1371/journal.pntd.0006197.g006>

proportion of mosquitoes infected with oocysts eight days post blood feed was significantly higher for the *S. mansoni* uninfected mic, compared to mice infected with *S. mansoni* (Fig 7C). Furthermore, oocyst numbers were significantly reduced in mosquitoes that had fed on mice infected with *S. mansoni*, compared to uninfected controls (Fig 7D).

Discussion

We have examined the impact of schistosome infection on malaria pathology and transmission capacity in a mouse model. Previous animal studies on the interactions between *Plasmodium* and *Schistosoma* have focused exclusively on the blood stage pathology of the malaria parasite infection, and have consistently shown that parasitaemia is enhanced in the presence of the helminth [10–13]. We have investigated the effect of helminth-malaria parasite coinfection on the growth of rodent malaria parasites in the liver, and show that *S. mansoni*-coinfection significantly reduces *Plasmodium* parasite numbers in the liver following sporozoite inoculation. *S. mansoni*-coinfection with malaria parasites resulted in a large reduction in the percentage of mice developing blood stage malaria parasite infection when mice were challenged with numbers of SPZ similar to those inoculated during natural inoculation through the bites of infected mosquitoes [36–39].

These results might suggest a possible explanation for the often contradictory results observed in epidemiological studies. Some prospective cohort studies in human populations have suggested a protective effect of *Schistosoma* infection on *Plasmodium* infection: Lyke et al. concluded that children infected with *Schistosoma haematobium* showed increased time to first clinical malaria infection and fewer malaria episodes over the follow-up period compared to children without *S. haematobium* infection [7]. Doumbo et al. also showed that people coinfecting with *S. haematobium* showed significant delays in time-to-first malaria episode [9]. Hürlimann et al. demonstrated that the *Plasmodium* parasitaemia incidence rate of children infected with *S. mansoni* increased after treatment with praziquantel; in contrast, children infected with hookworms had reduced *Plasmodium* parasitaemia rates following albendazole treatment [40]. Our results, which show that *S. mansoni* infection reduces the malaria parasite burden in the liver and the number of hosts developed blood stage infection following SPZ




Fig 7. Gametocyte infectivity. (A) Parasitemia. Female BALB/c mice were each inoculated with 1×10^6 *Plasmodium yoelii* parasitized erythrocytes intravenously with (n = 4 mice) or without (n = 5) pre-existing *Schistosoma mansoni* infection. Parasitaemia was determined by microscopic examination on days 3, 4, and 5 post-inoculation; day 3: Student's two-tailed t-test; $**P < 0.01$, $t = -4.906$, $df = 7$; Day5: $*P < 0.05$, $t = -2.922$, $df = 5$ (B) Gametocyte density. Gametocyte density was determined on days 3 and 4 post-inoculation of 1×10^6 *P. yoelii*-parasitized erythrocytes intravenously. Day 3: Student's two-tailed t-test; $**P < 0.01$, $t = 3.813$, $df = 5$; Day 4: $**P < 0.01$, $t = 3.608$, $df = 5$. Error bars show the geometric mean with 95% confidence intervals. (C) Percentage of mosquitoes with one or more oocysts present on the midgut eight days post-feeding on infected mice. A minimum of eight mosquitoes were allowed to feed on each individual mouse in the group per day $**P = 0.0003$, (2-way ANOVA, $F = 22.23$, $DFn = 1$, $DFd = 14$). Error bars mar the standard error of the mean per mouse group. (D) Oocyst numbers per mosquito. The numbers of oocysts present on mosquito midguts were determined eight days post-mosquito feeding; day 3: Student's two-tailed t-test, $**P < 0.01$, $t = 3.077$, $df = 25$. Error bars show the geometric mean with 95% confidence intervals. Data is representative of three independent experiments.

<https://doi.org/10.1371/journal.pntd.0006197.g007>

inoculation, may account for some of the observed protective effects of *Schistosoma* infection against malaria.

The intraportal inoculation of *S. mansoni* eggs reduced malaria parasite burden in the liver to the same extent as inoculation of cercariae. As shown in Fig 4, intraportal inoculation of *S. mansoni*-eggs induced an increase in the numbers of various lymphocytes, including both Th1 and Th2 cells, NK, NKT, and $\gamma\delta$ T cells in the liver without apparent fibrotic or granulomatous liver damage. This suggests that the suppression of the growth of malaria parasites in the liver is not mediated by physiological or morphological changes of the liver environment provoked by *S. mansoni*-eggs, but rather by changes in the hepatic immune microenvironment.

Previous animal studies have shown that interferon-mediated innate immune responses are important modulators of malaria parasite growth in the liver [41]; Miller et al. have shown that IFN- γ production from NKT cells (but not NK cells or T cells) is critical in reducing malaria parasite burden in the liver [35]. We initially hypothesised that malaria parasite liver burden

would increase in *S. mansoni*-coinfected mice as *S. mansoni* infection has been shown to induce robust Th2 type immune responses with a corresponding downregulation of Th1 responses [42]. However, contrary to our expectations, malaria parasite liver burden was significantly reduced in *S. mansoni*-coinfected mice. Both IFN- γ and IL-4 are required for this effect, and both the inoculation of *S. mansoni* eggs to the hepatic portal vein, and the infection of mice with cercariae were shown to increase innate immune cells and both Th1 and Th2 cells in the liver (S2 Fig). Taken together, this suggests that possible sources of IFN- γ and IL-4 include not only innate immune cells such as NKT cells but also helper T cells accumulated and activated by *S. mansoni* infection in the liver [43].

IFN- $\gamma^{-/-}$ and IL-4 $^{-/-}$ mice had lower numbers of malaria parasites in the liver after SPZ challenge compared with B6 WT mice even without *S. mansoni* infection. The experiment was repeated twice with similar results. These results are in agreement with previous studies with IL-4 $^{-/-}$ mice although the mechanisms are not fully understood. IL-4 $^{-/-}$ mice are known to be more resistant to sporozoite infection than wild-type mice owing to increased NK cell numbers and expression of inducible nitric oxide synthase in the liver [44]. Since the reduction of malaria parasite burden in the liver in co-infected mice was reversed in IFN- $\gamma^{-/-}$ and IL-4 $^{-/-}$ mice, we assume that the mechanism of protection is both IFN- γ and IL-4 dependent. To elucidate the mechanisms underlying this phenomenon, further research into the interactions between *Plasmodium* and *Schistosoma* are required.

We demonstrated that the infectivity of *Plasmodium* gametocytes to *Anopheles* mosquitoes dropped abruptly from a peak on day 3 to zero on day 4 in control mice not infected with *S. mansoni*. This phenomenon is known in animal models as malaria infection crisis and has previously been reported to occur in numerous *Plasmodium* species [45]. The mechanisms behind this loss of infectivity have not been fully explained, but previous animal studies suggest that cytokines such as tumor necrosis factor (TNF), IFN- γ , interleukin-6 (IL-6), and nitric oxide may inhibit the development of gametocytes [25–27]. The infectivity of *P. yoelii* gametocytes in *S. mansoni* infected mice was significantly reduced on day 3 post-inoculation which is the peak of infectivity in non-*S. mansoni* infected mice. Observing higher levels of IFN- γ in the sera of *S. mansoni* infected mice on day 3 and day 4 (S3 Fig), we assume that pre-existing *S. mansoni* infection induces activation of the host immune environment and this leads to an accelerated infection crisis. Although the mechanisms remain to be elucidated, this experimental transmission model suggests that *S. mansoni* infection may reduce malaria transmissibility from mosquitoes to mice.

In conclusion, the presence of *S. mansoni* infection dramatically reduced the number of rodent malaria parasites in the liver. This reduction leads to the inhibition of the development of blood stage malaria parasite infection following inoculation of biologically relevant numbers of sporozoites. We also demonstrate that *P. yoelii* gametocyte infectivity to mosquitoes is significantly reduced in the presence of *S. mansoni* in co-infected mice. These results imply that *S. mansoni* infection can reduce malaria transmission both from mosquitoes to mice and from mice to mosquitoes, and may explain some of the protective effects of *Schistosoma* infection against malaria described in previous epidemiological studies.

Supporting information

S1 Fig. Representative flow cytometry data of immune cells in the liver. Hepatic nonparenchymal cells were isolated from C57BL/6 mice at 1, 3 and 5 weeks post 3000 *S. mansoni* frozen eggs portal vein inoculation, or from BALB/c mice at 8 and 10 weeks post 50 *S. mansoni*-cercariae s.c. inoculation.

(PDF)

S2 Fig. The numbers of immune cells induced by 50 *Schistosoma mansoni*-cercariae infection. Female B6 mice (N = 5/group) were infected with 50 *S. mansoni*-cercariae. *P<0.05, **P<0.01, ***P<0.001, Student's two-tailed t-test.

(PDF)

S3 Fig. Concentration of IFN- γ in the serum. Female BALB/c mice were inoculated one million *Plasmodium yoelii*-parasitized erythrocytes intravenously with or without pre-existing *Schistosoma mansoni* infection. On day 3 and 4 after inoculation, the concentration of interferon-gamma (IFN- γ) was measured in the serum. Day 3:*P<0.05, t = -2.209, df = 6; Day 4:*P<0.05, t = -2.516, df = 4, Student's two-tailed t-test.

(PDF)

Acknowledgments

We thank Megumi Hamasaki and all laboratory members for technical support. This study was conducted at the Joint Usage/ Research Center on Tropical Disease, Institute of Tropical Medicine, Nagasaki University, Japan.

Author Contributions

Conceptualization: Taeko Moriyasu, Megumi Inoue, Richard Culleton, Shinjiro Hamano.

Data curation: Taeko Moriyasu.

Formal analysis: Taeko Moriyasu, Risa Nakamura.

Funding acquisition: Richard Culleton, Shinjiro Hamano.

Investigation: Taeko Moriyasu, Risa Nakamura, Richard Culleton.

Methodology: Taeko Moriyasu, Risa Nakamura, Sharmina Deloer, Masachika Senba, Richard Culleton.

Resources: Masato Kubo, Richard Culleton, Shinjiro Hamano.

Supervision: Richard Culleton, Shinjiro Hamano.

Validation: Richard Culleton, Shinjiro Hamano.

Visualization: Taeko Moriyasu.

Writing – original draft: Taeko Moriyasu.

Writing – review & editing: Richard Culleton, Shinjiro Hamano.

References

1. World Malaria Report 2016. Geneva: World Health Organization; 2016. Licence: CC BY-NC-SA 3.0 IGO. Available at: <http://apps.who.int/iris/bitstream/10665/252038/1/9789241511711-eng.pdf?ua=1>. Accessed 29 September 2017.
2. Degarege A, Degarege D, Veledar E, Erko B, Nacher M, Beck-Sague CM, et al. *Plasmodium falciparum* Infection Status among Children with *Schistosoma* in Sub-Saharan Africa: A Systematic Review and Meta-analysis. PLoS Negl Trop Dis 2016; 10(12):e0005193. <https://doi.org/10.1371/journal.pntd.0005193> PMID: 27926919
3. Sokhna C, Le Hesran J-Y, Mbaye PA, Akiana J, Camara P, Diop M, et al. Increase of malaria attacks among children presenting concomitant infection by *Schistosoma mansoni* in Senegal. Malar J 2004; 3:43. <https://doi.org/10.1186/1475-2875-3-43> PMID: 15544703
4. Mazigo HD, Waihenya R, Lwambo NJ, Mnyone LL, Mahande AM, Seni J, et al. Co-infections with *Plasmodium falciparum*, *Schistosoma mansoni* and intestinal helminths among schoolchildren in endemic

- areas of northwestern Tanzania. *Parasit Vectors* 2010; 3:44. <https://doi.org/10.1186/1756-3305-3-44> PMID: 20482866
5. Florey LS, King CH, Van Dyke MK, Muchiri EM, Mungai PL, Zimmerman PA, et al. Partnering parasites: evidence of synergism between heavy *Schistosoma haematobium* and *Plasmodium* species infections in Kenyan children. *PLoS Negl Trop Dis* 2012; 6(7):e1723. <https://doi.org/10.1371/journal.pntd.0001723> PMID: 22848765
 6. Briand V, Watier L, Le Hesran J-Y, Garcia A, Cot M. Coinfection with *Plasmodium falciparum* and *Schistosoma haematobium*: protective effect of schistosomiasis on malaria in Senegalese children? *Am J Trop Med Hyg* 2005; 72(6):702–7. PMID: 15964953
 7. Lyke Ke, Dicko A, Dabo A, Sangare L, Kone A, Coulibaly D, et al. Association of *Schistosoma haematobium* infection with protection against acute *Plasmodium falciparum* malaria in Malian children. *Am J Trop Med Hyg* 2005; 73(6):1124–30. PMID: 16354824
 8. Lemaitre M, Watier L, Briand V, Garcia A, Le Hesran JY, Cot M. Coinfection with *Plasmodium falciparum* and *Schistosoma haematobium*: additional evidence of the protective effect of Schistosomiasis on malaria in Senegalese children. *Am J Trop Med Hyg* 2014; 90(2):329–34. <https://doi.org/10.4269/ajtmh.12-0431> PMID: 24323515
 9. Doumbo S, Tran TM, Sangala J, Li S, Doumtabe D, Kone Y, et al. Co-infection of long-term carriers of *Plasmodium falciparum* with *Schistosoma haematobium* enhances protection from febrile malaria: a prospective cohort study in Mali. *PLoS Negl Trop Dis* 2014; 8(9):e3154. <https://doi.org/10.1371/journal.pntd.0003154> PMID: 25210876
 10. Helmby H, Kullberg M, Troye-Blomberg M. Altered Immune Responses in Mice with Concomitant *Schistosoma mansoni* and *Plasmodium chabaudi* Infections. *Infect Immun* 1998; 66(11):5167–74. PMID: 9784518
 11. Yoshida A, Maruyama H, Kumagai T, Amano T, Kobayashi F, Zhang M, et al. *Schistosoma mansoni* infection cancels the susceptibility to *Plasmodium chabaudi* through induction of type 1 immune responses in A/J mice. *Int Immunol* 2000; 12(8):1117–25. PMID: 10917886
 12. Legesse M, Erko B, Balcha F. Increased parasitaemia and delayed parasite clearance in *Schistosoma mansoni* and *Plasmodium berghei* co-infected mice. *Acta Trop* 2004; 91(2):161–6. <https://doi.org/10.1016/j.actatropica.2004.04.002> PMID: 15234665
 13. Sangweme D, Shiff C, Kumar N. *Plasmodium yoelii*. Adverse outcome of non-lethal *P. yoelii* malaria during co-infection with *Schistosoma mansoni* in BALB/c mouse model. *Exp Parasitol* 2009; 122(3):254–9. <https://doi.org/10.1016/j.exppara.2009.04.003> PMID: 19366621
 14. Wang M, Cao Y, Luo E, Zhang Y, Guo Y. Pre-existing *Schistosoma japonicum* infection alters the immune response to *Plasmodium berghei* infection in C57BL/6 mice. *Malar J* 2013; 12(1):322.
 15. Wang M, Feng Y, Pang W, Qi Z, Zhang Y, Guo Y, et al. Parasite densities modulate susceptibility of mice to cerebral malaria during co-infection with *Schistosoma japonicum* and *Plasmodium berghei*. *Malar J* 2014; 13(1):116.
 16. Amoani B, Ameyaw EO, Asante D-BB, Armah FA, Prah J, Kwesi Botchey CP, et al. Effect of pre-existing *Schistosoma haematobium* infection on *Plasmodium berghei* multiplications in imprinting control region mice. *Asian Pac J Trop Biomed* 2015; 5(6):488–92.
 17. Nyakundi RK, Nyamongo O, Maamun J, Akinyi M, Mulei I, Farah IO, et al. Protective effect of chronic schistosomiasis in baboons coinfecting with *Schistosoma mansoni* and *Plasmodium knowlesi*. *Infect Immun* 2016; 84(5):1320–30. <https://doi.org/10.1128/IAI.00490-15> PMID: 26883586
 18. Steinfelder S, Andersen JF, Cannons JL, Feng CG, Joshi M, Dwyer D, et al. The major component in schistosome eggs responsible for conditioning dendritic cells for Th2 polarization is a T2 ribonuclease (omega-1). *J Exp Med* 2009; 206(8):1681–90. <https://doi.org/10.1084/jem.20082462> PMID: 19635859
 19. Salazar-Castaon VH, Legorreta-Herrera M, Rodriguez-Sosa M. Helminth Parasites Alter Protection against *Plasmodium* Infection. *Biomed Res Int* 2014; 2014:913696. <https://doi.org/10.1155/2014/913696> PMID: 25276830
 20. Druilhe P, Tall A, Sokhna C. Worms can worsen malaria: towards a new means to roll back malaria? *Trends Parasitol* 2005; 21(8):359–62. <https://doi.org/10.1016/j.pt.2005.06.011> PMID: 15967721
 21. Belnoue E, Voza T, Costa FTM, et al. Vaccination with Live *Plasmodium yoelii* Blood stage Parasites under Chloroquine Cover Induces Cross-Stage Immunity against Malaria Liver Stage. *J Immunol* 2008; 181(12):8552–8. PMID: 19050274
 22. Nahrendorf W, Scholzen A, Sauerwein RW, Langhorne J. Cross-stage immunity for malaria vaccine development. *Vaccine* 2015; 33(52):7513–7. <https://doi.org/10.1016/j.vaccine.2015.09.098> PMID: 26469724

23. Gomes PS, Bhardwaj J, Rivera-Correa J, Freire-De-Lima CG, Morrot A. Immune Escape Strategies of Malaria Parasites. *Front Microbiol* 2016; 7:1617. <https://doi.org/10.3389/fmicb.2016.01617> PMID: [27799922](https://pubmed.ncbi.nlm.nih.gov/27799922/)
24. Butler NS, Schmidt NW, Vaughan AM, Aly AS, Kappe SHI, Harty JT, et al. Superior antimalarial immunity after vaccination with late liver stage-arresting genetically attenuated parasites. *Cell Host Microbe* 2011; 9(6):451–62. <https://doi.org/10.1016/j.chom.2011.05.008> PMID: [21669394](https://pubmed.ncbi.nlm.nih.gov/21669394/)
25. Naotunne BTDS, Karunaweera ND, Giudice G Del, Kularatne SMU, Grau GE, Carter R, et al. Cytokines Kill Malaria Parasites during Infection Crisis: Extracellular Complementary Factors Are Essential. *J Exp Med* 1991; 173(3):523–529. PMID: [1900073](https://pubmed.ncbi.nlm.nih.gov/1900073/)
26. Motard A, Landau I, Nussler A, Grau G, Baccam D, Mazier D, et al. The role of reactive nitrogen intermediates in modulation of gametocyte infectivity of rodent malaria parasites. *Parasite Immunol* 1993; 15(1):21–6. PMID: [8433851](https://pubmed.ncbi.nlm.nih.gov/8433851/)
27. Cao Y, Tsuboi T, Torii M. Nitric oxide inhibits the development of *Plasmodium yoelii* gametocytes into gametes. *Parasitol int* 1998; 47(2):157–166.
28. Inoue M, Tang J, Miyakoda M, Kaneko O, Yui K, Culleton R. The species specificity of immunity generated by live whole organism immunisation with erythrocytic and pre-erythrocytic stages of rodent malaria parasites and implications for vaccine development. *Int J Parasitol* 2012; 42(9):859–70. <https://doi.org/10.1016/j.ijpara.2012.07.001> PMID: [22846785](https://pubmed.ncbi.nlm.nih.gov/22846785/)
29. Inoue M, Culleton RL. The intradermal route for inoculation of sporozoites of rodent malaria parasites for immunological studies. *Parasite Immunol* 2011; 33(2):137–42. <https://doi.org/10.1111/j.1365-3024.2010.01263.x> PMID: [21226727](https://pubmed.ncbi.nlm.nih.gov/21226727/)
30. Abkallo HM, Liu W, Hokama S, Ferreira PE, Nakazawa S, Maeno Y, et al. DNA from pre-erythrocytic stage malaria parasites is detectable by PCR in the faeces and blood of hosts. *Int J Parasitol* 2014; 44(7):467–73. <https://doi.org/10.1016/j.ijpara.2014.03.002> PMID: [24704779](https://pubmed.ncbi.nlm.nih.gov/24704779/)
31. Stevenson MM, John J. Murine Malaria: Genetic Control of Resistance. *Infect Immun* 1982; 38(1):80–8. PMID: [7141699](https://pubmed.ncbi.nlm.nih.gov/7141699/)
32. Sayles PC, Wassom DL. Immunoregulation in murine malaria. Susceptibility of inbred mice to infection with *Plasmodium yoelii* depends on the dynamic interplay of host and parasite genes. *J Immunol* 1988; 141(1):241–248. PMID: [3379305](https://pubmed.ncbi.nlm.nih.gov/3379305/)
33. Pierrot C., Wilson S., Lallet H., Lafitte S., Jones FM., Daher W., Khalife J. Identification of a Novel Antigen of *Schistosoma mansoni* Shared with *Plasmodium falciparum* and Evaluation of Different Cross-Reactive Antibody Subclasses Induced by Human Schistosomiasis and Malaria. *Infect Immun* 2006; 74(6): 3347–3354. <https://doi.org/10.1128/IAI.01724-05> PMID: [16714563](https://pubmed.ncbi.nlm.nih.gov/16714563/)
34. Elbaz T, Esmat G. Hepatic and Intestinal Schistosomiasis: Review. *J Adv Res* 2013; 4(5):445–52. <https://doi.org/10.1016/j.jare.2012.12.001> PMID: [25685451](https://pubmed.ncbi.nlm.nih.gov/25685451/)
35. Miller JL, Sack BK, Baldwin M, Vaughan AM, Kappe SHI. Interferon-mediated innate immune responses against malaria parasite liver stages. *Cell Rep* 2014; 7(2):436–47. <https://doi.org/10.1016/j.celrep.2014.03.018> PMID: [24703850](https://pubmed.ncbi.nlm.nih.gov/24703850/)
36. Jin Y, Kebaier C, Vanderberg J. Direct microscopic quantification of dynamics of *Plasmodium berghei* sporozoite transmission from mosquitoes to mice. *Infect Immun* 2007; 75(11):5532–9. <https://doi.org/10.1128/IAI.00600-07> PMID: [17785479](https://pubmed.ncbi.nlm.nih.gov/17785479/)
37. Medica DL, Sinnis P. Quantitative Dynamics of *Plasmodium yoelii* Sporozoite Transmission by Infected Anopheline Mosquitoes. *Infect Immun* 2005; 73(7):4363–4369. <https://doi.org/10.1128/IAI.73.7.4363-4369.2005> PMID: [15972531](https://pubmed.ncbi.nlm.nih.gov/15972531/)
38. Kabiru EW, Mbogo CM, Muiruri SK, Ouma JH, Githure JL, Beier JC. Sporozoite loads of naturally infected Anopheles in Kilifi district, Kenya. *J Am Mosq Control Assoc* 1997; 13(3):259–62. PMID: [9383768](https://pubmed.ncbi.nlm.nih.gov/9383768/)
39. Churcher TS, Sinden RE, Edwards NJ, Poulton ID, Rampling TW, Brock PM, et al. Probability of Transmission of Malaria from Mosquito to Human Is Regulated by Mosquito Parasite Density in Naive and Vaccinated Hosts. *PLoS Pathog* 2017; 13(1):1–18.
40. Hürlimann E., Hougbedji CA., N'Dri PB., Bänninger D., Coulibaly JT., Yap P., et al. Effect of deworming on school-aged children's physical fitness, cognition and clinical parameters in a malaria-helminth co-endemic area of Côte d'Ivoire. *BMC Infectious Diseases*, 2014; 14(1): 411.
41. King T, Lamb T. Interferon- γ : The Jekyll and Hyde of Malaria. *PLoS Pathog* 2015; 11(10):8–13.
42. McKee AS, Pearce EJ. CD25⁺CD4⁺ cells contribute to Th2 polarization during helminth infection by suppressing Th1 response development. *J Immunol* 2004; 173(2):1224–31. PMID: [15240714](https://pubmed.ncbi.nlm.nih.gov/15240714/)
43. Adachi K, Osada Y, Nakamura R, Tamada K, Hamano S. Unique T cells with unconventional cytokine profiles induced in the livers of mice during *Schistosoma mansoni* infection. *PLoS One* 2013; 8(12):1–10.

44. Saeftel M., Krueger A., Arriens S., Heussler V., Racz P., Fleischer B., Hoerauf A. Mice Deficient in Interleukin-4 (IL-4) or IL-4 Receptor α Have Higher Resistance to Sporozoite Infection with *Plasmodium berghei*(ANKA) than Do Naive Wild-Type Mice. *Infect Immun* 2004; 72(1):322–331. <https://doi.org/10.1128/IAI.72.1.322-331.2004>
45. Bastien P, Landau I, Baccam D. Inhibition de l'infektivité des gamétocytes de Plasmodium par le sérum de l'hôte parasité. *Ann Parasitol Hum Comparée* 1987; 62(3):195–208.